

Supplement: Length-weight equations used to determine individual fish biomass for fish species caught in the early period (1977-1989). N = number of individuals used to determine the length-weight relationship for each species.

Species	Length-weight equation	n	Source	Study area and Notes
<i>Alepocephalus agassizii</i>	$TW(g) = 0.0035 \times (SLcm)^{3.2914}$	11	PSB trawl data	NE Atlantic
<i>Alepocephalus australis</i>	$TW(g) = 0.003 \times (SLcm)^{3.3582}$	3	John Gordon	NE Atlantic
<i>Alepocephalus bairdii</i>	$TW(g) = 0.0055 \times (SLcm)^{3.1403}$	38	PSB trawl data	NE Atlantic
<i>Alepocephalus productus</i>	$TW(g) = 0.0047 \times (SLcm)^{3.2293}$	5	PSB trawl data	NE Atlantic
<i>Alepocephalus rostratus</i>	$TW(g) = 0.0037 \times (SLcm)^{3.3155}$	35	PSB trawl data	NE Atlantic
<i>Antimora rostrata</i>	$TW(g) = 0.0015 \times (SLcm)^{3.4912}$	244	PSB trawl data	NE Atlantic
<i>Aphanopus carbo</i>	$TW(g) = 0.00003 \times (SLcm)^{3.8474}$	15	PSB trawl data	NE Atlantic
<i>Argentina silus</i>	$TW(g) = 0.0055 \times (SLcm)^{3.2242}$	421	Gordon & Hunter 1994	NE Atlantic
<i>Argentina sphyraena</i>	$TW(g) = 0.0053 \times (TLcm)^{3.053}$	157	Coull et al. 1989	NE Atlantic, $TL(cm) = 1.1027 \times (SLcm) + 0.2154$
<i>Bathypterois dubius</i> <i>Bathypterois longipes</i>	$TW(g) = 0.0074 \times (SLcm)^{2.982}$	123	PSB trawl data	<i>Bathypterois dubius</i> data, NE Atlantic
<i>Bathyraja richardsoni</i>	$TW(g) = 0.0081 \times (TLcm)^{2.9441}$	3	PSB trawl data	NE Atlantic
<i>Bathysaurus ferox</i>	$TW(g) = 0.0006 \times (SLcm)^{3.6252}$	21	PSB trawl data	NE Atlantic
<i>Bathysaurus mollis</i>	$TW(g) = 0.0009 \times (SLcm)^{3.5055}$	4	PSB trawl data	NE Atlantic

<i>Bathytroctes microlepis</i>	$TW(g) = 0.0156 \times (SLcm)^{2.8664}$	17	PSB trawl data	NE Atlantic
<i>Bathytroctes macrolepis</i>	$TW(g) = 0.0032 \times (SLcm)^{3.3948}$	11	PSB trawl data	NE Atlantic
<i>Bathytroctes michaelsarsi</i>	$TW(g) = 0.0165 \times (TLcm)^{2.8148}$	3	PSB trawl data	NE Atlantic
<i>Beryx decadactylus</i>	$TW(g) = 0.027 \times (FLcm)^{2.9738}$	NA	John Gordon	NE Atlantic
<i>Caelorinchus caelorhincus</i>	$TW(g) = 0.00219 \times (TLcm)^{3.106}$	25	Borges et al. 2003	S Portugal $TL(cm)=4.3706 \times (HLcm) + 1.1642$
<i>Caelorinchus labiatus</i>	$TW(g) = 0.145 \times (HLcm)^{3.0234}$	84	PSB trawl data	NE Atlantic
<i>Cataetyx laticeps</i> <i>Cataetyx alleni</i>	$TW(g) = 0.0045 \times (SLcm)^{3.1049}$	22	PSB trawl data	used <i>Cataetyx laticeps</i> data, NE Atlantic
<i>Centrophorus squamosus</i>	$TW(g) = 0.00007* \times (TLcm)^{3.9318}$	NA	John Gordon	NE Atlantic
<i>Centroscymnus coelolepis</i>	$TW(g) = 0.0043 \times (TLcm)^{3.1204}$	227	Gordon & Hunter 1994	NE Atlantic
<i>Chimaera monstrosa</i>	$TW(g) = 0.00014 \times (TLcm)^{3.475}$	22	Borges et al. 2003	S Portugal
<i>Conocara macropterum</i>	$TW(g) = 0.0022 \times (SLcm)^{3.3882}$	25	PSB trawl data	NE Atlantic
<i>Conocara murrayi</i>	$TW(g) = 0.0022 \times (SLcm)^{3.4581}$	23	PSB trawl data	NE Atlantic
<i>Conocara salmoneum</i>	$TW(g) = 0.0025 \times (SLcm)^{3.486}$	7	PSB trawl data	NE Atlantic
<i>Coryphaenoides armatus</i>	$TW(g) = 0.2926 \times (HLcm)^{3.4506}$	761	PSB trawl data	NE Atlantic
<i>Coryphaenoides brevibarbis</i>	$TW(g) = 0.4336 \times (HLcm)^{2.8262}$	69	PSB trawl data	NE Atlantic

<i>Coryphaenoides carapinus</i>	$TW(g) = 0.1734 \times (HLcm)^{3.6864}$	39	PSB trawl data	NE Atlantic
<i>Coryphaenoides guentheri</i>	$TW(g) = 0.8921 \times (HLcm)^{2.5784}$	1023	PSB trawl data	NE Atlantic
<i>Coryphaenoides leptolepis</i>	$TW(g) = 0.2947 \times (HLcm)^{3.1591}$	185	PSB trawl data	NE Atlantic
<i>Coryphaenoides mediterraneus</i>	$TW(g) = 0.4966 \times (HLcm)^{3.0369}$	60	PSB trawl data	NE Atlantic
<i>Coryphaenoides profundicolus</i>	$TW(g) = 0.2155 \times (HLcm)^{3.383}$	17	PSB trawl data	NE Atlantic
<i>Coryphaenoides rupestris</i>	$TW(g) = 0.231 \times (HLcm)^{3.2892}$	342	PSB trawl data	NE Atlantic
<i>Cottunculus thomsonii</i>	$TW(g) = 0.0156 \times (SLcm)^{3.0806}$	12	PSB trawl data	NE Atlantic
<i>Dalatias licha</i>	$TW(g) = 0.0008 \times (TLcm)^{3.433}$	30	John Gordon	NE Atlantic
<i>Deania calcea</i>	$TW(g) = 0.0012 \times (TLcm)^{3.2623}$	494	Gordon & Hunter 1994	NE Atlantic
<i>Dipturus nidarosiensis</i>	$TW(g) = 0.0108 \times (DWcm)^{3.079}$	NA	Coull et al. 1989	used <i>Dipturus batis</i> , NE Atlantic, DW (cm) = TL(cm) / 1.424
<i>Echiodon drummondii</i>	$TW(g) = 0.00059 \times (TLcm)^3$	NA	Pauly et al. 1998	NE Atlantic
<i>Epigonus telescopus</i>	$TW(g) = 0.0111 \times (SLcm)^{3.1601}$	10	PSB trawl data	NE Atlantic
<i>Etmopterus princeps</i>	$TW(g) = 0.0028 \times (TLcm)^{3.1476}$	30	Gordon & Hunter 1994	NE Atlantic
<i>Etmopterus spinax</i>	$TW(g) = 0.0018 \times (TLcm)^{3.2430}$	198	Gordon & Hunter 1994	NE Atlantic
<i>Gaidropsarus macrophthalmus macrophthalmus</i>	$TW(g) = 0.0041 \times (SLcm)^{3.2879}$	4	PSB trawl data	NE Atlantic

<i>Gaidropsarus argentatus</i>	$TW(g) = 0.0065 \times (SLcm)^{3.279}$	NA	John Gordon	NE Atlantic
<i>Galeus melastomus</i>	$TW(g) = 0.00166 \times (TLcm)^{3.145}$	311	Borges et al. 2003	S Portugal
<i>Galeus murinus</i>	$TW(g) = 0.0048 \times (SLcm)^{3.5928}$	NA	John Gordon	NE Atlantic
<i>Glyptocephalus cynoglossus</i>	$TW(g) = 0.0017 \times (TLcm)^{3.39}$	571	Coull et al. 1989	NE Atlantic
<i>Guttigadus latifrons</i>	$TW(g) = 0.0103 \times (SLcm)^{2.8844}$	15	PSB trawl data	NE used <i>Halargyreus johnsonii</i> , NE Atlantic
<i>Halargyreus johnsonii</i>	$TW(g) = 0.0103 \times (SLcm)^{2.8844}$	15	PSB trawl data	NE Atlantic
<i>Halosauropsis macrochir</i>	$TW(g) = 0.0138 \times (GPLcm)^{3.033}$	34	PSB trawl data	NE Atlantic
<i>Halosaurus johnsonianus</i>	$TW(g) = 0.0138 \times (GPLcm)^{3.033}$	34	PSB trawl data	used <i>Halosauropsis macrochir</i> , NE Atlantic
<i>Harriotta raleighana</i>	$TW(g) = 0.0007 \times (TLcm)^{3.38}$	NA	John Gordon	NE Atlantic
<i>Helicolenus dactylopterus dactylopterus</i>	$TW(g) = 0.0128 \times (SLcm)^{3.2604}$	6	PSB trawl data	NE Atlantic
<i>Histiobranchus bathybius</i>	$TW(g) = 0.0002 \times (TLcm)^{3.4447}$	55	PSB trawl data	NE Atlantic
<i>Hoplostethus atlanticus</i>	$TW(g) = 0.00735 \times (SLcm)^{2.7759}$	10	PSB trawl data	NE Atlantic
<i>Hoplostethus mediterraneus mediterraneus</i>	$TW(g) = 0.0417 \times (SLcm)^{2.9121}$	10	PSB trawl data	NE Atlantic
<i>Hydrolagus affinis</i>	$TW(g) = 0.0043 \times (TLcm)^{3.08}$	NA	John Gordon	NE Atlantic

<i>Hydrolagus mirabilis</i>	$TW(g) = 0.0152 \times (TLcm)^{2.2802}$	10	PSB trawl data	NE Atlantic
<i>Ilyophis arx</i>	$TW(g) = 0.000077 \times (SLcm)^{3.6349}$	297	Gordon & Duncan 1987	used <i>Synaphobranchus kaupii</i> , NE Atlantic
<i>Ilyophis blachei</i>	$TW(g) = 0.000077 \times (SLcm)^{3.6349}$	297	Gordon & Duncan 1987	used <i>Synaphobranchus kaupii</i> , NE Atlantic
<i>Ilyophis brunneus</i>	$TW(g) = 0.000077 \times (SLcm)^{3.6349}$	297	Gordon & Duncan 1987	used <i>Synaphobranchus kaupii</i> , NE Atlantic
<i>Lepidion eques</i>	$TW(g) = 0.0026 \times (SLcm)^{3.2909}$	657	PSB trawl data	NE Atlantic
<i>Lepidorhombus boscii</i>	$TW(g) = 0.00349 \times (TLcm)^{3.219}$	7	Borges et al. 2003	S Portugal
<i>Lepidorhombus whiffiagonis</i>	$TW(g) = 0.0008 \times (SLcm)^{3.1109}$	31	John Gordon	NE Atlantic
<i>Leucoraja circularis</i>	$TW(g) = 0.0024 \times (TLcm)^{3.233}$	NA	Dorel 1986	NE Atlantic
<i>Lophius piscatorius</i>	$TW(g) = 0.0063 \times (TLcm)^{3.284}$	47	Borges et al. 2003	S Portugal
<i>Lycodes terraenovae</i>	$TW(g) = 0.00005 \times (TLcm)^{4.1968}$	4	John Gordon	NE Atlantic
<i>Merluccius merluccius</i>	$TW(g) = 0.0047 \times (SLcm)^{3.099}$	NA	John Gordon	NE Atlantic
<i>Microchirus variegatus</i>	$TW(g) = 0.0089 \times (TLcm)^{3.0790}$	5	Coull et al. 1989	NE Atlantic
<i>Molva dypterygia</i>	$TW(g) = 0.0015 \times (TLcm)^{3.228}$	240	Gordon & Hunter 1994	NE Atlantic
<i>Molva macrophthalma</i>	$TW(g) = 0.0009 \times (TLcm)^{3.24}$	NA	John Gordon	NE Atlantic
<i>Mora moro</i>	$TW(g) = 0.0013 \times (SLcm)^{3.6134}$	20	PSB trawl data	NE Atlantic

<i>Narctes stomias</i>	$TW(g) = 0.0072 \times (SLcm)^{3.0679}$	27	PSB trawl data	NE Atlantic
<i>Nezumia aequalis</i>	$TW(g) = 0.8878 \times (HLcm)^{2.7835}$	193	PSB trawl data	NE Atlantic
<i>Notacanthus bonaparte</i>	$TW(g) = 5E-5 \times (TLcm)^{3.9591}$	28	PSB trawl data	NE Atlantic
<i>Notacanthus chemnitzii</i>	$\ln(Wg) = 2.910 * \ln \times (GPLcm)^{-2.979}$	97	Coggan et al. 1998	NE Atlantic
<i>Pachycara crassiceps</i> <i>Pachycara bulbiceps</i>	$TW(g) = 0.0004 \times (TLcm)^{3.673}$	3	PSB trawl data	used <i>Pachycara crassiceps</i> , NE Atlantic
<i>Paraliparis hystrix</i>	$TW(g) = 0.0011 \times (SLcm)^{3.7392}$	12	John Gordon	NE Atlantic
<i>Phycis blennoides</i>	$TW(g) = 0.012 \times (SLcm)^{2.9543}$	8	PSB trawl data	NE Atlantic
<i>Polyacanthonotus challengerii</i>	$TW(g) = 0.0051 \times (GPLcm)^{3.7161}$	5	PSB trawl data	NE Atlantic
<i>Polyacanthonotus rissoanus</i>	$TW(g) = 0.0005 \times (TLcm)^{3.1395}$	77	PSB trawl data	NE Atlantic
<i>Pseudonezumia flagellicauda</i>	$TW(g) = 0.8878 \times (HLcm)^{2.7835}$	193	PSB trawl data	used <i>Nezumia aequalis</i> , NE Atlantic
<i>Rajella bigelowi</i>	$TW(g) = 0.0154 \times (TLcm)^{2.6517}$	5	PSB trawl data	NE Atlantic
<i>Rajella fyllae</i>	$TW(g) = 0.02 \times (TLcm)^{2.5963}$	3	PSB trawl data	NE Atlantic
<i>Rinoctes nasutus</i>	$TW(g) = 0.0137 \times (SLcm)^{2.8874}$	10	PSB trawl data	NE Atlantic
<i>Rouleina attrita</i>	$TW(g) = 0.0137 \times (SLcm)^{2.8874}$	22	John Gordon	NE Atlantic
<i>Scymnodon ringens</i>	$TW(g) = 0.0043 \times (TLcm)^{3.1286}$	62	Gordon & Hunter 1994	NE Atlantic

<i>Spectrunculus grandis</i>	$TW(g) = 0.0083 \times (SLcm)^{2.9416}$	101	PSB trawl data	NE Atlantic
<i>Synaphobranchus kaupii</i>	$TW(g) = 9E-5 \times (TLcm)^{3.5584}$	1532	PSB trawl data	NE Atlantic
<i>Trachyrincus murrayi</i>	$TW(g) = 0.1503 \times (HLcm)^{2.9995}$	44	PSB trawl data	NE Atlantic
<i>Trachyrincus scabrus</i>	$TW(g) = 0.04444 \times (HLcm)^{3.525}$	10	PSB trawl data	NE Atlantic
<i>Trachyscorpia cristulata echinata</i>	$TW(g) = 0.0031 \times (SLcm)^{3.6574}$	4	PSB trawl data	NE Atlantic
<i>Venefica proboscidea</i>	$TW(g) = 0.0083 \times (SLcm)^{2.9416}$	1532	PSB trawl data	used <i>Synaphobranchus kaupii</i> , NE Atlantic

Borges, T.C., Olim, S. and Erzini, K.: Weight-length relationships for fish species discarded in commercial fisheries of the Algarve (southern Portugal), J. Appl. Ichthyol., 19, 394-396.

Coggan, R.A., Gorodn, J.D.M. and Merrett, N.R.: Abundance, distribution, reproduction and diet of notacanthid fishes from the north-east Atlantic, J. Fish Biol., 52, 1038-1057, 1998.

Coull, K.A., Jermyn, A.S., Newton, A.W., Henderson, G.I. and Hall, W.B.: Length/weight relationships for 88 species of fish encountered in the north east Atlantic, Scottish Fisheries Research Report, 43, 1-81, 1989.

Dorel, D.: Poissons de l'Atlantique nord-est relations taille-poids. Institut Francais de Recherche pour l'Exploitation de la Mer. Nantes, France. 165 pp, 1986.

Gordon, J.D.M. and Duncan, J.A.R.: Deep-sea bottom-living fishes at two repeat stations at 2200 and 2900 m in the Rockall Trough, northeastern Atlantic Ocean, Mar. Biol., 96, 309-325, 1987

Gordon, J.D.M. and Hunter, J.E.: Study of deep-water fish stocks to the west of Scotland, Final Report, Scottish Association for Marine Science, 1994.

Data referenced as John Gordon, is unpublished data collected by J.D.M. Gordon at the Scottish Association for Marine Science.