



Supplement of

CO₂ and nutrient-driven changes across multiple levels of organization in *Zostera noltii* ecosystems

B. Martínez-Crego et al.

Correspondence to: B. Martínez-Crego (bmcrego@ualg.pt)

Table S1. Full comparative description of the low- and high-nutrient meadows of *Zostera noltii* in June-August 2011, prior to the start of the experiment. Seagrass and *Ulva* spp. cover were measured within five quadrats of 0.5 m x 0.5 m placed every 5 m along two 25 m transects. Shoot density was measured within three 0.2 m x 0.2 m quadrats. Canopy height was measured in each quadrat ignoring the 20 % tallest leaves. The other seagrass traits were measured within cores of 10.5 cm diameter as described for the experimental samples. Water column pH (ranging from 8.0 to 8.2), salinity (ranging from 38 to 40 psu) and temperature (ranging from 23 to 26 °C) were also measured during low spring and neap tides; they were not significantly different between meadows. Values are means (\pm SE). Unpaired *t* test statistics and *p*-levels are shown. ^aSqrt-transformed data to meet normality. ^bMann-Whitney rank tests were conducted for variables that did not meet normality even after transformation and U statistics are shown.

	Low-nutrient meadow	High-nutrient meadow	<i>n</i>	<i>t/U</i>	<i>p</i> level
Leaf phenolics (mg (gDW) ⁻¹)	48 \pm 1.2	29 \pm 3.2	4	5.6	0.001
Leaf nitrogen (mg (gDW) ⁻¹)	21 \pm 0.8	25 \pm 0.6	4	-3.8	0.01
Leaf C : N	19 \pm 0.7	16 \pm 0.4	4	4.2	0.01
Rhizome nitrogen (mg (gDW) ⁻¹)	5.7 \pm 0.2	6.2 \pm 0.9	4	-0.5	0.63
Rhizome C : N	78 \pm 3.0	73 \pm 14	4	0.3	0.76
Rhizome sucrose (mg Glu (gDW) ⁻¹)	195 \pm 5.5	177 \pm 9.8	4	1.6	0.17
Rhizome starch (mg Glu (gDW) ⁻¹)	473 \pm 14	355 \pm 27	4	3.9	0.01
Rhizome TNC (mg Glu (gDW) ⁻¹)	668 \pm 18	532 \pm 28	4	4.1	0.01
Shoot area (cm ² shoot ⁻¹) ^a	7.8 \pm 0.6	4.5 \pm 0.7	20	3.8	<0.001
Shoot leaves (# of leaves shoot ⁻¹) ^b	3.1 \pm 0.1	2.8 \pm 0.2	20	141	0.053
Above:belowground biomass	1.4 \pm 0.3	1.9 \pm 0.6	4	-0.7	0.51
Rhizome length (cm)	0.7 \pm 0.1	1.0 \pm 0.1	20	-1.9	0.07
<i>Z. noltii</i> density (shoots m ⁻²)	5517 \pm 755	2664 \pm 411	3	3.3	0.03
Density of flowering shoots (# m ⁻²) ^b	0.0 \pm 0.0	11 \pm 11	4	6.0	0.69
<i>Z. noltii</i> canopy height (cm)	25 \pm 3.1	16 \pm 2.0	3	2.4	0.07
<i>Z. noltii</i> cover (% of sediment surface) ^b	96 \pm 2.2	18 \pm 9.8	9-10	0.0	<0.001
Epiphyte load (mg cm ⁻²)	0.4 \pm 0.1	0.8 \pm 0.2	4	-1.7	0.14
<i>Ulva</i> spp. cover (% of sediment surface) ^b	absent	38 \pm 12	9-10	10	0.001
Fish herbivory (% leaves with bite marks shoot ⁻¹) ^b	14 \pm 5.0	13 \pm 4.7	20	199	0.99
Seawater nitrate (μ M) ^b	< 0.01	1.1 \pm 0.2	4	0.0	0.03
Seawater ammonium (μ M)	0.7 \pm 0.2	3.0 \pm 0.4	11	-4.8	<0.001
Seawater phosphate (μ M)	0.5 \pm 0.1	1.2 \pm 0.1	11	-5.4	<0.001

Table S2. Seawater chemistry within the experimental mesocosms. Values are means (\pm SE). Nutrient samples were weekly collected ($n = 24$ nutrient-unfertilized; $n = 29$ nutrient-enriched). Temperature ($n = 44$) and pH were measured twice a week at approximately the same hour of the day using a Multimeter 340 (WTW, Weilheim, Germany) corrected for temperature and calibrated using NBS buffers ($n = 47$). Salinity was measured weekly using a refractometer ($n = 22$). Total alkalinity (TA, $n = 16$) was analyzed on samples collected the first and last week of the experiment and preserved in Winkler bottles poisoned with HgCl_2 saturated solution. TA was estimated by Gran titration with a Metrohm 794 Titroprocessor (accuracy 1%, checked using Certified Reference Material CRM batch #110; Scripps Institution of Oceanography, La Jolla, CA, USA). Total dissolved inorganic carbon (DIC) and its speciation into carbon dioxide (CO_2), bicarbonate (HCO_3^-) and carbonate (CO_3^{2-}) were calculated from mean values of TA, pH, temperature and salinity using the CO2SYS program (Pelletier et al. 1997) with the constants of Mehrbach refitted by Dickson & Millero (1987). Different superscript letters indicate significant differences tested with unpaired *t* tests. ^aMann-Whitney rank tests were conducted for variables that did not meet normality even after transformation.

	Nutrient-unfertilized	Nutrient-enriched
Ammonium (μM) ^a	$0.3 \pm 0.1^{\text{a}}$	$74 \pm 1.4^{\text{b}}$
Nitrate (μM) ^a	$2.3 \pm 0.2^{\text{a}}$	$44 \pm 1.7^{\text{b}}$
Phosphate (μM)	$0.3 \pm 0.02^{\text{a}}$	$3.9 \pm 0.1^{\text{b}}$
N:P molar ratio	$3.4 \pm 0.3^{\text{a}}$	$14 \pm 0.2^{\text{b}}$
	CO ₂ -unfertilized	CO ₂ -enriched
pH*	$8.02 \pm 0.01^{\text{a}}$	$7.86 \pm 0.004^{\text{b}}$
Temperature ($^{\circ}\text{C}$) ^a	24 ± 0.3	24 ± 0.3
Salinity (psu)	37 ± 0.1	37 ± 0.1
TA* ($\mu\text{mol (Kg seawater)}^{-1}$)	2668 ± 20	2659 ± 14
DIC ($\mu\text{mol (Kg seawater)}^{-1}$)	2422	2496
CO ₂ ($\mu\text{mol (Kg seawater)}^{-1}$)	20	30
HCO ₃ ⁻ ($\mu\text{mol (Kg seawater)}^{-1}$)	2212	2328
CO ₃ ²⁻ ($\mu\text{mol (Kg seawater)}^{-1}$)	189	138

References

- Pelletier, G., Lewis E., and Wallace, D.: CO2 sys.xls (version 1.0). A calculator for the CO₂ system in seawater for Microsoft Excel/VBA. Washington State Department of Ecology, Olympia, WA, 1997.
- Dickson, A. G. and Millero, F. J.: A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media. Deep-Sea Research, 34, 1733-1743, 1987.

SECTION S3. Full results of the response to CO₂ and nutrient additions of *Zostera noltii* plant-, community-, and ecosystem-level traits measured through time.

Table S3. Results of three-way RM ANOVA tests to assess the effects of CO₂ and nutrient additions (among-subject factors) on *Zostera noltii* plant-, community-, and ecosystem-level traits through time (within-subject factor).

Variable	LOW-NUTRIENT MEADOW					HIGH-NUTRIENT MEADOW				
	SS	df	MS	F	p level	SS	df	MS	F	p level
Shoot recruitment (%)										
Time	6402	4.0	1600	35	<0.0001	32991	4.0	8248	219	<0.0001
Time x CO ₂	139	4.0	35	0.8	0.57	1160	4.0	290	7.7	0.001
Time x Nutrients	350	4.0	87	1.9	0.16	198	4.0	49	1.3	0.31
Time x CO ₂ x Nut	553	4.0	138	3.0	0.05	852	4.0	213	5.7	0.005
Error (Time)	742	16	46			603	16	38		
CO ₂	33	1.0	33	0.1	0.77	2498	1.0	2498	6.5	0.06
Nutrients	3425	1.0	3425	10	0.03	867	1.0	867	2.2	0.21
CO ₂ x Nut	1741	1.0	1741	5.2	0.08	4.7	1.0	4.7	0.01	0.92
Error	1327	4.0	332			1549	4.0	387		
Leaf area index (m² m⁻²)^{a, c}										
Time	38	4.0	9.5	24	<0.0001	9	5.0	2	57	<0.0001
Time x CO ₂	3.6	4.0	0.9	2.3	0.11	0.1	5.0	0.03	0.9	0.50
Time x Nutrients	11	4.0	2.7	6.6	0.002	0.2	5.0	0.03	1.0	0.46
Time x CO ₂ x Nut	3.7	4.0	0.9	2.3	0.11	0.1	5.0	0.03	0.8	0.54
Error (Time)	6.5	16	0.4			0.7	20	0.03		
CO ₂	2.7	1.0	2.7	2.0	0.23	0.003	1.0	0.003	0.01	0.93
Nutrients	50	1.0	50	37	<0.005	0.2	1.0	0.2	0.5	0.50
CO ₂ x Nut	0.5	1.0	0.5	0.4	0.58	0.1	1.0	0.1	0.2	0.70
Error	5.4	4.0	1.3			1.2	4.0	0.3		
Density of flowering shoots (# m⁻²)^{b, c, d, e}										
Time	194	1.4	135	0.6	0.54	323	1.4	232	1.2	0.34
Time x CO ₂	194	1.4	135	0.6	0.54	202	1.4	145	0.8	0.46
Time x Nutrients	194	1.4	135	0.6	0.54	202	1.4	145	0.8	0.46
Time x CO ₂ x Nut	776	1.4	539	2.3	0.19	323	1.4	232	1.2	0.34
Error (Time)	1359	5.8	236			1051	5.6	188		
CO ₂	109	1.0	109	0.5	0.51	40	1.0	40	0.4	0.56
Nutrients	109	1.0	109	0.5	0.51	40	1.0	40	0.4	0.56
CO ₂ x Nut	12	1.0	12	0.1	0.82	162	1.0	162	1.6	0.27
Error	825	4.0	206			404	4.0	101		
Ulva cover (%)^{c, e}										
Time	-	-	-	-	-	17352	1.6	10863	255	<0.0001
Time x CO ₂	-	-	-	-	-	7	1.6	4.1	0.1	0.87
Time x Nutrients	-	-	-	-	-	137	1.6	86	2.0	0.21
Time x CO ₂ x Nut	-	-	-	-	-	6	1.6	4.0	0.1	0.87
Error (Time)	-	-	-	-	-	272	6.4	43		
CO ₂	-	-	-	-	-	3	1.0	2.5	0.1	0.76
Nutrients	-	-	-	-	-	10	1.0	10.1	0.4	0.54
CO ₂ x Nut	-	-	-	-	-	1	1.0	0.8	0.03	0.86
Error	-	-	-	-	-	90	4.0	23		
Meso-herbivory incidence(% leaves with bite marks shoot⁻¹)^{b, c, d, e}										
Time	247	1.5	162	2.9	0.14	40946	1.8	22538	40.3	0.0001
Time x CO ₂	124	1.5	81	1.4	0.30	356	1.8	196	0.4	0.70
Time x Nutrients	122	1.5	80	1.4	0.30	982	1.8	540	1.0	0.42
Time x CO ₂ x Nut	203	1.5	134	2.3	0.18	2215	1.8	1219	2.2	0.18
Error (Time)	346	6.1	57			4059	7.3	559		
CO ₂	29	1.0	29	0.7	0.46	122	1.0	122	0.1	0.75
Nutrients	41	1.0	41	0.9	0.39	232	1.0	232	0.2	0.66


Variable	LOW-NUTRIENT MEADOW					HIGH-NUTRIENT MEADOW				
	SS	df	MS	F	p level	SS	df	MS	F	p level
CO ₂ x Nut	79	1.0	79	1.8	0.25	3417	1.0	3417	3.4	0.14
Error	175	4.0	44			4050	4.0	1013		
# of leaves (# shoot ⁻¹) ^{c, e}										
Time	3.3	4.0	0.8	22	<0.0001	9	5.0	1.8	26	<0.0001
Time x CO ₂	0.8	4.0	0.2	5.3	0.01	0.3	5.0	0.1	0.9	0.49
Time x Nutrients	1.6	4.0	0.4	11	<0.001	0.7	5.0	0.1	1.9	0.15
Time x CO ₂ x Nut	0.4	4.0	0.1	2.9	0.054	1.0	5.0	0.2	3.0	0.04
Error (Time)	0.6	16	0.04			1.0	15	0.1		
CO ₂	0.2	1.0	0.2	2.4	0.19	0.2	1.0	0.2	31	0.01
Nutrients	1.9	1.0	1.9	29	0.01	0.1	1.0	0.1	17	0.03
CO ₂ x Nut	0.03	1.0	0.03	0.4	0.57	0.2	1.0	0.2	28	0.01
Error	0.3	4	0.1			0.02	3.0	0.01		
Detritus production (g FW day ⁻¹)										
Time	26	3.0	8.8	32	<0.0001	30	3.0	10	73	<0.0001
Time x CO ₂	2.1	3.0	0.7	2.5	0.11	2.5	3.0	0.8	6.1	0.01
Time x Nutrients	3.1	3.0	1.0	3.7	0.04	0.4	3.0	0.1	1.0	0.41
Time x CO ₂ x Nut	1.2	3.0	0.4	1.4	0.29	0.6	3.0	0.2	1.4	0.30
Error (Time)	3.4	12	0.3			1.7	12	0.1		
CO ₂	6.8	1.0	6.8	16	0.02	0.3	1.0	0.3	1.1	0.34
Nutrients	2.1	1.0	2.1	4.8	0.09	3.3	1.0	3.3	12	0.03
CO ₂ x Nut	0.01	1.0	0.01	0.02	0.90	0.6	1.0	0.6	2.0	0.23
Error	1.7	4.0	0.4			1.1	4.0	0.3		
Net community production (mmol O ₂ m ⁻² h ⁻¹) ^{b, c}										
Time	1904	1.6	1191	15	0.005	3578	1.7	2064	28	0.001
Time x CO ₂	548	1.6	343	4.4	0.07	352	1.7	203	2.8	0.13
Time x Nutrients	58	1.6	36	0.5	0.61	449	1.7	259	3.5	0.09
Time x CO ₂ x Nut	930	1.6	581	7.4	0.02	70	1.7	40	0.6	0.58
Error (Time)	502	6.4	78			508	6.9	73		
CO ₂	85	1.0	85	2.6	0.18	4	1.0	4	0.1	0.79
Nutrients	50	1.0	50	1.5	0.29	82	1.0	82	1.6	0.27
CO ₂ x Nut	3.0	1.0	3.0	0.1	0.78	166	1.0	166	3.3	0.14
Error	131	4.0	33			201	4.0	50		
Community respiration (mmol O ₂ m ⁻² h ⁻¹) ^c										
Time	87	4.0	22	4.3	0.01	25	1.7	14	2.1	0.19
Time x CO ₂	51	4.0	13	2.5	0.08	46	1.7	27	4.0	0.08
Time x Nutrients	66	4.0	17	3.3	0.04	5.4	1.7	3.1	0.5	0.62
Time x CO ₂ x Nut	37	4.0	9.3	1.9	0.17	3.1	1.7	1.8	0.3	0.74
Error (Time)	81	16	5.0			46	6.9	6.7		
CO ₂	0.6	1.0	0.6	0.1	0.81	8	1.0	7.9	2.8	0.17
Nutrients	20	1.0	20	2.4	0.20	0.1	1.0	0.1	0.03	0.86
CO ₂ x Nut	5.2	1.0	5.2	0.6	0.48	0.003	1.0	0.003	0.001	0.97
Error	34	4.0	8.6			11	4.0	2.8		

^a Data from the high-nutrient meadow were sqrt-transformed to meet normality.

^{b, c} Corrected significance levels from Greenhouse-Geisser adjustment were used when sphericity was not met in data from the low-nutrient (b) or high-nutrient (c) meadows.

^{d,e} Violation of normality was allowed on data from the low-nutrient (d) or high-nutrient (e) meadows. For those variables, the significance level was more restrictive ($p < 0.03$) to minimize the possibility of Type I error.

Fig. S3. *Zostera noltii* plant- and community-level traits from the low-nutrient and high-nutrient meadow showing no significant response to CO₂ and nutrient additions through time based on RM ANOVA tests. Symbols are means (+SE, *n*=2). Detailed RM ANOVA results are shown on Table S3.



* Variables that did not meet normality even after transformation.


SECTION S4. Full results of the response to CO₂ and nutrient additions of *Zostera noltii* plant-, community-, and ecosystem-level traits measured at the end of the experiment.

Table S4. Results of two-way ANOVA tests to assess the effects of CO₂ and nutrient additions (fixed crossed factors) on *Z. noltii* plant-, community-, and ecosystem-level traits measured at the end of the experiment. Restrictive significant *p* levels (*p* < 0.03) were selected to minimize the possibility of Type I error due to unequal variances and are highlighted in black. Significant Welch's *t* tests used to interpret significant interactions are shown. Biomass above:belowground allocation was not measured in the high-nutrient meadow due to the reduced number of shoots at the end of the experiment.

Variable	LOW-NUTRIENT MEADOW					<i>t</i> test	HIGH-NUTRIENT MEADOW					
	SS	df	MS	F	<i>p</i> level		SS	df	MS	F	<i>p</i> level	<i>t</i> test
Leaf nitrogen (mg (gDW) ⁻¹)												
CO ₂	0.6	1	0.6	0.1	0.80		37	1	37	2.4	0.19	
Nutrients	687	1	687	89	<0.001		260	1	260	17	0.01	
CO ₂ x Nut	0.002	1	0.002	0.0002	0.99		24	1	24	1.6	0.28	
Error	31	4	7.7				62	4	15			
Rhizome nitrogen (mg (gDW) ⁻¹)												
CO ₂	0.6	1	0.6	0.3	0.60		22	1	22	5.7	0.08	
Nutrients	47	1	47	27	<0.01		1.3	1	1.3	0.3	0.60	
CO ₂ x Nut	1.0	1	1.0	0.6	0.48		1.4	1	1.4	0.4	0.58	
Error	7.0	4	1.8				15	4	3.8			
Leaf C : N												
CO ₂	0.004	1	0.004	0.01	0.91		56	1	56	7.6	0.051	
Nutrients	77	1	77	276	<0.001		128	1	128	17	0.01	
CO ₂ x Nut	0.3	1	0.3	1.0	0.36		19	1	19	2.6	0.18	
Error	1.1	4	0.3				29	4	7.3			
Rhizome C : N												
CO ₂	0.5	1	0.5	0.1	0.83		315	1	315	15	0.02	
Nutrients	296	1	296	30	0.01		5.6	1	5.6	0.3	0.64	
CO ₂ x Nut.	0.9	1	0.9	0.1	0.78		1.5	1	1.5	0.1	0.80	
Error	39	4	10				87	4	22			
Rhizome sucrose (mg Glu (gDW) ⁻¹)												
CO ₂	66	1	66	1.1	0.36		175	1	175	0.3	0.61	
Nutrients	13	1	13	0.2	0.67		350	1	350	0.6	0.48	
CO ₂ x Nut	18	1	18	0.3	0.61		2	1	1.8	0.003	0.96	
Error	247	4	62				1661	3	554			
Rhizome starch (mg Glu (gDW) ⁻¹)												
CO ₂	1298	1	1298	5.2	0.08		9101	1	9101	1.5	0.31	
Nutrients	928	1	928	3.7	0.13		1374	1	1374	0.2	0.67	
CO ₂ x Nut	2092	1	2092	8.4	0.04		15893	1	15893	2.5	0.21	
Error	998	4	249				18783	3	6261			
Rhizome non-structural carbohydrates (mg Glu (gDW) ⁻¹)												
CO ₂	1951	1	1951	4.0	0.12		11805	1	11805	1.2	0.36	
Nutrients	1160	1	1160	2.4	0.20		3110	1	3110	0.3	0.62	
CO ₂ x Nut	2503	1	2503	5.1	0.09		15554	1	15554	1.5	0.30	
Error	1958	4	489				30626	3	10209			
Leaf phenolics (mg (gDW) ⁻¹)												
CO ₂	5	1	5	0.9	0.39		952	1	952	20	0.01	t= 4.5; p=0.13
Nutrients	75	1	75	13	0.02		450	1	450	9.3	0.04	t= 3.4; p=0.11
CO ₂ x Nut	6	1	6	1.1	0.36		632	1	632	13	0.02	t= 4.3; p=0.15
Error	23	4	5.6				194	4	49			

LOW-NUTRIENT MEADOW						HIGH-NUTRIENT MEADOW						
Variable	SS	df	MS	F	p level	t test	SS	df	MS	F	p level	t test
Above:Belowground biomass												
CO ₂	2.2	1	2.2	1.7	0.26		-	-	-	-	-	
Nutrients	2.5	1	2.5	1.9	0.24		-	-	-	-	-	
CO ₂ x Nut	0.9	1	0.9	0.7	0.45		-	-	-	-	-	
Error	5.1	4	1.3				-	-	-			
Rhizome length (cm)												
CO ₂	0.1	1	0.1	0.6	0.49		0.4	1	0.4	6.6	0.06	
Nutrients	0.1	1	0.1	1.3	0.32		0.7	1	0.7	13	0.02	
CO ₂ x Nut	0.2	1	0.2	1.7	0.27		0.6	1	0.6	11	0.03	
Error	0.4	4	0.1				0.2	4	0.1			
Total root length (cm)												
CO ₂	0.1	1	0.1	0.04	0.86		0.01	1	0.01	0.02	0.91	
Nutrients	0.3	1	0.3	0.2	0.70		0.8	1	0.8	0.9	0.41	
CO ₂ x Nut	0.3	1	0.3	0.3	0.66		3.4	1	3.4	4.1	0.14	
Error	2.7	2	1.4				2.5	3	0.8			
Epiphyte biomass (mg cm ⁻²)												
CO ₂	3.7	1	3.7	6.8	0.06	t= -9.5; p=0.05	16	1	16	24	<0.01	t= -8.7; p=0.01
Nutrients	0.7	1	0.7	1.2	0.33	t= -3.6; p=0.17	1.0	1	1.0	1.6	0.28	t= -2.3; p=0.20
CO ₂ x Nut	14	1	14	25	0.01	t= -5.7; p=0.11	5.1	1	5.1	7.8	0.049	t= -5.6; p=0.03
Error	2.2	4	0.6				2.6	4	0.7			
Sediment organic matter (% DW)												
CO ₂	0.18	1	0.18	2.2	0.21	t= 5.7; p=0.03	0.2	1	0.2	0.5	0.53	
Nutrients	0.12	1	0.12	1.5	0.29	t= 5.2; p=0.04	2.1	1	2.1	4.0	0.12	
CO ₂ x Nut	4.1	1	4.1	52	0.002	t= 2.2; p=0.20	1.2	1	1.2	2.4	0.20	
Error	0.3	4	0.1				2.1	4	0.5			

Fig. S4. *Zostera noltii* plant-level traits from the low-nutrient (grey bars) and high-nutrient meadow (black bars) showing no significant response (p level ≥ 0.03) to CO₂ and nutrient additions at the end of the experiment based on two-way ANOVA tests. Bars are means (\pm SE, $n=2$). Detailed ANOVA results are shown on Table S4. TNC refers to total non-structural carbohydrates.



SECTION S5. Results of principal component analyses.

Table S5. Variable loadings in the PCA of responsive variables from the low- and high-nutrient meadows at the end of the experiment (scaling 2, correlation biplot). Descriptors showing the highest correlation with each component ($r \geq 0.7$) were selected to interpret the components.

	pc-1	pc-2
Rhizome starch	0.37	-0.29
Leaf phenolics	0.66	-0.56
Leaf C : N	0.75	-0.52
Rhizome C : N	0.40	-0.60
Epiphyte biomass	-0.19	0.74
Sediment organic matter	-0.85	-0.44
Rhizome length	-0.24	0.40
Shoot mortality	0.89	0.22
Leaf area index	-0.74	-0.52
Meso-herbivory	0.92	0.27
Net community production	-0.93	-0.26
Detritus	-0.20	0.51
Epiphyte composition	-0.10	0.70
Shoot # of leaves	-0.71	-0.10

Table S5. Variable loadings in the PCAs of *Zostera noltii* plant- community and ecosystem- level responses to treatments through time (scaling 2, correlation biplot). Descriptors showing the highest correlation with each component ($r \geq 0.7$) were selected to interpret the components.

	Low-nutrient meadow		High-nutrient meadow	
	pc-1	pc-2	pc-1	pc-2
Detritus	0.01	0.30	0.85	-0.05
Mortality	0.92	0.00	0.93	0.24
Leaf area index	-0.91	0.07	-0.95	-0.18
Shoot # of leaves	-0.82	0.14	-0.86	0.05
Meso-herbivory	0.71	0.01	0.90	0.13
Net community production	-0.37	-0.72	-0.68	-0.05
Community respiration	0.45	-0.67	0.34	-0.79
Flowering shoots	-0.30	-0.70	-0.45	0.61
Ulva cover	-	-	-0.98	-0.07

Fig. S5. Ordination of treatments in the PCA of responsive *Zostera noltii* plant-, community-, and ecosystem-level traits at the end of the experiment. PCA was conducted using the mesocosms from low- and high-nutrient meadows as replicates in order to balance the number of cases and variables. The variable loadings (grey lines) reflect the correlation to the components and the angles between lines are proportional to their covariances. Symbols represent treatments according to the legend and symbol shading differentiates mesocosms from the low-nutrient (grey) and high-nutrient meadow (black). LAI refers to leaf area index, NCP to net community production, CR to community respiration, OM to organic matter and TNC to total non-structural carbohydrates. Epiphyte assemblage refers to the score on the axis II of the NMDS.

