

Biogeochemistry of the North Atlantic during Oceanic Anoxic Event 2: role of changes in ocean circulation and phosphorus input

I. Ruvalcaba Baroni¹, R. P. M. Topper¹, N. A. G. M. van Helmond¹, H. Brinkhuis^{1,2}, and C. P. Slomp¹

¹Utrecht University, Faculty of Geosciences, The Netherlands

²Royal Netherlands Institute for Sea Research (NIOZ), Den Burg, Texel, The Netherlands

December 4, 2013

Supplementary Information




Figure S1: Profiles of a) mean temperature and b) salinity and c) a temperature (T)-salinity (S) diagram for several regions of the proto-North Atlantic, where the basin average is also shown with a dashed line (Topper et al., 2011). The TS diagram highlights that the proto-North Atlantic was poorly stratified. Note that the temperatures may be underestimated since the model uses yearly averaged air temperature as its forcing.

Table S1: Compilation of particulate organic carbon (POC) contents and various proxies for redox conditions such as POC/P_{TOT} ratios, biomarkers, trace metals (TM) and indicators for bioturbation for pre-OAE2 and OAE2 sediments in the proto-North Atlantic.

Sites	Location	Depth (m)	POC (wt%) pre-OAE2	OAE2	POC/P _{TOT} pre-OAE2	OAE2	Biomarkers, TM pre-OAE2	OAE2
CENTRAL OPEN OCEAN								
<i>Rise hill</i>								
105		4100	~3 ^[25]	9.5 ^[3]	n.a.	n.a.	low Mo ^[30]	
603	Cape Hatteras	4000	0.5-1 ^[1]	3 ^[2]	n.a.	330 ^[2]	low Mo ^[30]	
<i>Hatteras Abyssal Plain</i>								
417		5500	n.d.	5.8 ^[3]	n.a.	n.a.		
638-641	Hatteras Abyssal Plain	>3000	<0.5 ^[1]	11.3 ^[4,126]	n.a.	n.a.		
551	Galicia Bank	1500	n.d.	6.5 ^[3]	n.a.	n.a.		
1276	Goban Spur	>2000	<0.5 ^[5,6]	5 ^[5,6]	n.a.	n.a.		
135	Newfoundland basin	>2000	n.d.	7 ^[7]	n.a.	n.a.		
138		>2000	n.d.	7 ^[7,8]	n.a.	n.a.		
386	Bermuda Rise	<3200	n.d.	>5 ^[33]	n.a.	n.a.		
398	Galicia Plateau	<700	~0.5 ^[1,22]	2.8 ^[1,22]	n.a.	n.a.		
<i>AVERAGE</i>								
144		3000	5 ^[4]	6 ^[3]	n.a.	n.a.	low isoren. ^[4]	
367	Cape Verde Basin	3700	6.4 ^[4]	21 ^[3,4]	227 ^[28]	552 ^[28]	high Mo ^[30]	
368	Cape Verde Rise	3700	n.d.	9.2 ^[3]	n.a.	n.a.	low isoren. ^[4]	
<i>AVERAGE</i>								
SOUTHERN OPEN OCEAN								
1258-1261		500-1500	7.6 ^[18]	15.6 ^[18]	114 ^[2]	730 ^[2]	high isoren., chlorobac., TM enrich, high lycoc./n-alkane ^[18]	
<i>SOUTHERN COAST</i>								
	Demerara Rise						high isoren., chlorobac.	
<i>AVERAGE</i>								
<i>S_n</i>								
	Tarfaya Basin	~250	5.5 ^[2,4,19,20]	8.5 ^[2,4,19]	160 ^[2,19]	458 ^[2,27,19]	high isoren., chlorobac.	
<i>Agadir Basin</i>								
<i>AVERAGE</i>								
WESTERN INTERIOR SEAWAY								
<i>Colorado</i>								
El Pueblo	Portland Colorado	<700	n.a.	1 ^[9]	n.a.	n.a.	250 ^[9]	
USGS	Rock Canyon	<700	n.a.	2 ^[10]	n.a.	n.a.		
<i>AVERAGE</i>								
NORTHERN COAST								
627, 635	Bahamas	<500	<0.5 ^[12]	1-5 ^[12]	n.a.	n.a.		
1050, 1052	Blake Nose	~500	<0.5 ^[13,14]	1-5 ^[13,14]	n.a.	n.a.		
	Bass River	30	0.6 ^[15,16,17]	1.1 ^[15,16,17]	45 ^[17]	100 ^[17]		

Sites	Location	Depth (m)	POC (wt%) pre-OAE2	continuation			Biomarkers, TM pre-OAE2	OAE2
				OAE2	POC/P _{TOT} pre-OAE2	OAE2		
AVERAGE								
	NORTH-EASTERN COAST							
549	Wunstorf Germany	~100	<0.1[21,22]	0.7[21,22]	10[21,22]	33[21,22]	Si/Al, K/Al, Mn/Al rise[21]	
	Eastborn England	~100	>0.1[23]	0.2[23]	n.a.	n.a.	fall of I/Ca ^[31]	
							and Mn, low Mo and Fe _T /Al ^[30]	
	Vocontian Basin France	~200	<0.1[22,23]	0.5-1[22,23]	n.a.	n.a.		
	Goban Spur	<700	<0.1[13,14]	<1[13,14]	n.a.	n.a.		
	AVERAGE		~0.1	~1	10	33		
	TETHYS GATEWAY							
	Coast of Spain	<700	<0.5[24]	0.5-1 ^[24]	n.a.	n.a.		
	Hedionda	<700	<0.5[24]	1-5 ^[24]	n.a.	n.a.		
	Maniwa	<700	0.5-1[9]	0.5-1 ^[9]	n.a.	750 ^[9]		
	Coast of N. Africa and Spain	<700	n.d.	0.31 ^[8]	n.a.	n.a.		
	AVERAGE		~0.5	~1	750			

n.a. = not available; n.d. = not detected; [1]: Thunow et al., 1988; [2]: Kraal et al., 2010b; [3]: Simminghe Damst  and K ster, 1998; [4]: Kuypers et al., 2002; [5]: Simminghe Damst  et al., 2007; [6]: Urquhart et al., 1980; [7]: Lancelot et al., 1980; [8]: Herbin et al., 1986; [9]: Mort et al., 2007; [10]: Meyers et al., 2001; [11]: Snow et al., 2005; [12]: Austin et al., 1986; [13]: Norris et al., 2001; [14]: Barker et al., 2001; [15]: Sugarman et al., 1999; [16]: Bowman and Bralower, 2005; [17]: this study; [18]: van Bentum et al., 2009; [19]: Nederbragt et al., 2004; [20]: Kolomic et al., 2005; [21]: H tzel et al., 2011; [22]: Arthur and Dean, 1988; [23]: Voigt, 2000; [24]: Rodr guez-Tovar et al., 2010; [25]: Kuypers et al., 2004; [26]: Kuhnt et al., 1990; [27]: Mort et al., 2008; [28]: M. M. Kuypers, personal communication; [29]: H tzel et al., 2009; [30]: Owens et al., 2012; [31]: van Bentum et al., 2012; [32]: Pancost et al., 2004; [33]: Kuhnt and Wiedmann, 1995

Table S2: Water cycle: Reservoirs and fluxes for pre-OAE2 and OAE2 conditions as implemented in the multi-box model. Note that several fluxes change direction from pre-OAE2 to OAE2 conditions.

Box	Water reservoir	Description	
Name	pre-OAE2	OAE2	units in Tm^3
W1s	1522.11	1554.12	Surface water of the central open ocean
W1i	8623.65	8805.83	Intermediate water of the central open ocean
W1b	38518.87	40583.26	Bottom water of the central open ocean
W2s	254.6	259.51	Surface water of the southern open ocean
W2i	1527.58	1557.09	Intermediate water of the southern open ocean
W2b	7056.33	7408.83	Bottom water of the southern open ocean
W3s	53.47	227.61	Surface water of the Western Interior Seaway
W3b	139.12	387.49	Shallow bottom water of the Western Interior Seaway
W4s	129.37	202.61	Surface water of the northern coast
W4b	340.46	461.35	Shallow bottom water of the northern coast
W5s	129.17	202.41	Surface water of the southern coast
W5b	340.21	461.1	Shallow bottom water of the southern coast
W6s	47.75	76.25	Surface water of the north-eastern coast
W6b	128.28	180.06	Shallow bottom water of the north-eastern coast
W7s	35.66	34.29	Surface water of the Tethys gateway
W7b	91.35	125.63	Shallow bottom water of the Tethys gateway
Name	Water flux (Sv)	Description	
	pre-OAE2	OAE2	($\times 3.1536$ for Tm^3 y^{-1})
F1	4.91	7.09	Net surface water flux from W1s to the Pacific Ocean
F3	9.26	15.15	Net intermediate water flux from W1i to the Pacific Ocean
F6	13.75	21.98	Net bottom water flux from the Pacific Ocean to W1b
F7	1.3	6.67	Net surface water flux from the Tethys gateway to W7s
F10	0.88	6.47	Net bottom water flux from W7b to the Tethys gateway
F11	1.32	1.32	Evaporation in W1s
F12	0.81	0.81	Precipitation in W1s
F13	6.57	7.46	Surface water flux from W4s to W1s
F14	0.59	3.36	Surface water flux from W1s to W6s
F15	0.65	4.15	Surface water flux from W7s to W1s
F16	10.6	17.13	Downwelling from W1s to W1i
F17	2.89	7.83	Upwelling from W1i to W1s
F18	1.02	1.17	Surface water flux from W5s to W2s (Changes direction during OAE2)
F19	1.75	2.49	Surface water flux from W1s to W3s
F20	10.35	9.25	Intermediate water flux from W1i to W3b
F21	5.64	13.77	Shallow bottom water flux from W4b to W1i
F22	0.61	2.89	Intermediate water flux from W1i to W6b
F23	0.12	8.64	Intermediate water flux from W1i to W7b
F24	4.65	2.73	Shallow bottom water flux from W5b to W2i
F25	0.12	5.86	Deep bottom water flux from W1b to W7b
F26	1.71	1.75	Downwelling water flux from W7s to W7b
F27	2.12	3.44	Upwelling water flux from bottom W7b to W7s
F28	0.75	1.31	Downwelling water flux from W5s to W5b
F29	4.38	5.74	Upwelling water flux from W5b to W5s

continuation

Name	Water flux (Sv) pre-OAE2	Water flux (Sv) OAE2	Description ($x 3.1536$ for Tm^3 y $^{-1}$)
F30	0.3	0.3	River water flux to W5st
F31	5.21	6.05	Downwelling water flux from W3s to W3b
F32	6.9	7.38	Upwelling water flux from W3b to W3s
F33	5.39	3.83	Downwelling water flux from W4s to W4b
F34	8.24	4.62	Upwelling water flux from W4b to W4s
F35	1.1	3.17	Downwelling water flux from W6s to W6b
F36	0.43	1.15	Upwelling water flux from W6b to W6s
F37	8.38	19.16	Downwelling water flux from W1i to W1b
F38	21.23	33.31	Upwelling water flux from W1b to W1i
F39	3.53	3.97	Surface water flux from W3s to W4s
F40	0.08	2.61	Surface water flux from W6s to W4s
F41	0.15	1.25	Surface water flux from W7s to W6s
F42	0.91	2.95	Surface water flux from W7s to W5s
F43	0.41	4.62	Shallow bottom water flux from W7b to W5b
F44	1.46	1.73	Shallow bottom water flux from W6b to W7b (Changes direction during OAE2)
F45	0.18	6.63	Shallow bottom water flux from W4b to W6b (Changes direction during OAE2)
F46	8.67	7.93	Shallow bottom water flux from W3b to W4b
F47	3.85	4.59	Surface water flux from W1s to W2s
F48	8.27	6.87	Surface water flux from W2s to W1s
F49	8.34	2.1	Intermediate water flux from W1i to W5b
F50	9.92	11	Intermediate water flux from W1i to W2i
F51	12.42	11.79	Intermediate water flux from W2i to W1i
F52	15.37	17.73	Deep bottom water flux from W1b to W2b
F53	14.6	15.76	Deep bottom water flux from W2b to W1b
F54	3.82	8.85	Surface water flux from W5s to W1s
F55	0.12	0.15	Downwelling water flux from W2s to W2i
F56	3.51	3.6	Upwelling water flux from W2i to W2s
F57	0.37	0.37	Downwelling water flux from W2i to W2b
F58	1.62	1.9	Upwelling water flux from bottom W2b to W2i
F59	0.48	0.45	Shallow bottom water flux from W5b to W2b (Changes direction during OAE2)
R3	0.1	0.15	River water flux to W3s
R4	0.1	0.1	River water flux to W4s
R6	0.01	0.01	River water flux to W6s

Table S3: Carbon cycle: Reservoir sizes and fluxes for pre-OAE2 and OAE2 conditions and pre-OAE2 biogeochemical fluxes as implemented in the multi-box model. Note that all OAE2 reservoirs and fluxes, including the ones in Table S4 and S5, are the ones obtained after equilibrium.

Name	C reservoir pre-OAE2	C reservoir OAE2	Description units in Tmol C
POC1s	4.84	4.54	Organic C in the surface water of the central open ocean
POC1i	9.14	8.58	Organic C in the intermediate water of the central open ocean
POC1b	8.17	7.67	Organic C in the bottom water of the central open ocean
POC2s	2.16	1.74	Organic C in the surface water of the southern open ocean
POC2i	6.48	5.23	Organic C in the intermediate water of the southern open ocean
POC2b	2.24	1.81	Organic C in the bottom water of the southern open ocean
POC3s	0.23	0.52	Organic C in the surface water of the Western Interior Seaway
POC3b	0.01	0.03	Organic C in the bottom water of the Western Interior Seaway
POC4s	0.69	0.49	Organic C in the surface water of the northern coast
POC4b	0.07	0.05	Organic C in the bottom water of the northern coast
POC5s	1.1	1.37	Organic C in the surface water of the southern coast
POC5b	0.29	0.36	Organic C in the bottom water of the southern coast
POC6s	0.05	0.07	Organic C in the surface water of the north-eastern coast
POC6b	0.01	0.02	Organic C in the bottom water of the north-eastern coast
POC7s	0.04	0.03	Organic C in the surface water of the Tethys gateway
POC7b	0.01	0.01	Organic C in the bottom water of the Tethys gateway
Name	C flux pre-OAE2	C flux OAE2	Description units in Tmol C y ⁻¹
BCF1	163.79	153.79	Primary productivity in the central open ocean
BCF2	142.73	134.01	Remineralization in the surface water of the central open ocean
BCF3	16.38	15.28	Remineralization in the intermediate water of the central open ocean
BCF4	4.67	4.38	Remineralization in the bottom water of the central open ocean
BCF5	0.02	0.015	Organic C burial in the central open ocean
BCF6	224	180.89	Primary productivity in the southern open ocean
BCF7	195.2	157.62	Remineralization in the surface water of the southern open ocean
BCF8	22.4	18.09	Remineralization in the intermediate water of the southern open ocean
BCF9	6.38	5.15	Remineralization in the bottom water of the southern open ocean
BCF10	0.03	0.02	Organic C burial in the southern open ocean
BCF11	45.76	105.16	Primary productivity in the Western Interior Seaway
BCF12	39.81	91.49	Remineralization in the surface water of the Western Interior Seaway
BCF13	5.83	13.39	Remineralization in the shallow bottom water of the Western Interior Seaway
BCF14	0.16*	0.28	Organic C burial in the Western Interior Seaway
BCF15	104.64	75.16	Primary productivity in the northern coast
BCF16	91.04	65.39	Remineralization in the surface water of the northern coast
BCF17	13.31	9.57	Remineralization in the shallow bottom water of the northern coast
BCF18	0.29*	0.2	Organic C burial in the northern coast
BCF19	168.3	210.01	Primary productivity in the southern coast
BCF20	146.42	182.7	Remineralization in the surface water of the southern coast
BCF21	21.25	26.52	Remineralization in the shallow bottom water of the southern coast
BCF22	0.63*	0.77	Organic C burial in the southern coast
BCF23	4.25	6.14	Primary productivity in the north-eastern coast
BCF24	3.7	5.34	Remineralization in the surface water of the north-eastern coast

continuation

Name	C flux pre-OAE2	OAE2	Description units in Tmol C y ⁻¹
BCF25	0.54	0.78	Remineralization in the shallow bottom water of the north-eastern coast
BCF26	0.01	0.02	Organic C burial in the north-eastern coast
BCF27	24	22.19	Primary productivity in the Tethys gateway
BCF28	20.88	19.3	Remineralization in the surface water of the Tethys gateway
BCF29	3.06	2.83	Remineralization in the shallow bottom water of the Tethys gateway
BCF30	0.069	0.057	Organic C burial in the Tethys gateway
BCF31	21.06	19.78	Organic C export to the intermediate water in the central open ocean
BCF32	4.68	4.4	Organic C export to the bottom water in the central open ocean
BCF33	28.8	23.26	Organic C export to the intermediate water in the southern open ocean
BCF34	6.4	5.17	Organic C export to the bottom water in the southern open ocean
PCF35	5.95	13.67	Organic C export to the shallow bottom water in the Western Interior Seaway
BCF36	13.60	9.77	Organic C export to the shallow bottom water in the northern coast
BCF37	21.88	27.3	Organic C export to the shallow bottom water in the southern coast
BCF38	0.55	0.8	Organic C export to the shallow bottom water in the north-eastern coast
BCF39	3.12	2.89	Organic C export to the shallow bottom water in the Tethys gateway

Table S4: Phosphorus cycle: Reservoir sizes and fluxes for pre-OAE2 and OAE2 conditions.

Name	P reservoir pre-OAE2	OAE2	Description units in Tmol P
SRP1s	0.68	0.64	Soluble reactive P in the surface water of the open ocean
SRP1i	13.6	10.48	Soluble reactive P in the intermediate water of the open ocean
SRP1b	59.7	52.86	Soluble reactive P in the bottom water of the open ocean
SRP2s	0.04	0.03	Soluble reactive P in the surface water of the southern open ocean
SRP2i	3.29	2.64	Soluble reactive P in the intermediate water of the southern open ocean
SRP2b	11.98	10.34	Soluble reactive P in the bottom water of the southern open ocean
SRP3s	0.06	0.13	Soluble reactive P in the surface water of the Western Interior Seaway
SRP3b	0.21	0.47	Soluble reactive P in the bottom water of the Western Interior Seaway
SRP4s	0.15	0.1	Soluble reactive P in the surface water of the northern coast
SRP4b	0.56	0.52	Soluble reactive P in the shallow bottom water of the northern coast
SRP5s	0.08	0.1	Soluble reactive P in the surface water of the southern coast
SRP5b	0.73	0.95	Soluble reactive P in the shallow bottom water of the southern coast
SRP6s	0.02	0.04	Soluble reactive P in the surface water of the north-eastern coast
SRP6b	0.13	0.17	Soluble reactive P in the shallow bottom water of the north-eastern coast
SRP7s	0.01	0.01	Soluble reactive P in the surface water of the Tethys gateway
SRP7b	0.09	0.15	Soluble reactive P in the shallow bottom water of the Tethys gateway
POP1s	0.046	0.04	Organic P in the surface water of the open ocean
POP1i	0.086	0.08	Organic P in the intermediate water of the open ocean
POP1b	0.077	0.07	Organic P in the bottom water of the open ocean
POP2s	0.02	0.02	Organic P in the surface water of the southern open ocean
POP2i	0.06	0.05	Organic P in the intermediate water of the southern open ocean
POP2b	0.02	0.02	Organic P in the bottom water of the southern open ocean
POP3s	0.002	0.005	Organic P in the surface water of the Western Interior Seaway
POP3b	0.0001	0.0003	Organic P in the shallow bottom water of the Western Interior Seaway
POP4s	0.006	0.005	Organic P in the surface water of the northern coast
POP4b	0.0007	0.0005	Organic P in the shallow bottom water of the northern coast
POP5s	0.01	0.01	Organic P in the surface water of the southern coast
POP5b	0.003	0.003	Organic P in the shallow bottom water of the southern coast
POP6s	0.0005	0.0007	Organic P in the surface water of the north-eastern coast
POP6b	0.0001	0.0002	Organic P in the shallow bottom water of the north-eastern coast
POP7s	0.0004	0.0003	Organic P in the surface water of the Tethys gateway
POP7b	0.0001	0.0001	Organic P in the shallow bottom water of the Tethys gateway
Name	P flux pre-OAE2	OAE2	Description units in Tmol P y ⁻¹
Fluxes coupled to the water cycle			
PF1	0.07	0.09	Surface [SRP] of the central open ocean times F1
PF3	0.46	0.57	Intermediate [SRP] of the central open ocean times F3
PF6	0.55	0.88	Bottom [SRP] of the Pacific Ocean times F6
PF7	0.004	0.02	Surface [SRP] of the Tethys gateway times F7
PF10	0.03	0.24	Bottom [SRP] of the Tethys gateway times F10
PF13	0.23	0.12	Surface [SRP] of the Tethys gateway times F13
PF14	0.008	0.04	Surface [SRP] of central open ocean times F14
PF15	0.008	0.05	Surface [SRP] of the Tethys gateway times F15
PF16	0.15	0.22	Surface [SRP] of the central open ocean times F16 (downwelling)

continuation

Name	P flux pre-OAE2	OAE2	Description
			units in Tmol P y ⁻¹
PF17	0.14	0.29	Intermediate [SRP] of the central open ocean times F17 (upwelling)
PF18	0.02	0.005	Surface [SRP] of the southern coast times F18 in pre-OAE2
			Surface [SRP] of the southern open ocean times F18 in OAE2
PF19	0.02	0.03	Surface [SRP] of the central open ocean times F19
PF20	0.51	0.35	Intermediate [SRP] of the central open ocean times F20
PF21	0.29	0.48	Bottom [SRP] of the northern coast times F21
PF22	0.03	0.11	Intermediate [SRP] of the central open ocean times F22
PF23	0.006	0.32	Intermediate [SRP] of the central open ocean times F23
PF24	0.32	0.18	Bottom [SRP] of the southern coast times F24
PF25	0.006	0.24	Bottom [SRP] of the central open ocean times F25
PF26	0.02	0.02	Surface [SRP] of the Tethys gateway times F26 (downwelling)
PF27	0.07	0.13	Bottom [SRP] of the Tethys gateway times F27 (upwelling)
PF28	0.02	0.02	Surface [SRP] of the southern coast times F28 (downwelling)
PF29	0.3	0.37	Bottom [SRP] of the southern coast times F29 (upwelling)
PF30	0.01	0.01	[SRP] of the South river times F30
PF31	0.18	0.11	Surface [SRP] of the Western Interior Seaway times F31 (downwelling)
PF32	0.33	0.28	Bottom [SRP] of the Western Interior Seaway times F32 (upwelling)
PF33	0.19	0.06	Surface [SRP] of the northern coast times F33 (downwelling)
PF34	0.43	0.16	Bottom [SRP] of the northern coast times F34 (upwelling)
PF35	0.02	0.05	Surface [SRP] of the north-eastern coast times F35 (downwelling)
PF36	0.01	0.03	Bottom [SRP] of the north-eastern coast times F36 (upwelling)
PF37	0.42	0.72	Intermediate [SRP] of the central open ocean times F37
PF38	1.04	1.37	Bottom [SRP] of the central open ocean times F38
PF39	0.12	0.07	Surface [SRP] of the Western Interior Seaway times F39
PF40	0.001	0.04	Surface [SRP] of the north-eastern coast times F40
PF41	0.002	0.02	Surface [SRP] of the Tethys gateway times F41
PF42	0.01	0.04	Surface [SRP] of the Tethys gateway times F42
PF43	0.01	0.17	Bottom [SRP] of the Tethys gateway times F43
PF44	0.05	0.07	Bottom [SRP] of the north-eastern coast times F44 in pre-OAE2
			Bottom [SRP] of the Tethys gateway times F44 in OAE2
PF45	0.01	0.19	Bottom [SRP] of the northern coast times F45
			Bottom [SRP] of the north-eastern coast times F45
PF46	0.42	0.3	Bottom [SRP] of the northern coast times F46
PF47	0.05	0.06	Surface [SRP] of the central open ocean times F47
PF48	0.04	0.03	Surface [SRP] of the southern open ocean times F48
PF49	0.41	0.08	Intermediate [SRP] of the central open ocean times F49
PF50	0.49	0.41	Intermediate [SRP] of the central open ocean times F50
PF51	0.84	0.63	Intermediate [SRP] of the southern open ocean times F51
PF52	0.75	0.73	Bottom [SRP] of the central open ocean times F52
PF53	0.78	0.69	Bottom [SRP] of the southern open ocean times F53
PF54	0.08	0.14	Surface [SRP] of the southern coast times F54
PF55	0.0006	0.0006	Surface [SRP] of the southern open ocean times F55 (downwelling)
PF56	0.24	0.19	Intermediate [SRP] of the southern open ocean times F56 (upwelling)
PF57	0.03	0.02	Intermediate [SRP] of the southern open ocean times F57
PF58	0.09	0.08	Bottom [SRP] of the southern open ocean times F58
PF59	0.03	0.02	Bottom [SRP] of the southern coast times F59 in pre-OAE2

continuation

Name	P flux pre-OAE2	OAE2	Description units in Tmol P y ⁻¹
PR3	0.0004	0.0006	Bottom [SRP] of the southern open ocean times F59 in OAE2
PR4	0.0018	0.0018	[SRP] of the North-West river times R3
PR6	0.00001	0.00001	[SRP] of the North river times R4
			[SRP] of the north-eastern river times R6
			Fluxes coupled to the C and O₂ cycles
BPF1	1.55	1.45	SRP uptake in the central open ocean
BPF2	1.35	1.26	SRP release in the surface water of the central open ocean
BPF3	0.155	0.15	SRP release in the intermediate water of the central open ocean
BPF4	0.04	0.04	SRP release in the bottom water of the central open ocean
BPF5	0.0001	0.0001	Organic P burial in the central open ocean
BPF6	2.11	1.71	SRP uptake in the southern open ocean
BPF7	1.84	1.49	SRP release in the surface water of the southern open ocean
BPF8	0.21	0.17	SRP release in the intermediate water of the southern open ocean
BPF9	0.06	0.05	SRP release in the bottom water of the southern open ocean
BPF10	0.0002	0.0002	Organic P burial in the southern open ocean
BPF11	0.43	0.99	SRP uptake in the Western Interior Seaway
BPF12	0.38	0.86	SRP release in the surface water of the Western Interior Seaway
BPF13	0.06	0.13	SRP release in the shallow bottom water of the Western Interior Seaway
BPF14	0.0005	0.001	Organic P burial in the Western Interior Seaway
BPF15	0.99	0.71	SRP uptake in the northern coast
BPF16	0.86	0.62	SRP release in the surface water of the northern coast
BPF17	0.13	0.09	SRP release in the shallow bottom water of the northern coast
BPF18	0.001	0.001	Organic P burial in the northern coast
BPF19	1.59	1.98	SRP uptake in the southern coast
BPF20	1.38	1.72	SRP release in the surface water of the southern coast
BPF21	0.20	0.26	SRP release in the shallow bottom water of the southern coast
BPF22	0.001	0.002	Organic P burial in the southern coast
BPF23	0.04	0.06	SRP uptake in the north-eastern coast
BPF24	0.03	0.05	SRP release in the surface water of the north-eastern coast
BPF25	0.005	0.007	SRP release in the shallow bottom water of the north-eastern coast
BPF26	0.00005	0.00007	Organic P burial in the north-eastern coast
BPF27	0.23	0.21	SRP uptake in the Tethys gateway
BPF28	0.2	0.18	SRP release in the surface water of the Tethys gateway
BPF29	0.03	0.03	SRP release in the shallow bottom water of the Tethys gateway
BPF30	0.0003	0.0003	Organic P burial in the Tethys gateway
BPF31	0.2	0.19	Organic P export to the intermediate water in the central open ocean
BPF32	0.04	0.04	Organic P export to the bottom water in the central open ocean
BPF33	0.27	0.22	Organic P export to the intermediate water in the southern open ocean
BPF34	0.06	0.05	Organic P export to the bottom water in the southern open ocean
BPF35	0.06	0.13	Organic P export to the shallow bottom water in the Western Interior Seaway
BPF36	0.13	0.09	Organic P export to the shallow bottom water in the northern coast
BPF37	0.21	0.26	Organic P export to the shallow bottom water in the southern coast
BPF38	0.005	0.008	Organic P export to the shallow bottom water in the north-eastern coast
BPF39	0.03	0.03	Organic P export to the shallow bottom water in the Tethys gateway
FePB1	0.00014	0.00016	Inorganic Fe-P burial in the central open ocean
CaPB1	0.00029	0.00029	Inorganic Ca-P burial in the central open ocean

continuation

Name	P flux		Description
	pre-OAE2	OAE2	units in Tmol P y ⁻¹
FePB2	0.00016	0.0002	Inorganic Fe-P burial in the southern open ocean
CaPB2	0.00037	0.00033	Inorganic Ca-P burial in the southern open ocean
FePB3	0.00043	0.00045	Inorganic Fe-P burial in the Western Interior Seaway
CaPB3	0.0009	0.0022	Inorganic Ca-P burial in the Western Interior Seaway
FePB4	0.0009	0.001	Inorganic Fe-P burial in the northern coast
CaPB4	0.0019	0.0018	Inorganic Ca-P burial in the northern coast
FePB5	0.0004	0.0004	Inorganic Fe-P burial in the southern coast
CaPB5	0.0017	0.0017	Inorganic Ca-P burial in the southern coast
FePB6	0.00005	0.00005	Inorganic Fe-P burial in the north-eastern coast
CaPB6	0.0001	0.00016	Inorganic Ca-P burial in the north-eastern coast
FePB7	0.0002	0.0003	Inorganic Fe-P burial in the Tethys gateway
CaPB7	0.0005	0.0005	Inorganic Ca-P burial in the Tethys gateway

Table S5: Oxygen cycle: Reservoir sizes and fluxes for pre-OAE2 and OAE2 conditions as implemented in the multi-box model.

Name	O ₂ reservoir pre-OAE2	Description units Tmol O ₂
OX1i	987.95	1323.70
OX1b	5777.83	6991.77
OX2i	54.28	130.72
OX2b	900.53	1175.8
OX3b	17.77	51.87
OX4b	39.6	64.38
OX5b	10.11	14.64
OX6b	20.18	30.8
OX7b	12.78	19.65
Name	O ₂ flux pre-OAE2	Description units in Tmol O ₂ y ⁻¹
Fluxes coupled to the water cycle		
OF3	33.44	71.82
OF6	90.76	145.14
OF10	3.9	31.9
OF16	66.51	107.51
OF17	10.46	37.11
OF20	37.4	43.86
OF21	20.67	60.59
OF22	2.2	13.67
OF23	0.45	40.95
OF24	4.37	2.73
OF25	0.58	31.84
OF26	10.78	11.04
OF27	9.36	16.95
OF28	4.62	8.07
OF29	4.1	5.75
OF31	32.87	38.18
OF32	27.77	31.13
OF33	33.97	24.17
OF34	30.22	20.32
OF35	7.22	20.8
OF36	2.12	6.23
OF37	30.27	90.81
OF38	100.43	180.96
OF43	1.82	22.78
OF44	7.26	8.53
OF45	0.66	35.73
OF46	34.92	33.47
OF49	30.14	9.94
OF50	35.84	52.13

continuation

Name	O ₂ flux pre-OAE2	OAE2	Description units in Tmol O ₂ y ⁻¹
OF51	13.91	31.22	Intermediate [O ₂] of the southern open ocean times F51
OF52	72.71	96.31	Bottom [O ₂] of the central open ocean times F52
OF53	58.76	78.86	Bottom [O ₂] of the southern open ocean times F53
OF55	0.72	0.94	Surface [O ₂] of the southern open ocean times F55 (downwelling)
OF56	3.94	9.5	Intermediate [O ₂] of the southern open ocean times F56 (upwelling)
OF57	0.41	0.99	Intermediate [O ₂] of the southern open ocean times F57
OF58	6.5	9.5	Bottom [O ₂] of the southern open ocean times F58
OF59	0.45	2.23	Bottom [O ₂] of the southern coast times F59 in pre-OAE2 Bottom [O ₂] of the southern open ocean times F59 in OAE2
Fluxes coupled to the C and P cycles			
BOF3	21.32	20.02	O ₂ consumption in the intermediate water of the central open ocean
BOF4	6.07	5.7	O ₂ consumption in the bottom water of the central open ocean
BOF8	29.16	23.55	O ₂ consumption in the intermediate water of the southern open ocean
BOF9	8.3	6.71	O ₂ consumption in the bottom water of the southern open ocean
BOF13	7.58	17.43	O ₂ consumption in the shallow bottom water in the Western Interior Seaway
BOF17	17.33	12.46	O ₂ consumption in the shallow bottom water in the northern coast
BOF21	27.67	34.54	O ₂ consumption in the shallow bottom water in the southern coast
BOF25	0.71	1.02	O ₂ consumption in the shallow bottom water in the north-eastern coast
BOF29	3.98	3.68	O ₂ consumption in the shallow bottom water in the Tethys gateway

Table S6: Linear equations as implemented in the model for fluxes in oxic conditions. The k_* is a rate constant (y^{-1}). SRP is the reservoir of soluble reactive P, POC and POP are the reservoirs for particulate organic C and P, respectively. $_s$: surface water; $_i$: intermediate water; $_b$: bottom water.

Flux	Linear equations
C cycle	(Tmol C y^{-1})
Primary productivity in surface ocean	$PP = k_{PP} \cdot SRP_s \cdot 106$ [1]
Inorganic C release in surface ocean	$CREL_s = k_{CREL_s} \cdot POC_s$ [1]
Inorganic C release in intermediate open ocean	$CREL_i = k_{CREL_i} \cdot POC_i$ [1]
Inorganic C release in shallow/deep bottom ocean	$CREL_b = k_{CREL_b} \cdot POC_b$ [1]
POC Export in surface to intermediate/shallow bottom ocean	$CEXP_s = POPEXP_s * 106$ [1]
POC Export in intermediate to deep bottom open ocean	$CEXP_i = POPEXP_i * 106$ [1]
POC burial in coastal ocean	$CBUR = k_{CBUR} \cdot CEXP_s$ [1]
POC burial in open ocean	$CBUR = k_{CBUR} \cdot (CEXP_s + CEXP_i)$ [2]
P cycle	(Tmol P y^{-1})
SRP uptake in surface ocean	$SRPU = PP/106$ [1]
SRP release in surface ocean	$PREL_s = CREL_s/106$ [1]
SRP release in intermediate open ocean	$PREL_i = CREL_i/106$ [1]
SRP release in shallow/deep bottom ocean	$PREL_b = k_{PREL_b} \cdot POP_b$ [1]
POP Export from surface to intermediate/shallow bottom	$POPEXP_s = k_{POPEXP_s} * POP_s$ [1]
POP Export from intermediate to deep bottom ocean	$POPEXP_i = k_{POPEXP_i} * POP_i$ [1]
POP burial in coastal ocean	$POPBUR = CBUR/200$ [3]
POP burial in open ocean	$POPBUR = CBUR/106$ [1]
Ca-P burial in shallow/deep bottom ocean	$CaPBUR = k_{CaPBUR} \cdot PREL_b$ [1]
Fe-P burial in shallow/deep bottom ocean	$FePBUR = FePBUR_{oxic} \cdot \frac{[O_2]}{[O_2]_{crit}}$ [1]
O₂ cycle	(Tmol O₂ y^{-1})
O ₂ consumption in intermediate open ocean	$OCON_i = CREL_i \cdot 138/106$ [1]
O ₂ consumption in shallow/deep bottom ocean	$OCON_b = CREL_b \cdot 138/106$ [1]

[1]Slomp and Van Cappellen, 2007, [2] This study, [3]Van Cappellen and Ingall, 1994

References

Arthur, M. A. and Dean, W. A. Pratt, L. M.: Geochemical and climatic effects of increased marine organic carbon burial at the Cenomanian-Turonian boundary, *Nature*, 335, 714–717, doi:10.1038/335714a0, 1988.

Austin, J. A., Schlager, W., Palmer, A. A., and et al.: Site 632 A/B, in Proceedings and initial reports (Part A) of the Ocean Drilling Program, Leg 101, College Station, Texas A&M University, p. 569, doi: 10.2973/odp.proc.ir.101.111, 1986.

Barker, C. E., Pawlewicz, M., and Cobabe, E. A.: Deposition of sedimentary organic matter in black shale facies indicated by the geochemistry and petrography of high-resolution samples, Blake Nose, western North Atlantic, Geological Society, London, Special Publications, 183, 49–72, doi:10.1144/GSL.SP.2001.183.01.03, 2001.

Bowman, A. R. and Bralower, T. J.: Paleoceanographic significance of high-resolution carbon isotope records across the Cenomanian-Turonian boundary in the Western Interior and New Jersey coastal plain, USA, *Marine Geology*, 217, 305–321, 2005.

Herbin, J. P., Montadert, L., Miiller, C., Gomez, R., Thurow, J., and Wiedmann, J.: Organic-rich sedimentation at

the Cenomanian-Turonian boundary in oceanic and coastal basins in the North Atlantic and Tethys, Geological Society Special Publication, 21, 389–422, 1986.

Hetzl, A., Böttcher, M. E., Wortmann, U. G., and Brumsack, H.-J.: Paleo-redox conditions during OAE 2 reflected in Demerara Rise sediment geochemistry (ODP Leg 207), *Palaeogeography Palaeoclimatology Palaeoecology*, 273, 302–328, 2009.

Hetzl, A., März, C., Vogt, C., and Brumsack, H.-J.: Geochemical environment of Cenomanian-Turonian black shale deposition at Wunstorf (northern Germany), *Cretaceous Research*, 32, 480–494, 2011.

Kolonic, S., Wagner, T., Forster, A., Sinninghe Damsté, J. S., Walsworth-Bell, B., Erba, E., Turgeon, S., Brumsack, H.-J., Hassane Chellai, E., Tsikos, H., Kuhnt, W., and Kuypers, M. M. M.: Black shale deposition on the northwest African Shelf during the Cenomanian-Turonian oceanic anoxic event: Climate coupling and global organic carbon burial, *Paleoceanography*, 20, 95–128, doi:10.1029/2003PA000950, 2005.

Kraal, P., Slomp, C. P., Forster, A., and Kuypers, M. M. M.: Phosphorus cycling from the margin to abyssal depths in the proto-Atlantic during oceanic anoxic event 2, *Palaeogeography Palaeoclimatology Palaeoecology*, 295, 42–54, 2010b.

Kuhnt, W. and Wiedmann, J.: Cenomanian–Turonian source rocks: paleobiogeographic and paleoenvironmental aspects, In: Huc, A. Y. (Ed.) : *Paleogeography, Paleoclimate and Source Rocks*. AAPG studies in Geology, 40, 213–231, 1995.

Kuhnt, W., Herbin, J. P., Thurow, J. W., and Wiedmann, J.: Distribution of Cenomanian-Turonian organic facies in the western Mediterranean and along the Adjacent Atlantic Margin, *AAPG Studies in Geology*, 30, 133–160, 1990.

Kuypers, M. M. M., Pancost, R. D., Nijenhuis, I. A., and Sinninghe Damsté, J. S.: Enhanced productivity led to increased organic carbon burial in the euxinic North Atlantic basin during the late Cenomanian oceanic anoxic event, *Paleoceanography*, 17, 1–13, 2002.

Kuypers, M. M. M., Lourens, L. J., Rijpstra, W. I. C., Pancost, R. D., Nijenhuis, I. A., and Sinninghe Damsté, J. S.: Orbital forcing of organic carbon burial in the proto-North Atlantic during oceanic anoxic event 2, *Earth and Planetary Science Letters*, 228, 465–482, 2004.

Lancelot, Y., Winterer, E. L., and et al.: Init. Rep. Deep Sea Drilling Project 50: Washington (U.S. Govt. Printing Office), 1980.

Meyers, S., Sageman, B., and Hinnov, L.: Integrated quantitative stratigraphy of the Cenomanian-Turonian Bridge Creek Limestone Member using evolutive harmonic analysis and stratigraphic modeling, *Journal of Sedimentary Research*, 71, 627–643, 2001.

Mort, H. P., Adatte, T., Föllmi, K. B., Keller, G., Steinmann, P., Matera, V., Berner, Z., and Stüben, D.: Phosphorus and the roles of productivity and nutrient recycling during oceanic anoxic event 2, *Geology*, 35, 483–486, 2007.

Mort, H. P., Adatte, T., Keller, G., Bartels, D., Föllmi, K. B., Steinmann, P., Berner, Z., and Chellai, E. H.: Organic carbon deposition and phosphorus accumulation during Oceanic Anoxic Event 2 in Tarfaya, Morocco, *Cretaceous Research*, 29, 1008–1023, 2008.

Nederbragt, A. J., Thurow, J., Vonhof, H., and Brumsack, H.-J.: Modelling oceanic carbon and phosphorus fluxes: implications for the cause of the late Cenomanian Oceanic Anoxic Event (OAE2), *Journal of the Geological Society of London*, 161, 721–728, doi:10.1144/0016-764903-075, 2004.

Norris, R. D., Kroon, D., and Klaus, A.: CretaceousPaleogene climatic evolution of the western North Atlantic: results from ODP Leg 171B, Blake Nose, In Kroon, D., Norris, R. D., and Klaus, A. (Eds.), *Proc. ODP, Sci. Results, 171B: College Station, TX (Ocean Drilling Program)*, 161, 1–10, doi:10.2973/odp.proc.sr.171b.101, 2001.

Owens, J. D., Lyons, T. W., Li, X., Macleod, K. G., Gordon, G., Kuypers, M. M. M., Anbar, A., Kuhnt, W., and Severmann, S.: Iron isotope and trace metal records of iron cycling in the proto-North Atlantic during the Cenomanian-Turonian oceanic anoxic event (OAE-2), *Palaeogeography*, 27, 1–13, doi:10.1029/2012PA002328, 2012.

Pancost, R. D., Crawford, N., Magness, S., Turner, A., Jenkyns, H. C., and Maxwell, J. R.: Further evidence for the development of photic-zone euxinic conditions during Mesozoic oceanic anoxic events, *Journal of the Geological Society of London*, 161, 353–364, doi:10.1144/0016764903-059, 2004.

Rodríguez-Tovar, F. J., Uchman, A., Payros, A., Orue-Etxebarria, X., Apellaniz, E., and Molina, E.: Sea-level dynamics and palaeoecological factors affecting trace fossil distribution in Eocene turbiditic deposits (Gorrondatxe section, N Spain), *Palaeogeography Palaeoclimatology Palaeoecology*, 285, 50–65, doi: 10.1016/j.palaeo.2009.10.022, 2010.

Sinninghe Damsté, J. S. and Köster, J.: A euxinic southern North Atlantic Ocean during the Cenomanian-Turonian oceanic anoxic event, *Earth and Planetary Science Letters*, 158, 165–173, doi:10.5194/cp-3-647-2007, 1998.

Sinninghe Damsté, J. S., van Bentum, E. C., Reichart, G.-J., Pross, P., and Schouten, S.: A CO₂ decrease-driven cooling and increased latitudinal temperature gradient during the mid-Cretaceous Oceanic Anoxic Event 2, *Earth and Planetary Science Letters*, 293, 97–103, doi:10.1016/j.epsl.2010.02.027, 2010.

Slomp, C. P. and Van Cappellen, P.: The global marine phosphorus cycle: sensitivity to oceanic circulation, *Biogeosciences*, 4, 155–171, 2007.

Snow, L. J., Duncan, R. A., and Bralower, T. J.: Trace element abundances in the Rock Canyon anticline, Pueblo, Colorado, marine sedimentary section and their relationship to Caribbean plateau construction and oxygen anoxic event 2, *Paleoceanography*, 20, doi:10.1029/2004PA001093, 2005.

Sugarman, P. J., Miller, K. G., Olsson, R. K., Browning, J. V., Wright, J. D., de Romero, L. M., White, T. S., Muller, F. L., and Uptegrove, J.: The Cenomanian-Turonian carbon burial event, Bass River, NJ, USA: Geochemical, paleoecological, and sea-level changes, *Journal of Foraminiferal Research*, 29, 438–452, 1999.

Thurow, J. W., Moullade, M., Brumsack, H.-J., Masure, E., Taugourdou, J., and Dunham, K.: The Cenomanian-Turonian Boundary Event (CTBE) at Leg 103/Hole 641A, *Proceedings of the Ocean Drilling Program Scientific Results*, 103, 587–634, 1988.

Topper, R. P. M., Trabucho Alexandre, J., Tuenter, E., and Meijer, P. Th.: A regional ocean circulation model for the mid-Cretaceous North Atlantic Basin: implications for black shale formation, *Climate of the past*, 7, 277–297, 2011.

Urquhart, E., Gardin, S., Leckie, R. M., Wood, S. A., Pross, J., Georgescu, M. D., Ladner, B., and Takata, H.: A Paleontological Synthesis of ODP Leg 210, Newfoundland Basin, In Tucholke, B.E and Sibuet, J. C and Klaus, A. (Eds.), *Proc. ODP, Sci. Results*, 210: College Station, TX (Ocean Drilling Program), pp. 1–53, 2007.

van Bentum, E. C., Hetzel, A., Brumsack, H.-J., Forster, A., Reichart, G.-J., and Sinninghe Damsté, J. S.: Reconstruction of water column anoxia in the equatorial Atlantic during the Cenomanian-Turonian oceanic anoxic event using biomarker and trace metal proxies, *Palaeogeography Palaeoclimatology Palaeoecology*, 280, 489–498, 2009.

van Bentum, E. C., Reichart, G.-J., and Sinninghe-Damsté, J. S.: Organic matter provenance, palaeoproductivity and bottom water anoxia during the Cenomanian/Turonian oceanic anoxic event in the Newfoundland Basin (northern proto North Atlantic Ocean), *Organic Geochemistry*, 50, 11–18, 2012.

Van Cappellen, P. and Ingall, E. D.: Benthic phosphorus regeneration, net primary production, and ocean anoxia: A model of the coupled marine biogeochemical cycles of carbon and phosphorus, *Paleoceanography*, 9, 677–692, 1994.

Voigt, S.: Cenomanian-Turonian composite $\delta^{13}\text{C}$ curve for Western and Central Europe: the role of organic and inorganic carbon fluxes, *Palaeogeography Palaeoclimatology Palaeoecology*, 160, 91–104, 2000.