



## N<sub>2</sub>O fluxes from the littoral zone of a Chinese reservoir

M. Yang<sup>1</sup>, X. M. Geng<sup>1</sup>, J. Grace<sup>2</sup>, Y. F. Jia<sup>1,a</sup>, Y. Z. Liu<sup>1</sup>, S. W. Jiao<sup>1</sup>, L. L. Shi<sup>1</sup>, C. Lu<sup>1</sup>, Y. Zhou<sup>1</sup>, and G. C. Lei<sup>1</sup>

<sup>1</sup>School of Nature Conservation, Beijing Forestry University, Beijing, China

<sup>2</sup>School of Geosciences, The University of Edinburgh EH9 3FF, Edinburgh, UK

<sup>a</sup>now at: Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing, China

Correspondence to: G. C. Lei (guangchun.lei@foxmail.com)

Received: 22 January 2015 – Published in Biogeosciences Discuss.: 2 April 2015

Revised: 10 July 2015 – Accepted: 19 July 2015 – Published: 6 August 2015

**Abstract.** There have been few studies of greenhouse gas emissions from reservoirs, despite the remarkable growth in the number of reservoirs in developing countries. We report a case study that focuses on the littoral zone of a major Chinese reservoir, where we established measurements of N<sub>2</sub>O fluxes using the static chamber technique at five different water levels (deep water, shallow water, seasonally flooded, control for seasonally flooded, and non-flooded). The “control for seasonally flooded” had similar vegetation to the “seasonally flooded” but was not actually flooded as it was on a higher piece of land. Seasonal, diurnal and spatial variations of N<sub>2</sub>O flux and environmental factors were monitored throughout the growing season which included a flood event during summer rains. The N<sub>2</sub>O flux ranged from  $-136.6$  to  $381.8 \mu\text{g m}^{-2} \text{h}^{-1}$  averaging  $6.8 \mu\text{g m}^{-2} \text{h}^{-1}$ . Seasonal and spatial variation was significant but diurnal variation was not. Non-flooded dry land emitted more N<sub>2</sub>O than flooded land, no matter whether it was permanently or seasonally flooded. Piecewise correlation was found between N<sub>2</sub>O flux, air temperature and soil nitrate concentration. Positive correlation was shown between N<sub>2</sub>O flux and dissolved oxygen in water. There were significantly higher emissions from farmland. We compared these results with our recently published study of CH<sub>4</sub> emissions, carried out simultaneously at the same site as those in the present study. Completely different patterns between the two gases are demonstrated. We conclude that the littoral zone is a hotspot for N<sub>2</sub>O emissions in the summer, especially when the shores of the lake are used for the farming of maize. But in terms of the overall greenhouse gas budget, the fluxes of N<sub>2</sub>O are not as important as those of CH<sub>4</sub>.

### 1 Introduction

Reservoirs are increasing rapidly in number and area, growing with the continuing demand for water and hydropower. In rapidly developing countries like China, India and Brazil this growth is likely to continue for many years (Yang and Lu, 2014; Kumar et al., 2011). There are several environmental impacts of reservoirs, particularly sediment accumulation and vegetation change. Moreover, when fertile agricultural lands are inundated by rising water there may be a strong enhancement of greenhouse gas emissions (Tranvik et al., 2009; L. Yang et al., 2014).

The pelagic zones of reservoirs have more often been studied (Beaulieu et al., 2014; Guérin et al., 2008; Huttunen et al., 2002; X. L. Liu et al., 2011) but few researchers have investigated the littoral zone, which could be a hotspot of N<sub>2</sub>O emissions (Wang et al., 2006). In the few cases where it has been studied, N<sub>2</sub>O emissions of the littoral zone in natural lakes have been observed to be higher than the pelagic zone even though the area differences had been taken into account (Huttunen et al., 2003).

Because of the strong gradients in water level and water level fluctuations, compared to the more or less stable pelagic zone and strictly terrestrial areas nearby (e.g. grassland and farmland), the environment of the littoral zone is more diverse and dynamic in terms of soil moisture, plant species and soil nutrients across scales of both space and time (Peng et al., 2011; Ahn et al., 2014; Trost et al., 2013). These variables may be expected to influence N<sub>2</sub>O production (Lu and Xu, 2014). Limited previous studies on N<sub>2</sub>O emissions of the littoral zone suggested significant spatio-temporal variations. But most of the studies just focus on a single water level


(with different communities sometimes), and they overlook the spatial variations between different water levels (Chen et al., 2011b; Y. Liu et al., 2011). Temporally, seasonal variation has been demonstrated but not diurnal variation (Chen et al., 2010; Huttunen et al., 2003). To match the diverse and dynamic environment of the littoral zone, we combined five water levels on a transect from water to dry land, three plant communities for each water level including both natural and cropped land, six times during the year and seven times of day. The improved sampling both in space and time was expected to provide representative data on N<sub>2</sub>O emission of the littoral zone, and to provide further insights into the nature of the underlying processes.

To be more specific, the objectives of this present study included (i) capturing the spatial and temporal variation of the N<sub>2</sub>O flux at the littoral zone of the Miyun Reservoir; (ii) finding the relationship between the observed flux and environmental factors; and (iii) evaluating the relative importance of N<sub>2</sub>O and CH<sub>4</sub> fluxes by comparing with our earlier report of the CH<sub>4</sub> fluxes made simultaneously from the same site (M. Yang et al., 2014). The over-arching hypothesis in this work is that the littoral zone is a hotspot of N<sub>2</sub>O emissions that is influenced by seasonal changes in the water level.

## 2 Methods

### 2.1 Study area


The research was carried out at Miyun Reservoir (40°29' N, 116°50' E), which is located in the northern mountainous area of Beijing, China. It was built in 1960 with a maximum water area of 188 km<sup>2</sup>. Its catchment is characterized by warm temperate semi-humid monsoonal climate with an annual average air temperature of 10.5 °C, maximum air temperature of 38 °C, and a minimum of −18 °C. The reservoir is normally covered by ice from the middle of November to the end of March. The growing season is from April to November. The annual average precipitation is close to 600 mm, of which 80 % is concentrated from July to August (Gao, 1989). Alongside the reservoir, higher land (sometimes just slightly higher) is nearly always used by local people for growing maize. This opportunistic agriculture is typically from May to September. Nitrogenous fertilizer is applied during sowing, and sometimes with further application in the middle of the growing season. This reservoir is mainly used as the domestic water supply for Beijing. The water quality is controlled to level II according to Environmental Quality Standards for Surface Water of People's Republic of China GB3838-2002 (levels are rated on a scale I to V, where level I is the cleanest, available at <http://kjs.mep.gov.cn/hjbhbz/index.htm>). The annual change in the water level is 1–5 m, reflecting the balance between rainfall, evaporation and usage. The water area between the points of highest and lowest water level (assessed from 1984



**Figure 1.** Experimental design. WL: water level. The difference between high WL and low WL was caused by summer flooding. m<sup>H</sup> indicates height (metres) in the horizontal; m<sup>V</sup> indicates metres in the vertical. The sites are grouped at different heights as follows. DW: deep water site; SW: shallow water site; SF: seasonally flooded site; SFC: “control site” for the seasonally flooded site, which had similar vegetation and soil moisture as site SF before it was flooded; NF: non-flooded site, which flooded once per several years and not flooded in the sampling year. A, B and C denote samples from different vegetation types within each height band; for species details see Table 1. There were 15 plots in total, four replicates in each plot, repeatedly sampled six times in the year to cover different seasons and covering the transition in and out of the flooding season. Also to capture diurnal variation, plots were repeatedly sampled seven times per day. For more details on water depth and other environmental parameters, see Figs. 2 and 3.

to 2005) was 84 km<sup>2</sup> (Cao et al., 2008). In the summer of 2012, when the work was carried out, unusual and continuous heavy rain in July caused a sudden water level increase of 0.8 m in 15 days, and part of the littoral vegetation was inundated. This provided us with a seasonal flooded area which made possible an exploration of the effects of summer flooding on greenhouse gas emissions.

We divided the littoral zone into five areas based on water level (Fig. 1). Sites were selected ranging from locations in open water to the dry area on higher ground, to provide five contrasting environments: (i) deep water area (DW); (ii) shallow water area (SW); (iii) seasonal (August and September) flooded area (SF); (iv) “seasonally flooded control area” (SFC), which was 500 m away from SF, had the same plant species as SF, but escaped the flood in August and September because of its slightly (about 1 m) higher elevation; and (v) an area which is seldom flooded (the last flooding was several years ago) which hereafter we call the non-flooded area (NF). Three typical plant communities in each water level were selected. At SW, SF, SFC and NF, land cropped with maize (*Zea mays*) was included as it is a typical practice, and allows some assessment of the impact of farming. Maize growing in SW and SF was abandoned by the local farmer after our first sampling campaign because of flooding. So these lands were colonized by wild plants after abandon-



**Figure 2.** Environmental characteristics (mean  $\pm$  SE) of each sampling area. Some SE bars are not visible. Days between dotted lines were the high water level period and thus the seasonal flooded site (SF) was under water. DW: deep water site; SW: shallow water site; SF: seasonally flooded site; SFC: “control site” for the seasonally flooded site; NF: non-flooded site. There were no soil water content data for July because of instrument malfunction.

ment. Dominant species of each month are shown in Table 1. Details of climate, biomass and soil–sediment parameters are shown in Figs. 2 and 3.

## 2.2 N<sub>2</sub>O flux measurements

Nitrous oxide flux was measured in November 2011, then May, July, August, September and October 2012. Measurements at site SFC were carried out just after the flooding and during the time when the water level dropped from August to October 2012. In order to reduce uncertainty in the average daily flux, a sampling protocol designed to capture any diurnal variation was performed at three-hourly intervals (local time: 6, 9, 12, 15, 18, 21 and 24 h). Each plot had four replicate chambers located within 3 m from each other. To eliminate trampling disturbance to the soil–sediment during sampling, wooden access platforms were built.

The static opaque chamber technique was used to determine the N<sub>2</sub>O flux. The chambers were made of stainless steel (volume: 125 L; surface area: 0.25 m<sup>2</sup>) and coated with polyethylene foam to minimize any warming effect inside the chamber. An extension chamber (volume: 200 L; surface area: 0.25 m<sup>2</sup>) was added whenever plants were especially

tall. Two fans were built into the chamber for air mixing. Four gas samples (200 mL each) were taken using 100 mL polypropylene syringes at 15 min intervals over a 45 min period after enclosure, and stored in 500 mL plastic and aluminum membrane gas sampling bags (Guangming Research and Design Institute of Chemical Industry, China). The concentration of N<sub>2</sub>O was analysed within 1 week by gas chromatography (7890A, Agilent, USA) equipped with a micro-electron capture detector ( $\mu$ -ECD). Gases were separated with a column (3 m, 3.2 mm) packed with Porapak Q (80/100 mesh). The temperatures of the oven, injector, and detector were 70, 20, and 330 °C, respectively. The flow rate of the carrier gas (N<sub>2</sub>) was 25 mL min<sup>-1</sup>. Standard N<sub>2</sub>O gas (310 ppb in air, China National Research Center for Certified Reference Materials, China) was used for precision verification for N<sub>2</sub>O concentrations. The coefficient of variation was below 1.5 %. The flux of N<sub>2</sub>O was calculated via the following relationship by Chen et al. (2011b):

$$F = \frac{M}{V_0} \times \frac{P}{P_0} \times \frac{T_0}{T} \times \frac{dC_t}{dt} \times H, \quad (1)$$

where  $F$  is the flux of N<sub>2</sub>O (mg m<sup>-2</sup> h<sup>-1</sup>);  $M$  is the molar mass of N<sub>2</sub>O (g mol<sup>-1</sup>);  $P$  (kPa) is the atmospheric pressure of the sampling site;  $T$  (K) is the absolute temperature of the sampling time;  $V_0$  (22.4 L),  $P_0$  (101.325 kPa) and  $T_0$  (273.15 K) are the molar volume, atmosphere pressure and absolute temperature, respectively, under standard conditions;  $dC_t/dt$  (ppm h<sup>-1</sup>) is the rate of concentration change; and  $H$  (m) is the chamber height over the water or soil surface.

Chambers were reset into new positions near the old positions each sampling month. All positions at each site were within an area of 20 m<sup>2</sup>, but not so close to each other to cause artefacts in the data through (for example) changes in the local hydrology.

## 2.3 Environmental factors

Weekly precipitation was accessed through the China Meteorological Data Sharing Service System (<http://www.esi.cn/metadata/page/index.html>). Average wind speed was recorded during the sampling period with a hand-held vane anemometer (4101, Testo, Germany), taking an average over the 45 min period during which gas was sampled. Air temperature was measured by a digital thermometer (JM624, Jinming, China) at the start and end of each gas sampling at every plot. Dissolved oxygen (DO) in water was measured during the gas sampling by a handheld multi-parameter meter (Professional Plus, YSI, USA). The aboveground biomass of every replicate in the chamber was weighed after drying at 80 °C to constant mass.

Water level was measured after gas sampling at DW, SW and SF (when SF had standing water in August and September 2012). At site SF (when there was no standing water in November 2011, May, July and October 2012) and SFC, a

**Table 1.** Dominant plant species at each plot in different months. DW: deep water site, SW: shallow water site, SF: seasonally flooded site, SFC: “control site” for seasonally flooded site, NF: non-flooded site. A, B and C indicate sample plot with different vegetation. Species with aerenchyma are denoted <sup>A</sup>; species that are emergent are denoted <sup>E</sup>.

Site	Nov 2011	May 2012	Jul 2012	Aug 2012	Sep 2012	Oct 2012
DW	A	<i>Echinochloa colonum</i> <sup>AE</sup>		<i>Myriophyllum</i> sp.	<i>Trapa</i> <sup>AE</sup> sp.	
	B	no vegetation				
	C	<i>Typha angustifolia</i> <sup>AE</sup>				
SW	A	<i>Xanthium sibiricum</i> <sup>E</sup>	<i>Scirpus planiculmis</i> <sup>AE</sup>	<i>Echinochloa colonum</i> <sup>AE</sup>		
	B	<i>Setaria viridis</i> <sup>E</sup>	<i>Bidens pilosa</i> <sup>E</sup>	<i>Echinochloa colonum</i> <sup>AE</sup>		
	C	<i>Zea mays</i> <sup>E</sup>	<i>Polygonum lapathifolium</i> <sup>E</sup>	<i>Typha angustifolia</i> <sup>AE</sup>		
SF	A	<i>Xanthium sibiricum</i>	<i>Cirsium setosum</i>	<i>Cirsium setosum</i> <sup>E</sup>	<i>Cirsium setosum</i>	
	B	<i>Setaria viridis</i>	<i>Hemarthria altissima</i>	<i>Hemarthria altissima</i> <sup>E</sup>	<i>Hemarthria altissima</i>	
	C	<i>Zea mays</i>	<i>Polygonum lapathifolium</i>	<i>Polygonum lapathifolium</i> <sup>E</sup>	<i>Polygonum lapathifolium</i>	
SFC	A	no data		<i>Cirsium setosum</i>		
	B	no data		<i>Hemarthria altissima</i>		
	C	no data		<i>Zea mays</i>		
NF	A	<i>Xanthium sibiricum</i>				
	B	<i>Setaria viridis</i>	<i>Artemisia argyi</i>			
	C	<i>Zea mays</i>				

1m PVC tube was inserted vertically into the soil under the chamber after all monthly gas sampling was complete, allowing 2 h for the water level to equilibrate before measuring the level. The water table of site NF was calculated according to the elevation measured by a Global Navigation Satellite System receiver (BLH-L90, Daheng International, China).

Soil water content (SWC) was measured every month after all gas sampling with a soil water sensor (UNI1000, Shunlong, China). Soil–sediment samples (0–30 cm) at site DW, SW, SF and NF were collected at each replicate location in November 2011, except site SFC in October 2012. Fresh soil–sediment samples were used for NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> analysis using a discrete analyser (Smartchem 300, AMS, Italy). After air-drying and grinding (passing through a 100 mesh sieve), pH of 1 : 5 soil-water extractions was measured using a pH meter (IQ160, Hach, USA) while soil total carbon (TC) and nitrogen (TN) were analysed using an elemental analyser (vario MACRO cube, Elementar, Germany). Soil bulk density was measured following Chinese national standards NY/T 1121.4-2006 (MAPRC, 2007).

## 2.4 Statistical analysis

Flux differences were tested using a three-way ANOVA, and then using LSD for multiple comparisons (Table 2 and Fig. 4). A one-sample *t* test was used for testing whether the negative fluxes were statistically significantly different from zero. A log<sub>10</sub> transformation was used to explore the correlation between N<sub>2</sub>O flux and environmental variables (air temperature and soil NO<sub>3</sub><sup>-</sup>); where appropriate, a piecewise function (a two segment line) was calculated (SigmaPlot 11.0,

SYSTAT, USA). Spearman’s rank correlation was used to test for correlations between flux and environmental factors. Figures 5, 6, 7 and Table 3 were made using daily average fluxes. All the analyses above were performed using IBM SPSS Statistics (version 19.0, IBM, USA). Charts were made using SigmaPlot (version 11.0, SYSTAT, USA).


## 3 Results

### 3.1 Environmental characteristics

Precipitation occurred from March to November. The highest rainfall was in July, which accounted for one-fourth of the total (Fig. 2a). Water levels rose rapidly after the summer monsoon, and then declined after August (Fig. 2d). Temperature peaked during summer (Fig. 2c). The diurnal range in temperature was about 10 °C. The non-flooded site was very dry before the rains began, having fallen to only 10 %, but rose to 35 % after rain (Fig. 2e).

### 3.2 N<sub>2</sub>O fluxes

The mean flux from the littoral zone of the Miyun Reservoir was 6.8 μg m<sup>-2</sup> h<sup>-1</sup> (0.15 μmol m<sup>-2</sup> h<sup>-1</sup>), ranging from -136.6 to 381.8 μg m<sup>-2</sup> h<sup>-1</sup>. Negative flux was observed in about one-third of all the cases (*n* = 739, *p* < 0.001). In ANOVA (Table 2), both time of year and position on the transect had statistically significant effects (both *p* < 0.001), but time of day was not significant (*p* = 0.97). N<sub>2</sub>O emission from the non-flooded area (NF) was 17.0 ± 2.3 μg m<sup>-2</sup> h<sup>-1</sup>, which was significantly higher




**Figure 3.** Physicochemical properties (mean  $\pm$  SE) of soil–sediment of each site. Some SE bars are not visible because they are too small. DW: deep water site; SW: shallow water site; SF: seasonally flooded site; SFC: “control site” for the seasonally flooded site; NF: non-flooded site. (a), (b) and (c) denote samples from different vegetation types within each height band.

( $p < 0.001$ ) than the other four areas. There was no statistical difference ( $p = 0.91$ ) between emissions from the seasonal flooded area (SF) and its control site (SFC): fluxes were  $4.4 \pm 0.7$  and  $4.2 \pm 0.7 \mu\text{g m}^{-2} \text{h}^{-1}$ , respectively. For SW, SF, SFC and NF, the average emission of non-farmland plots was  $2.6 \mu\text{g m}^{-2} \text{h}^{-1}$  but the land growing maize during the sampling summer or the last summer reached 24.0 and  $8.4 \mu\text{g m}^{-2} \text{h}^{-1}$ , respectively (Fig. 4). Especially high emissions ( $43.7 \mu\text{g m}^{-2} \text{h}^{-1}$ ) were observed on farmland of NF (Fig. 4). Besides SF, where the highest emission occurred in late autumn, other high emissions were observed in the warm season, July and August in particular (Fig. 5).

### 3.3 Relationships between flux and environmental parameters

Rank correlation analysis was carried out between N<sub>2</sub>O flux and environmental parameters, but the coefficients were no higher than 0.38 (Table 3). For more information, correlation analysis was also carried out separately at each water level. The correlations were different among water levels and higher coefficients were shown between flux and air temperature in several cases (Table S1 in Supplement). Linear correlations can hide important non-linear features and so scatterplots are also shown, where log<sub>10</sub> flux was plotted against air temperature and soil NO<sub>3</sub><sup>−</sup> (Fig. 6). As fluxes were often negative (and significantly less than zero, implying a sink for N<sub>2</sub>O), we carried out a separate analysis of negative fluxes. Piecewise correlations were found between log<sub>10</sub> flux and



**Figure 4.** N<sub>2</sub>O flux (mean ± SE) at different water levels, months and times of day. Farmland included four plots, i.e. SW-C, SF-C, SFC-C and NF-C, which grew maize in the sampling growing season or the last growing season. Non-farmland included other 11 spots (see Table 1 for details of vegetation). DW: deep water site; SW: shallow water site; SF: seasonally flooded site; SFC: “control site” for the seasonally flooded site; NF: non-flooded site. Bars with different letters indicate a significant difference at  $p < 0.05$ . Difference analysis of bars with capital letters and small letters was done separately.

**Table 2.** ANOVA table to test the effects of water level, sampling month and time of day on N<sub>2</sub>O flux. The category of farmland included four plots, i.e. SW-C, SF-C, SFC-C and NF-C, which grew maize in the year of the study or the previous year. The category of non-farmland included other 11 spots (see Table 1 for details of vegetation).

	Effect	Type III SS	df	MS	F	p
All	Water level	65 808	4	16 452	25.3	< 0.001
	Month	65 546	5	13 109	20.2	< 0.001
	Time (of day)	918	6	153	0.2	0.965
	Water level × month	176 351	17	10 374	16.0	< 0.001
	Water level × time (of day)	4901	24	204	0.3	0.999
	Month × time (of day)	7277	30	243	0.4	0.999
	Water level × month × time	31 728	102	311	0.5	1.000
	Error	1347 885	2073	650		
Non-farmland	Water level	2982	4	745	5.9	< 0.001
	Month	3525	5	705	5.6	< 0.001
	Time (of day)	668	6	111	0.9	0.505
	Water level × month	11 830	17	696	5.5	< 0.001
	Water level × time (of day)	3087	24	129	1.0	0.431
	Month × time (of day)	4657	30	155	1.2	0.179
	Water level × month × time	14 385	102	141	1.1	0.198
	Error	186 701	1485	126		
Farmland (or use to be)	Water level	145 935	3	48 645	48.8	< 0.001
	Month	214 645	5	42 929	43.1	< 0.001
	Time (of day)	1286	6	214	0.2	0.972
	Water level × month	490 401	12	40 867	41.0	< 0.001
	Water level × time (of day)	6406	18	356	0.4	0.994
	Month × time (of day)	16 766	30	559	0.6	0.972
	Water level × month × time	46 388	72	644	0.6	0.988
	Error	439 735	441	997		

air temperature (Fig. 6). For positive fluxes, there was a negative correlation ( $p = 0.03$ ,  $n = 65$ ) when the air temperature was from 5.2 to 18.7 °C but a positive correlation ( $p < 0.01$ ,  $n = 175$ ) when air temperature was from 18.7 to 31.1 °C. For negative fluxes, there was a positive correlation ( $p < 0.01$ ,  $n = 43$ ) when the air temperature was from 5.2 to 17.6 °C


and an insignificant negative correlation ( $p = 0.12$ ,  $n = 41$ ) when air temperature was from 17.6 to 31.1 °C.

We present the relationship between nitrate and N<sub>2</sub>O emission. For positive flux, the soil NO<sub>3</sub><sup>-</sup> seemed to accelerate N<sub>2</sub>O emission when its concentration was higher than 7.1 mg kg<sup>-1</sup> ( $p < 0.01$ ,  $n = 122$ ), but it did not influence emission rate when lower than this “knot point” ( $p = 0.30$ ,



**Table 3.** Spearman's rank correlation ( $r$ ) between flux and environmental variables, included in the table are data from M. Yang et al. (2014) on the flux of CH<sub>4</sub>, collected at the same time as the N<sub>2</sub>O. \*\* indicates significant correlation ( $p < 0.01$ ); \* indicates significant correlation ( $p < 0.05$ ). SWC: soil water content, DO: dissolved oxygen, TC: total carbon, TN: total nitrogen. Daily average fluxes were used in the correlation analysis;  $n$  is from 84 to 324. #: data of DW were not included in the analysis since there was no contract sampling of farmland and non-farmland.

	N <sub>2</sub> O flux	N <sub>2</sub> O flux non-farmland#	N <sub>2</sub> O flux farmland#	CH <sub>4</sub> flux	Wind speed	Air temp	Water depth	SWC	Water DO	Biomass	Bulk density	Soil pH	Soil TC	Soil TN	Soil NH <sub>4</sub> <sup>+</sup>	Soil NO <sub>3</sub> <sup>-</sup>
N <sub>2</sub> O flux	1															
CH <sub>4</sub> flux	-0.10			1												
Wind speed	0.14*	0.06	-0.01	0.03	1											
Air temp	0.19**	0.05	0.38**	0.25**	0.30**	1										
Water depth	-0.02	-0.21**	-0.11	0.75**	0.06	0.16**	1									
SWC	-0.12*	-0.33**	-0.04	0.70**	0.03	0.29**	0.87**	1								
Water DO	0.35**	0.04	0.14	-0.28**	0.43**	-0.15	0.24**	0.00	1							
Biomass	-0.08	0.11	0.11	-0.26**	-0.15**	-0.34**	-0.38**	-0.52**	-0.48**	1						
Bulk density	0.00	0.17*	0.13	-0.53**	-0.01	-0.05	-0.78**	-0.67**	-0.26**	0.35**	1					
Soil pH	0.08	0.21**	0.19	-0.17**	-0.02	-0.03	-0.25**	-0.18**	-0.14	0.06	0.35**	1				
Soil TC	-0.04	-0.06	-0.08	0.62**	0.01	0.05	0.81**	0.74**	0.13	-0.35**	-0.77**	-0.26**	1			
Soil TN	0.03	-0.01	-0.06	0.56**	0.03	0.05	0.76**	0.67**	0.15	-0.33**	-0.73**	-0.21**	0.96**	1		
Soil NH <sub>4</sub> <sup>+</sup>	0.01	-0.13	0.03	0.18**	-0.14*	0.02	0.06	0.23**	-0.21**	-0.16**	-0.12*	-0.02	0.08	0.06	1	
Soil NO <sub>3</sub> <sup>-</sup>	0.25**	0.09	0.25*	-0.02	0.04	-0.01	0.09	0.10	0.28**	-0.07	-0.20**	0.27**	0.17**	0.19**	-0.11*	1




**Figure 5.** Monthly N<sub>2</sub>O flux (mean ± SE) of each site. Days between dotted lines were the high water level period and thus the seasonal flooded site (SF) was under water. DW: deep water site; SW: shallow water site; SF: seasonally flooded site; SFC: “control site” for the seasonally flooded site; NF: non-flooded site. (a), (b) and (c) denote samples from different vegetation types within each height band. Superscript F indicates farmland during the whole/part sampling time.

$n = 118$ ). Piecewise analysis was not attempted between negative flux and nitrate because of the very narrow nitrate concentrations (almost no data when soil NO<sub>3</sub><sup>-</sup> higher than 10 mg kg<sup>-1</sup>).

## 4 Discussion

### 4.1 N<sub>2</sub>O flux

Variations of N<sub>2</sub>O fluxes were compared at different spatial and temporal scales (Fig. 4 and Table 2). Whilst significant




**Figure 6.** Relationship between flux, air temperature and soil NO<sub>3</sub><sup>-</sup>. DW: deep water site; SW: shallow water site; SF: seasonally flooded site; SFC: “control site” for the seasonally flooded site; NF: non-flooded site. The result of piecewise correlation was plotted using flux data after log<sub>10</sub> transformation. Dashed lines indicate insignificant correlations while solid lines indicate significant correlations. See text for details.

differences were observed among water levels and sampling months, there were no differences among times of day, just as reported by Xia et al. (2013) in a polluted riverine system. We would expect soil microbes to respond to temperature, and given a diurnal range in air temperature of about 10 °C we would expect a detectable diurnal pattern in the N<sub>2</sub>O flux. We assume that the reason for a lack of response is that the microbial population is mostly deep in the soil–sediment–water system, where temperature variations are much smaller.

The mean flux from the littoral zone of the Miyun Reservoir was 6.8, from -136.6 to 381.8 μg m<sup>-2</sup> h<sup>-1</sup>. Negative fluxes were observed in about one-third of the cases, demonstrating a process of N<sub>2</sub>O consumption to be occurring. It

**Table 4.** Comparison of N<sub>2</sub>O and CH<sub>4</sub> emission from reservoir and farmland (both expressed as CO<sub>2</sub> equivalent, see text). Flux was transformed into CO<sub>2</sub> equivalent according to the global-warming potential (Stocker et al., 2013), i.e. 1 N<sub>2</sub>O = 298 CO<sub>2</sub>, 1 CH<sub>4</sub> = 34 CO<sub>2</sub>. Superscripts are \* the N<sub>2</sub>O flux was equivalent to 0.87 mg CO<sub>2</sub> m<sup>-2</sup> h<sup>-1</sup> while CH<sub>4</sub> flux was equivalent to 60.2 mg CO<sub>2</sub> m<sup>-2</sup> h<sup>-1</sup> when farmlands were excluded, i.e. SW-C, SF-C, SFC-C and NF-C; #, just SFC-C and NF-C were used for the calculation, where maize grew over the whole sampling time; \*\* unpublished data. Notes: Hubei is the province where part of the Three Gorges Reservoir is situated. Beijing is the city which includes the Miyun Reservoir. Maize, rice and wheat are the first three crops in terms of area in China.

	Study area	Zone or crop	N <sub>2</sub> O (mg CO <sub>2</sub> m <sup>-2</sup> h <sup>-1</sup> )	CH <sub>4</sub> (mg CO <sub>2</sub> m <sup>-2</sup> h <sup>-1</sup> )	Sum (mg CO <sub>2</sub> m <sup>-2</sup> h <sup>-1</sup> )	Data source
Reservoir	Three Gorges Reservoir	littoral zone	9.2	227.8	237	Chen et al. (2009, 2010)
		pelagic zone	4.2	8.8	13	D. Zhu et al. (2013), Chen et al. (2011a)
	Miyun Reservoir	littoral zone	2.0*	44.2*	46.2	This study, M. Yang et al. (2014)
		pelagic zone	no data	10.2	10.2	Yang et al. (2011)
Farmland	China-IPCC		2.5–16.7	no data	2.5–16.7	Xu et al. (2014), Smith et al. (2002)
	Hubei-DNDC typical farmland near Three Gorges Reservoir-observed	rice	26.8	85	111.8	Li et al. (2003)
		rice and rape	24.1	100.6	124.7	Zhang et al. (2012)
	Beijing-DNDC typical farmland near Miyun Reservoir-observed	rice and rape	33.7	47.6	81.3	Zhang et al. (2012)
		wheat	17.9	6.8	24.7	Li et al. (2003)
		maize	4.8	0.4	5.2	Hu et al. (2013)
	maize	24.1	0.5	24.6	Hu et al. (2013)	
	maize	9.1#	-0.3#**	8.8	This study	



**Figure 7.** Relationship between flux and water DO (mean  $\pm$  SE). DW: deep water site; SW: shallow water site; SF: seasonally flooded site.

is generally acknowledged that under certain conditions the capacity of soil to be a sink for N<sub>2</sub>O can, through denitrification, exceed its capacity to emit N<sub>2</sub>O (Baggs and Pilippot, 2010).

How do these fluxes compare to those reported from elsewhere? Our fluxes are comparable to those from the littoral zone of temperate-zone lakes, for example, a shallow lake in eastern Austria (Soja et al., 2014). However, in most of the cases, our fluxes were lower, as shown by the following comparisons. One similar-latitude lake, Lake Baiyangdian, had nearly 10 times higher N<sub>2</sub>O emissions, averaging 58  $\mu\text{g m}^{-2} \text{h}^{-1}$  (Yang et al., 2012). Higher emissions have also been reported in the littoral zone of lower-latitude sites, for example the Three Gorges Reservoir (Table 4). The seriously eutrophic Lake Taihu (latitude: 30° N) had a broader extent ranging from -278 to 2101  $\mu\text{g m}^{-2} \text{h}^{-1}$  in the littoral zone (Wang et al., 2007). Greenhouse gas emissions from low-latitude ecosystems are found to be higher than the corresponding ecosystems at high latitude because of the temperature effects (D. Zhu et al., 2013). The average N<sub>2</sub>O emission found in the present research was lower than that reported for boreal and Antarctic lakes (Huttunen et al., 2003; Y. Liu et al., 2011). The low N<sub>2</sub>O emission of Miyun Reservoir might be, in part, the consequence of relatively good water quality or high soil pH (Van den Heuvel et al., 2011).

N<sub>2</sub>O emissions from the littoral zone have been reported to be greater than for the pelagic zone (e.g. Huttunen et al., 2003, and see Table 4). We did not examine N<sub>2</sub>O fluxes



from the pelagic zone in this research, but we can compare our fluxes with pelagic data from elsewhere, as follows. The N<sub>2</sub>O emission in this study is slightly higher than those from five perialpine and alpine reservoirs ( $1.56 \mu\text{g m}^{-2} \text{h}^{-1}$ ) in Switzerland (Diem et al., 2012), while it is much lower than a same-latitude fluvial reservoir ( $84 \mu\text{g m}^{-2} \text{h}^{-1}$ ) located in an agricultural landscape near Indianapolis, USA (Jacinthe et al., 2012b). It should be noted that the comparison between littoral zone and pelagic zone of different reservoirs includes uncertainties, for example differences of elevation, nutrients input and the influence of topography on microclimate.

#### 4.2 Relative greenhouse gas effect: comparison with CH<sub>4</sub>

Elsewhere, we presented data on methane emissions from this reservoir (M. Yang et al., 2014). The global-warming potential (GWP) of N<sub>2</sub>O over a 100-year time span is 298 while CH<sub>4</sub> is 34 (Stocker et al., 2013). We can use the GWPs to calculate the emissions as CO<sub>2</sub>-equivalent emissions, and thus compare the warming effect of the two gases. The mean N<sub>2</sub>O emission in this study was  $2.0 \text{ mg CO}_2\text{-equivalent m}^{-2} \text{h}^{-1}$ . The CH<sub>4</sub> emission was  $44.2 \text{ mg CO}_2\text{-equivalent m}^{-2} \text{h}^{-1}$  (M. Yang et al., 2014), which is 22.1 times that of N<sub>2</sub>O. This contrasts with our previous findings, where the warming ratio of CH<sub>4</sub>:N<sub>2</sub>O was 1.5 (Li et al., 2014). But in our earlier report, N<sub>2</sub>O variation was investigated with a water recession process. Significant increases (nearly up to 1000 times) were observed after sediment exposure of 5 months. The high emissions may be the result of soil water content declining to 60–90 % (Ciarlo et al., 2007). In this research, the soil water content was not in this range at all, and that may have biased the comparison. In general, the flux ratio of CH<sub>4</sub> to N<sub>2</sub>O in aquatic environments varies considerably. For example, the CH<sub>4</sub>:N<sub>2</sub>O ratio of permanent flooded areas at Poyang Lake was 1.1 (Liu et al., 2013) while the ratio was 0.6 for the pelagic zone of a fluvial reservoir in central Indiana (Jacinthe et al., 2012b). In a study which monitored the flux of both littoral and pelagic zone of a temperate lake, the average CH<sub>4</sub>:N<sub>2</sub>O ratio is 7.2 (Soja et al., 2014). For a freshwater marsh at northeast of China, it was found to be as high as 66.5 (Yang et al., 2013). Although the ratio varies greatly, there is nevertheless a considerable contribution of N<sub>2</sub>O emission from aquatic ecosystems to global warming, whose importance may have been somewhat understated in relation to the large CH<sub>4</sub> emission.

#### 4.3 Environmental controls

##### 4.3.1 Flooding

Unlike the specific influence of flooding on CH<sub>4</sub> emission (M. Yang et al., 2014), flooding effects on N<sub>2</sub>O emission were not very clear in this study. The N<sub>2</sub>O flux of seasonal flooded area SF was as high as its control area SFC which

escaped flooding because of higher elevation (Fig. 4). Inundation nearly always causes a drop of N<sub>2</sub>O emissions (Yang et al., 2013). Standing water could inhibit N<sub>2</sub>O emission through slowing down the diffusive transportation of gas, causing anoxia, activating a different component of the microbiota, leading to the reduction of N<sub>2</sub>O to N<sub>2</sub> (Lienggaard et al., 2013; Pilegaard, 2013). Our results did not reject those possibilities when looking into the seasonal variation of N<sub>2</sub>O flux of seasonal flooded sites, but they did not completely support that hypothesis either (Fig. 5c). After flooding, the fluxes of two sites (SF-A and SF-B) were no higher than before flooding and no higher than their control sites. However, a single extraordinary observation showed the highest emission was during flooding (Fig. 5c, SF-C). A somewhat similar result was also observed at an artificial wetland (Hernandez and Mitsch, 2006). An incubation study showed both increasing N<sub>2</sub>O emission and stable emission during flooding at different treatments, i.e. N<sub>2</sub>O emission of residue-incorporated soils, increased remarkably from the 6th to 30th days of flooding and decreased to lower level than before flooding afterward. However, the N<sub>2</sub>O emission of the soils with residues on the surface was stable before and during flooding (Zschornack et al., 2011). It suggested that other factors would influence N<sub>2</sub>O emission responses to flooding. Even though there are uncertainties about the mechanisms, this study implied that flooding introduces a complex set of processes that influence N<sub>2</sub>O flux, when compared to non-flooded areas whose fluxes were all more or less coordinated with temperature variation (Fig. 5a, b and d).

Besides, floods may influence N<sub>2</sub>O production both in the long term and short term (Jacinthe et al., 2012a). Quick response of N<sub>2</sub>O flux after flooding was shown at a coastal marsh; that is, N<sub>2</sub>O emission decreased in 2.5–5 h after flooding but then increased to the original level after flooding for 7.5 h (Sun et al., 2014). The possibility of emissions occurring in discrete pulses, especially by ebullition, should be kept in mind when interpreting results from flux chambers. It also emphasizes the importance of continuous high-frequency monitoring to reveal flooding effects with lower uncertainties.

##### 4.3.2 Other environmental conditions

Positive correlations between N<sub>2</sub>O emission and temperature were reported in previous studies (e.g. Wang et al., 2014). But in this study we found both positive and negative fluxes, and decided to fit a piecewise regression to the log-transformation data (Fig. 6). This complex and non-linear picture might explain the low coefficients in the correlation analysis (Table 3).

N<sub>2</sub>O production is generally caused by several processes, for example denitrification, nitrification, nitrate ammonification and nitrifier denitrification. N<sub>2</sub>O consumption has been much less studied (Baggs and Pilippot, 2010). Some studies have found denitrification to be the main contributor in

N<sub>2</sub>O emission while some others pointed out that several processes occurred simultaneously with a shifting dominance of processes caused by environmental limitations, for instance soil moisture and O<sub>2</sub> availability (Kool et al., 2011; X. Zhu et al., 2013). Controlled studies showed that N<sub>2</sub>O production via a single process always changes according to temperature, if not exceeded by biotic tolerance (Sierra, 2002; Veraart et al., 2011). Our complex N<sub>2</sub>O response to temperature supported the latter notion, i.e. multi-processes occurring and competing during our sampling campaigns. Furthermore, it demonstrated that the response of N<sub>2</sub>O production and consumption to temperature was at different rates (Xie et al., 2003). As some chambers within a treatment showed efflux whilst others showed influx, we may presume that the substrate is patchy, over scales of a few metres, reflecting an underlying heterogeneity possibly the result of the distribution of underlying decaying vegetation.

Negative relationships between N<sub>2</sub>O flux and O<sub>2</sub> are reported in both laboratory experiments and field studies (Rosamond et al., 2012; Rubol et al., 2012; Zhao et al., 2014). This is explained by the fact that denitrification, which is activated in anoxic environments, is likely to be controlling N<sub>2</sub>O emissions (Xia et al., 2013). Our present result contradicted those previous conclusions because a significantly positive correlation was observed between N<sub>2</sub>O flux and water DO (Fig. 7). N<sub>2</sub>O accumulation in the water column has been shown to depend not only on production rate but also on the extent of N<sub>2</sub>O reduction to N<sub>2</sub> by reductase enzymes (Zhao et al., 2014). An incubation study showed that denitrifying activity decreased along with rise of DO concentration, but the N<sub>2</sub>O producing activity increased because of less N<sub>2</sub>O reduction to N<sub>2</sub> (Senga et al., 2002). Furthermore, Senga's study also pointed out that N<sub>2</sub>O produced by nitrification could also be reduced to N<sub>2</sub> via denitrification. That might have happened in our sampling field, i.e. along with increasing of water DO, decreasing of N<sub>2</sub>O reduction to N<sub>2</sub> allowing more N<sub>2</sub>O to be released at water–air interface, no matter in which processes the N<sub>2</sub>O was produced. Further study should focus on responses of both N<sub>2</sub>O production and reduction to water DO and factors determining which process is dominant.

Soil NO<sub>3</sub><sup>-</sup> is an important anion in N cycle (Butterbach-Bahl et al., 2013). Positive correlations between N<sub>2</sub>O flux and nitrate have been reported (Soja et al., 2014; Y. Liu et al., 2011; X. L. Liu et al., 2011). It is therefore not surprising to find the highest emission where highest soil NO<sub>3</sub><sup>-</sup> occurred. However, in this research when soil NO<sub>3</sub><sup>-</sup> was less than the threshold value of 7.1 mg kg<sup>-1</sup> there was no relationship with NO<sub>3</sub><sup>-</sup>. In agricultural studies the NO<sub>3</sub><sup>-</sup> concentrations are generally much higher, but even then a threshold phenomenon has been reported (Bao et al., 2012). This implies that substrate constraint might be a reason for the weak correlations between N<sub>2</sub>O flux and other environmental factors. In the present study, no significant correlation was shown between N<sub>2</sub>O flux and NH<sub>4</sub><sup>+</sup>, although NH<sub>4</sub><sup>+</sup> is

also important in the N cycle. An N fertilizer experiment in a temperate forest found that the N<sub>2</sub>O emissions were only significantly correlated with soil NO<sub>3</sub><sup>-</sup> and temperature, but not soil NH<sub>4</sub><sup>+</sup> (Bai et al., 2014). A global review study found that among the five chemical forms of N fertilizer assessed (including NH<sub>4</sub><sup>+</sup>), NO<sub>3</sub><sup>-</sup> showed the strongest stimulation of N<sub>2</sub>O emission, approximately 2 to 3 times higher than the others (Liu and Greaver, 2009).

Based on the above discussion and discussion in a previous paper (M. Yang et al., 2014), the influence of environmental factors on N<sub>2</sub>O and CH<sub>4</sub> emission was summarized as follows. The emissions of these two gases are influenced by different factors and in different ways (Table 3), depending on soil conditions, meteorology and vegetation. Methane shows relatively strong correlation with environmental variables while the correlations are always rather weak in N<sub>2</sub>O, reflecting the number and complexity of the microbial processes governing the flux of N<sub>2</sub>O. The variables likely to be associated with anoxia (soil water depth, soil water content, water DO) were important for both N<sub>2</sub>O (see above discussion) and CH<sub>4</sub> (Serrano-Silva et al., 2014) but acted in converse ways. Soil nutrients also influence both of the two gases, but it seems through different parameters (Table 3). The soil water status in the natural environment controls anoxia and influences soil temperature and soil nutrients, implying a fundamental role of soil water levels acting on N<sub>2</sub>O and CH<sub>4</sub> emissions. Therefore, we conclude that water level is the most important factor determining N<sub>2</sub>O and CH<sub>4</sub> emission in littoral zone.

#### 4.4 Comparison with farmland

Reclamation of the shore by local farmers, to supplement their income, is not rare. In this research we compared the N<sub>2</sub>O emission of natural and farm-related area in the littoral zone. Significantly higher emissions were observed at sites cropped with maize in the sampling season or the last growing season. The emission was 24.0 and 8.4 μg m<sup>-2</sup> h<sup>-1</sup>, respectively, while the emission of natural sites was 2.6 μg m<sup>-2</sup> h<sup>-1</sup>. As discussed in the above section, soil NO<sub>3</sub><sup>-</sup> might partly explain the flux difference between farm-related land and natural land. Besides, tillage might also influencing N<sub>2</sub>O emission through soil aeration (Buchkina et al., 2013).

Reservoirs are being developed, in part, for “clean energy”, and reports of high greenhouse gas emissions from reservoirs have already led some authors to question the “clean” concept, especially in relation to the mitigation of climate change (Gunkel, 2009). To evaluate the role that reservoirs play in climate change, their greenhouse gas emissions ought to be compared with those of the prior ecosystem (Tremblay et al., 2005). Farmland is one of the several ecosystems which are lost by flooding during reservoir construction in China. Total emission of N<sub>2</sub>O and CH<sub>4</sub> in the littoral zone was higher than in farmland (Table 4). The range

of soil water content of most farmland soils is relatively narrow and even. Crops, with the exception of rice, do not tolerate flooding or drought. But soil moisture of the littoral zone is patchy and ranges from flooded to seasonally dry. The littoral zone is therefore more precarious in terms of N<sub>2</sub>O or CH<sub>4</sub> emissions than farming (Groffman et al., 2009). Even though the emission from the littoral zone was higher, considering its small area and the low emission of the pelagic zone, N<sub>2</sub>O and CH<sub>4</sub> emissions from reservoirs are likely to be lower than farmland. It is worth noting that N<sub>2</sub>O and CH<sub>4</sub> emissions may vary with the type of crop, and so there could be exceptions to this generalization.

## 5 Conclusions

Finally, we return to our original hypothesis: the littoral zone is a hotspot of N<sub>2</sub>O emissions that is influenced by seasonal changes in the water level. We find that the littoral zone is indeed a hotspot for N<sub>2</sub>O in the summer, especially when the shores of the lake are used for opportunistic farming of maize. But in terms of the overall greenhouse gas budget, the fluxes of N<sub>2</sub>O from the littoral zone are not as important as those of CH<sub>4</sub>.

**The Supplement related to this article is available online at doi:10.5194/bg-12-4711-2015-supplement.**

*Acknowledgements.* This study was financially supported by State Forestry Administration of China under grant 200804005 and China Scholarship Council. We thank Yi Zhu, Lei Guan, Yamian Zhang, Nana Li, Jialin Lei, Rui Li, Duoduo Feng, Hairui Duo, Lei Jing, Qing Zeng, Chu Lang, Xu Luo, Jiayuan Li, Yonghong Gao, Defeng Bai, Siyu Zhu, and Jianing Xu for their great support during the course of this study. We also thank Beijing North Miyun Reservoir Eco-agriculture Co. Ltd for granting us permission to conduct the study on its land. We sincerely appreciate suggestions and help from editors and reviewers on this paper.

Edited by: T. J. Battin

## References

- Ahn, J. H., Choi, M. Y., Kim, B. Y., Lee, J. S., Song, J., Kim, G. Y., and Weon, H. Y.: Effects of water-saving irrigation on emissions of greenhouse gases and Prokaryotic Communities in rice paddy soil, *Microb. Ecol.*, 68, 271–283, 2014.
- Bai, E., Li, W., Li, S. L., Sun, J. F., Peng, B., Dai, W. W., Jiang, P., and Han, S. J.: Pulse increase of soil N<sub>2</sub>O emission in response to N addition in a temperate forest on Mt Changbai, northeast China, *Plos One*, 9, e102765, doi:10.1371/journal.pone.0102765, 2014.
- Baggs, E. and Pilippot, L.: Microbial Terrestrial Pathways to Nitrous Oxide, in: *Nitrous Oxide and Climate Change*, edited by: Smith, K. A., Earthscan, London, 4–35, 2010.
- Bao, Q., Ju, X., Gao, B., Qu, Z., Christie, P., and Lu, Y.: Response of nitrous oxide and corresponding bacteria to managements in an agricultural soil, *Soil Sci. Soc. Am. J.*, 76, 130–141, 2012.
- Beaulieu, J. J., Smolenski, R. L., Nietch, C. T., Townsend-Small, A., Elovitz, M. S., and Schubauer-Berigan, J. P.: Denitrification alternates between a source and sink of nitrous oxide in the hypolimnion of a thermally stratified reservoir, *Limnol. Oceanogr.*, 59, 495–506, 2014.
- Buchkina, N. P., Rizhiya, E. Y., Pavlik, S. V., and Balashov, E. V.: Soil Physical Properties and Nitrous Oxide Emission from Agricultural Soils, in: *Advances in Agrophysical Research*, edited by: Grundas, S. and Stepniewski, A., 194–196, 2013.
- Butterbach-Bahl, K., Baggs, E. M., Dannenmann, M., Kiese, R., and Zechmeister-Boltenstern, S.: Nitrous oxide emissions from soils: how well do we understand the processes and their controls?, *Philos. T. R. Soc. B.*, 368, 20130122, doi:10.1098/rstb.2013.0122, 2013.
- Cao, R., Li, C. J., Liu, L. J., Wang, J. H., and Yan, G. J.: Extracting Miyun reservoirs water area and monitoring its change based on a revised normalized different water index, *Sci. Surv. Mapp.*, 33, 158–160, 2008.
- Chen, H., Wu, Y. Y., Yuan, X. Z., Gao, Y. H., Wu, N., and Zhu, D.: Methane emissions from newly created marshes in the drawdown area of the Three Gorges Reservoir, *J. Geophys. Res. Atmos.*, 114, D18301, doi:10.1029/2009jd012410, 2009.
- Chen, H., Yuan, X., Gao, Y., Wu, N., Zhu, D., and Wang, J.: Nitrous oxide emissions from newly created littoral marshes in the drawdown area of the Three Gorges Reservoir, China, *Water, Air Soil Pollut.*, 211, 25–33, 2010.
- Chen, H., Yuan, X., Chen, Z., Wu, Y., Liu, X., Zhu, D., Wu, N., Zhu, Q. a., Peng, C., and Li, W.: Methane emissions from the surface of the Three Gorges Reservoir, *J. Geophys. Res. Atmos.*, 116, D21306, doi:10.1029/2011JD016244, 2011a.
- Chen, H., Wang, M., Wu, N., Wang, Y., Zhu, D., Gao, Y., and Peng, C.: Nitrous oxide fluxes from the littoral zone of a lake on the Qinghai-Tibetan Plateau, *Environ. Monit. Assess.*, 182, 545–553, 2011b.
- Ciarlo, E., Conti, M., Bartoloni, N., and Rubio, G.: The effect of moisture on nitrous oxide emissions from soil and the N<sub>2</sub>O/(N<sub>2</sub>O+ N<sub>2</sub>) ratio under laboratory conditions, *Biol. Fertility Soils*, 43, 675–681, 2007.
- Diem, T., Koch, S., Schwarzenbach, S., Wehrli, B., and Schubert, C. J.: Greenhouse gas emissions (CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O) from several perialpine and alpine hydropower reservoirs by diffusion and loss in turbines, *Aquat. Sci.*, 74, 619–635, 2012.
- Gao, Z. Q.: Beijing chronicles of water conservancy, volume III, Editing Committee of the Beijing Chronicles of Water Conservancy, Beijing, 1989.
- Groffman, P. M., Butterbach-Bahl, K., Fulweiler, R. W., Gold, A. J., Morse, J. L., Stander, E. K., Tague, C., Tonitto, C., and Vidon, P.: Challenges to incorporating spatially and temporally explicit phenomena (hotspots and hot moments) in denitrification models, *Biogeochemistry*, 93, 49–77, 2009.
- Guérin, F., Abril, G., Tremblay, A., and Delmas, R.: Nitrous oxide emissions from tropical hydroelectric reservoirs, *Geophys. Res. Lett.*, 35, L06404, doi:10.1029/2007GL033057, 2008.

- Gunkel, G.: Hydropower – a green energy? Tropical reservoirs and greenhouse gas emissions, *Clean-Soil Air Water*, 37, 726–734, 2009.
- Hernandez, M., and Mitsch, W.: Influence of hydrologic pulses, flooding frequency, and vegetation on nitrous oxide emissions from created riparian marshes, *Wetlands*, 26, 862–877, 2006.
- Hu, X. K., Su, F., Ju, X. T., Gao, B., Oenema, O., Christie, P., Huang, B. X., Jiang, R. F., and Zhang, F. S.: Greenhouse gas emissions from a wheat-maize double cropping system with different nitrogen fertilization regimes, *Environ. Pollut.*, 176, 198–207, 2013.
- Huttunen, J. T., Vaisanen, T. S., Hellsten, S. K., Heikkinen, M., Nykanen, H., Jungner, H., Niskanen, A., Virtanen, M. O., Lindqvist, O. V., Nenonen, O. S., and Martikainen, P. J.: Fluxes of CH<sub>4</sub>, CO<sub>2</sub>, and N<sub>2</sub>O in hydroelectric reservoirs Lokka and Porttipahta in the northern boreal zone in Finland, *Global Biogeochem. Cy.*, 16, 1003, doi:10.1029/2000GB001316, 2002.
- Huttunen, J. T., Juutinen, S., Alm, J., Larmola, T., Hammar, T., Silvola, J., and Martikainen, P. J.: Nitrous oxide flux to the atmosphere from the littoral zone of a boreal lake, *J. Geophys. Res.*, 108, 4421, doi:10.1029/2002JD002989, 2003.
- Jacinte, P. A., Bills, J. S., Tedesco, L. P., and Barr, R. C.: Nitrous oxide emission from riparian buffers in relation to vegetation and flood frequency, *J. Environ. Qual.*, 41, 95–105, 2012a.
- Jacinte, P. A., Filippelli, G. M., Tedesco, L. P., and Raftis, R.: Carbon storage and greenhouse gases emission from a fluvial reservoir in an agricultural landscape, *Catena*, 94, 53–63, 2012b.
- Kool, D. M., Dolfing, J., Wrage, N., and Van Groenigen, J. W.: Nitrifier denitrification as a distinct and significant source of nitrous oxide from soil, *Soil Biol. Biochem.*, 43, 174–178, 2011.
- Kumar, A., Schei, T., Ahenkorah, A., Rodriguez, R. C., Devernay, J. M., Freitas, M., Hall, D., A. Killingtveit, and Liu, Z.: Hydropower, in: *IPCC Special Report on Renewable Energy Sources and Climate Change Mitigation* edited by: Edenhofer, O., Pichs-Madruga, R., Sokona, Y., Seyboth, K., Matschoss, P., Kadner, S., Zwickel, T., Eickemeier, P., Hansen, G., Schlömer, S., and von Stechow, C., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 443–466, 2011.
- Li, C., Xiao, X. M., Frolking, S., Moore, B., Salas, W., Qiu, J., Zhang, Y., Zhuang, Y., Wang, X., and Dai, Z.: Greenhouse gas emissions from croplands of China, *Quaternary Sci.*, 23, 493–404, 2003.
- Li, H., Yang, M., Lei, T., Zhang, M., Bridgewater, P., Lu, C., Geng, X., and Lei, G.: Nitrous oxide emission from the littoral zones of the Miyun Reservoir near Beijing, China, *Hydrol. Res.*, in press, doi:10.2166/nh.2014.095, 2014.
- Lienggaard, L., Nielsen, L. P., Revsbech, N. P., Prieme, A., Elberling, B., Enrich-Prast, A., and Kuhl, M.: Extreme emission of N<sub>2</sub>O from tropical wetland soil (pantanal, South America), *Front Microbiol.*, 3, 433, doi:10.3389/fmicb.2012.00433, 2013.
- Liu, L. L. and Greaver, T. L.: A review of nitrogen enrichment effects on three biogenic GHGs: the CO<sub>2</sub> sink may be largely offset by stimulated N<sub>2</sub>O and CH<sub>4</sub> emission, *Ecol. Lett.*, 12, 1103–1117, 2009.
- Liu, L., Xu, M., Lin, M., and Zhang, X.: Spatial variability of greenhouse gas effluxes and their controlling factors in the Poyang Lake in China, *Pol. J. Environ. Stud.*, 22, 749–758, 2013.
- Liu, X. L., Liu, C. Q., Li, S. L., Wang, F. S., Wang, B. L., and Wang, Z. L.: Spatiotemporal variations of nitrous oxide (N<sub>2</sub>O) emissions from two reservoirs in SW China, *Atmos. Environ.*, 45, 5458–5468, 2011.
- Liu, Y., Zhu, R., Ma, D., Xu, H., Luo, Y., Huang, T., and Sun, L.: Temporal and spatial variations of nitrous oxide fluxes from the littoral zones of three alga-rich lakes in coastal Antarctica, *Atmos. Environ.*, 45, 1464–1475, 2011.
- Lu, Y. and Xu, H.: Effects of soil temperature, flooding, and organic matter addition on N<sub>2</sub>O emissions from a soil of Hongze Lake wetland, China, *The Scientific World Journal*, 2014, 272684–272684, 2014.
- MAPRC: Ministry of agriculture of People's Republic of China, available at: <http://cx.spsp.gov.cn/index.aspx> (last access: 10 November 2013), 2007.
- Peng, S., Hou, H., Xu, J., Mao, Z., Abudu, S., and Luo, Y.: Nitrous oxide emissions from paddy fields under different water managements in southeast China, *Paddy Water Environ.*, 9, 403–411, 2011.
- Pilegaard, K.: Processes regulating nitric oxide emissions from soils, *Philos. T. R. Soc. B.*, 368, 20130126, doi:10.1098/rstb.2013.0126, 2013.
- Rosamond, M. S., Thuss, S. J., and Schiff, S. L.: Dependence of riverine nitrous oxide emissions on dissolved oxygen levels, *Nature Geosci.*, 5, 715–718, 2012.
- Rubol, S., Silver, W. L., and Bellin, A.: Hydrologic control on redox and nitrogen dynamics in a peatland soil, *Sci. Total Environ.*, 432, 37–46, 2012.
- Senga, Y., Mochida, K., Okamoto, N., Fukumori, R., and Seike, Y.: Nitrous oxide in brackish Lake Nakaumi, Japan II: the role of nitrification and denitrification in N<sub>2</sub>O accumulation, *Limnology*, 3, 21–27, 2002.
- Serrano-Silva, N., Sarria-Guzman, Y., Dendooven, L., and Luna-Guido, M.: Methanogenesis and methanotrophy in soil: a review, *Pedosphere*, 24, 291–307, 2014.
- Sierra, J.: Nitrogen mineralization and nitrification in a tropical soil: effects of fluctuating temperature conditions, *Soil Biol. Biochem.*, 34, 1219–1226, 2002.
- Smith, K. A., Bouwman, L., and Braatz, B.: N<sub>2</sub>O: direct emissions from agricultural soils. In: *Background Papers: IPCC Meetings on Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories*, Inst. for Global Environmental Strategies, Kanagawa, Japan, 361–380, 2002.
- Soja, G., Kitzler, B., and Soja, A. M.: Emissions of greenhouse gases from Lake Neusiedl, a shallow steppe lake in Eastern Austria, *Hydrobiologia*, 731, 125–138, 2014.
- Stocker, T. F., Qin, D., Plattner, G. K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M.: *Climate Change 2013: The physical science basis. contribution of working group I to the fifth assessment report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- Sun, Z., Wang, L., Mou, X., Jiang, H., and Sun, W.: Spatial and temporal variations of nitrous oxide flux between coastal marsh and the atmosphere in the Yellow River estuary of China, *Environ. Sci. Pollut. R.*, 21, 419–433, 2014.
- Tranvik, L. J., Downing, J. A., Cotner, J. B., Loiselle, S. A., Striegl, R. G., Ballatore, T. J., Dillon, P., Finlay, K., Fortino, K., Knoll, L.

- B., Kortelainen, P. L., Kutser, T., Larsen, S., Laurion, I., Leech, D. M., McCallister, S. L., McKnight, D. M., Melack, J. M., Overholt, E., Porter, J. A., Prairie, Y., Renwick, W. H., Roland, F., Sherman, B. S., Schindler, D. W., Sobek, S., Tremblay, A., Vanni, M. J., Verschoor, A. M., von Wachenfeldt, E., and Weyhenmeyer, G. A.: Lakes and reservoirs as regulators of carbon cycling and climate, *Limnol. Oceanogr.*, 54, 2298–2314, 2009.
- Tremblay, A., Varfalvy, L., Roehm, C., and Garneau, M.: Greenhouse gas emissions-fluxes and processes, *Hydroelectric Reservoirs and Natural Environments*, Springer, Berlin, 21–60, 2005.
- Trost, B., Prochnow, A., Drastig, K., Meyer-Aurich, A., Ellmer, F., and Baumecker, M.: Irrigation, soil organic carbon and N<sub>2</sub>O emissions: a review, *Agron. Sustain. Dev.*, 33, 1–17, 2013.
- Van den Heuvel, R. N., Bakker, S. E., Jetten, M. S. M., and Hefting, M. M.: Decreased N<sub>2</sub>O reduction by low soil pH causes high N<sub>2</sub>O emissions in a riparian ecosystem, *Geobiology*, 9, 294–300, 2011.
- Veraart, A. J., De Klein, J. J., and Scheffer, M.: Warming can boost denitrification disproportionately due to altered oxygen dynamics, *Plos One*, 6, e18508, doi:10.1371/journal.pone.0018508, 2011.
- Wang, H., Wang, W., Yin, C., Wang, Y., and Lu, J.: Littoral zones as the “hotspots” of nitrous oxide (N<sub>2</sub>O) emission in a hyper-eutrophic lake in China, *Atmos. Environ.*, 40, 5522–5527, 2006.
- Wang, H. J., Yang, L. Y., Wang, W. D., Lu, J. W., and Yin, C. Q.: Nitrous oxide (N<sub>2</sub>O) fluxes and their relationships with water-sediment characteristics in a hyper-eutrophic shallow lake, China, *J. Geophys. Res. Biogeo.*, 112, G01005, doi:10.1029/2005jg000129, 2007.
- Wang, Y. S., Cheng, S. L., Fang, H. J., Yu, G. R., Xu, M. J., Dang, X. S., Li, L. S., and Wang, L.: Simulated nitrogen deposition reduces CH<sub>4</sub> uptake and increases N<sub>2</sub>O emission from a subtropical plantation forest soil in southern China, *Plos One*, 9, e93571, doi:10.1371/journal.pone.0093571, 2014.
- Xia, Y., Li, Y., Li, X., Guo, M., She, D., and Yan, X.: Diurnal pattern in nitrous oxide emissions from a sewage-enriched river, *Chemosphere*, 92, 421–428, 2013.
- Xie, S., Zhang, X., and Wang, Z.: Temperature effect on aerobic denitrification and nitrification, *J. Environ. Sci. (China)*, 15, 669–673, 2003.
- Xu, X., He, P., Qiu, S., Pampolino, M. F., Zhao, S., Johnston, A. M., and Zhou, W.: Estimating a new approach of fertilizer recommendation across small-holder farms in China, *Field Crops Res.*, 163, 10–17, 2014.
- Yang, J., Liu, J., Hu, X., Li, X., Wang, Y., and Li, H.: Effect of water table level on CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O emissions in a freshwater marsh of Northeast China, *Soil Biol. Biochem.*, 61, 52–60, 2013.
- Yang, L., Lu, F., Zhou, X., Wang, X., Duan, X., and Sun, B.: Progress in the studies on the greenhouse gas emissions from reservoirs, *Acta Ecol. Sinica*, 34, 204–212, 2014.
- Yang, M., Li, H. L., Lei, T., Zhou, Y., and Lu, C.: Spatial-temporal variation of CH<sub>4</sub> flux and its environmental factors at Miyun Reservoir, *Wetland Sci.*, 9, 191–197, 2011.
- Yang, M., Geng, X., Grace, J., Lu, C., Zhu, Y., Zhou, Y., and Lei, G.: Spatial and seasonal CH<sub>4</sub> flux in the littoral zone of Miyun Reservoir near Beijing: the effects of water level and its fluctuation, *Plos One*, 9, e94275, doi:10.1371/journal.pone.0094275, 2014.
- Yang, X. and Lu, X.: Drastic change in China’s lakes and reservoirs over the past decades, *Sci. Rep.*, 4, 6041, doi:10.1038/srep06041, 2014.
- Yang, Z., Zhao, Y., and Xia, X.: Nitrous oxide emissions from *Phragmites australis*-dominated zones in a shallow lake, *Environ. Pollut.*, 166, 116–124, 2012.
- Zhang, J. K., Jiang, C. S., Hao, Q. J., Tang, Q. W., Cheng, B. H., Li, H., and Chen, L. H.: Effects of tillage-cropping systems on methane and nitrous oxide emissions from agro-ecosystems in a purple paddy soil, *Huan Jing Ke Xue*, 33, 1979–1986, 2012.
- Zhao, Y., Xia, Y., Li, B., and Yan, X.: Influence of environmental factors on net N<sub>2</sub> and N<sub>2</sub>O production in sediment of freshwater rivers, *Environ. Sci. Pollut. R.*, 21, 9973–9982, 2014.
- Zhu, D., Chen, H., Yuan, X., Wu, N., Gao, Y., Wu, Y., Zhang, Y., Peng, C., Zhu, Q., Yang, G., and Wu, J.: Nitrous oxide emissions from the surface of the Three Gorges Reservoir, *Ecol. Eng.*, 60, 150–154, 2013.
- Zhu, X., Burger, M., Doane, T. A., and Horwath, W. R.: Ammonia oxidation pathways and nitrifier denitrification are significant sources of N<sub>2</sub>O and NO under low oxygen availability, *P. Natl. Acad. Sci. USA*, 110, 6328–6333, 2013.
- Zschornack, T., Bayer, C., Zanatta, J. A., Vieira, F. C. B., and Anghinoni, I.: Mitigation of methane and nitrous oxide emissions from flood-irrigated rice by no incorporation of winter crop residues into the soil, *Rev. Bras. Cienc. Solo.*, 35, 623–634, 2011.