



## Corrigendum to “Coastal-ocean uptake of anthropogenic carbon” published in Biogeosciences, 13, 4167–4185, 2016

Timothée Bourgeois<sup>1</sup>, James C. Orr<sup>1</sup>, Laure Resplandy<sup>2</sup>, Jens Terhaar<sup>1</sup>, Christian Ethé<sup>3</sup>, Marion Gehlen<sup>1</sup>, and Laurent Bopp<sup>1</sup>

<sup>1</sup>Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA-CNRS-UVSQ, Université Paris-Saclay, 91191 Gif-sur-Yvette, France


<sup>2</sup>Scripps Institution of Oceanography, University of California San Diego, La Jolla, CA, USA

<sup>3</sup>Institut Pierre Simon Laplace, 4 Place Jussieu, 75005 Paris, France

Correspondence to: Timothée Bourgeois (timothee.bourgeois@lsce.ipsl.fr)

Published: 11 August 2016

We chose to remove the panel (b) of the Fig. 6 during the manuscript revision. However, the caption of the Fig. 6 has not been updated accordingly in the manuscript. The new caption of Fig. 6 should be read as follows:



**Figure 6.** Simulated vs. observed MARCATS sea-to-air flux of total carbon in  $\text{mol C m}^{-2} \text{yr}^{-1}$ . Vertical error bars show the standard deviation from the 1993–2012 interannual variability for model results and the horizontal bars correspond to the 1990–2011 variability from computational methods used in LA14 for observation-based estimates. Here, regression line (grey dotted) has y intercept forced to 0. All MARCATS regions have been used except the Black Sea, the Persian Gulf (no data estimate), and the Sea of Okhotsk (see text).