



Urbanisation-related land use change from forest and pasture into turf grass modifies soil nitrogen cycling and increases N₂O emissions

Lona van Delden, David W. Rowlings, Clemens Scheer, and Peter R. Grace

Institute for Future Environments (IFE), Queensland University of Technology (QUT), Brisbane, QLD, Australia

Correspondence to: Lona van Delden (lona.vandelden@qut.edu.au)

Received: 19 May 2016 – Published in Biogeosciences Discuss.: 31 May 2016

Revised: 6 September 2016 – Accepted: 17 October 2016 – Published: 7 November 2016

Abstract. Urbanisation is becoming increasingly important in terms of climate change and ecosystem functionality worldwide. We are only beginning to understand how the processes of urbanisation influence ecosystem dynamics, making peri-urban environments more vulnerable to nutrient losses. Brisbane in South East Queensland has the most extensive urban sprawl of all Australian cities. This research estimated the environmental impact of land use change associated with urbanisation by examining soil nitrogen (N) turnover and subsequent nitrous oxide (N₂O) emissions using a fully automated system that measured emissions on a sub-daily basis. There was no significant difference in soil N₂O emissions between the native dry sclerophyll eucalypt forest and an extensively grazed pasture, wherefrom only low annual emissions were observed amounting to 0.1 and 0.2 kg N₂O ha⁻¹ yr⁻¹, respectively. The establishment of a fertilised turf grass lawn increased soil N₂O emissions 18-fold (1.8 kg N₂O ha⁻¹ yr⁻¹), with highest emissions occurring in the first 2 months after establishment. Once established, the turf grass lawn presented relatively low N₂O emissions for the rest of the year, even after fertilisation and rain events. Soil moisture was significantly higher, and mineralised N accumulated in the fallow plots, resulting in the highest N₂O emissions (2.8 kg N₂O ha⁻¹ yr⁻¹) and significant nitrate (NO₃⁻) losses, with up to 63 kg N ha⁻¹ lost from a single rain event due to reduced plant cover removal. The study concludes that urbanisation processes creating peri-urban ecosystems can greatly modify N cycling and increase the potential for losses in the form of N₂O and NO₃⁻.

1 Introduction

Global urbanisation processes are becoming increasingly important in terms of global warming and ecosystem functionality. Urban populations worldwide have not only exceeded rural populations but are also predicted to account for all future population growth (United Nations, 2008). Urban sprawl and increasing population densities are causing severe land use changes from intact biomes and commercially focused agriculture into smaller residential properties with introduced species. This transition from rural to semi-rural, i.e. peri-urban, and urban environments is increasingly associated with development and construction processes and the extensive establishment of turf grass for residential backyards, public parks and sports grounds, and golf courses (IPCC, 2006). How these urbanisation processes influence ecosystem dynamics in biogeochemical cycling, and therefore contribute to ecosystem vulnerability and global warming, is only beginning to be understood.

The consequences of land use change from native vegetation to agriculture have been identified by several studies and include a loss in soil quality (structure and nutrient losses) and quantity (erosion), increased greenhouse gas (GHG) emissions, and a reduced potential for soil carbon (C) sequestration (Livesley et al., 2009; Grover et al., 2012). On the other hand, changing soils from agricultural to residential use in temperate climates has shown the potential to improve critical ecosystem services by (i) providing stormwater treatment, (ii) acting as a sink for atmospheric nitrogen (N), and (iii) sequestering C (Golubiewski, 2006; Raciti et al., 2011).

Studies on the impact of these land use changes on climate change are few but suggest that urbanisation will alter biogeochemical cycling of C and N and associated nutrient turnover (Grimm et al., 2008). These biogeochemi-

cal alterations induced by land use change interact with urban effects (Betts, 2007) such as the creation of heat islands through vegetation replacement and surface sealing, and they also increased local carbon dioxide concentration of over 500 ppm around cities compared to 390 ppm in natural environments (Pataki et al., 2007; IPCC, 2013). These changing local climatic conditions and their feedback effects onto natural ecosystems make peri-urban environments more vulnerable to nutrient losses and potential sources of GHG emissions. With peri-urban areas expanding worldwide it is most likely that these changing local climates will increasingly have an impact on global climate change, making the examination of GHG emissions from peri-urban land uses all the more urgent.

Nitrous oxide (N_2O), along with carbon dioxide (CO_2) and methane (CH_4), is one of the major greenhouse gases with a global warming potential (GWP) nearly 300 times that of CO_2 (IPCC, 2013). Nitrous oxide is produced principally by microorganisms during nitrification and denitrification processes from mineral N (NH_4^+ and NO_3^-) in the soil. The production of N_2O is influenced by a number of soil parameters including substrate availability; temperature; and the availability of oxygen, which is dependent on water content and texture of the soil (Rowlings et al., 2015). With predicted climatic changes, Australia's ancient and fragile soils will most likely be affected in their balance between GHG gas emissions and consumption (Baldock et al., 2012). Management practices such as fertilisation and irrigation enhance N_2O production in the soil by increasing the mineral N content and limiting the oxygen availability (Scheer et al., 2008; Rowlings et al., 2013). Turf grass is the most highly managed land use of peri-urban environments in terms of fertilisation, irrigation, and frequent mowing, and it therefore has a high potential for N_2O emissions.

Research on urban and peri-urban areas in temperate zones suggests that urbanisation can result in emissions comparable to agriculture, with the intensive management of these peri-urban areas expanding rapidly worldwide (Milesi et al., 2005; Groffman and Pouyat, 2009). More than half the world's 7.2 billion population currently occupies 2.4 % of the global terrestrial land surface in urban areas (Potere and Schneider, 2007; United Nations, 2013). While peri-urban environments are often considered too small to be of consequence, the rapid growth of peri-urban areas has resulted in over $160\,000\text{ km}^{-2}$ being converted to turf grass lawn in the USA alone, 3 times more than any other irrigated crop in the country (Milesi et al., 2005). In Australia about 60 % of all anthropogenic N_2O emissions come from cropped and grazed soils (AGO, 2010), and the first GHG estimations from turf grass establishment support the emission intensity reported from temperate zones (van Delden et al., 2016). The area under turf grass is consistently growing in Australia with up to 17 320 ha in turf sales with an approximate gross value production of AUD 240 million per annum (ABS, 2012; Turf Australia, 2012). Detailed estimates of turf grass cover, how-

ever, currently do not exist for the Australian continent and other subtropical regions like South-east Asia, China, India, or Mexico. Urbanisation is currently neglected in modelled IPCC climate scenarios, mainly due to limited data on C and N processes in urban and peri-urban environments (IPCC, 2006, 2013).

This study therefore aims to identify the impact of those land use changes associated with urbanisation on annual N_2O emissions and their driving parameters in subtropical peri-urban environments. Following a short-term (80-day) GHG sampling campaign focussing on lawn establishment (van Delden et al., 2016), a fully automated closed static chamber system was used to continuously monitor N_2O fluxes together with soil biogeochemical processes over a full year to determine the seasonal impact of construction work, fallows, and conversion from extensively grazed pasture to turf grass lawn. This study's high-resolution flux measurements and supporting soil N mineralisation illustrate the vulnerability of ecosystems to urbanisation processes and the potential impact on N cycling and N_2O emissions.

2 Materials and method

2.1 Site description

The study was conducted at the Samford Ecological Research Facility (SERF) in the Samford Valley, 20 km from Brisbane in South East Queensland (SEQ), Australia. The Samford Valley covers an area of approximately 82 km^2 and is surrounded by mountains to the north, west, and south. Mostly cleared in the early 1900s, the valley was developed in the 1960s for dairy and beef cattle as well as intensive agriculture including banana and pineapple. Samford's population density has increased rapidly, almost doubling from 1996 to 2006, causing land use change from predominately rural to residential properties (Moreton Bay Regional Council, 2011). As a result, SERF contains the last remnant forest of the valley floor. The valley is influenced by a humid subtropical climate with seasonal summer rain. The long-term mean annual precipitation is 1110 mm with mean annual minimum and maximum temperatures of 13 and 25.6°C respectively (BOM, 2015). The soil at the experimental site is characterised by a strong texture contrast between the A and B horizon and is classified as brown Chromosol according to the Australian soil classification (Isbell, 2002) and Planosol according to the World Reference Base (WRB, 2015).

2.2 Experimental design

This study examined the impact of land use change from a native forest to well-established pasture, turf grass lawn, and fallow soil without plant cover using the same sampling campaign setup as van Delden et al. (2016). Each land use treatment included three replicated plots, 2 m wide by 10 m long and separated by 0.5 m of pasture as a buffer zone. The turf

grass lawn and fallow treatments were established within the well-established pasture to create a randomised plot design, 50 m from the native forest. The SERF native forest (dry sclerophyll eucalypt forest) was used as the baseline for historical land use and was unmanaged. The well-established *Chloris gayana* pasture represents rural development in the area and has been extensively grazed for the last 15 years. Livestock were excluded over the course of the study, and the pasture grass was slashed 5 times to ensure it did not exceed the maximum height of the GHG measurement chamber.

The turf grass lawn was established from the well-established pasture by removing 5 cm of topsoil with grass roots. The soil was rotary-hoed twice to a depth of 15 cm, and blue couch (*Digitaria didactyla*) turf rolls were planted with 50 kg N ha⁻¹ fertilisation (13 June 2013) to aid in establishment. Over the experimental year the turf grass lawn was fertilised twice more (26 October 13, 6 March 2014) with 50 kg N ha⁻¹ and irrigated, in all 150 kg N ha⁻¹ yr⁻¹ of Pro-lific Blue AN fertiliser (12.0 % nitrogen, 5.2 % phosphorus, 14.1 % potassium, 1.2 % magnesium) with two-thirds of the N content by mass in the ammonium form. The turf grass lawn was irrigated only enough to ensure its survival, with a total of 30 mm applied during drier months as well as after fertilisation. The turf grass was mowed with the clippings removed as soon as the grass grew to the maximum chamber height – once in spring, twice in summer, and twice in autumn – and kept free of weeds manually at all times. Fertilisation rates were based on half the local industry practice recommendation. Infrequent mowing represents the normal management for residential properties in this region and is normally in response to increased growth in the wetter and warmer summer months.

The fallow treatment simulated the impact of transitional processes such as construction work and plant cover replacement. In the fallow treatment, the grass cover was removed and the bare soil was rotary-hoed twice to a depth of 15 cm. The fallow treatment was kept free from plant cover over the full experimental year with a non-selective herbicide (Biac-tive, 360 g L⁻¹ glyphosate) and a broadleaf herbicide (Double Time, 340 g L⁻¹ MCPA + 80 g L⁻¹ dicamba). During the experimental year, high-resolution sub-daily N₂O flux measurements were combined with mineral N analysis and site-specific climate and soil moisture measurements.

2.3 N₂O flux measurements

Nitrous oxide fluxes were determined from mid-June 2013 to mid-June 2014 using an automated sampling system as detailed by Scheer et al. (2014), extending the turf grass establishment phase documented by van Delden et al. (2016) into a full measurement year. The pneumatically operated 50 cm × 50 cm × 15 cm high, clear acrylic glass chambers were secured to stainless steel bases, permanently inserted 10 cm into the ground. The chambers were moved each week between two bases per treatment plot to minimise the influ-

ence of the chamber microclimate, while measurements were analysed continuously. The chambers were connected to an automated sampling system and an in situ gas chromatograph (SRI GC8610, Torrance, CA, USA) equipped with a ⁶³Ni electron capture detector (ECD) for N₂O. One replicate chamber from each of the four treatments was closed for 1 h, and four headspace gas concentrations measured at 15 min intervals, followed a known calibration standard (0.5 ppm N₂O, Air Liquide, Houston, TX, USA). This process was repeated for the remaining two replicate chambers over a full cycle of 3 h, allowing eight flux measurements to be calculated per day for each of the 12 chambers.

2.4 Auxiliary measurements

Soil samples were taken fortnightly from all replicated treatment plots over the experimental year and divided into two depths (0–10, 10–20 cm). NH₄⁺ and NO₃⁻ were extracted from the soil using a 1 : 5 KCl solution with 20 g of fresh soil with additional soil moisture determination at 105 °C to identify the dry soil weight for the mineral N calculation as described by Carter and Gregorich (2007). The extract was analysed for mineral N (NH₄⁺ and NO₃⁻) with an AQ2+ discrete analyser (SEAL Analytical WI, USA). The net mineralisation rate was determined from differences in mineral N content between sampling dates (Hart et al., 1994). Soil moisture and temperature for each treatment were collected using a time-domain reflectometer (TDR) probe (HydroSense CD 620 CSA) and a PT100 probe (IMKO Germany). Soil moisture was then converted with the treatment-specific bulk density (BD) to water-filled pore space (WFPS). Soil samples were taken for site characterisation with a hydraulic soil corer to 1 m depth, air-dried, and sieved to 2 mm. Particle size analysis for soil texture as well as BD, pH, and electrical conductivity (EC) analysis was undertaken according to Carter and Gregorich (2007). The cation exchange capacity (CEC) was determined based on Rayment and Higginson (1992). Total C and N content of air-dried soil and plant material was determined by dry combustion (CNS-2000, LECO Corporation, St. Joseph, MI, USA) from ground samples.

2.5 Flux calculations and statistical analysis

Fluxes were calculated from the slope of the linear increase or decrease of the four concentrations measured over the closure time and corrected for chamber temperature and atmospheric pressure using the procedure outlined by Knowles and Singh (2003) and Scheer et al. (2014). The linear regression coefficient (r^2) was calculated and used as a quality check for fluxes above the detection limit to assure linearity of the gas concentration increase. Flux rates were discarded when r^2 was < 0.85 for N₂O fluxes (Scheer et al., 2013). Daily fluxes from the automated chambers were calculated by averaging sub-daily measurements for each chamber over

Table 1. SERF site characteristics.

Parameters											
Longitude						152°52'37.3" E					
Latitude						27°23'22.211" S					
Altitude						60 m					
Slope						2°					
Mean annual min temp.						13 °C ^a					
Mean annual max temp.						25.6 °C ^a					
Mean annual rain						1110 mm ^a					
Soil profile	Horizons ^b	Depths (cm)	Sand (%)	Silt (%)	Clay (%)	BD (g cm ⁻³)	pH	EC (μS)	CEC (meq + 100 g ⁻¹)	Total C (%)	Total N (%)
Pasture	A1	0–17	70	24	6	1.4	5.4	46	4.0	1.5	0.12
	A2	17–45	74	18	8	1.6	6.0	10	0.9	0.9	0.07
	B2	45–92	9	18	73	1.8	6.1	31	11.8	0.4	0.03
Forest	A1	0–20	75	18	7	1.4	5.5	29	2.8	1.8	0.14
	A2	20–47	78	15	7	1.5	5.6	30	0.9	1.1	0.08
	B2	47–70	41	7	52	1.7	5.6	30	11.8	0.2	0.02

^a Long-term means by Commonwealth Bureau of Meteorology, Australian Government (BOM, 2015). ^b According to the Australian soil classification.

Table 2. Seasonal and cumulative rain, number of rain events and seasonal and annual averages of minimum and maximum temperatures of the experimental year.

	Sum rain (mm)	Number of rain events*	Avg temperature (°C)	
			Min	Max
Winter	51.2	0	11.5	22.6
Spring	248.2	5	16.7	28.2
Summer	137.2	3	20.7	30.2
Autumn	303.2	3	17.6	27.5
	739.8	11	16.7	27.1

* Rain event if sum > 10 mm per day.

the 24 h period. The detection limit determined for the gas sampling system was $\pm 1.2 \text{ g N}_2\text{O ha}^{-1} \text{ d}^{-1}$. Gaps in the data set were filled by linear interpolation across missing days.

Statistical analysis was undertaken using SPSS Statistics 21.0 (IBM Corp., Armonk, NY). Non-normal distribution meant all cumulative data were log-transformed for analysis of variance (ANOVA) using Games–Howell as the post hoc test. Daily N_2O flux differences between treatments were interpreted by plotting 95 % confidence intervals using R studio. A significant difference of $p < 0.05$ between treatments was assumed in case the confidence intervals of all treatments were not overlapping. A Spearman's rho correlation analysis was used to examine relationships between gas fluxes, soil chemistry, soil moisture, and temperature. The significance value (p) is shown for each analysis, as well as the correlation coefficient (r) with its significance level ($p < 0.05^*$, $p < 0.01^{**}$).

3 Results


3.1 Site characteristics

The site received 740 mm of rain during the experimental year, substantially less than the long-term average (Table 1). Wet-season rainfall was delayed compared to the historic average, with less than half the rainfall in summer (December to February) compared to autumn (March to May; Table 2). Substantial out-of-season rain also fell in the spring with over 200 mm in November alone. Rainfall was highly episodic, with the highest daily rain event of 108.8 mm in March 2014. The mean annual minimum and maximum temperatures for the experimental year were 16.7 and 27.1 °C respectively, and light ground frost occurred twice in August. The turf grass and fallow treatment were established within the pasture and therefore share the same soil profile with its characteristics, except for bulk density in the A1 horizon, which changed after the turf grass establishment from 1.4 to 1.2 g cm⁻³. The CEC of the sandy topsoil was very low, and slightly higher in the A1 compared to the A2 horizon due to the higher soil organic matter as indicated by the total C and N content. Nutrient removal in turf grass clippings added up to 1.8 t C ha⁻¹ yr⁻¹ and 30 kg N ha⁻¹ yr⁻¹ lost from the system during the experiment year respectively. The turf's biomass production was approximately 6.3 kg C ha⁻¹ d⁻¹ and 0.13 kg N ha⁻¹ d⁻¹ in dry matter respectively but varied widely depending on fertilisation and available water with a maximum growth of 10.4 kg C ha⁻¹ d⁻¹.

Table 3. Annual mineral N averages of NH_4^+ -N and NO_3^- -N at 0–20 cm soil depth, WFPS, and daily maximum and average N_2O fluxes from all treatments with their cumulative annual fluxes over the experimental year with their standard error.

	NH_4^+ -N (kg ha^{-1})	NO_3^- -N (kg ha^{-1})	WFPS (%)	Max daily flux ($\text{g N}_2\text{O ha}^{-1} \text{d}^{-1}$)	Avg daily flux ($\text{g N}_2\text{O ha}^{-1} \text{d}^{-1}$)	Annual flux ($\text{kg N}_2\text{O ha}^{-1} \text{yr}^{-1}$)
Forest	$13.7^a \pm 1.2$	$3.9^a \pm 0.6$	23 ^a	8.1	$0.4^a \pm 0.1$	$0.1^a \pm 0.03$
Pasture	$17.4^b \pm 1.4$	$1.1^b \pm 0.3$	42 ^b	18.3	$0.6^a \pm 0.1$	$0.2^a \pm 0.2$
Turf grass	$21.9^{bc} \pm 2.4$	$8.9^c \pm 2.5$	43 ^b	83.0	$4.9^b \pm 0.6$	$1.8^b \pm 0.3$
Fallow	$26.0^c \pm 1.9$	$35.2^d \pm 5.6$	55 ^c	123.8	$7.7^b \pm 1.0$	$2.8^b \pm 1.0$

abcd Different letters indicate significant differences between treatments based on $p < 0.05$.

**Figure 1.** Annual soil NO_3^- (a) and NH_4^+ (b) content variations from forest, pasture, turf grass, and fallow averaged across replicates ($n = 3$) and summed for separate analysed soil depths of 0–10 and 10–20 cm with the climatic conditions (c) for the experimental year 2013/2014 as well as fertilisation and irrigation indication for the turf grass treatment.

3.2 Environmental parameters

The lowest WFPS recorded during the experimental year was 13 % in the forest, with the highest occurring in the pasture, which briefly reached saturation in March 2014 (Fig. 3). In all treatments, the lowest WFPS occurred in spring and sum-

mer with an average of 33 and 32 % respectively, together with the highest average maximum daily temperatures of 28 and 30 °C. While the highest seasonal WFPS for all treatments occurred in winter, the maximum WFPS occurred during autumn after the heavy rain in March 2014. The forest had significantly lower WFPS throughout the experimental year than all other treatments ($p < 0.01$, Table 3), while the fallow had significantly higher WFPS ($p < 0.01$). No significant difference in WFPS was observed between pasture and turf grass ($p > 0.05$), although during spring, summer, and autumn turf grass had lower minimum and maximum values than the pasture. The fallow had significantly higher and forest significantly lower WFPS than pasture and turf grass ($p < 0.01$) throughout the experimental year.

3.3 Temporal variability of mineral nitrogen

Averaged over the experimental year, the fallow treatment had the highest NH_4^+ and NO_3^- content across the top 20 cm soil profile, followed by turf grass, pasture, and forest (Table 3). These differences in mineral N were significant for all treatments ($p < 0.01$) except between pasture and forest ($p > 0.05$). The 0–10 cm depth had higher average total mineral N than the 10–20 cm depth for all treatments, with significant differences between all treatments ($p < 0.01$). In the 10–20 cm soil depth only the fallow had significantly higher total mineral N, NH_4^+ , and NO_3^- contents ($p < 0.01$). Soil NH_4^+ showed relatively little temporal variation and remained consistently above $3 \text{ kg NH}_4^+ \text{ ha}^{-1}$, while NO_3^- decreased substantially after rain events and fell below detection limit several times in all treatments but the fallow (Fig. 1).

Total mineral N in the forest ranged from 8 to $40.1 \text{ kg N ha}^{-1} 20 \text{ cm}^{-1}$ throughout the year with marginally higher mineral N content at 0–10 than 10–20 cm, with 9.7 and 8 kg N ha^{-1} respectively. Total mineral N in the pasture ranged from 5.1 to $42.4 \text{ kg N ha}^{-1} 20 \text{ cm}^{-1}$, with a comparable distribution in depth to the 0–10 and 10–20 cm forest soil with 10.7 and 7.8 kg N ha^{-1} respectively. Total mineral N in the turf grass soil ranged from 9.1 to $127.6 \text{ kg N ha}^{-1} 20 \text{ cm}^{-1}$. The turf grass had twice as much mineral N at 0–10 than 10–20 cm depths, with 20.7 and $10.1 \text{ kg N ha}^{-1}$ respectively. A short-term increase in both NH_4^+ and NO_3^- content




Figure 2. Daily N₂O flux averages (max eight fluxes per day for three replicates each) with standard errors ($N = 3$) over the experimental year 2013/2014 for forest (a), pasture (b), turf grass (c), and fallow (d) with the treatment-specific water-filled pore space (WFPS).

in the soil was evident after fertilisation in June, October, and March, which decreased to the background levels after approximately 1 month. Total mineral N contents in the fallow soil ranged from 19.7 to 160.7 kg N ha⁻¹ 20 cm⁻¹, with about two-thirds of the mineral N being located in the upper 10 cm. All main changes in the fallow's total mineral N content were caused by variations in NO₃⁻ rather than NH₄⁺. The NO₃⁻ content increased in the fallow until the major rain event in March, when it dropped from 95.5 to 32.8 kg N ha⁻¹ 20 cm⁻¹. From the linear increase in mineral N content within the upper 10 cm between January and March 2014 a soil mineralisation rate of 0.6 kg N ha⁻¹ d⁻¹ was estimated.

3.4 Temporal variation of N₂O fluxes

Daily N₂O fluxes across all treatments ranged from extended periods of close to 0 to over 123 g N₂O ha⁻¹ d⁻¹ from the fallow when WFPS was highest after heavy-rain events (Fig. 2).





Figure 3. Cumulative daily N₂O fluxes ($n = 3$) for forest, pasture, turf grass, and fallow with rainfall for the experimental year 2013/2014.

The N₂O fluxes from turf grass were more often significantly different on a daily basis than any other treatment with 76 days (21 %) of the experimental year; 80 % of this difference occurred in the first 2 months after establishment. This was followed by the forest with 65 days (18 %), fallow with 58 days (16 %), and pasture with 29 days (8 %). Daily N₂O fluxes from the forest soil showed no substantial temporal variation throughout the experimental year, with minor emission peaks up to 8.1 g N₂O ha⁻¹ d⁻¹ after large rain events (> 60 mm) in November and March. From September until October one of the two bases in one pasture replicate emitted substantially more N₂O than the other replicates; however, the exact cause of this is unknown. Without these spatially variable emissions, the annual flux would have been about 40 % lower and therefore comparable to the forest N loss of 0.09 kg N ha⁻¹ yr⁻¹. During the initial emission peak between June and August the daily average N₂O flux from the turf grass was 24 g N₂O ha⁻¹ d⁻¹, reaching a maximum of 83 g N₂O ha⁻¹ d⁻¹. Excluding this initial emission peak, daily N₂O fluxes from the turf grass averaged 1.2 g N₂O ha⁻¹ d⁻¹. The highest annual N₂O flux was measured in the fallow from only three large peaks over 19, 10, and 44 consecutive days after rain events, which together accounted for 85 % of the total N losses. Over a third of the significantly high daily N₂O fluxes in the fallow occurred from the heavy-rain event in March 2014.

Annual N₂O losses were highest in the fallow and turf grass treatments, totalling 1.78 and 1.15 kg N ha⁻¹ yr⁻¹ respectively, compared to the pasture and forest losses of 0.15 and 0.09 kg N ha⁻¹ yr⁻¹ ($p < 0.01$, Table 3). About 80 % of the annual N₂O losses in the turf occurred in the first 8 weeks after establishment (Fig. 3). Mineral N fertiliser input of 150 kg N ha⁻¹ yr⁻¹ and the yearly N₂O–N losses from the turf grass lawn corrected for background emissions (zero N fertilisation) from the pasture resulted in an emission factor (EF) of 0.7 % (Kroeze et al., 1997).

Table 4. Spearman's rho correlation coefficient between N₂O fluxes and mineral N, WFPS, and temperature for each treatment.

	N ₂ O			Mineral N	
	Mineral N	WFPS	Temperature	WFPS	Temperature
Forest	−0.39	0.61*	0.40*	0.07	−0.17
Pasture	0.49*	0.30*	−0.46*	0.39*	−0.41*
Turf grass	0.47*	0.78*	−0.45*	0.47*	−0.50*
Fallow	−0.0	0.43*	0.20*	−0.61*	0.72*

* Correlation coefficient significant with $p < 0.01$.**Figure 4.** Linear relationship of log-transformed N₂O emissions with mineral N content at 20 cm soil depth for each replicate of forest, pasture, turf grass, and fallow land use during the establishment phase (a) and the rest of the year (b), with the coefficient of determination R^2 .

3.5 Environmental parameters influencing N₂O fluxes

Mineral N contents in the forest and fallow soils were not significantly correlated with N₂O fluxes on a daily basis (Table 4). However, the linear regression shown in Fig. 4 identified a clear increase of N₂O emissions with increasing annual mineral N contents for all treatments during the establishment phase as well as during the rest of the year. This relationship is supported by the substantial N₂O emissions peaks from the fallow and simultaneous decrease in NO₃[−] after the two biggest rain events in November 2013 and March 2014 when WFPS reached above 70 %. The separate linear regression for all land uses with plant cover – i.e. forest, pasture, and turf grass – identified an even stronger relationship of mineral N and N₂O. Forest and turf grass N₂O fluxes were strongly, and fluxes from the pasture and fallow moderately, correlated with their WFPS. Temperature was moderately negatively correlated with N₂O fluxes as well as mineral N for pasture and turf grass. In the fallow temperature strongly affected mineral N contents but not N₂O fluxes. Mineral N in the fallow soil was strongly negatively correlated with its WFPS mostly because of the strong negative correlation of NO₃[−] with WFPS, with $r = -0.56^{**}$.

4 Discussion

This study combines the first high-frequency estimates of subtropical N₂O fluxes and annual mineral N cycling from dry sclerophyll forests, unfertilised pastures, and turf grass lawns: the most common land uses associated with urban and peri-urban environments. The lack of high-frequency field measurements in urban and peri-urban environments makes accurate assumptions and mitigation strategies difficult. Conventional gas sampling methods most likely result in an over- or underestimation of emissions, as the production and release of N₂O can differ in time (Mosier et al., 1998). This research gap, together with the strong temporal variability of subtropical heavy-rain events, underlines the importance of automated high-frequency measurements to capture representative soil–atmosphere gas exchange. The subtropical climatic zone represents an often-neglected area of research, despite the subtropics covering 3.26 M ha in Australia alone, as well as large areas on the North and South American continents, Africa, and Asia. Differences between other climates and the humid subtropics are the heavy summer rains leading to high soil moisture and temperatures favourable for high soil microbial activity. Therefore this study identified mineral N content and WFPS as the main parameters driving N₂O

production in the soil, while studies from temperate zones report temperature as the main driver (Butterbach-Bahl and Kiese, 2005; Fest et al., 2009).

4.1 Mineral N

Mineralised N in the form of NH_4^+ and NO_3^- determines the production and loss of N via N_2O and depends on climatic parameters like temperature as well as substrate and oxygen availability. Nutrient mineralisation is often faster in sandy soils, but the rapid infiltration and low nutrient holding capacity of the A horizon of the Chromosol decreased the highly mobile NO_3^- content substantially after heavy-rain events. This NO_3^- is not only lost for plant uptake but can pollute groundwater and open waterways, resulting in eutrophication. In this study mineral N contents from the forest and pasture treatments were driven by annual variation in temperature and moisture contents, whereas turf grass lawn and fallow were dominated by management. The negative correlation of temperature in the pasture and turf grass can most likely be explained by the higher plant productivity during the warmer summer and spring resulting in higher plant NO_3^- and water uptake with increasing temperatures, subsequently reducing soil moisture conditions.

Soil mineral N in the SERF forest was generally low and dominated by NH_4^+ , and while less seasonally variable throughout the year than NO_3^- , it still responded to rainfall. The overall mineral N reported from temperate eucalypt forest soils was double the annual SERF average of $17.6 \text{ kg N ha}^{-1}$, with up to $38.1 \text{ kg N ha}^{-1}$ reported (Fest et al., 2009, 2015; Livesley et al., 2009). However, the greater proportion of NO_3^- in the sandy SERF soil of 3.9 compared to 0.8 kg N ha^{-1} of temperate sandy forest soils (Livesley et al., 2009) indicates a higher mineral N availability in the subtropics. Average NO_3^- contents reported from other dry sclerophyll forest are even lower at $0.02 \text{ kg N ha}^{-1}$ (Fest et al., 2015). The higher N availability is most likely due to faster soil organic matter turnover in the subtropical climate with higher temperatures in combination with the main annual rainfall. While in temperate summers it is mostly dry during the periods of high temperatures, which limits microbial activity, the humid summers in SEQ accelerate not only N turnover but also water and N uptake by plants, therefore reducing potential N losses. Subtropical rainforests, on the other hand, present with up to 6-times-higher mineral N contents ($> 97 \text{ kg N ha}^{-1}$) than the SERF soil, suggesting a lower N turnover associated with the low net primary productivity (NPP) of the dry sclerophyll forests (Rowlings et al., 2012). Overall $\text{NH}_4^+:\text{NO}_3^-$ ratios from Australian forests indicate higher NO_3^- availability in subtropical forest soils (3–4) than in temperate zones (28–125; Livesley et al., 2009; Rowlings et al., 2012; Fest et al., 2015). These differences in N availability suggest that N cycling in forest soils is mainly regulated by the climate as opposed to soil type and NPP.

Ammonium was the dominant mineral N form in the SERF pasture, similar to the forest and in line with other subtropical pastures in Australia (Rowlings et al., 2015). The SERF soil reflects the overall minor annual variability of NH_4^+ compared to NO_3^- across most climates in Australia. The overall mineral N content at the SERF pasture soil was at the lower end of the reported values from both temperate and subtropical pastures, which is most likely explained by the lower clay contents at the site which fixes NH_4^+ and higher N inputs by legumes (Livesley et al., 2009; Rowlings et al., 2015). For example, in other extensively grazed subtropical pastures NH_4^+ annual values did not drop below 55 kg N ha^{-1} (Rowlings et al., 2015), 3 times higher than the SERF annual NH_4^+ average. While NH_4^+ at SERF is comparable to temperate Australian pastures, NO_3^- in the SERF pasture soil is at the lower end (Livesley et al., 2009). This indicates an efficient system from tied-up N in organic material to the plant uptake of NO_3^- , which supports the hypothesis of an efficient N cycle within well-established land use.

Annual mineral N variations in the SERF turf grass were mainly controlled by the fertilisation events but rapidly fell back to background levels after each application. The fertiliser mineral N peak was particularly emphasised after the first application, where soil NO_3^- was more than twice as much as after subsequent fertilisation events. This is possibly due to the undeveloped root system and therefore less N uptake as well as additional plant available N in the added turf grass rolls. These NO_3^- peaks together with irrigation, which is particularly needed during turf grass establishment, imply a high N leaching potential, with up to $80 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ reported in other Australian sandy soils (Barton et al., 2006). With the high potential of heavy-rain events in the subtropics, fertiliser rates and timing need to be considered carefully to avoid excessive N losses in the form of NO_3^- displacement.

The fallow soil had the highest WFPS content throughout the year due to plant cover removal and therefore no further water uptake by the roots, creating favourable conditions for soil mineralisation and subsequent losses (Robertson and Groffman, 2007). The moist conditions together with the temperatures of the warmer season resulted in accelerated N turnover, and without plant uptake substantial amounts of NO_3^- accumulated in the soil. Despite the fact that mineral N in the fallow soil never dropped back to 0, substantial amounts of NO_3^- were lost from the topsoil after heavy-rain events, not only as N_2O emissions but also through NO_3^- displacement into deeper soil layers. The low CEC of the sandy topsoil highlights the minor nutrient holding capacity of these soils. These potential N losses after heavy-rain events demonstrate the significant impact of plant cover removal and soil disturbance in peri-urban ecosystems.

4.2 N_2O fluxes

The study illustrates that land use change associated with urbanisation can significantly alter soil N turnover, resulting in

elevated soil N_2O emissions and increased N losses from the soil. During the experimental year of this study, autumn was the wettest season and therefore had the highest N_2O emissions from all treatments but with different intensity from the different land use systems. Soil N_2O emissions were significantly different between the investigated land use systems with the temporal variations in daily N_2O fluxes and primarily controlled by WFPS. However, the linear increase of N_2O emissions with increasing NO_3^- content in the soil may be the result of higher denitrification than nitrification rates in the SERF soil. The high surface sand content of the Chromosol, combined with the moderate slope, prevents excessive water logging over long periods of time, which limits N_2O gaseous losses from denitrification in saturated soil conditions.

The daily N_2O average of $0.4 \text{ g N}_2\text{O ha}^{-1} \text{ d}^{-1}$ from this study's subtropical dry sclerophyll forest is lower than the averages of $<1.2 \text{ g N}_2\text{O ha}^{-1} \text{ d}^{-1}$ reported from temperate Australian dry sclerophyll forests (Fest et al., 2009; Livesley et al., 2009). This might be explained by the overall low total C and N and the below-average rainfall during the experimental year. Considering the positive correlation of N_2O emissions and NO_3^- content in the soil, it was expected that the higher NO_3^- availability in the SERF forest compared to the temperate dry sclerophyll forest also causes higher N_2O emissions. The low WFPS, which was $>40\%$ for most of the year, inhibited denitrification processes and therefore caused lower N_2O emissions than in the temperate zones as well as increased NO_3^- uptake during the humid subtropical summer. This efficient N cycling together with the low NPP of the dry sclerophyll forest and low clay content at SERF also cause lower N_2O losses compared to subtropical rainforests (Rowlings et al., 2012). This study supports the general hypothesis that forest soils are minor contributors to the global N_2O budget, although other N_2O emission studies of Australian forest soils provide only a limited comparison of temporal N_2O variability due to infrequent or short-term measurements (Fest et al., 2009, 2015; Page et al., 2011).

The annual N_2O emissions from the SERF pasture are comparable to other reported extensive pastures across Australia ($1\text{--}2 \text{ kg N ha}^{-1} \text{ yr}^{-1}$) but substantially lower than unfertilised pasture in the Northern Hemisphere (Dalal et al., 2003). Annual emissions from other studies on subtropical Australian pastures have been reported to be up to $3.4 \text{ kg N}_2\text{O ha}^{-1} \text{ yr}^{-1}$ and highly inter-annually variable depending on rainfall (Rowlings et al., 2015). This exceeded the annual N_2O emissions at SERF by nearly 17 times, which may have been limited by the dry year and high sand content.

The first N_2O emission peak after the turf grass's establishment caused the majority of the annual N_2O emissions and was not repeated after two additional fertilisation events. This initial N_2O peak can be explained by the underdeveloped root system and consequently a reduced NO_3^- uptake by the turf grass, which together with the irrigation stimulated nitrification and denitrification and consequential N_2O emis-

sions. The high N demand from the highly productive turf grass later on results in the immediate uptake of mineral N and therefore minor N_2O emissions. The annual N_2O emissions from the SERF's turf grass are more than double the N_2O emissions from extensive Australian pastures reported in the literature (Dalal et al., 2003). The SERF turf grass lawn emitted on average about 3.3-times-more N_2O daily during the experimental year than native pasture from the temperate zones, but only half of the reported values for urban turf grass in the USA which were comparable to intensive agriculture (Kaye et al., 2004). However, compared to Australian intensively managed pastures, N_2O emissions from the SERF turf grass were 50 % lower (Scheer et al., 2011). Differences between reported values and the SERF turf grass are most likely explained by differences in texture and the total N content in the SERF soil being nearly 4 times lower. Reported EFs from temperate pastures also vary substantially between experimental years due to differences in received rainfall (Jones et al., 2005). It could therefore be expected that the SERF's turf grass EF will increase in wetter years. However, in subtropical systems it has been proven that the total amount of annual rainfall received is not as decisive for annual N_2O emissions as rainfall patterns and intensities (Rowlings et al., 2015). These differences between temperate and subtropical N cycling make short-term N_2O flux measurements difficult to compare, and further investigation is needed in the global subtropics.

Significantly higher NO_3^- contents occurred 3 months after plant cover removal in the fallow soil, but only during the warm and wet summer season were substantial N_2O emissions observed. The two significant N_2O emission peaks from the fallow were most likely caused by denitrification processes from the accumulated NO_3^- and soil moisture conditions after major rain events. These N_2O emission peaks mirror the NO_3^- decrease from the soil after those rain events but cannot completely account for it, suggesting that most NO_3^- was leached below 20 cm soil depths or lost via other gases such as N_2 . All other treatments, including the fertilised turf grass, prevented potential N_2O production in the soil by rapid NO_3^- uptake from plants. Therefore, plant cover removal makes ecosystems undergoing land use change most vulnerable to substantial N losses under humid subtropical climate conditions.

4.3 Effect of land use change associated with urbanisation

This study determined that urbanisation-related land use change results in an accumulation of NO_3^- in fallow topsoil and elevated N_2O emissions, mainly after heavy-rain events. The results presented here verify that subtropical N_2O emissions positively correlate with mineral N content in the soil and therefore indicate that land use change increases N_2O emissions from the soil, especially after plant cover removal and establishment of fertilised turf grass lawn. The annual

variation in daily N_2O fluxes confirms that, despite soil moisture as the strongest climatic parameter influencing N_2O emissions, the individual land use is the main influence on the soil–atmosphere gas exchange. Extended periods of fallow soil in particular should be avoided during urbanisation processes as bare soil is highly vulnerable to N losses due to plant cover removal. Turf grass lawn, as a fertilised and highly managed land cover, leads to significantly changed soil conditions compared to the forest and pasture land use types. However, this turf grass lawn in the subtropical climate of SEQ has lower emissions against expectations based on the high-emission findings from temperate zones (Kaye et al., 2004; Tratalos et al., 2007; Grimm et al., 2008). Substantial N_2O emissions were only observed within the first 2 months after turf grass establishment, while over the remaining 10 months only minor fluxes occurred even after further fertilisation events. While N_2O emissions from the turf grass were reduced substantially over time, emissions from the fallow increased with time due to more available NO_3^- . Therefore, the N_2O emissions of well-established turf grass lawns need to be considered separately to their production and establishment phase as well as potential N losses from fallow land targeted for the entire duration of land use change, which should be kept as short as possible (Barton et al., 2006; van Delden et al., 2016).

Research from temperate zones suggests a C sequestration potential from the higher productivity of turf grass lawns (Golubiewski, 2006; Lorenz and Lal, 2009; Raciti et al., 2011). Others argue that the positive effect of C sequestration can easily be offset by the high N demand together with irrigation, resulting in increased N_2O emissions and overall nutrient losses caused by management practices like mowing and clipping removal (Conant et al., 2005; Wang et al., 2014). Australian ecosystems with highly weathered soils, however, are generally low in nutrient stocks and often limited in their C sequestration potential (Livesley et al., 2009). The SERF turf grass, however, presented relatively low N_2O emissions when excluding the establishment phase, which implies the potential to balance emissions with C sequestration. A full life cycle assessment needs to determine if turf grass lawn in the subtropics is increasing or decreasing the GWP of peri-urban environments by balancing C sequestration and GHG emissions, not only from the soil but also through the production; distribution; and use of fertiliser, fuel, and chemicals (Selhorst and Lal, 2011).

5 Conclusions

This study provides evidence that land use change associated with urbanisation accelerates N turnover and increases N_2O emissions from soils by presenting the first high-temporal-frequency data set on peri-urban soils in the subtropics for a full year after land use change. These findings demonstrate that GHG emissions from peri-urban areas should be in-

cluded into future IPCC climate change scenarios, and rural-to-urban land development guidelines need to be established for GHG emission mitigation. Three main factors need to be considered to target N_2O losses from soils during land use change associated with urbanisation: (i) previous land use, (ii) duration of development process, and (iii) new land use purpose that it is being changed into, i.e. public or private. The dry sclerophyll forest in this study supports the general hypothesis that forest soils are low N_2O emitters, contrary to expectation that the humid subtropical summer conditions would increase emissions compared to temperate forest soils. The accumulation of NO_3^- in fallow soil increases the potential for N_2O emissions, which may be amplified considering future predictions of rising temperatures and more frequent heavy-rain events worldwide. Increased fertiliser application may be required to compensate for these N losses after land use change to keep land uses, such as turf grass, highly productive while altering N cycling in peri-urban environments. The outcomes of this study highlight the substantial NO_3^- accumulation in soils during land use change, which consequently increases N_2O emissions and should be accounted for in global climate forecasts as urbanisation processes are predicted to increase worldwide with increasing population growth.

6 Data availability

The data set “Greenhouse gas emissions from peri-urban land use at SERF, SEQ, 2013–2015” can be found online at the N_2O network at <http://www.N2O.net.au/knb/metacat/vandelden.3.3/html>.

Acknowledgements. This study was undertaken at the Samford Ecological Research Facility (SERF) one of the supersites in the Terrestrial Ecosystem Research Network (TERN). The study was supported by the Institute for Future Environments (IFE) of the Queensland University of Technology (QUT).

Edited by: A. Ibrom

Reviewed by: two anonymous referees

References

- ABS: 7106.0 – Australian Farming in Brief, edited by: Australian Bureau of Statistics, Canberra, 2012.
- AGO: National Inventory Report 2008, National Greenhouse Account, Australian Greenhouse Office, Commonwealth of Australia, Canberra, 2010.
- Baldock, J. A., Wheeler, I., McKenzie, N., and McBratney, A.: Soils and climate change: potential impacts on carbon stocks and greenhouse gas emissions, and future research for Australian agriculture, *Crop Pasture Sci.*, 63 3, 269–283, doi:10.1071/cp11170, 2012.

- Barton, L., Wan, G. G. Y., and Colmer, T. D.: Turfgrass (*Cynodon dactylon* L.) sod production on sandy soils, II. Effects of irrigation and fertiliser regimes on N leaching, *Plant Soil*, 284, 147–164, 2006.
- Betts, R.: Implications of land ecosystem-atmosphere interactions for strategies for climate change adaptation and mitigation, *Tellus B*, 59, 602–615, 2007.
- BOM: Commonwealth Bureau of Meteorology, Australian Government, 2015.
- Butterbach-Bahl, K. and Kiese, R.: Significance of forests as sources for N_2O and NO , in: *Tree species effects on soils: implications for global change*, Springer, 173–191, 2005.
- Carter, R. and Gregorich, E. G.: *Soil Sampling and Methods of Analysis*, Second Edition, edited by: Canadian Society of Soil Science, Taylor & Francis, 2007.
- Conant, R. T., Paustian, K., Del Grosso, S. J., and Parton, W. J.: Nitrogen pools and fluxes in grassland soils sequestering carbon, *Nutr. Cycl. Agroecosys.*, 71, 239–248, doi:10.1007/s10705-004-5085-z, 2005.
- Dalal, R. C., Wang, W. J., Robertson, G. P., and Parton, W. J.: Nitrous oxide emission from Australian agricultural lands and mitigation options: a review, *Aust. J. Soil Res.*, 41, 165–195, doi:10.1071/sr02064, 2003.
- Fest, B. J., Livesley, S. J., Drösler, M., van Gorsel, E., and Arndt, S. K.: Soil-atmosphere greenhouse gas exchange in a cool, temperate *Eucalyptus delegatensis* forest in south-eastern Australia, *Agr. Forest Meteorol.*, 149, 393–406, doi:10.1016/j.agrformet.2008.09.007, 2009.
- Fest, B. J., Livesley, S. J., von Fischer, J. C., and Arndt, S. K.: Repeated fuel reduction burns have little long-term impact on soil greenhouse gas exchange in a dry sclerophyll eucalypt forest, *Agr. Forest Meteorol.*, 201, 17–25, doi:10.1016/j.agrformet.2014.11.006, 2015.
- Golubiewski, N. E.: Urbanization increases grassland carbon pools: Effects of landscaping in Colorado's front range, *Ecol. Appl.*, 16, 555–571, doi:10.1890/1051-0761(2006)016[0555:uigepe]2.0.co;2, 2006.
- Grimm, N. B., Foster, D., Groffman, P., Grove, J. M., Hopkinson, C. S., Nadelhoffer, K. J., Pataki, D. E., and Peters, D. P. C.: The changing landscape: ecosystem responses to urbanization and pollution across climatic and societal gradients, *Front. Ecol. Environ.*, 6, 264–272, 2008.
- Groffman, P. M. and Pouyat, R. V.: Methane Uptake in Urban Forests and Lawns, *Environ. Sci. Technol.*, 43, 5229–5235, doi:10.1021/es803720h, 2009.
- Grover, S. P. P., Livesley, S. J., Hutley, L. B., Jamali, H., Fest, B., Beringer, J., Butterbach-Bahl, K., and Arndt, S. K.: Land use change and the impact on greenhouse gas exchange in north Australian savanna soils, *Biogeosciences*, 9, 423–437, doi:10.5194/bg-9-423-2012, 2012.
- Hart, S. C., Stark, J. M., Davidson, E. A., and Firestone, M. K.: Nitrogen mineralization, immobilization, and nitrification, *Methods of Soil Analysis: Part 2 – Microbiological and Biochemical Properties*, 985–1018, 1994.
- IPCC: Chapter 8: Settlements in Guidelines for National Greenhouse Gas Inventories: Agriculture, Forestry and Other Land Use, edited by: Ginzo, H. D., Jenkins, J. C., Ogle, S. M., Verchot, L. V., Handa, M., and Tsunekawa, A., 4, available at: http://www.ipcc-nggip.iges.or.jp/public/2006gl/pdf/4_Volume4/V4_08_Ch8_Settlements.pdf (last access: 1 January 2016), 2006.
- IPCC: Climate Change 2013: The Physical Science Basis, Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Stocker, T. F., Qin, Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp., doi:10.1017/CBO9781107415324, 2013.
- Isbell, R.: The Australian soil classification, CSIRO publishing: Collingwood, 4, 2002.
- Jones, S. K., Rees, R. M., Skiba, U. M., and Ball, B. C.: Greenhouse gas emissions from a managed grassland, *Global Planet. Change*, 47, 201–211, doi:10.1016/j.gloplacha.2004.10.011, 2005.
- Kaye, J. P., Burke, I. C., Mosier, A. R., and Guerschman, J. P.: Methane and nitrous oxide fluxes from urban soils to the atmosphere, *Ecol. Appl.*, 14, 975–981, doi:10.1890/03-5115, 2004.
- Kaye, J. P., Groffman, P. M., Grimm, N. B., Baker, L. A., and Pouyat, R. V.: A distinct urban biogeochemistry?, *Trends Ecol. Evol.*, 21, 192–199, 2006.
- Knowles, T. A. and Singh, B.: Carbon storage in cotton soils of northern New South Wales, *Aust. J. Soil Res.*, 41, 889–903, doi:10.1071/sr02023, 2003.
- Kroeze, C., Mosier, A., Nevison, C., Oenema, O., Seitzinger, S., and van Cleemput, O.: Revised 1996 IPCC guidelines for national greenhouse gas inventories: Intergovernmental Panel on Climate Change, 1997.
- Livesley, S. J., Kiese, R., Miehe, Weston, C. J., Butterbach-Bahl, K., and Arndt, S. K.: Soil-atmosphere exchange of greenhouse gases in a *Eucalyptus marginata* woodland, a clover-grass pasture, and *Pinus radiata* and *Eucalyptus globulus* plantations, *Glob. Change Biol.*, 15, 425–440, doi:10.1111/j.1365-2486.2008.01759.x, 2009.
- Lorenz, K. and Lal, R.: Biogeochemical C and N cycles in urban soils, *Environ. Int.*, 35, 1–8, 2009.
- Milesi, C., Running, S. W., Elvidge, C. D., Dietz, J. B., Tuttle, B. T., and Nemani, R. R.: Mapping and modeling the biogeochemical cycling of turf grasses in the United States, *Environ. Manage.*, 36, 426–438, doi:10.1007/s00267-004-0316-2, 2005.
- Moreton Bay Regional Council: Community profile, Samford valley area, <http://profile.id.com.au/Default.aspx?id=311&pg=101&gid=560&type=enum> (last access: 1 January 2016), 2011.
- Mosier, A., Kroeze, C., Nevison, C., Oenema, O., Seitzinger, S., and Van Cleemput, O.: Closing the global N_2O budget: nitrous oxide emissions through the agricultural nitrogen cycle, *Nutr. Cycl. Agroecosys.*, 52, 225–248, 1998.
- Page, K. L., Dalal, R. C., and Raison, R. J.: The impact of harvesting native forests on vegetation and soil C stocks, and soil CO_2 , N_2O and CH_4 fluxes, *Aust. J. Bot.*, 59, 653–668, doi:10.1071/bt11207, 2011.
- Pataki, D. E., Xu, T., Luo, Y. Q., and Ehleringer, J. R.: Inferring biogenic and anthropogenic carbon dioxide sources across an urban to rural gradient, *Oecologia*, 152, 307–322, 2007.
- Potere, D. and Schneider, A.: A critical look at representations of urban areas in global maps, *GeoJournal*, 69, 55–80, 2007.
- Raciti, S. M., Groffman, P. M., Jenkins, J. C., Pouyat, R. V., Fahey, T. J., Pickett, S. T. A., and Cadenasso, M. L.: Accumulation of carbon and nitrogen in residential soils with different land-use

- histories, *Ecosystems*, 14, 287–297, doi:10.1007/s10021-010-9409-3, 2011a.
- Rayment, G. E. and Higginson F. R.: Australian laboratory handbook of soil and water chemical methods: Inkata Press Pty Ltd, 1992.
- Robertson, G. P. and Groffman, P.: Nitrogen transformations, edited by: Paul, E. A. C. F., *Soil Microbiology, Ecology, and Biochemistry*, Oxford, UK, Elsevier Academic Press, 341–364, 2007.
- Rowlings, D. W., Grace, P. R., Kiese, R., and Weier, K. L.: Environmental factors controlling temporal and spatial variability in the soil-atmosphere exchange of CO₂, CH₄ and N₂O from an Australian subtropical rainforest, *Glob. Change Biol.*, 18, 726–738, doi:10.1111/j.1365-2486.2011.02563.x, 2012.
- Rowlings, D. W., Grace, P. R., Scheer, C., and Kiese, R.: Influence of nitrogen fertiliser application and timing on greenhouse gas emissions from a lychee (*Litchi chinensis*) orchard in humid subtropical Australia, *Agr. Ecosyst. Environ.*, 179, 168–178, 2013.
- Rowlings, D. W., Grace, P. R., Scheer, C., and Liu, S.: Rainfall variability drives interannual variation in N₂O emissions from a humid, subtropical pasture, *Sci. Total Environ.*, 512, 8–18, 2015.
- Scheer, C., Wassmann, R., Kienzler, K., Ibragimov, N., and Eschanov, R.: Nitrous oxide emissions from fertilized, irrigated cotton (*Gossypium hirsutum* L.) in the Aral Sea Basin, Uzbekistan: Influence of nitrogen applications and irrigation practices, *Soil Biol. Biochem.*, 40, 290–301, 2008.
- Scheer, C., Grace, P. R., Rowlings, D. W., Kimber, S., and Van Zwieten, L.: Effect of biochar amendment on the soil-atmosphere exchange of greenhouse gases from an intensive subtropical pasture in northern New South Wales, Australia, *Plant Soil*, 345, 47–58, doi:10.1007/s11104-011-0759-1, 2011.
- Scheer, C., Grace, P. R., Rowlings, D. W., and Payero J.: Soil N₂O and CO₂ emissions from cotton in Australia under varying irrigation management, *Nutr. Cycl. Agroecosys.*, 95, 43–56, doi:10.1007/s10705-012-9547-4, 2013.
- Scheer, C., Rowlings, D. W., Firrel, M., Deuter, P., Morris, S., and Grace, P. R.: Impact of nitrification inhibitor (DMPP) on soil nitrous oxide emissions from an intensive broccoli production system in sub-tropical Australia, *Soil Biol. Biochem.*, 77, 243–251, 2014.
- Selhorst, A. L. and Lal, R.: Carbon budgeting in golf course soils of Central Ohio, *Urban Ecosystems*, 14, 771–781, doi:10.1007/s11252-011-0168-5, 2011.
- Tratalos, J., Fuller, R. A., Warren, P. H., Davies, R. G., and Gaston, K. J.: Urban form, biodiversity potential and ecosystem services, *Landscape Urban Planning* 83, 308–317, 2007.
- Turf Australia: Newsletter 2012, available at: <https://www.turfaustralia.com.au/> (last access: 1 January 2016), 2012.
- United Nations: World urbanization prospects: The 2007 Revision, edited by: Economics and Social Affairs, United Nations (NY): United Nations Population Division, 2008.
- United Nations: World Population Prospects: The 2012 Revision, Highlights and Advance Tables, edited by Economics and Social Affairs, United Nations (NY), United Nations Population Division, 2013.
- van Delden, L.: Greenhouse gas emissions from peri-urban land use at SERF, SEQ, Australia, 2013–2015, Research Data Australia: N₂O Network, available at: <http://www.N2O.net.au/knb/metacat/vandelden.3.3/html>, 2015.
- van Delden, L., Larsen, E., Rowlings, D., Scheer, C., and Grace, P.: Establishing turf grass increases soil greenhouse gas emissions in peri-urban environments, *Urban Ecosystems*, 19, 749–762, 2016.
- Wang, Y., Tu, C., Li, C., Tredway, L., Lee, D., Snell, M., Zhang, X., and Hu, S.: Turfgrass Management Duration and Intensities Influence Soil Microbial Dynamics and Carbon Sequestration., *Int. J. Agr. Biol.*, 16, 139–145, 2014.
- WRB: IUSS Working Group, World Reference Base for Soil Resources 2014, update 2015, World Soil Resources Reports No. 106, International soil classification system for naming soils and creating legends for soil maps, FAO, Rome, 2015.