

Supplement of Biogeosciences, 14, 2133–2149, 2017
<http://www.biogeosciences.net/14/2133/2017/>
doi:10.5194/bg-14-2133-2017-supplement
© Author(s) 2017. CC Attribution 3.0 License.



Biogeosciences  Open Access

Supplement of

The oxic degradation of sedimentary organic matter 1400 Ma constrains atmospheric oxygen levels

Shuichang Zhang et al.

Correspondence to: Shuichang Zhang (sczhang@petrochina.com.cn)

The copyright of individual parts of the supplement might differ from the CC-BY 3.0 licence.

The oxic degradation of sedimentary organic matter 1.4 Ga constrains atmospheric oxygen levels

Supplementary Information

Table S1. Repeated analysis of standards by ICP-MS for V, Mo and U; by XRF for Al and Fe and by HHXRF for Fe.

No	Sample	V (ppm)	Mo (ppm)	U (ppm)	No	Sample	Al (wt%)	Fe (wt%)	No	Sample	HHXRF Fe (wt%)
1	GBW07309	99.6	0.62	2.48	1	GBW07104	8.52	2.62	1	PACS -3	4.08
2	GBW07309	98.4	0.63	2.51	2	GBW07104	8.54	2.61	2	PACS -3	4.07
3	GBW07309	93.3	0.67	2.57	3	GBW07104	8.50	2.61	3	PACS -3	3.99
4	GBW07309	93.7	0.62	2.68	4	GBW07104	8.51	2.61	4	PACS -3	4.07
5	GBW07309	99.9	0.61	2.50	5	GBW07104	8.54	2.61	5	PACS -3	3.93
6	GBW07309	94.7	0.66	2.70	6	GBW07104	8.54	2.62	6	PACS -3	3.95
7	GBW07309	96.5	0.67	2.48	7	GBW07104	8.54	2.62	7	PACS -3	3.98
8	GBW07309	95.3	0.61	2.57	8	GBW07104	8.54	2.62	8	PACS -3	3.96
9	GBW07309	93.9	0.64	2.56	9	GBW07104	8.55	2.62	9	PACS -3	3.95
10	GBW07309	101.0	0.62	2.73	10	GBW07104	8.54	2.63	10	PACS -3	3.90
					11	GBW07104	8.54	2.61	11	PACS -3	3.87
									12	PACS -3	3.95
									13	PACS -3	3.94
									14	PACS -3	3.95
average		96.6	0.64	2.58			8.53	2.62			3.97
std dev		2.9	0.02	0.09			0.02	0.01			0.06
Accepted		97.0	0.64	2.60			8.57	2.65			4.11
1	GBW07310	104.0	1.21	2.04	1	GBW07407	15.51	10.14	1	XML Z4	5.97
2	GBW07310	105.0	1.18	2.02	2	GBW07407	15.48	10.14	2	XML Z4	5.94
3	GBW07310	104.0	1.26	2.21	3	GBW07407	15.45	10.12	3	XML Z4	5.87
4	GBW07310	107.0	1.23	2.10					4	XML Z4	5.90

5	GBW07310	110.0	1.24	2.07
6	GBW07310	109.0	1.23	2.15
7	GBW07310	107.0	1.19	2.12
8	GBW07310	105.0	1.26	2.00
9	GBW07310	109.0	1.16	2.14
10	GBW07310	106.0	1.17	2.09

5	XML Z4	5.86
6	XML Z4	5.87
7	XML Z4	5.87
8	XML Z4	5.92
9	XML Z4	5.85
10	XML Z4	5.80
11	XML Z4	5.91

average	106.6	1.21	2.09	15.48	10.13	5.89
std dev	2.2	0.04	0.06	0.03	0.01	0.05
Accepted	107.0	1.200	2.10	15.51	10.13	5.83

Table S2. Geochemical data and sediment description for outcrop samples.

Depth	Sediment	Org-C	$\delta^{13}\text{C}$	Al	Fe	Fe/Al	V	V/Al	Mo	Mo/Al	U	U/Al	HI
M		wt%	‰	wt%	wt%		ppm		ppm		ppm		mg/gTOC
0	black	1.14	-34.1	5.77	5.39	0.93	80	13.9	1.54	0.27	2.6	0.45	138
3.60	green	0.27	-31.7	5.57	3.54	0.64	60	10.8	0.62	0.11	2.0	0.37	111
4.60	gray	0.20	-32.9	6.76	2.95	0.44	74	10.9	0.67	0.10	2.8	0.41	100
6.10	black	2.25		6.90	3.26	0.47	112	16.2	2.2	0.32	3.0	0.43	318
7.60	black	3.32	-33.4	6.95	3.31	0.48	163	23.4	5.28	0.76	4.0	0.58	373
9.10	black	2.65		6.73	3.42	0.51	203	30.2	4.63	0.69	3.8	0.56	353
10.10	black	2.21	-33.4	6.72	3.27	0.49	126	18.8	3.32	0.49	3.1	0.46	313
11.10	gray	0.09	-33.3	7.80	2.03	0.26	86	11.0	1.06	0.14	2.9	0.38	67
12.10	black	3.37	-32.8	7.57	3.22	0.43	221	29.2	5.48	0.72	4.7	0.63	353
14.10	gray	0.94		4.54	1.95	0.43	44	9.8	1.21	0.27	2.0	0.44	220
14.60	black	3.86	-33.0	9.06	2.28	0.25	239	26.4	3.25	0.36	4.9	0.54	338
15.60	gray	0.23	-32.7	10.10	2.22	0.22	132	13.1	0.83	0.08	4.0	0.40	43
15.80	gray	0.20		10.09	2.57	0.25	109	10.8	0.91	0.09	3.6	0.36	55
17.10	gray	0.26	-32.3	9.49	2.07	0.22	114	12.0	1.29	0.14	4.4	0.46	62
18.60	gray	0.24		9.20	2.95	0.32	89	9.7	1.88	0.20	3.6	0.39	71
20.10	gray	0.18		9.54	2.73	0.29	112	11.7	1.33	0.14	3.4	0.36	72
21.60	black	3.92	-32.4	8.18	4.20	0.51	245	29.9	7.22	0.88	5.0	0.61	319
22.50	gray	0.30	-32.2	9.35	2.62	0.28	106	11.3	1.46	0.16	3.9	0.41	93
22.60	gray	0.15	-31.6	9.07	2.62	0.29	103	11.4	1.83	0.20	3.8	0.42	80
26.10	black	3.53		7.64	4.53	0.59	223	29.2	5.34	0.70	4.6	0.60	310
26.60	gray	0.40	-32.0	8.60	2.73	0.32	108	12.6	2.22	0.26	4.1	0.47	113
27.60	black	3.64	-32.7	8.27	3.64	0.44	200	24.2	4.48	0.54	5.6	0.68	336
28.40	gray	0.43		8.65	2.50	0.29	119	13.7	1.94	0.22	4.6	0.53	128
28.50	black	3.91	-32.2	7.76	4.18	0.54	259	33.4	7.72	0.99	6.6	0.85	287
29.00	gray	0.24		8.78	1.90	0.22	105	12.0	1.03	0.12	3.7	0.42	104
29.10	black	3.28		8.15	3.85	0.47	271	33.2	7.04	0.86	6.3	0.78	241
29.50	gray	0.30		9.09	2.15	0.24	112	12.3	1.55	0.17	4.4	0.48	107

29.60	black	2.41		7.81	3.56	0.46	132	16.9	4.36	0.56	5.3	0.67	256
30.10	gray	0.19	-32.8	9.29	1.72	0.19	100	10.8	0.62	0.07	3.4	0.37	89
31.10	black	2.84		8.07	4.19	0.52	169	20.9	4	0.50	5.9	0.72	205
34.10	black	3.02		7.96	3.70	0.47	148	18.6	3.91	0.49	4.3	0.54	343
35.60	gray	0.35	-32.5	8.36	1.49	0.18	110	13.2	1.07	0.13	3.2	0.39	120
36.60	gray	0.53		8.34	1.63	0.20	114	13.7	1.29	0.15	3.4	0.40	160
37.60	black	3.42	-33.0	7.93	2.08	0.26	216	27.2	5.5	0.69	5.2	0.66	430
39.00	gray	0.30	-32.7	8.71	1.84	0.21	122	14.0	0.89	0.10	4.1	0.47	127
39.10	gray	0.19	-31.7	8.37	1.72	0.20	77	9.2	0.66	0.08	3.1	0.37	100
40.60	gray	0.15		8.11	1.93	0.24	80	9.9	0.68	0.08	3.0	0.37	73
42.10	gray	0.26	-32.4	8.14	2.52	0.31	114	14.0	1.33	0.16	4.3	0.53	135
43.70	black	2.51	-33.7	6.70	4.32	0.64	197	29.4	7.93	1.18	4.8	0.71	314
43.75	gray	0.28		7.85	2.33	0.30	101	12.9	1.17	0.15	3.8	0.49	139
43.80	black	3.68	-32.9	7.39	3.71	0.50	189	25.6	5.58	0.75	3.9	0.53	390
43.90	gray	0.31	-31.9	7.55	1.82	0.24	104	13.8	1.07	0.14	4.1	0.54	145
43.94	black	3.10		7.40	2.46	0.33	176	23.8	5.21	0.70	5.3	0.72	342
43.95	gray	0.25		7.30	1.31	0.18	93	12.8	1.06	0.15	3.8	0.52	136
44.00	black	3.17		7.09	4.09	0.58	249	35.1	10	1.41	5.6	0.78	302
44.10	black	3.51	-33.3	7.03	3.21	0.46	246	35.0	7.65	1.09	6.0	0.85	403
44.40	black	3.40	-33.6	7.04	3.56	0.51	238	33.8	7.58	1.08	6.2	0.88	385
44.70	black	2.48	-33.8	6.69	3.53	0.53	178	26.6	6.75	1.01	4.7	0.70	363

Table S3. Fe speciation results from fresh core material

Depth	TOC	FeT	FeCARB	FeOX	FeMAG	FePy	FePy/FeHR	FeHR/FeT	FeT method
M	wt%	wt%	wt%	wt%	wt%	wt%			
15.00	2.14	3.72	0.37	0.19	0.24	0.41	0.34	0.33	HCl
15.71	0.21	2.71	0.05	0.16	0.03	0.19	0.44	0.16	HCl
16.66	0.31	4.31	0.19	0.05	0.17	0.51	0.55	0.22	HCl
17.85	0.36	4.13	0.34	0.05	0.10	0.77	0.61	0.30	XRF
18.33	0.31	3.22	0.22	0.11	0.13	0.50	0.52	0.30	XRF
18.8	0.39	4.71	0.47	0.09	0.23	0.78	0.49	0.33	XRF
19.21	3.55	2.50	0.35	0.04	0.02	1.50	0.78	0.76	HHXRF
19.75	0.41	4.14	0.47	0.06	0.14	0.72	0.52	0.34	HCl
20.94	0.39	3.61	0.16	0.03	0.06	0.75	0.75	0.28	HCl
21.89	0.53	4.00	0.24	0.03	0.09	0.97	0.73	0.33	Wet
22.13	1.66	3.82	0.65	0.16	0.24	0.58	0.35	0.43	XRF
22.84	0.38	3.51	0.15	0.03	0.06	0.71	0.74	0.27	HCl
23.79	0.26	3.98	0.41	0.06	0.22	0.36	0.34	0.27	HCl
24.74	0.18	4.62	0.24	0.05	0.26	0.33	0.38	0.19	HCl
24.98	0.51	2.86	0.15	0.10	0.08	0.60	0.65	0.32	XRF
25.46	3.92	5.60	1.26	0.15	0.31	1.46	0.46	0.57	HHXRF
25.69	0.27	3.71	0.11	0.05	0.11	0.57	0.67	0.23	HCl
25.93	0.79	5.65	0.18	0.04	0.15	1.09	0.74	0.26	HHXRF
26.41	0.86	6.10	0.21	0.04	0.15	0.61	0.60	0.17	HHXRF
26.64	0.29	3.64	0.20	0.05	0.09	0.83	0.71	0.32	HCl
26.88	0.47	3.75	0.13	0.03	0.12	0.73	0.72	0.27	HHXRF
27.36	0.93	5.35	0.14	0.04	0.12	0.52	0.64	0.15	HHXRF
27.59	0.45	3.74	0.12	0.04	0.06	0.69	0.76	0.24	HCl
27.83	0.47	2.88	0.08	0.06	0.03	0.73	0.81	0.31	XRF
28.30	2.91	3.56	0.18	0.04	0.15	1.43	0.79	0.51	HHXRF

28.54	0.99	3.58	0.17	0.04	0.09	1.08	0.78	0.39	HCl
28.78	1.10	2.71	0.10	0.04	0.09	0.79	0.77	0.37	HHXRF
29.26	1.02	3.07	0.23	0.08	0.14	1.23	0.74	0.54	HHXRF
29.49	0.42	3.76	0.18	0.04	0.09	0.97	0.76	0.34	HCl
29.73	0.67	2.85	0.21	0.05	0.12	0.79	0.68	0.41	HHXRF
29.96	0.47	3.21	0.18	0.04	0.03	0.94	0.79	0.37	HCl
30.21	0.51	2.68	0.08	0.04	0.09	0.68	0.76	0.34	HHXRF
30.44	0.32	2.97	0.10	0.03	0.04	0.46	0.72	0.21	HCl
30.68	2.52	3.72	0.70	0.11	0.10	0.63	0.41	0.42	XRF
31.16	3.59	3.37	0.66	0.14	0.19	0.74	0.43	0.51	HHXRF
31.39	0.43	3.33	0.30	0.04	0.08	0.93	0.69	0.41	HCl
31.63	0.44	2.57	0.07	0.05	0.05	0.67	0.80	0.33	HHXRF
32.11	1.08	3.41	0.27	0.05	0.09	1.38	0.77	0.53	HHXRF
32.34	0.34	2.94	0.14	0.04	0.04	0.59	0.74	0.27	HCl
32.58	0.71	2.84	0.12	0.05	0.05	1.11	0.84	0.47	HHXRF
33.06	0.72	3.01	0.33	0.06	0.09	0.93	0.66	0.47	HHXRF
33.29	0.87	4.88	0.37	0.05	0.08	2.21	0.82	0.56	HCl
33.53	3.36	3.20	0.55	0.08	0.03	1.73	0.72	0.75	XRF
34.01	3.41	3.48	0.46	0.07	0.07	2.03	0.77	0.75	HHXRF
34.48	4.20	3.50	0.60	0.11	0.08	2.08	0.72	0.82	HHXRF
34.96	3.50	3.39	0.59	0.09	0.12	1.82	0.69	0.77	HHXRF
35.43	3.16	3.68	1.76	0.11	0.01	1.29	0.41	0.86	HHXRF
35.91	4.77	3.06	0.40	0.06	0.04	1.78	0.78	0.74	HHXRF
36.38	4.79	3.38	1.38	0.08	0.08	0.80	0.34	0.69	XRF
36.86	3.95	3.04	0.43	0.07	0.04	1.80	0.77	0.77	HHXRF
37.33	4.94	3.04	0.54	0.07	0.04	1.85	0.74	0.82	HHXRF
37.81	4.01	3.00	0.40	0.06	0.04	1.80	0.78	0.77	HHXRF
38.28	3.52	3.20	0.51	0.07	0.04	1.92	0.76	0.79	HHXRF
38.76	3.11	2.90	0.49	0.07	0.06	1.40	0.69	0.70	HHXRF

39.23	2.61	3.60	0.00	0.03	0.01	2.16	0.98	0.61	XRF
39.71	4.45	3.20	0.41	0.06	0.04	1.94	0.79	0.76	HHXRF
40.18	4.07	2.89	0.38	0.05	0.04	1.51	0.76	0.69	HHXRF
40.66	4.03	3.41	0.42	0.05	0.03	2.38	0.83	0.85	HHXRF
41.13	3.74	3.48	0.60	0.07	0.04	2.24	0.76	0.85	HHXRF
41.61	4.70	3.49	0.66	0.08	0.04	2.50	0.76	0.94	HHXRF
42.08	1.68	1.16	0.11	0.04	0.04	0.11	0.38	0.25	XRF
42.56	2.98	2.73	0.39	0.03	0.06	1.74	0.78	0.81	HHXRF
43.03	4.40	3.13	0.47	0.05	0.03	1.97	0.78	0.81	HHXRF
43.51	4.20	3.28	0.51	0.05	0.03	2.24	0.79	0.86	HHXRF
43.98	3.16	3.08	0.48	0.02	0.03	1.95	0.79	0.81	HHXRF
44.46	4.83	3.08	0.53	0.05	0.03	2.19	0.78	0.91	HHXRF
44.93	3.93	2.39	0.54	0.04	0.01	1.16	0.66	0.73	XRF

Table S4. Parameters used to calculate O₂ exposure time and OC burial efficiencies

site	Depth	O ₂ pen.	Sed rate	OrgC	O ₂ uptake rate	O ₂ exposure time	Burial eff.	ref
	m	cm	cm y ⁻¹	wt%	mmol O ₂ cm ⁻² y ⁻¹	years	%	
Cape lookout Bight	10	0.30	5.4x10 ⁰	3.0	2.1x10 ⁰	5.6x10 ⁻²	83.0	(Canfield, 1989)
Aarhus Bay	16	0.60	1.7x10 ⁻¹	2.0	6.5x10 ⁻¹	3.6x10 ⁰	25.0	(Jørgensen, 1996; Thamdrup et al., 1994)
Kattegat/Skagerrak Sta 2	25	0.25	1.3x10 ⁻¹	1.5	6.3x10 ⁻¹	1.9x10 ⁰	16.6	(Jørgensen and Revsbech, 1989)
Young Sound, Greenland	36	1.00	1.2x10 ⁻¹	1.3	2.4x10 ⁻¹	8.5x10 ⁰	29.6	(Rysgaard et al., 1998)
Katt/Skagerrak Sta 6	43	0.42	2.6x10 ⁻¹	1.4	3.9x10 ⁻¹	1.6x10 ⁰	37.6	(Jørgensen and Revsbech, 1989)
Katt/Skagerrak Sta 9	65	0.55	2.6x10 ⁻¹	1.7	4.4x10 ⁻¹	2.1x10 ⁰	39.0	(Jørgensen and Revsbech, 1989)
Katt/Skagerrak Sta 10	73	0.30	2.6x10 ⁻¹	1.8	8.3x10 ⁻¹	1.2x10 ⁰	27.0	(Jørgensen and Revsbech, 1989)
Svalbard Sta 3	115	0.96	3.6x10 ⁻¹	1.8	4.8x10 ⁻¹	2.6x10 ⁰	46.7	(Glud et al., 1998; Sagemann et al., 1998)
Svalbard Sta 2	155	0.55	6.1x10 ⁻¹	1.5	6.0x10 ⁻¹	9.0x10 ⁻¹	49.5	(Glud et al., 1998; Sagemann et al., 1998)
Svalbard Sta 5	175	1.10	1.8x10 ⁻¹	2.5	3.2x10 ⁻¹	6.0x10 ⁰	47.7	(Glud et al., 1998; Sagemann et al., 1998)
Katt/Skagerrak Sta 12	200	0.32	2.6x10 ⁻¹	2.4	5.0x10 ⁻¹	1.2x10 ⁰	44.4	(Jørgensen and Revsbech, 1989)
California slope Sta K	1000	0.50	6.2x10 ⁻³	3.9	4.3x10 ⁻²	8.0x10 ¹	26.6	(Reimers et al., 1992)
Chilean coast Sta 40	1015	0.60	3.1x10 ⁻²	2.0	2.1x10 ⁻¹	1.9x10 ¹	16.0	(Thamdrup and Canfield, 1996)
Northeast Atlantic Sta B	1023	1.60	1.6x10 ⁻³	0.3	5.4x10 ⁻²	1.0x10 ³	0.6	(Lohse et al., 1998)
California rise Sta D	1440	1.00	1.0x10 ⁻³	1.7	3.5x10 ⁻²	9.6x10 ²	3.1	(Reimers et al., 1992)
California rise Sta L	1890	0.90	3.0x10 ⁻³	1.2	5.3x10 ⁻²	3.0x10 ²	4.2	(Reimers et al., 1992)
San Clemente Basin	1900	0.80	2.0 x10 ⁻²	2.0	4.5 x10 ⁻²	4.1x10 ¹	36.0	(Bender et al., 1989)
Chilean coast Sta 41	2000	0.60	3.5x10 ⁻²	2.2	1.7x10 ⁻¹	1.7x10 ¹	20.9	(Thamdrup and Canfield, 1996)
Borderland Basins-SCI	2053	0.70	2.0x10 ⁻²	2.6	4.0 x10 ⁻²	3.6x10 ¹	45.0	(Berelson et al., 1996; Reimers, 1987)
California rise Sta H	3580	1.50	3.6x10 ⁻³	2.0	6.2x10 ⁻²	4.1x10 ²	7.0	(Reimers et al., 1992)
Borderland Basins-PE	3707	2.00	3.9x10 ⁻²	1.1	1.4x10 ⁻²	5.1x10 ²	17.0	(Berelson et al., 1996)
Northeast Atlantic Sta III	3719	5.50	2.1x10 ⁻³	0.4	5.7x10 ⁻²	2.6x10 ³	0.9	(Lohse et al., 1998)
California rise Sta M	3730	1.60	1.3 x10 ⁻²	2.8	6.4x10 ⁻²	1.2x10 ²	26.7	(Reimers, 1987; Reimers et al., 1992)
Northeast Atlantic Sta E	4486	10.00	2.3x10 ⁻³	0.4	3.6x10 ⁻²	4.3x10 ³	1.6	(Lohse et al., 1998)

Table S5. O₂ penetration depths, bottom water O₂ levels and sediment O₂ uptake rates for modern sediments.

Site	Water depth (m)	O ₂ pen. (cm)	BW O ₂ (μM)	O ₂ uptake mmol cm ⁻² y ⁻¹	reference
Cape lookout Bight	10	0.30	250	2.050	(Chanton et al., 1987)
Western Pac sta I	4765	60.00	250	0.005	(Murray and Grundmanis, 1980)
Western Pac Sta J	5361	60.00	250	0.005	(Murray and Grundmanis, 1980)
Western Pac StaK	5469	20.00	250	0.008	(Murray and Grundmanis, 1980)
Kattegat/Skagerrak Sta3	15	0.20	250	0.800	(Jørgensen and Revsbech, 1989)
Kattegat/Skagerrak Sta 8	17	0.45	250	0.550	(Jørgensen and Revsbech, 1989)
Kattegat/Skagerrak Sta 2	25	0.25	250	0.630	(Jørgensen and Revsbech, 1989)
Kattegat/Skagerrak Sta 6	43	0.42	250	0.390	(Jørgensen and Revsbech, 1989)
Kattegat/Skagerrak Sta 9	65	0.55	250	0.440	(Jørgensen and Revsbech, 1989)
Kattegat/Skagerrak Sta 10	73	0.30	250	0.830	(Jørgensen and Revsbech, 1989)
Kattegat/Skagerrak Sta 12	200	0.32	250	0.500	(Jørgensen and Revsbech, 1989)
Central Calif slope-rise	791	0.40	250	0.037	(Reimers et al., 1992)
Skagerrak S4	190	0.70	250	0.580	(Canfield et al., 1993)
Skagerrak S9	695	1.60	250	0.400	(Canfield et al., 1993)
Chile sta 40	1015	0.60	250	0.210	(Thamdrup and Canfield, 1996)
Chile sta 41	2000	0.60	250	0.170	(Thamdrup and Canfield, 1996)
Svalbard Sta 3	115	0.96	250	0.480	(Glud et al., 1998)
Svalbard sta 4	138	0.93	250	0.440	(Glud et al., 1998)
Svalbard sta 5	175	1.10	250	0.320	(Glud et al., 1998)
Svalbard Sta 2	155	0.55	250	0.600	(Glud et al., 1998)
S. Atlantic Geob 1713	604	0.40	250	0.340	(Glud et al., 1994)
S. Atlantic Geob 1719	1015	1.20	250	0.410	(Glud et al., 1994)
S. Atlantic Geob 1703	1747	1.10	250	0.566	(Glud et al., 1994)
S. Atlantic Geob 1711	1947	2.50	250	0.110	(Glud et al., 1994)
S. Atlantic Geob 1721	3095	4.10	250	0.153	(Glud et al., 1994)
S. Atlantic Geob 1702	3107	3.50	250	0.066	(Glud et al., 1994)

Aarhus Bay	16	0.60	250	0.650	(Thamdrup et al., 1994)
Arctic ocean-19/050	170	0.50	250	0.409	(Hulth et al., 1994)
Arctic ocean-19/143	188	0.60	250	0.143	(Hulth et al., 1994)
Arctic ocean-19/134	273	3.80	250	0.117	(Hulth et al., 1994)
Arctic ocean-19/070	318	0.50	250	0.335	(Hulth et al., 1994)
Arctic ocean-19/082	326	0.80	250	0.340	(Hulth et al., 1994)
Arctic ocean-19/119	486	2.50	250	0.092	(Hulth et al., 1994)
Arctic ocean-19/101	530	3.00	250	0.104	(Hulth et al., 1994)
Arctic ocean-19/086	550	1.70	250	0.210	(Hulth et al., 1994)
Arctic ocean-19/100	867	1.70	250	0.149	(Hulth et al., 1994)
Arctic ocean-19/112	1010	3.70	250	0.067	(Hulth et al., 1994)
Arctic ocean-19/078	2010	1.70	250	0.134	(Hulth et al., 1994)
Arctic ocean-19/108	2490	6.00	250	0.110	(Hulth et al., 1994)
Arctic ocean-19/098	2577	5.00	250	0.097	(Hulth et al., 1994)
Amazon shelf RMT-1	9	0.15	250	0.440	(Aller et al., 1996)
Amazon shelf OST-1	15	0.50	250	0.400	(Aller et al., 1996)
Amazon shelf RMT-2	18	0.15	250	0.580	(Aller et al., 1996)
Amazon shelf OST-2	20	0.50	250	0.390	(Aller et al., 1996)
Amazon shelf OST-3	40	0.15	250	0.470	(Aller et al., 1996)
Amazon shelf RMT-3	44	0.45	250	0.580	(Aller et al., 1996)
Patton Escarpment, E.N. PAC	3750	3.00	250	0.065	(Reimers et al., 1986; Reimers and Suess, 1983)
Young Sound, Greenland	36	1.00	250	0.235	(Rysgaard et al., 1998)
JGOFS 02S	4300	5.00	250	0.020	(Hammond et al., 1996)
JGOFS 02N	4400	6.00	250	0.019	(Hammond et al., 1996)
JGOFS 05N	4500	10.00	250	0.011	(Hammond et al., 1996)
JGOFS 00		6.00	250	0.018	(Hammond et al., 1996)
Northeast Atlantic StaI	668	0.80	250	0.086	(Lohse et al., 1998)
Northeast Atlantic StaB	1023	1.60	250	0.054	(Lohse et al., 1998)
Northeast Atlantic StaII	1442	2.50	250	0.065	(Lohse et al., 1998)
Northeast Atlantic StaC	1989	3.50	250	0.039	(Lohse et al., 1998)

Northeast Atlantic Sta III	3719	5.50	250	0.057	(Lohse et al., 1998)
Northeast Atlantic StaE	4486	10.00	250	0.036	(Lohse et al., 1998)
Borderland Basins-PE	3707	2.00	131	0.014	(Berelson et al., 1996)
Central Calif Sta M	3730	1.60	129	0.064	(Reimers et al., 1992)
East N. Pac NH14	114	0.40	127	0.670	(Devol and Christensen, 1993)
Central Calif Sta H	3580	1.50	125	0.062	(Reimers et al., 1992)
East N. Pac NH 18	146	0.30	106	0.530	(Devol and Christensen, 1993)
Central Calif Sta F	2000	1.10	75	0.056	(Reimers et al., 1992)
Central Calif Sta L	1890	0.90	72	0.053	(Reimers et al., 1992)
San Clemente Basin	1900	0.80	58	0.045	(Bender et al., 1989)
Borderland Basins-SCI	2053	0.70	58	0.040	(Berelson et al., 1996)
Central Calif Sta E	1570	0.80	50	0.048	(Reimers et al., 1992)
Central Calif Sta D	1440	1.00	33	0.035	(Reimers et al., 1992)
Central Calif Sta C	1200	0.40	26	0.048	(Reimers et al., 1992)
Central Calif Sta K	1000	0.50	20	0.043	(Reimers et al., 1992)
Central Calif Sta A	585	0.30	17	0.050	(Reimers et al., 1992)

References

- Aller, R. C., Blair, N. E., and Rude, P. D.: Remineralization rates, recycling, and storage of carbon in Amazon shelf sediments, *Cont Shelf Res*, 16, 753-786, 1996.
- Bender, M., Jahnke, R., Weiss, R., Martin, W., Heggie, D. T., Orchardo, J., and Sowers, T.: Organic carbon oxidation and benthic nitrogen and silica dynamics in San Clemente Basin, a continental borderland site, *Geochimica et Cosmochimica Acta*, 53, 685-697, 1989.
- Berelson, W. M., McManus, J., Coale, K. H., Johnson, K. S., Kilgore, T., Burdige, D., and Pilskaln, C.: Biogenic matter diagenesis on the sea floor: A comparison between two continental margin transects., *J Mar Res*, 54, 731-762, 1996.
- Canfield, D. E.: Sulfate reduction and oxic respiration in marine sediments: implications for organic carbon preservation in euxinic environments, *Deep-Sea Research*, 36, 121-138, 1989.
- Canfield, D. E., Jørgensen, B. B., Fossing, H., Glud, R., Gundersen, J., Ramsing, N. B., Thamdrup, B., Hansen, J. W., Nielsen, L. P., and Hall, P. O. J.: Pathways of organic carbon oxidation in three continental margin sediments, *Mar Geol*, 113, 27-40, 1993.
- Chanton, J. P., Martens, C. S., and Goldhaber, M. B.: Biogeochemical cycling in an organic-rich coastal marine basin. 8. A sulfur isotopic budget balanced by differential diffusion across the sediment-water interface, *Geochim Cosmochim Acta*, 51, 1201-1208, 1987.
- Devol, A. H. and Christensen, J. P.: Benthic fluxes and nitrogen cycling in sediments of the continental margin of the eastern North Pacific, *Journal of Marine Research*, 51, 345-372, 1993.
- Glud, R. N., Gundersen, J. K., Jørgensen, B. B., Revsbech, N. P., and Schultz, H. D.: Diffusive and total oxygen uptake of deep-sea sediments in the eastern South Atlantic Ocean: *in situ* and laboratory measurements, *Deep-Sea Research I*, 41, 1767-1788, 1994.
- Glud, R. N., Holby, O., Hoffmann, F., and Canfield, D. E.: Benthic mineralization and exchange in Arctic sediments (Svalbard, Norway), *Marine Ecology-Progress Series*, 173, 237-251, 1998.
- Hammond, D. E., McManus, J., Berelson, W. M., Kilgore, T. E., and Pope, R. H.: Early diagenesis of organic material in equatorial Pacific sediments: stoichiometry and kinetics, *Deep-sea research II*, 43, 1365-1412, 1996.
- Hulth, S., Blackburn, T. H., and Hall, P. O. J.: Arctic sediments (Svalbard): consumption and microdistribution of oxygen, *Mar Chem*, 46, 293-316, 1994.
- Jørgensen, B. B.: Case study-Aarhus Bay. In: *Eutrophication in Coastal Marine Ecosystems*, Jørgensen, B. B. and Richardson, K. (Eds.), American Geophysical Union, Washington DC, 1996.
- Jørgensen, B. B. and Revsbech, N. P.: Oxygen uptake, bacterial distribution, and carbon-nitrogen-sulfur cycling in sediments from the Baltic Sea-North Sea transition, *Ophelia*, 31, 29-49, 1989.
- Lohse, L., Helder, W., Epping, E. H. G., and Balzer, W.: Recycling of organic matter along shelf-slope transect across the N. W. European Continental Margin (Goban Spur), *Prog Oceanogr*, 42, 77-110, 1998.
- Murray, J. W. and Grundmanis, V.: Oxygen consumption in pelagic marine sediment, *Science*, 209, 1527-1530, 1980.
- Reimers, C. E.: An *in situ* microprofiling instrument for measuring interfacial porewater gradients: methods and oxygen profiles from the North Pacific ocean, *Deep-sea research*, 34, 2019-2035, 1987.

- Reimers, C. E., Fischer, K. M., Merewether, R., Smith, K. L., and Jahnke, R. A.: Oxygen microprofiles measured in situ in deep sea sediments, *Nature*, 320, 741-744, 1986.
- Reimers, C. E., Jahnke, R. A., and McCorkle, D. C.: Carbon fluxes and burial rates over the continental slope and rise off central California with implications for the global carbon cycle, *Global Biogeochem Cy*, 6, 199-224, 1992.
- Reimers, C. E. and Suess, E.: The partitioning of organic carbon fluxes and sedimentary organic matter decomposition rates in the ocean, *Mar Chem*, 13, 141-168, 1983.
- Rysgaard, S., Thamdrup, B., Risgaard-Petersen, N., Fossing, H., Berg, P., Christensen, P. B., and Dalsgaard, T.: Seasonal carbon and nutrient mineralization in a high-Arctic coastal marine sediment, Young Sound, Northeast Greenland, *Marine Ecology-Progress Series*, 175, 261-276, 1998.
- Sagemann, J., Jørgensen, B. B., and Greeff, O.: Temperature dependence and rates of sulfate reduction in cold sediments of Svalbard, Arctic Ocean, *Geomicrobiology Journal*, 15, 85-100, 1998.
- Thamdrup, B. and Canfield, D. E.: Pathways of carbon oxidation in continental margin sediments off central Chile, *Limnology and Oceanography*, 41, 1629-1650, 1996.
- Thamdrup, B., Fossing, H., and Jørgensen, B. B.: Manganese, iron, and sulfur cycling in a coastal marine sediment, Aarhus Bay, Denmark, *Geochim Cosmochim Acta*, 58, 5115-5129, 1994.