

Supplement of Biogeosciences, 14, 3387–3400, 2017
<https://doi.org/10.5194/bg-14-3387-2017-supplement>
© Author(s) 2017. This work is distributed under
the Creative Commons Attribution 3.0 License.



Supplement of

Ba incorporation in benthic foraminifera

Lennart J. de Nooijer et al.

Correspondence to: Lennart J. de Nooijer (ldeNooijer@nioz.nl)

The copyright of individual parts of the supplement might differ from the CC BY 3.0 License.

Table 1: measured calcitic El/Ca, calculated partition coefficients and experimental/ field conditions. The selected publications reflect the range in known ratios, but is not aimed to encompass all published element ratios. Particularly for Mg/Ca and Sr/Ca, the listed ratios only represent a small selection of published values.

Low-Mg species are marked in green (darker shade for benthics, light one for planktonics), high Mg-species are in red (milliolid species are in light shade). Reported element/Ca ratios are reported as a range of values (min-max) when reported in the original study, in other cases average values are reported. ‘Study type’ refers to core-top/ sediment trap calibrations (1) or culture experiment (2). ‘n.d.’ means not determined or not reported. Data from this study are referenced as ‘t.s.’. Numbers between parentheses refer to the standard deviation. For a number of field studies, (conservative) element concentrations are not measured, but are here calculated (**) to obtain a partition coefficient. Assumed concentrations at salinity of 35.0 are 10.3 mmol/kg for Ca, 24.5 $\mu\text{mol/kg}$ for Li, 416 $\mu\text{mol/kg}$ for B, 0.469 mol/kg for Na, 52.8 mmol/kg for Mg, 90.6 $\mu\text{mol/kg}$ for Sr and 109 nmol/kg for Ba.

| Li/Ca | | | | | | | |
|---------------------------------|---------------------|-------------------------|-----------------------------------|--------------------------|-------------|------------|---------------|
| calcite ($\mu\text{mol/mol}$) | seawater (mmol/mol) | D_{El} (*1000) | species | T ($^{\circ}\text{C}$) | salinity | study type | ref |
| 3.5 – 5.5 | 2.39* | 1.5 – 2.3 | <i>Hoeglundina elegans</i> | 4 – 16.5 | n.d. | 1 | ²⁴ |
| 2.7 – 4.6 | 2.39* | 1.1 – 1.9 | <i>Hoeglundina elegans</i> | 5.8 – 19.0 | 34.9 – 36.8 | 1 | ²² |
| 12 - 20 | 2.39* | 5.0 – 8.4 | <i>Cibicidoides</i> spp. | 4 – 16.5 | n.d. | 1 | ²⁴ |
| 11 - 14 | 2.39* | 4.6 – 5.9 | <i>Cibicidoides pachyderma</i> | 5.8 – 18.6 | 34.9 – 36.8 | 1 | ²² |
| 12 – 14 | 2.39* | 5.0 – 5.9 | <i>Cibicidoides wuellerstorfi</i> | 2.9 – 3.4 | n.d. | 1 | ²⁶ |
| 14 – 18 | 2.52 | 5.6 – 7.1 | <i>Oridorsalis umbonatus</i> | 1.1-3.6 | n.d. | 1 | ¹ |
| 12 - 15 | 2.39* | 5.0 – 6.3 | <i>Planulina ariminensis</i> | 7.0 – 12.1 | 34.9 – 36.8 | 1 | ²² |
| 11 - 17 | 2.39* | 4.6 – 7.1 | <i>Planulina foveolata</i> | 11.0 – 17.8 | 34.9 – 36.8 | 1 | ²² |
| 13 - 19 | 2.39* | 5.4 – 7.6 | <i>Uvigerina</i> spp. | 3.6 – 22.5 | 34.9 – 36.2 | 1 | ²⁵ |
| 15 – 18 | 2.39* | 6.3 – 7.1 | <i>Uvigerina peregrina</i> | 5.8 – 17.2 | 34.9 – 36.8 | 1 | ²² |
| | | | <i>Amphistegina lessonii</i> | 25 | n.d. | 2 | ³¹ |
| 14-18 | 2.39* | 5.9 - 7.5 | <i>Amphistegina lessonii</i> | 18-33 | 35 | 2 | ³² |
| 13-16 | 2.39* | 5.4 – 6.7 | <i>Amphistegina lobifera</i> | 24 | 35 | 2 | ³² |
| 12-20 | 2.39* | 5.0 – 8.4 | <i>Globorotalia inflata</i> | 6 -13 | 35.0 – 35.8 | 1 | ²³ |
| 13 – 17 | 2.39* | 5.3 – 7.0 | <i>Globigerinoides ruber</i> | n.d. | n.d. | 1 | ²⁸ |

| | | | | | | | |
|--|---------------------|----------------------------------|-------------------------------------|-------------|-------------|------------|---------------|
| 12 - 14 | 2.39* | 4.8 – 5.6 | <i>Globigerinoides sacculifer</i> | 26 | n.d. | 1 | ²⁴ |
| 0 - 14 | 2.39* | ≤5.6 | <i>Globigerinoides sacculifer</i> | n.d. | n.d. | 1 | ²⁸ |
| 17 - 20 | 2.39* | 7.1 – 8.4 | <i>Neogloboquadri na pachyderma</i> | -2 - 0 | 32.5 – 33.5 | 1 | ¹⁸ |
| 10 - 11 | 2.39* | 4.2 – 4.5 | <i>Orbulina universa</i> | 26 | n.d. | 1 | ²⁴ |
| 10-20 | 1-5 | 4-10 | <i>Globigerinoides sacculifer</i> | 30 | n.d. | 2 | ³⁴ |
| Maximum range in D_{Li} | | 1 - 9* 10⁻³ | | | | | |
| 55.7 | 1.8 | 31 | <i>Operculina ammonoides</i> | 22 | n.d. | 1 | ¹⁰ |
| 53-62 | 2.1 | 25 - 30 | <i>Operculina ammonoides</i> | 19-27 | 37 – 38 | 2 | ¹⁰ |
| Maximum range in D_{Li} | | 25 – 31 * 10⁻³ | | | | | |
| B/Ca | | | | | | | |
| calcite (μmol/mol) | seawater (mmol/mol) | D _{EI} (*1000) | species | T (°C) | salinity | study type | ref |
| 30 - 60 | 40* | 0.75 – 1.5 | <i>Hoeglundina elegans</i> | -1 - 4 | n.d. | 1 | ²¹ |
| 172 | 40* | 5.1 | <i>Cibicidoides mundulus</i> | 5.3 | 35 | 1 | ²⁷ |
| 100 - 175 | 40* | 2.5 – 4.4 | <i>Cibicidoides mundulus</i> | -1 - 4 | n.d. | 1 | ²¹ |
| 125 - 250 | 40* | 3.1 – 6.2 | <i>Cibicidoides wuellerstorfi</i> | -1 - 4 | n.d. | 1 | ²¹ |
| 202 - 218 | 40* | 5.1 - 5.5 | <i>Cibicidoides wuellerstorfi</i> | 2.9 – 3.4 | n.d. | 1 | ²⁶ |
| 239 | 40* | 6.0 | <i>Cibicidoides wuellerstorfi</i> | 3.7 | 35 | 1 | ²⁷ |
| 20 – 60 | 40* | 1.5 – 1.5 | <i>Oridorsalis umbonatus</i> | 1.1-3.6 | n.d. | 1 | ¹ |
| 15 - 50 | 40* | 0.4 – 1.3 | <i>Uvigerina spp.</i> | -1 - 4 | n.d. | 1 | ²¹ |
| 113 – 212 | 40* | 2.8 – 5.3 | <i>Globigerinoides ruber</i> | 24.0 – 29.3 | 33 - 40 | 2 | ¹² |
| 90 - 160 | 43.8* | 2.1 – 3.7 | <i>Globigerinoides ruber</i> | 19-27 | 36.2 – 37.0 | 1 | ¹⁶ |
| 112 - 121 | 40* | 2.8 – 3.0 | <i>Globigerinoides ruber</i> | 19.5 – 30.4 | 34.7 – 36.8 | 1 | ²⁷ |
| 103-122 | 40* | 2.6 – 3.1 | <i>Globigerinoides sacculifer</i> | 28 | 36 | 1 | ¹² |
| 79 - 145 | 40* | 2.0 – 3.6 | <i>Globigerinoides sacculifer</i> | 24.0 – 29.3 | 33 - 40 | 2 | ¹² |
| 73 - 98 | 40* | 1.8 – 2.5 | <i>Globigerinoides sacculifer</i> | 23.8 – 28.0 | 34.8 – 35.9 | 1 | ²⁷ |
| 48 - 62 | 40* | 1.2 – 1.6 | <i>Neogloboquadri na dutertrei</i> | 18.7 – 24.2 | 34.8 – 36.7 | 1 | ²⁷ |

| | | | | | | | |
|--|------------------------|------------------------------------|---|----------------|----------------|---------------|---------------|
| 45 - 77 | 40* | 1.1 – 1.9 | <i>Neogloboquadri na pachyderma</i> | -2 - 0 | 32.6 – 33.6 | 1 | ¹⁸ |
| 53 - 92 | 38 - 43 | 0.6 – 3.0 | <i>Orbulina universa</i> | 17.7 – 26.5 | 29.9 – 35.4 | 2 | ¹¹ |
| 66 - 76 | 40* | 1.7 – 1.9 | <i>Orbulina universa</i> | 28 | 36 | 1 | ¹² |
| Maximum range in D_B | | 0.4 – 6 * 10⁻³ | | | | | |
| 1200 - 6360 | 392 | 3.1 - 16 | <i>Amphistegina lessonii</i> | 25 | 32 | 2 | ⁸ |
| Maximum range in D_B | | 3 – 16 * 10⁻³ | | | | | |
| Na/Ca | | | | | | | |
| calcite (mmol/mol) | seawater (mol/mol) | D _{EI} (*1000) | species | T (°C) | salinity | study type | ref |
| 6.12 (±0.33) | 47.8 (±0.7) | 0.13 | <i>Ammonia tepida</i> | 20.0 (±0.2) | 32.5 (±0.2) | 2 | ³ |
| 5.9 – 7.6 | 45.5* | 0.13 – 0.17 | <i>Globigerinoides ruber</i> | n.d. | n.d. | 1 | ²⁸ |
| 5.1 – 6.4 | 45.5* | 0.11 – 0.14 | <i>Globigerinoides sacculifer</i> | n.d. | n.d. | 1 | ²⁸ |
| 5.5 – 6.0 | 45.5* | 0.12 – 0.13 | <i>Globigerinoides sacculifer</i> | n.d. | n.d. | 1 | ²⁹ |
| 4.5 – 5.2 | 45.5* | 0.099 – 0.11 | <i>Neogloboquadri na dutertrei</i> | n.d. | n.d. | 1 | ²⁹ |
| 4.0-6.0 | 30-50 | 0.1-0.15 | <i>Globigerinoides sacculifer</i> | 30 | n.d | 2 | ³⁴ |
| Maximum range in D_{Na} | | 0.1 – 0.2 * 10⁻³ | | | | | |
| 24 | 41.6 | 0.58 | <i>Operculina ammonoides</i> | 24 | 37 | 2 | ¹⁰ |
| Maximum range in D_{Na} | | 0.6 * 10⁻³ | | | | | |
| Mg/Ca | | | | | | | |
| calcite (mmol/mol) | seawater (mmol/mol) | D _{EI} (*1000) | species | T (°C) | salinity | study type | ref |
| 2 - 7 | 5158 | 0.39 – 1.4 | <i>Ammonia tepida</i> | 25 (±0.5) | 32.2 (±0.2) | 2 | ² |
| 1.3 – 2.2 | 5080 (±30) | 0.26 – 0.43 | <i>Ammonia tepida</i> | 20.0 (±0.2) | 32.5 (±0.2) | 2 | ³ |
| 1 - 3 | 5100 – 5300 | 0.31 – 0.70 | <i>Ammonia tepida</i> | 18.0 (±0.5) | 35.0 (±0.3) | 2 | ⁴ |
| 40 - 60 | 5200 | 7.7 - 12 | <i>Amphistegina lessonii</i> | 24 | 35 | 2 | ⁷ |
| 68 - 86 | 5126* | 13 - 17 | <i>Amphistegina lessonii</i> | 21 - 29 | n.d. | 1 | ³⁰ |
| 50 - 70 | 5200 | 9.6 - 13 | <i>Amphistegina lobifera</i> | 24 | 35 | 2 | ⁷ |
| 0.98 – 1.40 | 5126* | 0.19 – 0.27 | <i>Cibicidoides wuellerstorfi</i> | 2.9 – 3.4 | n.d. | 1 | ²⁶ |

| | | | | | | | |
|--|---------------------|-----------------------------------|-------------------------------------|--------------|--------------|------------|-----|
| 1 – 3 | 5300 | 0.19 – 0.57 | <i>Oridorsalis umbonatus</i> | 1.1-3.6 | n.d. | 1 | 1 |
| 0.75 – 2.5 | 5126* | 0.15 – 0.49 | <i>Uvigerina</i> spp. | 1.6 - 20 | n.d. | 1 | 6 |
| 3.5 - 5.5 | 5126* | 0.68 – 1.1 | <i>Globigerinoides ruber</i> | 20-26 | n.d. | 1 | 16 |
| 5.2 | 5126* | 1.0 | <i>Globigerinoides sacculifer</i> | 26 | 36 | 2 | 17 |
| 0.75 – 1.05 | 5126* | 0.15 – 0.20 | <i>Neogloboquadri na pachyderma</i> | -2 - 0 | 32.6 – 33.6 | 1 | 18 |
| Maximum range in D_{Mg} | | 0.2 – 17 * 10⁻³ | | | | | |
| 110 - 140 | 5200 – 6200 | 25– 28 | <i>Heterostegina depressa</i> | 18.0 (±0.5) | 35.0 (±0.1) | 2 | 4 |
| 214 - 267 | 5126* | 42 - 52 | <i>Neorotalia calcar</i> | 21 - 29 | n.d. | 1 | 30 |
| 141 (± 2) | 5330 | 27 | <i>Operculina ammonoides</i> | 24 | 37 | 2 | 10 |
| 138 - 144 | 5050 (± 50) | 27 - 29 | <i>Planoglabratella opercularis</i> | 23.1 (± 0.2) | 35 | 2 | 9 |
| 136 | 5050 (± 50) | 27 | <i>Quinqueloculina yabei</i> | 24.5 (± 1.5) | 36.5 | 2 | 9 |
| 95 -103 | 5126* | 19 - 20 | <i>Triloculina triincarta</i> | 18.3 | n.d. | 1 | 20 |
| 90 – 100 | 5126* | 18 - 20 | <i>Spiroloculina subimpressa</i> | 18.3 | n.d. | 1 | 20 |
| 80 -105 | 5126* | 16 - 20 | <i>Pyrgo sarsi</i> | 18.3 | n.d. | 1 | 20 |
| 224 - 256 | 5126* | 44 - 50 | <i>Amphisorus hemprichii</i> | 21 – 29 | n.d. | 1 | 30 |
| 213 - 255 | 5126* | 42 - 50 | <i>Marginopora vertebralis</i> | 21 - 29 | n.d. | 1 | 30 |
| 116 – 132 | 5126* | 23 – 26 | <i>Archaias ungulatus</i> | 23.9 (± 0.1) | 31.8 (± 0.2) | 2 | 33 |
| Maximum range in D_{Mg} | | 16 – 52 * 10⁻³ | | | | | |
| Sr/Ca | | | | | | | |
| calcite (mmol/mol) | seawater (mmol/mol) | D _{EI} | species | T (°C) | salinity | study type | ref |
| 1.2 – 1.9 | 9.47 | 0.13 – 0.20 | <i>Ammonia tepida</i> | 25 (±0.5) | 32.2 (±0.2) | 2 | 2 |
| 1.4 – 2.0 | 9.27 (±0.15) | 0.15 - 0.22 | <i>Ammonia tepida</i> | 20.0 (±0.2) | 32.5 (±0.2) | 2 | 3 |
| 1.35 (±0.03) | 4.6 – 15.6 | 0.16 – 0.17 | <i>Ammonia tepida</i> | 18.0 (±0.5) | 35.0 (±0.3) | 2 | 4 |
| 1.6 – 1.9 | 8.83* | 0.18 – 0.22 | <i>Amphistegina lessonii</i> | 21 - 29 | n.d. | 1 | 30 |
| 1.29 – 1.36 | 8.83* | 0.15 | <i>Cibicidoides wuellerstorfi</i> | 2.6 – 3.4 | n.d. | 1 | 26 |
| 0.8 – 1.00 | 8.72 | 0.09 – 0.11 | <i>Oridorsalis umbonatus</i> | 1.1-3.6 | n.d. | 1 | 1 |

| | | | | | | | |
|--|---------------------|------------------|-------------------------------------|-------------|-------------|------------|------|
| 1.35 | 8.83* | 0.15 | <i>Globigerinoides sacculifer</i> | 26 | 36 | 2 | 17 |
| 1.36 – 1.40 | 8.83* | 0.15 – 0.16 | <i>Neogloboquadri na pachyderma</i> | -2 - 0 | 32.6 – 33.6 | 1 | 18 |
| Maximum range in D_{Sr} | | 0.1 – 0.2 | | | | | |
| 2.56 | 4.8 – 17.8 | 0.27 – 0.33 | <i>Heterostegina depressa</i> | 18.0 (±0.5) | 35.0 (±0.1) | 2 | 4 |
| 1.9 – 2.2 | 8.83* | 0.22 – 0.25 | <i>Neorotalia calcar</i> | 21 - 29 | n.d. | 1 | 30 |
| 2.56 | 8.42 | 0.30 | <i>Operculina ammonoides</i> | 24 | 37 | 2 | 10 |
| 1.8 – 1.9 | 8.83* | 0.20 – 0.22 | <i>Amphisorus hemprichii</i> | 21 - 29 | n.d. | 1 | 30 |
| 0.6 – 1.8 | 8.83* | 0.08 – 0.20 | <i>Marginopora vertebralis</i> | 21 - 29 | n.d. | 1 | 30 |
| Maximum range in D_{Sr} | | 0.1 – 0.3 | | | | | |
| Ba/Ca | | | | | | | |
| calcite (μmol/mol) | seawater (μmol/mol) | D _{EI} | species | T (°C) | salinity | study type | ref |
| 10 - 40 | 50 - 90 | 0.32 | <i>Amphistegina lessonii</i> | 25 | 32.5 | 2 | t.s. |
| 2.2 – 5.0 | 4.8 – 7.3 | 0.39 - 0.41 | <i>Cibicidoides kullenbergi</i> | n.d. | n.d. | 1 | 5 |
| 1.8 – 4.4 | 4.5 – 13.5 | 0.35 – 0.37 | <i>Cibicidoides wuellerstorfi</i> | n.d. | n.d. | 1 | 5 |
| 1.9 – 4.7 | 4.6 - 13.1 | 0.32 – 0.34 | <i>Uvigerina</i> spp. | n.d. | n.d. | 1 | 5 |
| 0.5 - 0.8 | 3.7 | 0.14 – 0.22 | <i>Globigerina bulloides</i> | 22 | 33.7 | 2 | 13 |
| 1.2 – 3.1 | 10.6* | 0.11 – 0.29 | <i>Globigerinoides ruber</i> | n.d. | n.d. | 1 | 28 |
| 0.7 – 1.0 | 3.3 – 4.0 | 0.17 – 0.19 | <i>Globigerinoides ruber</i> | n.d. | n.d. | 1 | 15 |
| 0.6 – 0.9 | 3.3 – 4.0 | 0.17 – 0.19 | <i>Globigerinoides sacculifer</i> | n.d. | n.d. | 1 | 15 |
| 0.65 – 2.17 | 4.4 - 15 | 0.14 - 0.15 | <i>Globigerinoides sacculifer</i> | 22 - 29 | 36.7 | 2 | 14 |
| 1.0 – 1.8 | 10.6* | 0.094 – 0.17 | <i>Globigerinoides sacculifer</i> | n.d. | n.d. | 1 | 28 |
| 1.4 – 1.6 | 5.4 – 5.5 | 0.25 – 0.30 | <i>Neogloboquadri na pachyderma</i> | 0 | n.d. | 1 | 19 |
| 0.5 - 0.8 | 3.7 | 0.14 – 0.22 | <i>Orbulina universa</i> | 18 - 26 | 33.7 | 2 | 13 |
| 0.7 – 2.3 | 4.4 - 15 | 0.14 – 0.15 | <i>Orbulina universa</i> | 22 - 29 | 36.7 | 2 | 14 |
| 0.6 – 0.9 | 3.3 – 4.0 | 0.17 – 0.19 | <i>Orbulina universa</i> | n.d. | n.d. | 1 | 15 |
| Maximum range in D_{Ba} | | 0.1 – 0.4 | | | | | |

| | | | | | | | |
|--|---------|------------------|-------------------------------|----|------|---|---------------|
| 30 - 90 | 50 - 90 | 0.81 | <i>Heterostegina depressa</i> | 25 | 32.5 | 2 | t.s. |
| 0.3 – 13.5 | 15 - 19 | 0.53 – 0.71 | <i>Operculina ammonoides</i> | 24 | 37 | 2 | ¹⁰ |
| Maximum range in D_{Ba} | | 0.5 – 0.8 | | | | | |

References

- ¹Dawber, C.F., Tripathi, A., 2012. Relationships between bottom water carbonate saturation and element/Ca ratios in coretop samples of the benthic foraminifera *Oridorsalis umbonatus*. *Biogeosciences* 9, 3029-3045.
- ²De Nooijer, L.J., Hathorne, E.C., Reichart, G.J., Langer, G., Bijma, J., 2014a. Variability in calcitic Mg/Ca and Sr/Ca ratios in clones of the benthic foraminifer *Ammonia tepida*. *Marine Micropaleontology* 107, 33-43.
- ³Wit, J.C., De Nooijer, L.J., Wolthers, M., Reichart, G.J., 2013. A novel salinity proxy based on Na incorporation into foraminiferal calcite. *Biogeosciences* 10, 6375-6387.
- ⁴Dueñas-Bohórquez, Raitzsch, M., De Nooijer, L.J., Reichart, G.J., 2011. Independent impacts of calcium and carbonate ion concentration on Mg and Sr incorporation in cultured benthic foraminifera. *Marine Micropaleontology* 81, 122-130.
- ⁵Lea, D., Boyle, E., 1989, Barium content of benthic foraminifera controlled by bottom-water composition. *Nature* 338, 751-753.
- ⁶Elderfield, H., Yu, J., Anand, P., Kiefer, T., Nyland, B., 2006. Calibrations for benthic foraminiferal Mg/Ca paleothermometry and the carbonate ion hypothesis. *Earth Planet. Sci. Lett.* 250, 633-649.
- ⁷Segev, E., Erez, J., 2006. Effect of Mg/Ca ratio in seawater on shell composition in shallow benthic foraminifera. *Geochem. Geophys. Geosyst.* 7, GC000969.
- ⁸Kaczmarek, K., Langer, G., Nehrke, G., Horn, I., Misra, S., Janse, M., Bijma, J., 2015. Boron incorporation in the foraminifer *Amphistegina lessonii* under a decoupled carbon chemistry. *Biogeosciences* 12, 1753-1763.

- ⁹Toyofuku, T., Kitazato, H., Kawahata, H., 2000. Evaluation of Mg/Ca thermometry in foraminifera: Comparison of experimental results and measurements in nature. *Paleoceanography* 15, 456-464.
- ¹⁰Evans, D., Erez, J., Oron, S., Müller, W., 2015. Mg/Ca-temperature and seawater-test chemistry relationships in the shallow-dwelling large benthic foraminifera *Operculina ammonoides*. *Geochim. Cosmochim. Acta* 148, 325-342.
- ¹¹Allen, K.A., Hönisch, B., Eggins, S.M., Yu, J., Spero, H.J., Elderfield, H., 2011. Controls on Boron incorporation in cultured tests of the planktic foraminifer *Orbulina universa*. *Earth Planet. Sci. Lett.* 309, 291-301.
- ¹²Allen, K.A., Hönisch, B., Eggins, S.M., Rosenthal, Y., 2012. Environmental controls on B/Ca in calcite tests of the tropical planktic foraminifer species *Globigerinoides ruber* and *Globigerinoides sacculifer*. *Earth Planet. Sci. Lett.* 351-352, 270-280.
- ¹³Hönisch, B., Allen, K.A., Russell, A.D., Eggins, S.M., Bijma, J., Spero H.J., Lea, D.W., Yu, J., 2011. Planktic foraminifers as recorders of seawater Ba/Ca. *Marine Micropaleontology* 79, 52-57.
- ¹⁴Lea, D.W., Spero, H.J., 1994. Assessing the reliability of paleochemical tracers: Barium uptake in the shells of planktonic foraminifera. *Paleoceanography* 9, 445-452.
- ¹⁵Lea, D.W., Boyle, E.A. 1991. Barium in planktonic foraminifera. *Geochim. Cosmochim. Acta* 55, 3321-3331.
- ¹⁶Babila, T.L., Rosenthal, Y., Conte, M.H., 2014. Evaluation of the biogeochemical controls on B/Ca of *Globigerinoides ruber* from the Ocean Flux Program, Bermuda. *Earth Planet. Sci. Lett.* 404, 67-76.
- ¹⁷Dueñas-Bohórquez, A., Da Rocha, R., Kuroyanagi, A., Bijma, J., Reichert, G.J., 2009. Effect of salinity and seawater calcite saturation state on Mg and Sr incorporation in cultured planktonic foraminifera. *Marine Micropaleontology* 73, 178-189.

- ¹⁸Hendry, K.R., Rickaby, R.E.M., Meredith, M.P., Elderfield, H., 2009. Controls on stable isotope and trace metal uptake in *Neogloboquadrina pachyderma* (sinistral) from an Antarctic sea-ice environment. *Earth Planet. Sci. Lett.* 278, 67-77.
- ¹⁹Hall, J.M., Chan, L.-H., 2004b. Ba/Ca in *Neogloboquadrina pachyderma* as an indicator of deglacial meltwater discharge into the western Arctic Ocean. *Paleoceanography* 19, PA000910.
- ²⁰Sadekov, A., Bush, F., Kerr, J., Ganeshram, R., Elderfield, H., 2014. Mg/Ca composition of benthic foraminifera *Miliolacea* as a new tool of paleoceanography. *Paleoceanography* 29, 990-1001.
- ²¹Yu, J., Elderfield, H., 2007. Benthic foraminiferal B/Ca ratios reflect deep water carbonate saturation state. *Earth and Planetary Science Letters* 258, 73-86.
- ²²Bryan, S.P., Marchitto, T.M., 2008. Mg/Ca-temperature proxy in benthic foraminifera: New calibrations from the Florida Straits and a hypothesis regarding Mg/Li. *Paleoceanography* 23, PA2220.
- ²³Hathorne, E.C., James, R.H., Lampitt, R.S., 2009. Environmental versus biomineralization controls on the intratest variation in the trace element composition of the planktonic foraminifera *G. inflata* and *G. scitula*. *Paleoceanography* 24, PA4204.
- ²⁴Hall, J.M., Chan, L.H., 2014b. Li/Ca in multiple species of benthic and planktonic foraminifera: Thermocline, latitudinal, and glacial-interglacial variation. *Geochimica et Cosmochimica Acta* 68, 529-545.
- ²⁵Marriott, C.S., Henderson, G.M., Crompton, R., Staubwasser, M., Shaw, S., 2004. Effect of mineralogy, salinity, and temperature on Li/Ca and Li isotope composition of calcium carbonate. *Chemical Geology* 212, 5-15.

- ²⁶Yu, J., Day, J., Greaves, M., Elderfield, H., 2005. Determination of multiple element/ calcium ratios in foraminiferal calcite by quadrupole ICP-MS. *Geochemistry, Geophysics, Geosystems* 6, Q08P01.
- ²⁷Foster, G.L., 2008. Seawater pH, $p\text{CO}_2$ and $[\text{CO}_3^{2-}]$ variations in the Caribbean Sea over the last 130 kyr: A boron isotope and B/Ca study of planktonic foraminifera. *Earth and Planetary Science Letters* 271, 254-266.
- ²⁸Ni, Y., Foster, G.L., Bailey, T., Elliott, T., Schmidt, D.N., Pearson, P., Haley, B., Coath, C., 2007. A core top assessment of proxies for the ocean carbonate system in surface-dwelling foraminifera. *Paleoceanography* 22, PA3212.
- ²⁹Bian, N., Martin, P.A., 2010. Investigating the fidelity of Mg/Ca and other element data from reductively cleaned planktonic foraminifera. *Paleoceanography* 25, PA2215.
- ³⁰Raja, R., Saraswati, P.K., Rogers, K., Iwao, K., 2005. Magnesium and strontium compositions of recent symbiont-bearing benthic foraminifera. *Marine Micropaleontology* 58, 31-44.
- ³¹Langer, G., Sadekov, A., Thoms, S., Mewes, A., Nehrke, G., Greaves, M., Misra, S., Bijma, J., Elderfield, H., 2015. Li partitioning in the benthic foraminifer *Amphistegina lessonii*. *Geochemistry, Geophysics, Geosystems* 16, 4275-4279.
- ³²Vigier, N., Rollion-Bard, C., Levenson, Y., Erez, J., 2015. Lithium isotopes in foraminifera shells as a novel proxy for the ocean dissolved inorganic carbon (DIC). *C.R. Geoscience* 347, 43-51.
- ³³Knorr, P.O., Robbins, L.L., Harries, P.J., Hallock, P., Wynn, J., 2015. Response of the miliolid *Archaias unguulatus* to simulated ocean acidification. *Journal of Foraminiferal Research* 45, 109-127.

³⁴Delaney, M.L., Bé, A.W.H., Boyle E.A., 1985. Li, Sr, Mg and Na in foraminiferal calcite shells from laboratory culture, sediment traps, and sediment cores. *Geochimica et Cosmochimica Acta* 49, 1327-1341.