

Supplement of Biogeosciences, 14, 3445–3459, 2017
<https://doi.org/10.5194/bg-14-3445-2017-supplement>
© Author(s) 2017. This work is distributed under
the Creative Commons Attribution 3.0 License.



Supplement of

Boreal coniferous forest density leads to significant variations in soil physical and geochemical properties

Carole Bastianelli et al.

Correspondence to: Carole Bastianelli (carole.bastianelli@agroparistech.fr)

The copyright of individual parts of the supplement might differ from the CC BY 3.0 License.




Figure S1: Methods for sampling individual soil horizons



Figure S2: Cross-correlations heatmap between environment characteristics and chemical elements in B horizons. Correlations were computed on fully standardized variables using Pearson's coefficient method and tested by associated t-tests. Asterisks indicate statistical significance with *: p-value<0.05, **: p-value<0.01, ***: p-value<0.001.




Figure S3: Correlation heatmap of (a) chemical elements in the FH organic horizons and (b) cross-correlation matrix with environment characteristics. Correlations were computed on fully standardized variables using Pearson's coefficient method and tested by associated t-tests. Asterisks indicate statistical significance with *: p-value<0.05, **: p-value<0.01, ***: p-value<0.001.

Table S1: Characteristics of C horizons in mf and lw plots. lw plots = plots covered by lichen woodlands, mf plots = plots covered by moss forests. Values are given as means \pm standard deviations. Asterisks indicate statistical significance of mean comparison tests with *: p-value<0.05, **: p-value<0.01, ***: p-value<0.001.

	Horizon C			Metal species	Horizon C		
	lw	mf	p-values		lw	mf	p-values
%C	0.16 \pm 0.13	0.35 \pm 0.16	0.001***	Al _{pyro} (cmol.kg ⁻¹)	1.31 \pm 0.57	1.52 \pm 0.69	0.262
%N	0.01 \pm 0	0.01 \pm 0	0.083	Al _{oxa} (cmol.kg ⁻¹)	2.53 \pm 1.57	2.23 \pm 1.09	0.430
C:N	22.69 \pm 8.63	43.46 \pm 16.09	0.004**	Al _{dit} (cmol.kg ⁻¹)	1.93 \pm 1.26	2.04 \pm 1	0.921
CEC (cmol.kg ⁻¹)	4.17 \pm 3.63	8.64 \pm 6.91	0.032*	Al _{SRO} (cmol.kg ⁻¹)	1.23 \pm 1.02	0.71 \pm 0.71	0.077
Base saturation	0.11 \pm 0.06	0.1 \pm 0.08	0.926	Fe _{pyro} (cmol.kg ⁻¹)	0.38 \pm 0.36	0.77 \pm 0.47	0.104
P (cmol.kg ⁻¹)	53.34 \pm 33.69	21.75 \pm 12.14	0.000***	Fe _{oxa} (cmol.kg ⁻¹)	2.61 \pm 0.96	2.4 \pm 0.95	0.751
pH	5.61 \pm 0.15	5.41 \pm 0.12	0.000***	Fe _{dit} (cmol.kg ⁻¹)	2.53 \pm 1.36	3.34 \pm 1.68	0.323
				Fe _{SRO} (cmol.kg ⁻¹)	2.22 \pm 1.05	1.63 \pm 0.9	0.464
				Fe _{CRl} (cmol.kg ⁻¹)	-0.05 \pm 1.34	0.94 \pm 1.13	0.123

Table S2: Crystallinity index of mineral soil horizons in MF and LW sites. CI (Fe) accounts for iron crystallinity index of iron (Fe_{dit}:Fe_{oxa}). CI (Al) accounts for aluminium index (Al_{dit}:Al_{oxa}). Site mean comparisons were achieved by Mann-Whitney tests. Values are given as means \pm standard deviations.

Site	type	CI (Fe)	
		B horizon	C horizon
Lake Adele	LW	1.29 \pm 0.16	1.48 \pm 0.30
Lake Prisca	LW	1.09 \pm 0.11	0.65 \pm 0.21
Lake Mundi	LW	1.13 \pm 0.17	1.09 \pm 0.32
Lac des trotteurs	MF	1.09 \pm 0.11	1.05 \pm 0.26
Lake Arthur	MF	1.12 \pm 0.12	1.47 \pm 0.33
Lake Freeze	MF	1.16 \pm 0.24	1.88 \pm 0.63

Table S3: Results of the permutation test for homogeneity of multivariate dispersions of individual plot values around the centroids (MF vs LW).

Response: Distances	df	MS	F	p-value	Number of perm.
Groups	1	0.398	0.0203	0.895	999
Residuals	52	19.606			

Table S4: Stocks of Fe and Al species contained in the B horizon. Values are given as means \pm standard deviations.

Total stock at the site scale (t.ha⁻¹)		
Metal species	LW sites	MF sites
Fe _{pyro} (g.kg ⁻¹)	14.06 \pm 5.02	18.74 \pm 4.57
Fe _{oxa} (g.kg ⁻¹)	33.13 \pm 6.08	29.09 \pm 13.25
Fe _{dit} (g.kg ⁻¹)	39.12 \pm 9.79	31.83 \pm 13.60
Fe _{SRO} (g.kg ⁻¹)	19.07 \pm 4.87	10.36 \pm 8.75
Al _{pyro} (g.kg ⁻¹)	27.51 \pm 3.86	22.60 \pm 5.46
Al _{oxa} (g.kg ⁻¹)	57.93 \pm 14.51	28.18 \pm 8.45
Al _{dit} (g.kg ⁻¹)	41.95 \pm 11.55	29.63 \pm 8.32
Al _{SRO} (g.kg ⁻¹)	30.42 \pm 16.46	5.57 \pm 4.96