

Supplement of Biogeosciences, 14, 5647–5662, 2017  
<https://doi.org/10.5194/bg-14-5647-2017-supplement>  
© Author(s) 2017. This work is distributed under  
the Creative Commons Attribution 3.0 License.




*Supplement of*

## **Temporal variability of chlorophyll distribution in the Gulf of Mexico: bio-optical data from profiling floats**

**Orens Pasqueron de Fommervault et al.**

*Correspondence to:* Orens Pasqueron de Fommervault (orens@cicese.mx)


The copyright of individual parts of the supplement might differ from the CC BY 3.0 License.



**Figure S1. Basin-scale, monthly climatological mean and standard deviation of: (top) the mixed layer depth, (middle) mean surface chlorophyll concentration, (bottom) integrated content of chlorophyll over the 0-350 m layer, according a criterion based on the SSH (mean value  $\pm$  7cm). Red (blue) shows statistics for all profiles in anticyclonic (cyclonic) structures.**



**Figure S2. Float time-series of the (top) mixed layer depth, (middle) mean surface bbp concentration and (bottom) integrated content of bbp over the 0-350 m layer.**



**Figure S3.** Comparison of the mean monthly profiles of [CHL], bbp and bbp/[CHL] ratio, only for the months when the water column is well stratified. The adjusted depth (i.e., depth normalized by the depth of the DCM (Hense and Beckmann, 2008)) has been selected to account for