

Seasons	Stations	<i>T</i> (°C)	<i>S</i> (psu)	Chl <i>a</i> (µg L ⁻¹)	Copepod abundance (ind. m ⁻³)	Dominant species	% of dominant species
Winter February 2015	1	7.4	1.4	1.2	65	<i>Acartia hudsonica</i>	45
	2	9.1	8.0	2.2	551	<i>Acartia hudsonica</i>	89
	3	9.3	23.7	2.3	944	<i>Acartia omorii</i>	33
	4	8.5	28.2	1.6	1614	<i>Acartia omorii</i>	46
	5	8.4	29.9	1.0	2888	<i>Acartia omorii</i>	49
	6	9.1	27.4	0.8	2123	<i>Acartia omorii</i>	32
	7	8.7	26.9	1.3	1673	<i>Paracalanus parvus</i>	41
	8	9.0	26.9	1.1	2159	<i>Paracalanus parvus</i>	30
	9	9.0	26.1	1.0	2690	<i>Paracalanus parvus</i>	35
Spring May 2015	1	19.8	0.0	5.0	2265	<i>Pseudodiaptomus koreanus</i>	88
	2	19.8	4.7	2.0	175	<i>Pseudodiaptomus koreanus</i> ; <i>Tortanus dextrilobatus</i>	46
	3	19.0	11.2	0.9	324	<i>Acartia omorii</i>	53
	4	17.4	27.2	6.2	326	<i>Corycaeus affinis</i>	38
	5	17.0	30.1	2.2	266	<i>Corycaeus affinis</i>	52
	6	17.0	32.2	6.8	358	<i>Corycaeus affinis</i> ; <i>Calanus sinicus</i>	41
	7	18.0	32.7	5.3	148	<i>Corycaeus affinis</i>	73
	8	16.5	32.9	3.8	139	<i>Corycaeus affinis</i>	41
	9	16.5	32.8	2.7	150	<i>Acartia omorii</i>	81
Summer August 2015	1	26.8	0.4	1.0	53	<i>Tortanus dextrilobatus</i>	79
	2	27.4	10.6	4.3	3220	<i>Tortanus dextrilobatus</i>	58
	3	27.1	20.5	4.5	784	<i>Acartia ohtsukai</i>	71
	4	25.8	28.8	1.6	1401	<i>Acartia ohtsukai</i>	62
	5	23.7	32.2	2.8	366	<i>Acartia ohtsukai</i>	37
	6	23.9	32.2	2.9	129	<i>Acartia erythraea</i>	67
	7	24.1	32.3	2.3	79	<i>Labidocera rotunda</i>	60
	8	24.5	32.4	1.6	124	<i>Acartia erythraea</i>	93
	9	24.2	32.5	2.4	81	<i>Acartia erythraea</i>	55
Autumn November 2015	1	8.8	0.0	0.2	17	<i>Sinocalanus tenellus</i>	78
	2	9.9	4.7	0.1	22	<i>Paracalanus parvus</i>	32
	3	11.3	15.0	0.5	33	<i>Paracalanus parvus</i>	32
	4	12.1	20.9	0.4	32	<i>Corycaeus affinis</i>	65
	5	15.4	31.3	0.4	18	<i>Corycaeus affinis</i>	62
	6	14.6	31.3	0.4	41	<i>Corycaeus affinis</i>	55
	7	14.7	31.8	0.9	113	<i>Corycaeus affinis</i>	71
	8	14.8	32.3	1.1	118	<i>Corycaeus affinis</i>	30
	9	14.2	32.1	0.4	23	<i>Corycaeus affinis</i>	56