



Supplement of

A model based on Rock-Eval thermal analysis to quantify the size of the centennially persistent organic carbon pool in temperate soils

Lauric Cécillon et al.

Correspondence to: Lauric Cécillon (lauric.cecillon@irstea.fr)

The copyright of individual parts of the supplement might differ from the CC BY 4.0 License.

10 Table S1: Basic characteristics of the four long-term agronomic experimental sites used for the reference soil sample set. Abbreviation: *LTBF*, long-term bare fallow.

Site	Latitude, longitude	Mean annual temperature (°C)	Mean annual precipitation (mm)	Land use before experiment	Sampling depth (cm)	Soil texture (%, clay/silt/sand)
Versailles, France	48°48N, 2°08E	10.7	628	grassland	25	17/57/26
Rothamsted, United Kingdom	51°82N, 0°35E	9.5	712	grassland	23	25/62/13
Ultuna, Sweden	59°49N, 17°38E	5.5	533	arable	20	36/41/23
Grignon, France	48°51N, 1°55E	10.7	649	grassland	25	16/54/30



Figure S1: Error propagation scheme in the random forests regression model (adapted from Coulston et al. (2016) and modified).



Figure S2: Standard deviation of the CP_{SOC} proportion as a function of the CP_{SOC} proportion in the reference soil sample set (n = 5 118).

Supplementary reference

Coulston, J.W., Blinn, C.E., Thomas, V.A., and Wynne, R.H.: Approximating prediction uncertainty for random forest regression models. Photogramm. Eng. Remote Sens., 82, 189–197, 2016.