

Species	Ca content in aboveground parts			Ca content in belowground parts		
	Range (g kg <sup>-1</sup> )	Mean ± SE (g kg <sup>-1</sup> )	Correlation coefficient	Range (g kg <sup>-1</sup> )	Mean ± SE (g kg <sup>-1</sup> )	Correlation coefficient
<i>Smilax china</i>	5.77–36.35	18.5 ± 12.24	0.302	3.11–8.61	5.89 ± 2.75	0.931
<i>Aster baccharoides</i>	16.16–24.03	20.00 ± 3.60	0.418	6.20–12.02	8.91 ± 2.58	0.315
<i>Vitex negundo</i>	5.53–26.31	18.03 ± 7.44	0.198	2.83–8.17	5.59 ± 2.02	-0.116
<i>Sanguisorba officinalis</i>	17.68–27.77	24.01 ± 4.47	0.995**	13.41–40.14	32.25 ± 12.71	0.996**
<i>Themeda japonica</i>	2.15–9.23	5.51 ± 2.45	0.963**	0.42–7.91	3.88 ± 2.70	0.488
<i>Pyracantha fortuneana</i>	9.16–29.84	19.61 ± 8.46	0.240	17.08–31.86	21.43 ± 7.02	-0.189
<i>Loropetalum chinense</i>	10.33–33.44	27.25 ± 7.29	-0.203	13.62–27.69	19.69 ± 7.09	0.542
<i>Serissa japonica</i>	9.69–33.66	23.26 ± 9.90	-0.027	4.27–20.51	12.01 ± 7.81	0.838
<i>Indigofera tinctoria</i>	10.18–40.24	24.17 ± 11.49	0.215	3.39–9.83	5.98 ± 2.33	-0.289
<i>Digitaria sanguinalis</i>	4.75–9.80	6.67 ± 2.73	0.257	1.36–5.33	3.37 ± 1.98	-0.915
<i>Abelia chinensis</i>	5.07–29.64	18.08 ± 10.12	-0.163	0.87–7.12	4.10 ± 2.16	0.070
<i>Artemisia carvifolia</i>	15.34–19.39	17.37 ± 1.42	0.400	6.39–14.07	9.18 ± 3.07	0.028
<i>Glochidion puberum</i>	11.13–26.99	20.49 ± 7.04	0.357	5.33–13.64	10.45 ± 4.48	0.775
<i>Miscanthus sinensis</i>	4.34–7.60	5.61 ± 1.44	0.000	2.88–13.10	5.82 ± 4.87	0.118
<i>Rhus chinensis</i>	10.52–28.16	19.93 ± 6.43	0.076	8.92–20.38	14.13 ± 4.13	0.336
<i>Dendranthema indicum</i>	20.97–24.96	22.54 ± 1.86	0.666	2.97–7.39	5.39 ± 1.70	0.877*
<i>Castanea henryi</i>	12.99–38.74	22.4 ± 8.17	0.151	20.52–31.37	25.28 ± 3.92	0.963**