

Supplement of Biogeosciences, 15, 3975–4001, 2018  
<https://doi.org/10.5194/bg-15-3975-2018-supplement>  
© Author(s) 2018. This work is distributed under  
the Creative Commons Attribution 4.0 License.




*Supplement of*

## **A 1500-year multiproxy record of coastal hypoxia from the northern Baltic Sea indicates unprecedented deoxygenation over the 20th century**

**Sami A. Jokinen et al.**

*Correspondence to:* Sami A. Jokinen ([sami.jokinen@utu.fi](mailto:sami.jokinen@utu.fi))


The copyright of individual parts of the supplement might differ from the CC BY 4.0 License.



1

2 **Figure S1: Crossplots for Mo content against  $C_{org}$  content and Pr/Ph ratio. For Mo vs.  $C_{org}$ , a linear**  
 3 **regression for the samples post-dating the MCA is shown, while for Mo vs. Pr/Ph a regression for the**  
 4 **entire data set is presented.**


5



6

7 **Figure S2: Crossplot between C/N ratio and BIT index.**


8



9  
10 **Figure S3: Comparison between Mo/Al and raw Mo content profiles.**

11

12



13

14 **Figure S4: Near-bottom water (1 m above seabed) dissolved oxygen concentrations at the study site in**  
 15 **August from 1988 to 2015 AD (data from HERTTA database: [www.syke.fi/avoindata](http://www.syke.fi/avoindata)).** The values at  
 16 **0.1 mg L<sup>-1</sup> represent concentrations below the detection limit (0.2 mg L<sup>-1</sup>).**