

Supplement of Biogeosciences, 15, 4405–4429, 2018
<https://doi.org/10.5194/bg-15-4405-2018-supplement>
© Author(s) 2018. This work is distributed under
the Creative Commons Attribution 4.0 License.



Supplement of

Modeling seasonal and vertical habitats of planktonic foraminifera on a global scale

Kerstin Kretschmer et al.

Correspondence to: Kerstin Kretschmer (kkretschmer@marum.de)

The copyright of individual parts of the supplement might differ from the CC BY 4.0 License.

Contents of this file

5 **Figure S1** (page 2): 50-year (left panel) and 300-year (right panel) time series of the year-to-year difference (in mmol m^{-3}) of the modeled carbonate ion concentration (CO_3^{2-} ; grey), dissolved inorganic nitrate (NO_3^- ; light blue), small phytoplankton concentration (orange), zooplankton concentration (magenta), and the concentration of *N. pachyderma* (black) at the surface ocean, 105 m, 250 m, and 530 m water depth. Note that the left panel only shows a zoom for the latter three mentioned concentrations.

10 **Figure S2** (page 3): Modeled peak timing (top row) and/or modeled peak amplitude (bottom row) vs. annual mean temperature (in $^{\circ}\text{C}$) averaged over the top 55 m of the water column for (a) *N. pachyderma*, (b) *N. incompta*, (c) *G. bulloides*, (d) *G. ruber* (white), and (e) *T. sacculifer*. The color coding corresponds to latitude. Modeled peak timing is given in months and modeled peak amplitudes have been log-transformed. Note that peak timings of each species from the southern hemisphere have been transformed to northern hemisphere equivalents by adding or subtracting 6 months. For a better visualization the peak timing data has been offset along the ordinate axis to avoid that overlapping points plot on top of each other (this has been achieved by adding a small amount of white noise to the peak timing data). The
15 grey shadings in the top row panels show the data density, i.e., where most of the data points occur.

20 **Figure S3** (pages 4-14): Comparison of export planktonic foraminiferal shell fluxes in sediment traps (in $\log_{10}[\# \text{m}^{-2} \text{day}^{-1}]$; grey triangles) with the residuals (i.e., the deviation from the mean) of the volume integrated modeled biomass (in $\text{mmol C m}^{-3} \times 10^{-4}$; light blue squares). The respective location of each sediment trap is given in Table S1.

25 **Figure S4** (pages 15-26): Comparison of the vertical distribution of live specimens in plankton tows (in $\# \text{m}^{-3}$; grey bars) with modeled concentrations over depth (in mmol C m^{-3} ; light blue profiles). Dashed dark grey and blue lines indicate average living depth (in m) and vertical dispersion calculated for the plankton tows ($\text{ALD}_{\text{tow}} \pm \text{VD}_{\text{tow}}$) and PLAFOM2.0 ($\text{ALD}_{\text{mod}} \pm \text{VD}_{\text{mod}}$), respectively. The respective location of each plankton tow sample is given in Table S2.

30 **Figure S5** (page 27): Difference in potential temperature (in $^{\circ}\text{C}$) between the Community Earth System Model, version 1.2.2 (CESM1.2) and the World Ocean Atlas 2013 (WOA13; Locarnini et al., 2013) averaged over the top (a) 55 m and (b) 250 m of the water column.

Table S1 (page 28): Information on sediment trap data.

Table S2 (pages 29-30): Information on plankton tow data.

35 **Table S3** (pages 31-32): (a) Peak season (i.e., season of maximum production) and (b) peak amplitude (i.e., maximum in production divided by the annual mean) for each planktonic foraminiferal species at the locations of the sediment traps shown in Figure 1b in the main text. Empty cells indicate absence of species in either the sediment trap data or the model output.

Table S4 (page 33): Average living depths for each planktonic foraminiferal species at the locations of the plankton tows shown in Figure 1b in the main text. Empty cells indicate if species has been absent in either the plankton tow data or the model output.

40 **References** (pages 34-36)

Figure S1.




Figure S2.






Figure S3.



Site GS2





Site OG5





Site NB6/7





Site PAC50







Site PAPA







Site SA





Site KNOT





Site WCT6





Site WCT2





Site WCT7





Site WCT1





Site SBB





Site SPB




Site JGQFS34




Site L1





Site BATS





Site WAST





Site EA1





Site EA2





Site EA3





Site EA4





Site WA1






Site NCR



Site SCR



Site CP



Site WS34






Figure S4.


Station 93-36




Station PS78-25




Station PS78-44




Station PS78-75




Station PS55-025




Station PS55-043




Station PS55-063




Station MN116




Station MN2




Station MN323




Station MN314




Station PAPA




Station 101




Station 79




Station KNOT




Station #B




Station #b




Station #A




Station #E




Station POS383-165




Station POS383-175




Station POS247-1389




Station MOC1-38




Station MOC1-28




Station MOC1-23




Station 310




Station 920




Station 313




Station 917




Station MOC63




Station MOC65




Station MOC12




Station MOC66




Station MOC15




Station MOC69




Station MOC20




Station MOC71




Station MOC72




Station SO225-21-3




Station TNO57-16




Station TNO57-13




Station AN98-O



Station AN99-O



Station AN00-O



Station AN01-O




Figure S5.




Table S1.

#	Site	Latitude (°N)	Longitude (°E)	Water Depth (m)	Trap Depth (m)	Deployment Time (day/month/year)	Duration (days)	Species	Fraction (μm)	Source
1	GS2	75.00	0.00	3720	300	03/06/1994 to 11/05/1995	342	Np	63-500	Jensen (1998)
2	OG5	72.40	-7.70	2624	500	06/08/1991 to 10/07/1992	339	Np	63-500	Jensen (1998)
3	NB6/7	69.69	0.47	3273	500	06/08/1991 to 02/10/1993	780	Np, Ni	63-500	Jensen (1998)
4	PAC50	50.01	165.03	5570	3260	01/12/1997 to 10/06/2001	1091	Np, Ni, Gb	>125	Kuroyanagi et al. (2002)
5	PAPA	50.00	-145.00	4240	3800	23/09/1982 to 30/08/1986	1122	Np, Ni, Gb	>125	Sautter and Thunell (1989)
6	SA	49.00	-174.00	5406	4812	23/08/1990 to 03/08/1999	2702	Np, Gb	>125	Asahi and Takahashi (2007)
7	KNOT	43.97	155.06	5370	2957	01/12/1997 to 12/05/2000	799	Np, Ni, Gb, Gr	>125	Kuroyanagi et al. (2002)
8	WCT6	42.00	155.34	5578	1091	15/08/1999 to 31/08/2000	382	Np, Ni, Gb, Gr	>125	Mohiuddin et al. (2005)
9	WCT2	39.00	147.00	5356-5322	1371; 1586	19/11/1997 to 10/08/1999	608	Ni, Gb, Gr	>125	Mohiuddin et al. (2002)
10	WCT7	36.68	154.94	5578	5034	19/08/1999 to 29/08/2000	376	Gb, Ts	>125	Mohiuddin et al. (2004)
11	WCT1	25.00	136.99	4905-5308	917; 1388	07/12/1997 to 12/08/1999	560	Ni, Gr, Ts	>125	Mohiuddin et al. (2002)
12	SBB	34.23	-120.03	650	590; 470	12/08/1993 to 26/06/1999	1015	Ni, Gb, Gr	>125	Kincaid et al. (2000) Darling et al. (2003)
13	SPB	33.55	-118.50	880	500	07/01/1988 to 26/07/1988	199	Ni, Gb, Gr	>125	Sautter and Thunell (1991)
14	JGOFS34	34.00	-21.00	n.a.	2000	03/04/1989 to 16/04/1990	378	Ni, Gb, Ts	>150	Wolfteich (1994)
15	L1	33.00	-22.00	5300	3000	24/02/2002 to 01/04/2004	764	Ni, Gb, Gr, Ts	>125	Storz et al. (2009)
16	BATS	32.08	-64.25	4200	3200	06/04/1978 to 17/05/1984	1848	Gb, Gr, Ts	>125	Deuser et al. (1981) Deuser and Ross (1989)
17	WAST	16.32	60.47	4016	3026	10/05/1986 to 21/10/1987	506	Gb, Gr, Ts	>150	Curry et al. (1992)
18	EA1	3.17	-11.25	4524	984	13/04/1991 to 29/11/1991	230	Gb, Gr, Ts	>150	Fischer and Wefer (1996)
19	EA2	1.78	-11.25	4399	953	13/04/1991 to 29/11/1991	230	Gb, Gr, Ts	>150	Fischer and Wefer (1996)
20	EA3	0.08	-10.77	4141	1097	13/04/1991 to 29/11/1991	230	Gb, Gr, Ts	>150	Fischer and Wefer (1996)
21	EA4	-2.19	-10.09	3906	1068	13/04/1991 to 29/11/1991	230	Gb, Gr, Ts	>150	Fischer and Wefer (1996)
22	WA1	-4.00	-25.57	5530	652	17/10/1992 to 21/03/1993	155	Gr, Ts	>150	Fischer and Wefer (1996)
23	NCR	-42.70	178.63	1500	1000	14/09/1996 to 15/05/1997	243	Np, Ni, Gb, Gr, Ts	>150	King and Howard (2001)
24	SCR	-44.62	178.62	1500	1000	09/06/1996 to 15/05/1997	340	Np, Ni, Gb, Gr, Ts	>150	King and Howard (2001)
25	CP	-52.62	174.15	n.a.	442; 362	14/05/1998 to 13/07/1999	368	Np, Ni, Gb	>150	Northcote and Neil (2005)
26	WS34	-64.90	-2.60	5053	360	16/01/1988 to 26/02/1990	745	Np	>125	Donner and Wefer (1994)

Np – *N. pachyderma*; Ni – *N. incompta*; Gb – *G. bulloides*; Gr – *G. ruber* (white); Ts – *T. sacculifer*
n.a. – not available (i.e., not given in data set)

Table S2.

#	Station	Latitude (°N)	Longitude (°E)	Water Depth (m)	Depth Intervals	Date (season)	Species	Fraction (μm)	Source
1	93-36	80.36	-10.14	n.a.	0-20, 20-40, 40-60, 60-100, 100-150, 150-200, 200-250, 250-275	27/07/1993 (summer)	Np	n.a.	Kohfeld et al. (1996)
2	PS78-25	78.83	7.00	1465	0-50, 50-100, 100-200, 200-300, 300-500	26/06/2011 (summer)	Np, Ni, Gb	100-250	Pados and Spielhagen (2014)
3	PS78-44	78.83	0.08	2636	0-50, 50-100, 100-200, 200-300, 300-500	29/06/2011 (summer)	Np, Ni, Gb	100-250	Pados and Spielhagen (2014)
4	PS78-75	78.83	-3.92	1978	0-50, 50-100, 100-200, 200-300, 300-500	04/07/2011 (summer)	Np, Ni, Gb	100-250	Pados and Spielhagen (2014)
5	PS55-025	75.00	-10.58	3084	0-50, 50-100, 100-200, 200-300, 300-500	11/07/1999 (summer)	Np, Ni, Gb	125-250	Stangeew (2001)
6	PS55-043	75.00	0.36	3789	0-50, 50-100, 100-200, 200-300, 300-500	14/07/1999 (summer)	Np, Ni, Gb	125-250	Stangeew (2001)
7	PS55-063	75.00	10.65	2542	0-50, 50-100, 100-200, 200-300, 300-500	16/07/1999 (summer)	Np, Ni, Gb	125-250	Stangeew (2001)
8	MN116	75.00	-7.31	3393	0-50, 50-100, 100-150, 150-300	21/08/1994 (summer)	Np	125-250	Simstich et al. (2003)
9	MN2	70.00	3.40	3261	0-50, 50-100, 100-500, 500-1000, 1000-2000	10/07/1994 (summer)	Np	125-250	Simstich et al. (2003)
10	MN323	69.69	0.47	3290	0-20, 20-40, 40-60, 60-80, 80-100, 100-200, 200-300, 300-500, 500-700, 700-1000, 1000-1500, 1500-2000, 2000-2500	07/07/1992 (summer)	Np	125-250	Simstich et al. (2003)
11	MN314	67.54	5.58	1438	0-20, 20-40, 40-60, 60-80, 80-100, 100-200, 200-300, 300-500, 500-700	28/06/1992 (summer)	Np	125-250	Simstich et al. (2003)
12	PAPA	49.98	-144.97	4253	0-50, 50-100, 100-150, 150-200, 200-300	16/08/2015 (summer)	Np, Ni, Gb	> 100	Iwasaki et al. (2017)
13	101	47.00	-174.95	5790	0-50, 50-100, 100-150, 150-200, 200-300	05/08/2015 (summer)	Np, Ni, Gb	> 100	Iwasaki et al. (2017)
14	79	46.98	166.73	5957	0-50, 50-100, 100-150, 150-200, 200-300	31/07/2015 (summer)	Np, Ni, Gb	> 100	Iwasaki et al. (2017)
15	KNOT	44.08	154.98	5335	0-50, 50-100, 100-150, 150-200, 200-300	25/07/2015 (summer)	Np, Ni, Gb	> 100	Iwasaki et al. (2017)
16	#B	41.57	141.90	1000	0-20, 20-40, 40-60, 60-80, 80-120, 120-160, 160-200	03/06/2002 (summer)	Np, Ni, Gb, Gr, Ts	125-1000	Kuroyanagi and Kawahata (2004)
17	#b	41.15	143.38	2077	0-20, 20-40, 40-60, 60-80, 80-120, 120-160, 160-200	04/06/2002 (summer)	Np, Ni, Gb, Gr, Ts	125-1000	Kuroyanagi and Kawahata (2004)
18	#A	36.02	141.78	2220	0-20, 20-40, 40-60, 60-80, 80-120, 120-160, 160-200	25/05/2002 (spring)	Np, Ni, Gb, Gr, Ts	125-1000	Kuroyanagi and Kawahata (2004)
19	#E	32.17	133.88	2660	0-20, 20-40, 40-60, 60-80, 80-120, 120-160, 160-200	27/05/2002 (spring)	Np, Ni, Gb, Gr, Ts	125-1000	Kuroyanagi and Kawahata (2004)
20	POS383-165	34.00	-22.00	5288	0-100, 100-200, 200-300, 300-500, 500-700	23/04/2009 (spring)	Np, Ni, Gb, Gr, Ts	> 100	Rebotim et al. (2017)
21	POS383-175	33.15	-22.00	5232	0-100, 100-200, 200-300, 300-500, 500-700	26/04/2009 (spring)	Np, Ni, Gb, Gr, Ts	> 100	Rebotim et al. (2017)
22	POS247-1389	33.08	-22.00	5226	0-20, 20-40, 40-60, 60-80, 80-100, 100-200, 200-300, 300-500, 500-700	24/01/1999 (winter)	Np, Ni, Gb, Gr, Ts	> 100	Rebotim et al. (2017)
23	MOC1-38	38.92	-67.90	n.a.	0-25, 25-50, 50-75, 75-100, 100-125, 125-150, 150-175, 175-200	11/1975 (fall)	Gb, Gr, Ts	n.a.	Fairbanks et al. (1980)
24	MOC1-28	33.91	-71.78	n.a.	0-25, 25-50, 50-75, 75-100, 100-125, 125-150, 150-175, 175-200	11/1975 (fall)	Gr, Ts	n.a.	Fairbanks et al. (1980)
25	MOC1-23	32.73	-71.16	n.a.	0-25, 25-50, 50-75, 75-100, 100-125, 125-150, 150-175, 175-200	11/1975 (fall)	Gr, Ts	n.a.	Fairbanks et al. (1980)
26	310	16.02	52.73	n.a.	0-10, 10-25, 25-50, 50-75, 75-100, 100-150, 150-200, 200-300, 300-500	20/08/1992 (summer)	Np, Gb, Gr, Ts	> 125	Peeters and Brummer (2002)
27	920	16.09	52.70	n.a.	0-10, 10-25, 25-50, 50-75, 75-100, 100-150, 150-200, 200-300, 300-500	27/02/1993 (winter)	Np, Gb, Gr, Ts	> 125	Peeters and Brummer (2002)
28	313	15.91	53.02	n.a.	0-10, 10-25, 25-50, 50-75, 75-100, 100-150, 150-200, 200-300, 300-500	21/08/1992 (summer)	Np, Gb, Gr, Ts	> 125	Peeters and Brummer (2002)
29	917	15.89	52.97	n.a.	0-10, 10-25, 25-50, 50-75, 75-100, 100-150, 150-200, 200-300, 300-500	25/02/1993 (winter)	Np, Gb, Gr, Ts	> 125	Peeters and Brummer (2002)
30	MOC63	2.92	-140.20	n.a.	0-10, 10-20, 20-40, 40-60,	22/08/1992	Gb, Gr, Ts	> 150	Watkins et al. (1998)

Continued on next page

#	Station	Latitude (°N)	Longitude (°E)	Water Depth (m)	Depth Intervals	Date (season)	Species	Fraction (μm)	Source
					60-80, 80-100, 100-150, 150-200	(summer)			
31	MOC65	2.05	-141.49	n.a.	10-20, 40-60, 60-80, 80-100, 100-150, 150-200	26/08/1992 (summer)	Gb, Gr, Ts	> 150	Watkins et al. (1998)
32	MOC12	2.01	-139.88	n.a.	0-5, 5-10, 10-30, 30-50, 50-70, 70-90, 90-145, 145-200	17/02/1992 (winter)	Gb, Gr, Ts	> 150	Watkins et al. (1996)
33	MOC66	1.13	-140.01	n.a.	0-10, 10-20, 20-40, 40-60, 60-80, 80-100, 100-150, 150-200	27/08/1992 (summer)	Gb, Gr, Ts	> 150	Watkins et al. (1998)
34	MOC15	0.00	-140.07	n.a.	0-10, 10-20, 20-40, 40-60, 60-100, 100-150	23/02/1992 (winter)	Gr, Ts	> 150	Watkins et al. (1996)
35	MOC69	-1.05	-139.97	n.a.	0-10, 10-20, 20-40, 40-60, 60-80, 100-150, 150-200	01/09/1992 (fall)	Gb, Gr, Ts	> 150	Watkins et al. (1998)
36	MOC20	-2.02	-140.16	n.a.	0-10, 10-20, 20-40, 40-60, 60-80, 80-100, 100-150, 150-200	29/02/1992 (winter)	Gr, Ts	> 150	Watkins et al. (1996)
37	MOC71	-2.33	-140.32	n.a.	0-10, 10-20, 20-40, 40-60, 60-80, 80-100, 100-150, 150-200	04/09/1992 (fall)	Gb, Gr, Ts	> 150	Watkins et al. (1998)
38	MOC72	-3.21	-140.25	n.a.	0-10, 10-20, 20-40, 40-60, 60-80, 80-100, 100-150, 150-200	06/09/1992 (fall)	Gb, Gr, Ts	> 150	Watkins et al. (1998)
39	SO225-21-3	-3.05	-165.06	5188	0-50, 50-100, 100-200, 200-300, 300-500	08/12/2012 (winter)	Gb, Gr, Ts	> 150	Rippert et al. (2016)
40	TNO57-16	-50.12	5.75	3761	0-30, 30-60, 60-75, 75-85, 85-125, 125-235, 235-300, 300-440	24/02/1996 (winter)	Np, Ni, Gb	> 150	Mortyn and Charles (2003)
41	TNO57-13	-53.18	5.13	2851	0-25, 25-50, 50-75, 75-100, 100-150, 150-200, 200-300, 300-400, 400-500	21/02/1996 (winter)	Np, Ni, Gb	> 150	Mortyn and Charles (2003)
42	AN98/O	-63.25	177.25	4100	0-50, 50-90, 90-130	20/01/1998 (winter)	Np, Ni, Gb	> 100	Bergami et al. (2009)
43	AN99/O	-63.40	178.05	4074	0-35, 35-70	09/01/1999 (winter)	Np, Ni, Gb	> 100	Bergami et al. (2009)
44	AN00/O	-63.53	178.38	3548	0-60, 60-150, 150-220	11/01/2000 (winter)	Np, Ni, Gb	> 100	Bergami et al. (2009)
45	AN01/O	-63.43	178.10	3964	0-90, 90-150	09/01/2001 (winter)	Np, Ni, Gb	> 100	Bergami et al. (2009)

Np – *N. pachyderma*; Ni – *N. incompta*; Gb – *G. bulloides*; Gr – *G. ruber* (white); Ts – *T. sacculifer*
n.a. – not available (i.e., not given in data set)
Here the season refers to those of the Northern Hemisphere.

Table S3a.

Province	Sediment Trap Details			<i>N. pachyderma</i>		<i>N. incompta</i>		<i>G. bulloides</i>		<i>G. ruber</i> (white)		<i>T. sacculifer</i>	
	Site	Latitude (°N)	Longitude (°E)	Trap	PLAFOM2.0	Trap	PLAFOM2.0	Trap	PLAFOM2.0	Trap	PLAFOM2.0	Trap	PLAFOM2.0
Polar	GS2	75.00	0.00	Jun-Sep	Jul-Sep	-	-	-	-	-	-	-	-
	OG5	72.40	-7.70	Aug-Sep	Jul-Sep	-	-	-	-	-	-	-	-
	NB6/7	69.69	-0.47	Jun-Nov	Apr-Jun	Jul-Nov	Jul-Oct	-	-	-	-	-	-
Subpolar	PAC50	50.01	165.03	Mar-May Sep-Nov	Feb-May Jul-Sep	Mar-May	Aug-Nov	Apr-Jun	Aug-Nov	-	-	-	-
	PAPA	50.00	-145.00	Mar-May Nov-Dec	Mar-May	Apr-Jun	Aug-Nov	Apr-Jun Nov-Dec	Apr-May	-	-	-	-
	SA	49.00	-174.00	Apr-Jul Aug-Oct	Mar-May	-	-	Apr-May Jul-Sep	Apr-Jun	-	-	-	-
Transitional	KNOT	43.97	155.06	Mar-May	Mar-May	none	Jul-Oct	Mar-May Oct-Dec	Apr-Jun	Sep-Nov	Aug-Oct	-	-
	WCT6	42.00	155.34	Mar-May Jul-Sep	Feb-May	Jun-Oct	Mar-May	Mar-May Oct-Nov	Mar-May	Sep-Nov	Aug-Oct	-	-
	WCT2	39.00	147.00	-	-	none	Jan-Apr	none	Jan-Apr	Sep-Dec	Jul-Oct	-	-
	WCT7	36.68	154.94	-	-	-	-	May-Jun	Mar-May	-	-	Aug-Dec	Aug-Nov
Subtropics	SBB ^a	34.23	-120.03	-	-	Aug-Oct	Jul-Sep	Jun-Aug	Jul-Sep	Jul-Nov	Aug-Nov	-	-
	JGOF34	34.00	-21.00	-	-	Feb-Apr Aug-Sep	May-Aug	Mar-May	May-Sep	-	-	Apr-Jun	Jul-Oct
	SPB	33.55	-118.50	-	-	Mar-May	May-Sep	May-Jun	May-Sep	Apr-Jul	Aug-Nov	-	-
	L1	33.00	-22.00	-	-	Feb-Apr	May-Sep	Mar-May	May-Sep	Mar-May	Jul-Nov	Mar-May	Jul-Oct
	BATS	32.08	-64.25	-	-	-	-	Jan-May	Feb	none	May-Jul	May-Jul	Jul-Nov
WCT1	25.00	136.99	-	-	none	Feb-Jun	-	May-Sep	Jun-Aug	Jan-Mar	Jun-Aug	Jun-Oct	
Tropics	WAST	16.32	60.47	-	-	-	-	Jan-Feb Jul-Oct	Mar-May Oct-Nov	Dec-Feb May-Aug	Jan-Mar Aug-Sep	Jun-Sep	Apr-May
	EA1	3.17	-11.25	-	-	-	-	Sep-Nov	Feb-Apr	none	Jul-Sep	none	Jun-Aug
	EA2	1.78	-11.25	-	-	-	-	Jul-Oct	Mar-Apr	none	Jul-Nov	May-Jul	Jan-Feb
	EA3	0.08	-10.77	-	-	-	-	Jul-Sep	Aug-Sep	May-Jun	Sep-Jan	May-Jun	Sep-Nov
	EA4 WA1	-2.19 -4.00	-10.09 -25.57	-	-	-	-	Apr-May	Feb-Apr	Apr-Aug none	Aug-Dec Aug-Dec	Apr-Aug none	Dec-May Aug-Dec
Transitional	NCR	-42.70	178.63	Sep-Dec	Apr-May Aug-Nov	Sep-Oct	Jan-Feb Aug-Oct	Apr-May Sep-Nov	Aug-Oct	none	Jan-Apr	Oct-Dec	Jan-Apr
	SCR	-44.62	178.62	Sep-Oct	Aug-Oct	Feb-Apr Sep-Nov	Aug-Sep Dec-Feb	Sep-Oct	Aug-Oct Dec-Jan	-	-	-	-
Sub-polar	CP	-52.62	174.15	Mar-May	Aug-Nov	Mar-May	Feb-Apr Aug-Oct	Nov-May	Feb-Mar Aug-Oct	-	-	-	-
Polar	WS34	-64.90	-2.60	Mar-May	Feb-Apr Oct-Dec	-	-	-	-	-	-	-	-

^a The nearest model grid point for site SBB fell onto land. Therefore, we used the nearest model grid point in the ocean to perform a consistent model-data-comparison.

Table S3b.

Province	Sediment Trap Details			<i>N. pachyderma</i>		<i>N. incompta</i>		<i>G. bulloides</i>		<i>G. ruber</i> (white)		<i>T. sacculifer</i>	
	Site	Latitude (°N)	Longitude (°E)	Trap	PLAFOM2.0	Trap	PLAFOM2.0	Trap	PLAFOM2.0	Trap	PLAFOM2.0	Trap	PLAFOM2.0
Polar	GS2	75.00	0.00	0.78	0.26	-	-	-	-	-	-	-	-
	OG5	72.40	-7.70	0.64	0.19	-	-	-	-	-	-	-	-
	NB6/7	69.69	-0.47	0.80	0.42	0.95	0.57	-	-	-	-	-	-
Sub-polar	PAC50	50.01	165.03	0.75	0.20	0.77	0.36	0.66	0.36	-	-	-	-
	PAPA	50.00	-145.00	1.07	0.28	1.20	0.04	1.10	0.22	-	-	-	-
	SA	49.00	-174.00	0.95	0.29	-	-	0.94	0.21	-	-	-	-
Transitional	KNOT	43.97	155.06	0.69	0.36	0.79	0.16	0.66	0.33	0.76	0.70	-	-
	WCT6	42.00	155.34	0.42	0.33	0.47	0.15	0.62	0.25	0.73	0.59	-	-
	WCT2	39.00	147.00	-	-	0.64	0.17	0.69	0.22	0.74	0.44	-	-
	WCT7	36.68	154.94	-	-	-	-	0.55	0.19	-	-	0.57	0.34
Subtropics	SBB ^a	34.23	-120.03	-	-	0.68	0.04	0.67	0.06	0.87	0.19	-	-
	JGOF34	34.00	-21.00	-	-	0.85	0.05	0.76	0.07	-	-	0.69	0.14
	SPB	33.55	-118.50	-	-	0.73	0.04	0.85	0.06	0.77	0.18	-	-
	L1	33.00	-22.00	-	-	1.28	0.05	0.91	0.06	0.70	0.09	0.59	0.13
	BATS	32.08	-64.25	-	-	-	-	0.72	0.05	0.37	0.07	0.96	0.12
	WCT1	25.00	136.99	-	-	0.42	0.06	-	-	0.77	0.10	0.88	0.07
Tropics	WAST	16.32	60.47	-	-	-	-	0.77	0.02	0.70	0.14	0.66	0.02
	EA1	3.17	-11.25	-	-	-	-	0.48	0.03	0.36	0.15	0.47	0.07
	EA2	1.78	-11.25	-	-	-	-	0.52	0.02	0.33	0.13	0.59	0.03
	EA3	0.08	-10.77	-	-	-	-	0.81	0.02	0.57	0.10	0.47	0.05
	EA4	-2.19	-10.09	-	-	-	-	0.83	0.02	0.60	0.10	0.50	0.03
	WA1	-4.00	-25.57	-	-	-	-	-	-	0.67	0.15	0.58	0.06
Transitional	NCR	-42.70	178.63	0.91	0.47	0.78	0.11	0.73	0.16	0.77	0.46	0.85	0.54
	SCR	-44.62	178.62	0.87	0.14	0.57	0.08	0.89	0.16	-	-	-	-
Sub-polar	CP	-52.62	174.15	1.29	0.28	1.12	0.08	1.10	0.11	-	-	-	-
Polar	WS34	-64.90	-2.60	1.12	0.09	-	-	-	-	-	-	-	-

^a The nearest model grid point for site SBB fell onto land. Therefore, we used the nearest model grid point in the ocean to perform a consistent model-data-comparison.

Table S4.

Province	Plankton Tow Details			<i>N. pachyderma</i>		<i>N. incompta</i>		<i>G. bulloides</i>		<i>G. ruber</i> (white)		<i>T. sacculifer</i>	
	Site	Latitude (°N)	Longitude (°E)	Tow ^a (m)	PLAFOM2.0 ^a (m)	Tow ^a (m)	PLAFOM2.0 ^a (m)	Tow ^a (m)	PLAFOM2.0 ^a (m)	Tow ^a (m)	PLAFOM2.0 ^a (m)	Tow ^a (m)	PLAFOM2.0 ^a (m)
Polar	93-36	80.36	-10.14	85±35	55±35	-	-	-	-	-	-	-	-
	PS78-25	78.83	7.00	85±55	50±35	-	-	-	-	-	-	-	-
	PS78-44	78.83	0.08	80±40	45±35	-	-	-	-	-	-	-	-
	PS78-75	78.83	-3.92	70±40	45±35	-	-	-	-	-	-	-	-
	PS55-025	75.00	-10.58	90±70	45±35	-	-	-	-	-	-	-	-
	PS55-043	75.00	0.36	60±40	45±30	-	-	-	-	-	-	-	-
	PS55-063	75.00	10.65	85±65	45±30	-	-	55±25	25±15	-	-	-	-
	MN116	75.00	-7.31	150±40	45±30	-	-	-	-	-	-	-	-
	MN2	70.00	3.40	170±215	60±50	-	-	-	-	-	-	-	-
	MN323	69.69	0.47	140±155	55±45	-	-	-	-	-	-	-	-
	MN314	67.54	5.58	125±60	65±55	-	-	-	-	-	-	-	-
Sub-polar	PAPA	49.98	-144.97	230±30	80±35	105±85	60±35	60±45	70±40	-	-	-	-
	101	47.00	-174.95	95±50	60±40	140±40	55±35	65±30	50±35	-	-	-	-
	79	46.98	166.73	110±55	55±40	150±50	25±15	70±35	30±25	-	-	-	-
Transitional	KNOT	44.08	154.98	90±45	70±45	75±55	20±10	45±30	30±25	-	-	-	-
	#B ^b	41.57	141.90	125±40	80±40	105±45	70±45	100±55	75±45	122±40	5±5	-	-
	#b	41.15	143.38	85±40	80±40	35±25	70±45	40±30	75±45	-	-	-	-
	MOC1-38 #A	38.92 36.02	-67.90 141.78	- -	- -	- 25±20	- 65±35	- 65±35	65±55 25±25	75±40 65±35	30±20 20±20	35±20 25±20	35±20 25±20
Subtropics	POS383-165	34.00	-22.00	-	-	85±60	80±45	170±70	75±40	65±25	30±25	185±85	50±30
	MOC1-28	33.91	-71.78	-	-	-	-	-	-	80±35	50±30	60±35	50±30
	POS383-175	33.15	-22.00	-	-	85±55	80±40	95±50	75±40	65±25	35±25	190±65	50±30
	POS247-1389	33.08	-22.00	-	-	30±0	80±45	30±0	80±45	55±25	50±30	40±35	55±30
	MOC1-23	32.73	-71.16	-	-	-	-	140±0	70±35	115±0	45±25	95±50	50±30
	#E	32.17	133.88	-	-	60±30	80±40	-	-	70±45	30±20	45±40	35±25
Tropics	920	16.09	52.70	-	-	-	-	65±65	70±40	40±45	40±20	20±20	45±25
	310	16.02	52.73	-	-	-	-	30±35	50±35	180±100	10±5	180±85	15±10
	313	15.91	53.02	-	-	-	-	70±80	50±35	30±35	10±5	30±30	10±5
	917	15.89	52.97	-	-	-	-	90±65	70±40	75±50	40±20	20±15	45±25
	MOC63	2.92	-140.20	-	-	-	-	15±10	95±50	20±15	45±25	25±15	45±25
	MOC65	2.05	-141.49	-	-	-	-	25±15	90±50	35±25	40±20	25±15	40±25
	MOC12	2.01	-139.88	-	-	-	-	65±25	80±45	45±25	45±25	45±25	40±25
	MOC66	1.13	-140.01	-	-	-	-	55±25	85±45	45±20	35±20	45±25	35±20
	MOC15	0.00	-140.07	-	-	-	-	-	-	20±10	35±20	25±15	35±20
	MOC69	-1.05	-139.97	-	-	-	-	25±15	80±45	25±15	20±15	25±15	40±20
	MOC20	-2.02	-140.16	-	-	-	-	-	-	35±15	45±25	40±20	40±25
	MOC71	-2.33	-140.32	-	-	-	-	45±25	90±45	35±25	45±25	35±25	45±25
	SO225-21-3	-3.05	-165.06	-	-	-	-	145±90	55±30	65±35	55±30	75±45	55±30
	MOC72	-3.21	-140.25	-	-	-	-	40±20	90±50	35±15	50±25	35±20	50±25
Sub-polar	TNO57-16	-50.12	5.75	70±10	60±35	70±10	25±15	80±5	30±15	-	-	-	-
	TNO57-13	-53.18	5.13	85±60	55±35	-	-	-	-	-	-	-	-
Polar	AN98-O	-63.25	177.25	55±30	60±35	-	-	-	-	-	-	-	-
	AN99-O	-63.40	178.05	25±15	60±40	-	-	-	-	-	-	-	-
	AN01-O	-63.43	178.10	120±0	60±40	-	-	-	-	-	-	-	-
	AN00-O	-63.53	178.38	95±50	60±40	-	-	-	-	-	-	-	-

^aALD±VD (in m) of the planktonic foraminiferal species calculated after Rebotim et al. (2017) for the plankton tow samples and for PLAFOM2.0 (obtained at the nearest model grid points of the given plankton tow locations). Note that the values have been rounded to the nearest 5 m.

^b The nearest model grid point for site #B fell onto the shelf. Therefore, we used the nearest model grid point in the open ocean to perform a consistent model-data-comparison.

References

- Asahi, H. and Takahashi, K.: A 9-year time-series of planktonic foraminifer fluxes and environmental change in the Bering sea and the central subarctic Pacific Ocean, 1990-1999, *Progress in Oceanography*, 72, 343–363, <https://doi.org/10.1016/j.pocean.2006.03.021>, 2007.
- Bergami, C., Capotondi, L., Langone, L., Giglio, F., and Ravaoli, M.: Distribution of living planktonic foraminifera in the Ross Sea and the Pacific sector of the Southern Ocean (Antarctica), *Marine Micropaleontology*, 73, 37–48, <https://doi.org/10.1016/j.marmicro.2009.06.007>, 2009.
- Curry, W. B., Ostermann, D. R., Guptha, M. V. S., and Ittekkot, V.: Foraminiferal production and monsoonal upwelling in the Arabian Sea: evidence from sediment traps, Geological Society, London, Special Publications, 64, 93–106, <https://doi.org/10.1144/GSL.SP.1992.064.01.06>, 1992.
- 10 Darling, K. F., Kucera, M., Wade, C. M., von Langen, P., and Pak, D.: Seasonal distribution of genetic types of planktonic foraminifer morphospecies in the Santa Barbara Channel and its paleoceanographic implications, *Paleoceanography*, 18, 1032, <https://doi.org/10.1029/2001PA000723>, 2003.
- Deuser, W. G. and Ross, E. H.: Seasonally abundant planktonic foraminifera of the Sargasso Sea; succession, deep-water fluxes, isotopic compositions, and paleoceanographic implications, *Journal of Foraminiferal Research*, 19, 268–293, 1989.
- 15 Deuser, W. G., Ross, E. H., Hemleben, C., and Spindler, M.: Seasonal changes in species composition, numbers, mass, size, and isotopic composition of planktonic foraminifera settling into the deep Sargasso Sea, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 33, 103–127, [https://doi.org/10.1016/0031-0182\(81\)90034-1](https://doi.org/10.1016/0031-0182(81)90034-1), 1981.
- Donner, B. and Wefer, G.: Flux and stable isotope composition of *Neogloboquadrina pachyderma* and other planktonic foraminifers in the Southern Ocean (Atlantic sector), *Deep-Sea Research I*, 41, 1733–1743, 1994.
- 20 Fairbanks, R. G., Wiebe, P. H., and Bé, A. W. H.: Vertical Distribution and Isotopic Composition of Living Planktonic Foraminifera in the Western North Atlantic, *Science*, 207, 61–63, 1980.
- Fischer, G. and Wefer, G.: Long-term observation of particle fluxes in the eastern Atlantic: seasonality, changes of flux with depth and comparison with the sediment record, in: *The South Atlantic: Present and Past Circulation*, edited by Wefer, G., Berger, W. H., Siedler, G., and Webb, D., pp. 325–344, Springer, Berlin, Heidelberg, New York, 1996.
- 25 Iwasaki, S., Kimoto, K., Kuroyanagi, A., and Kawahata, H.: Horizontal and vertical distributions of planktic foraminifera in the subarctic Pacific, *Marine Micropaleontology*, 130, 1–14, <https://doi.org/10.1016/j.marmicro.2016.12.001>, 2017.
- Jensen, S.: Planktische Foraminiferen im Europäischen Nordmeer: Verbreitung und Vertikalfluß sowie ihre Entwicklung während der letzten 15000 Jahre, *Berichte Sonderforschungsbereich 313, Univ. Kiel*, 75, 1–105, 1998.
- Kincaid, E., Thunell, R. C., Le, J., Lange, C. B., Weinheimer, A. L., and Reid, F. M. H.: Planktonic foraminiferal fluxes in the Santa Barbara Basin: response to seasonal and interannual hydrographic changes, *Deep-Sea Research Part II*, 47, 1157–1176, [https://doi.org/10.1016/S0967-0645\(99\)00140-X](https://doi.org/10.1016/S0967-0645(99)00140-X), 2000.
- 30 King, A. L. and Howard, W. R.: Seasonality of foraminiferal flux in sediment traps at Chatham rise, SW Pacific: Implications for paleotemperature estimates, *Deep-Sea Research Part I*, 48, 1687–1708, [https://doi.org/10.1016/S0967-0637\(00\)00106-0](https://doi.org/10.1016/S0967-0637(00)00106-0), 2001.
- Kohfeld, K. E., Fairbanks, R. G., Smith, S. L., and Walsh, I. D.: *Neogloboquadrina pachyderma* (sinistral coiling) as paleoceanographic tracers in polar oceans: Evidence from Northeast Water Polynya plankton tows, sediment traps, and surface sediments, *Paleoceanography*, 11, 679–699, 1996.
- 35 Kuroyanagi, A. and Kawahata, H.: Vertical distribution of living planktonic foraminifera in the seas around Japan, *Marine Micropaleontology*, 53, 173–196, <https://doi.org/10.1016/j.marmicro.2004.06.001>, 2004.
- Kuroyanagi, A., Kawahata, H., Nishi, H., and Honda, M. C.: Seasonal changes in planktonic foraminifera in the northwestern North Pacific Ocean: sediment trap experiments from subarctic and subtropical gyres, *Deep-Sea Research Part II*, 49, 5627–5645, [https://doi.org/10.1016/S0967-0645\(02\)00202-3](https://doi.org/10.1016/S0967-0645(02)00202-3), 2002.
- 40

- Locarnini, R. A., Mishonov, A. V., Antonov, J. I., Boyer, T. P., Garcia, H. E., Baranova, O. K., Zweng, M. M., Paver, C. R., Reagan, J. R., Johnson, D. R., Hamilton, M., and Seidov, D.: World Ocean Atlas 2013, Volume 1: Temperature, Silver Spring, MD, <https://doi.org/10.1182/blood-2011-06-357442>, 2013.
- 5 Mohiuddin, M. M., Nishimura, A., Tanaka, Y., and Shimamoto, A.: Regional and interannual productivity of biogenic components and planktonic foraminiferal fluxes in the northwestern Pacific Basin, *Marine Micropaleontology*, 45, 57–82, [https://doi.org/10.1016/S0377-8398\(01\)00045-7](https://doi.org/10.1016/S0377-8398(01)00045-7), 2002.
- Mohiuddin, M. M., Nishimura, A., Tanaka, Y., and Shimamoto, A.: Seasonality of biogenic particle and planktonic foraminifera fluxes: Response to hydrographic variability in the Kuroshio Extension, northwestern Pacific Ocean, *Deep-Sea Research Part I*, 51, 1659–1683, <https://doi.org/10.1016/j.dsr.2004.06.002>, 2004.
- 10 Mohiuddin, M. M., Nishimura, A., and Tanaka, Y.: Seasonal succession, vertical distribution, and dissolution of planktonic foraminifera along the Subarctic Front: Implications for paleoceanographic reconstruction in the northwestern Pacific, *Marine Micropaleontology*, 55, 129–156, <https://doi.org/10.1016/j.marmicro.2005.02.007>, 2005.
- Mortyn, P. G. and Charles, C. D.: Planktonic foraminiferal depth habitat and $\delta^{18}\text{O}$ calibrations: Plankton tow results from the Atlantic sector of the Southern Ocean, *Paleoceanography*, 18, 1037, <https://doi.org/10.1029/2001PA000637>, 2003.
- 15 Northcote, L. C. and Neil, H. L.: Seasonal variations in foraminiferal flux in the Southern Ocean, Campbell Plateau, New Zealand, *Marine Micropaleontology*, 56, 122–137, <https://doi.org/10.1016/j.marmicro.2005.05.001>, 2005.
- Pados, T. and Spielhagen, R. F.: Species distribution and depth habitat of recent planktic foraminifera in Fram Strait, Arctic Ocean, *Polar Research*, 33, 22483, <https://doi.org/10.3402/polar.v33.22483>, 2014.
- Peeters, F. J. C. and Brummer, G.-J. A.: The seasonal and vertical distribution of living planktic foraminifera in the NW Arabian Sea, Geological Society, London, Special Publications, 195, 463–497, <https://doi.org/10.1144/GSL.SP.2002.195.01.26>, 2002.
- 20 Rebotim, A., Voelker, A. H. L., Jonkers, L., Waniek, J. J., Meggers, H., Schiebel, R., Fraile, I., Schulz, M., and Kucera, M.: Factors controlling the depth habitat of planktonic foraminifera in the subtropical eastern North Atlantic, *Biogeosciences*, 14, 827–859, <https://doi.org/10.5194/bg-14-827-2017>, 2017.
- Rippert, N., Nürnberg, D., Raddatz, J., Maier, E., Hathorne, E., Bijma, J., and Tiedemann, R.: Constraining foraminiferal calcification depths in the western Pacific warm pool, *Marine Micropaleontology*, 128, 14–27, <https://doi.org/10.1016/j.marmicro.2016.08.004>, 2016.
- 25 Sautter, L. R. and Thunell, R. C.: Seasonal succession of planktonic foraminifera, results from a four-year time-series sediment trap experiment in the Northeast Pacific, *Journal of Foraminiferal Research*, 19, 253–267, <https://doi.org/10.2113/gsjfr.19.4.253>, 1989.
- Sautter, L. R. and Thunell, R. C.: Planktonic foraminiferal response to upwelling and seasonal hydrographic conditions: sediment trap results from San Pedro Basin, Southern California Bight, *Journal of Foraminiferal Research*, 21, 347–363, <https://doi.org/10.2113/gsjfr.21.4.347>, 30 1991.
- Simstich, J., Sarnthein, M., and Erlenkeuser, H.: Paired $\delta^{18}\text{O}$ signals of *Neoglobobulimina pachyderma* (s) and *Turborotalita quinqueloba* show thermal stratification structure in Nordic Seas, *Marine Micropaleontology*, 48, 107–125, [https://doi.org/10.1016/S0377-8398\(02\)00165-2](https://doi.org/10.1016/S0377-8398(02)00165-2), 2003.
- Stangeew, E.: Distribution and Isotopic Composition of Living Planktonic Foraminifera *N. pachyderma* (sinistral) and *T. quinqueloba* in the High Latitude North Atlantic, Ph.D. thesis, Christian-Albrechts-Universität zu Kiel, 2001.
- 35 Storz, D., Schulz, H., Waniek, J. J., Schulz-Bull, D. E., and Kucera, M.: Seasonal and interannual variability of the planktic foraminiferal flux in the vicinity of the Azores Current, *Deep-Sea Research I*, 56, 107–124, <https://doi.org/10.1016/j.dsr.2008.08.009>, 2009.
- Watkins, J. M., Mix, A. C., and Wilson, J.: Living planktic foraminifera: tracers of circulation and productivity regimes in the central equatorial Pacific, *Deep Sea Research II*, 43, 1257–1282, 1996.
- 40 Watkins, J. M., Mix, A. C., and Wilson, J.: Living planktic foraminifera in the central tropical Pacific Ocean: Articulating the equatorial 'cold tongue' during La Niña, 1992, *Marine Micropaleontology*, 33, 157–174, [https://doi.org/10.1016/S0377-8398\(97\)00036-4](https://doi.org/10.1016/S0377-8398(97)00036-4), 1998.

Wolfeich, C. M.: Satellite-Derived Sea Surface Temperature, Mesoscale Variability, and Foraminiferal Production in the North Atlantic, M.Sc., Cambridge, MS, 1994.