

Species	<i>N. pachyderma</i>	<i>N. incompta</i>	<i>G. bulloides</i>	<i>G. ruber</i> (white)	<i>T. sacculifer</i>
$P_{F, 0}$	– (–)	– (–)	2.6 (–)	2.6 (–)	2.6 (–)
α_{PI}	– (–)	– (–)	0.012 (–)	0.01 (–)	0.07 (–)
$p_{\%}$	– (–)	– (–)	0.3 (–)	0.3 (–)	0.4 (–)
T_{thres}	18.0 (24.0)	3.0 (-0.3)	3.0 (-0.3)	10.0 (5.0)	15.0 (15.0)
$cl_{N.pachyderma, j}$	– (–)	0.2 (0.2)	0 (0)	0 (0)	0 (0)
$cl_{N.incompta, j}$	– (–)	– (–)	0.1 (0.1)	0.2 (0.8)	0 (0)
$cl_{G.bulloides, j}$	– (–)	0.8 (0.5)	– (–)	0.8 (0.8)	0.8 (0.8)
$cl_{G.ruber(white), j}$	– (–)	0.2 (0.8)	0.1 (0.5)	– (–)	0.2 (0.2)
$cl_{T.sacculifer, j}$	– (–)	0 (0)	0.1 (0.5)	0.2 (0.2)	– (–)