

Supplement of Biogeosciences, 15, 721–732, 2018
<https://doi.org/10.5194/bg-15-721-2018-supplement>
© Author(s) 2018. This work is distributed under
the Creative Commons Attribution 4.0 License.



Supplement of

Tracking the direct impact of rainfall on groundwater at Mt. Fuji by multiple analyses including microbial DNA

Ayumi Sugiyama et al.

Correspondence to: Kenji Kato (kato.kenji@shizuoka.ac.jp)

The copyright of individual parts of the supplement might differ from the CC BY 4.0 License.

Table S1. Summary of sample analysis.

sampling date	rainfall event (amount of rainfall)	SP-0m, Shibakawa, spring water, 726m a.s.l.				GW-42m, Yodoshi, groundwater, 150m a.s.l., 42m depth				GW-550m, Aoki, groundwater, 175m a.s.l., 550m depth				R1, Go-gome, rainwater, 2,364m a.s.l.				R2, Kokuyurin, rainwater, 1,431m a.s.l.				R3, Ni-gome, rainwater, 1,081m a.s.l.				R4, Asagiri, rainwater, 850m a.s.l.				R5, Shibakawa, rainwater, 723m a.s.l.					
		E	B	DG	NGS	E	B	DG	NGS	E	B	DG	NGS	E	B	DG	NGS	E	B	DG	NGS	E	B	DG	NGS	E	B	DG	NGS	E	B	DG	NGS		
2012/6/15	■ Event1 (30mm)	○	○																																
2012/10/18		○	○																													○	○		
2012/11/8		○	○																																
2012/11/22		○	○																																
2013/5/31		○																																	
2013/6/17		○	○	○										○	○					○	○					○	○				○	○			
2013/7/2								○	○	○		○	○	○					○	○															
2013/7/10		○	○	○	○																										○	○			
2013/8/6		○	○	○												○	○										○	○			○	○			
2013/8/7								○	○	○		○	○	○							○	○													
2013/8/27																○	○									○	○				○	○			
2013/9/4		○	○	○				○	○	○		○	○	○																	○	○			
2013/9/9	■ Event2 (>300mm)	○	○	○												○	○									○	○			○	○				
2013/9/19		○	○	○	○			○	○	○		○	○	○	○			○	○	○					○	○	○	○	○	○	○	○	○		
2013/10/17		○	○					○	○	○		○	○																						
2014/3/28		○	○	○				○	○	○		○	○													○	○			○	○				
2014/5/14																																			
2014/5/31																																			
2014/6/27		○	○	○				○	○		○	○																							
2014/6/28	■ Event3 (100mm)																														○	○			
2014/7/18		○	○	○				○	○		○	○				○	○									○	○			○	○				
2014/8/14		○	○	○				○	○		○	○				○	○									○	○			○	○				
2014/9/16		○	○	○				○	○		○	○				○	○									○	○			○	○				
2014/10/16	■ Event4 (>300mm)	○	○	○				○	○		○	○				○	○									○	○			○	○				
2014/10/30		○	○	○				○	○		○	○															○	○			○	○			
2014/11/20		○	○	○				○	○		○	○															○	○			○	○			
2015/3/10																																	○		

E: environmental parameters, stable isotopic analysis, Ion analysis and dissolved silica,
 B: number of prokaryote(TDC), Bacteria (CARD-FISH) and Archaea (CARD-FISH),
 DG: DGGE analysis,
 NGS: next generation sequence analysis.






Figure S1. Fluctuation of the Amount of discharge at the spring of SP-0m.


SP-0m (726 m a.s.l.)




GW-42m (150 m a.s.l.)




GW-550m (175 m a.s.l.)




R1 (2,364 m a.s.l.)




R2 (1,431 m a.s.l.)



R3 (1,081 m a.s.l.)



R4 (850 m a.s.l.)



R5 (723m a.s.l.)




Figure S2. Hexadiagram shows major eight ions dissolved in water by their relative abundance. Anions are shown in the right side of the figure, while cations are in the left. The shape of diagram suggests characteristics of water examined. Thus that of groundwater (left panels) and precipitation (right panels). Observation period is from June 2012 to November 2014. Blue arrow indicates high concentration of Na^+ at GW-550m.

Table S2. Chemical character of spring, groundwater and rainwater.

(meq L⁻¹)

Site ID	n	Na ⁺	K ⁺	Mg ²⁺	Ca ²⁺	Cl ⁻	NO ₃ ⁻	SO ₄ ²⁻	HCO ₃ ⁻
SP-0m	20	0.254	0.032	0.199	0.428	0.114	0.056	0.149	0.654
GW-42m	13	0.313	0.046	0.308	0.578	0.086	0.122	0.104	1.005
GW-550m	13	0.627	0.015	0.178	0.354	0.088	0.008	0.242	0.815
R1	1	0.002	0.002	0.001	0.025	0.002	0.004	0.000	0.059
R2	8	0.007	0.007	0.007	0.014	0.009	0.011	0.019	0.053
R3	11	0.034	0.002	0.013	0.011	0.040	0.019	0.024	0.035
R4	15	0.029	0.004	0.015	0.016	0.034	0.025	0.040	0.021
R5	14	0.016	0.006	0.009	0.012	0.019	0.020	0.025	0.095

Methods for denaturing gradient gel electrophoresis (DGGE) analysis

A nested-PCR approach was employed with Bacteria-specific primers Bac27F(5'-AGA GTT TGA TCM TGG CTC AG-3')-Uni1492R(5'-GGY TAC CTT GTT ACG ACT T-3') (DeLong, 1992) and 341F-GC(5'- CGC CCG CCG CGC GCG GCG GGC GGG GCG GGG GCA CGG GGG GCC TAC GGG AGG CAG CAG -3')-534R(5'- ATT ACC GCG GCT GCT GG -3') (Muyzer et al., 1993) or Archaea-specific primers ARC344F(5'- ACG GGG YGC AGC AGG CGC GA -3')-ARC915R(5'- GTG CTC CCC CGC CAA TTC CT -3') (Vetriani et al., 1999) and 344F-GC(5'-CGC CCG CCG CGC CCC GCG CCC GTC CCG CCG CCC CCG CCC GAC GGG GYG CAG CAG GCG CGA-3')-518R(5'- ATT ACC GCG GCT GCT GG -3') (Muyzer et al., 1993) was employed to amplify variable V3 region of bacterial and archaeal 16S rRNA gene. Primary and secondary amplification reactions were performed in a 25 μ L PCR mixture consisted of 2.5 μ L 10 \times PCR buffer (TaKaRa Bio Inc., Shiga, Japan), 250 μ M dNTPs (TaKaRa Bio Inc.) 2.0 μ L, 1 U of ExTaq polymerase (TaKaRa Bio Inc.) 0.125 μ L, 1.5 M of each primer and 50 ng template. The products were amplified in Thermal cycler Wako WK-0232 (Wako Pure Chemical Industries Ltd.) under the following conditions: for *Bacteria*, 94°C for 5 min, then 20 cycles for primary reactions and 40 cycles for secondary reactions of 94°C for 30 sec, 55°C for 30 sec, and 72°C for 40 sec, a final extension of 72°C for 10 min (Jiang et al., 2014), for Archaea, 94°C for 5 min, then 20 cycles for primary reactions and 40 cycles for secondary reactions of 94°C for 30 sec, 48°C for 30 sec, and 72°C for 30 sec, a final extension of 72°C for 10 min (Vetriani et al., 1999). PCR products confirmed by 2% (w/v) agarose gel electrophoresis, stained with ethidium bromide (EB), and visualized under UV light. Obtained PCR products were stored at -20°C for subsequent DGGE analysis.

DGGE was performed by Dcode Universal Mutation Detection System (BIO-RAD Laboratories Inc., CA, USA) according to Muyzer et al. (1993). A 35%-60% vertical denaturing gradient polyacrylamide gel was formed mixing 0% denaturant solution (40% Acrylamide/Bisacrylamide 5 mL, 50 \times TAE Buffer 250 μ L, MilliQ 19.75 mL) and 100% denaturant solution (40% Acrylamide/Bisacrylamide 3 mL, 50 \times TAE Buffer 150 μ L, formamide 6 mL, Urea 6.3 g, MilliQ 200 μ L) with Model 475 gradient former. PCR products were applied directly onto polyacrylamide gel. Electrophoresis was performed at a constant voltage of 75 V and at constant temperature of 60°C for 12 hours in 0.5 \times TAE buffer. Following electrophoresis, the gel was stained by SYBR Gold Nucleic Acid Gel Stain (SYBR Gold 0.5 μ L, Thermo Fisher Scientific Inc. 1 \times TAE 5 mL) for 30 min in dark and photographed with UV transillumination by AE-6932GXES-U (ATTO, Tokyo, Japan).

DNA fragments from the major DGGE bands were extracted and purified using

AxyPrep PCR Clean-up kit (Axygen Scientific Inc., CA, USA) for sequencing. The sequences of PCR products were determined by the capillary DNA sequencer (CEQ8000 DNA Analysis System, Beckman Coulter Inc. CA, USA or Applied Biosystems 3730xl, Thermo Fisher Scientific Inc.). Sequences were aligned using Basic Local Alignment Search Tool (BLAST; Altschul et al., 1997) in the DNA Data Bank of Japan (DDBJ; Thompson et al., 1994).

Reference:

- Altschul, S. F., Madden, T. L., Schaffer, A. A., Zhang, J. H., Zhang, Z., Miller, W. and Lipman, D. J.: Gapped BLAST and PSI-BLAST: a new generation of protein database search programs, *Nucleic Acids Res*, 25, 3389-3402, 1997.
- DeLong, E. F., Archaea in coastal marine environments. *Proceedings of the National Academy of Sciences of the United States of America*, 89, 5685-5689, 1992.
- Jiang, Z., Li, P., Wang, Y. H., Li, B., Deng, Y. M. and Wang, Y. X.: Vertical distribution of bacterial populations associated with arsenic mobilization in aquifer sediments from the Hetao plain, Inner Mongolia, *Environ Earth Sci*, 71, 311-318, 2014.
- Muyzer, G., Dewaal, E. C. and Uitterlinden, A. G.: Profiling of complex microbial-populations by denaturing gradient gel-electrophoresis analysis of polymerase chain reaction-amplified genes-coding for 16S ribosomal-RNA, *Appl Environ Microb*, 59, 695-700, 1993.
- Thompson, J. D., Higgins, D. G. and Gibson, T. J.: Clustal-W –Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice, *Nucleic Acids Res*, 22, 4673-4680, 1994 (<http://www.ddbj.nig.ac.jp/>).
- Vetriani, C., Jannasch, H. W., MacGregor, B. J., Stahl, D. A. and Reysenbach, A. L.: Population structure and phylogenetic characterization of marine benthic archaea in deep-sea sediments, *Appl Environ Microb*, 65, 4375-4384, 1999.




Figure S3. Bacterial community composition of groundwater (SP-0m, GW-42m and GW-550m) and rainwater (R2, R3, R4 and R5). Black and open arrows indicate the rainfall event; Event 2, Event 3 and Event 4. Black arrows particularly indicate the torrential rainfall.

Table S3. Similarity of sequences from DGGE bands to those of most related strains.

Band name	Band No.	Class	Most relative strain in DDBJ related species	Match	Rain					Groundwater	
					R2	R3	R4	R5	SP-0m	GW-42m	GW-550m
Rain_1	2, 4, 6	<i>Sphingobacteriia</i>	<i>Mucilaginibacter</i> sp. 001	168/170 98%	+	+	+				
Rain_2	10, 11, 18	<i>Sphingobacteriia</i>	<i>Mucilaginibacter</i> sp. KJ029	153/155 98%	+	+	+	+			
Rain_3	15, 17	<i>Nostocales</i>	<i>Tolypothrix</i> sp. CNP3-B1-C1	68/75 90%			+	+			
Rain_4	7	<i>Betaproteobacteria</i>	<i>Herbaspirillum</i> sp. MMD15	174/174 100%			+		+		
	1	<i>Sphingobacteriia</i>	<i>Mucilaginibacter</i> sp. Aws5	164/171 95%	+						
	3	<i>Gammaproteobacteria</i>	<i>Pseudomonas oryzihabitans</i>	150/156 96%	+						
	5	<i>Sphingobacteriia</i>	<i>Mucilaginibacter</i> sp. KJ029	162/170 95%			+				
	8	<i>Alphaproteobacteria</i>	<i>Caulobacter</i> sp. JM6	149/150 99%			+				
	9	<i>Sphingobacteriia</i>	<i>Mucilaginibacter</i> sp. KJ029	112/112 100%				+			
	12	<i>Sphingobacteriia</i>	<i>Mucilaginibacter</i> sp. PAMC 26640	153/155 98%		+					
	13	<i>Betaproteobacteria</i>	<i>Variovorax paradoxus</i>	161/162 99%		+					
	14	<i>Alphaproteobacteria</i>	<i>Sphingomonas</i> sp. PXM	127/127 100%		+					
	16	<i>Gammaproteobacteria</i>	<i>Pseudomonas fluorescens</i>	48/49 97%			+				
SP-0m_1	S1, S4	<i>Betaproteobacteria</i>	<i>Herbaspirillum</i> sp. MMD15	174/177 98%			+		+		
SP-0m_2	S2, S6	<i>Betaproteobacteria</i>	<i>Sphaerotilus</i> sp. IMCC12769	176/176 100%					+		
	S3	<i>Betaproteobacteria</i>	<i>Oxalobacteraceae</i> bacterium AKB-2008-RN12	153/155 98%					+		
	S5	<i>Bacilli</i>	<i>Paenibacillus</i> sp. PAMC26516	170/175 97%					+		
GW-42m_1	Y2, Y3, Y4	<i>Nitrospirales</i>	<i>Nitrospiraceae</i> bacterium vj1_c7	94/96 97%						+	
	Y1	<i>Sphingobacteriia</i>	<i>Flaviumibacter</i> sp. 7B-231	162/162 100%						+	
GW-550m_1	A1, A5, A7, A8	<i>Betaproteobacteria</i>	<i>Gallionella</i> sp. JA52	164/172 95%							+
GW-550m_2	A2, A6, A11	<i>Nitrospirales</i>	<i>Nitrospiraceae</i> bacterium vj1_c7	159/175 90%							+
	A3	<i>Nitrospirales</i>	<i>Nitrospiraceae</i> bacterium vj1_c7	159/175 90%							+
	A4	<i>Nitrospirales</i>	<i>Nitrospiraceae</i> bacterium vj1_c7	159/168 94%							+
	A8	<i>Gammaproteobacteria</i>	<i>Acinetobacter</i> sp.	176/177 99%							+
	A9	<i>Alphaproteobacteria</i>	<i>Sphingomonas</i> sp.	150/150 100%							+
	A10	<i>Alphaproteobacteria</i>	<i>Brevundimonas diminuta</i>	149/149 100%							+

Color indicates the DGGE band were obtained from more than two samples.

Table S4. Sequences affiliated with thermophilic bacteria retrieved from deep groundwater (GW-550m).

Order	GW-550m-1	GW-550m-2	GW-550m-3
<i>Firmicutes/Clostridia/Thermoanaerobacterales</i>	0.00	1.25	0.00
<i>Actinobacteria/Thermoleophilia/Gaiellales</i>	0.21	0.61	0.27
<i>Deinococcus-Thermus/Deinococci/Deinococcales</i>	0.01	1.36	0.05

*Number indicates relative percentage to whole reads which were removed chimeric sequence.