Corrigendum to Biogeosciences, 16, 3439–3455, 2019 https://doi.org/10.5194/bg-16-3439-2019-corrigendum © Author(s) 2020. This work is distributed under the Creative Commons Attribution 4.0 License.



Corrigendum to "Insights into architecture, growth dynamics, and biomineralization from pulsed Sr-labelled *Katelysia rhytiphora* shells (Mollusca, Bivalvia)" published in Biogeosciences, 16, 3439–3455, 2019

Laura M. Otter¹, Oluwatoosin B. A. Agbaje¹, Matt R. Kilburn², Christoph Lenz^{3,4}, Hadrien Henry^{1,3}, Patrick Trimby⁵, Peter Hoppe⁶, and Dorrit E. Jacob^{1,3}

¹Department of Earth and Planetary Science, Macquarie University, Sydney, New South Wales 2109, Australia ²Centre for Microscopy, Characterisation and Analysis, University of Western Australia, Perth, WA 6009, Australia ³Australian Research Council Centre of Excellence for Core to Crust Fluid Systems (CCFS)/GEMOC, Australia ⁴Institute of Mineralogy and Crystallography, University of Vienna, Althanstr. 14, 1090 Vienna, Austria ⁵Oxford Instruments NanoAnalysis, High Wycombe, HP12 3SE, UK

⁶Particle Chemistry Department, Max Planck Institute for Chemistry, Hahn-Meitner-Weg 1, 55128 Mainz, Germany

Correspondence: Laura M. Otter (laura.otter@mq.edu.au)

Published: 15 January 2020

A mistake was introduced during the production. A wrong figure file was used for Fig. 2. Please see the correct version here. This mistake only concerns the PDF version of the paper. The full-text HTML version is correct as is.



Figure 2. FEG-SEM BSE images showing polished cross sections of the oOSL (**a**) and iOSL (**b**) of a Sr-labelled *K. rhytiphora* shell (specimen ID: K2-06) overlain with NanoSIMS 88 Sr/ 40 Ca maps. Shell layers grown in ambient seawater 88 Sr/ 40 Ca ratios are depicted in blue, while shell formed during Sr-enriched incubations are shown in pink. White arrows identify the general growth direction of the shell, while the Sr-labelled shell layers from the underlying BSE image visualize the local growth directions for each ultrastructure. A schematic of the shell tip shows the exact location of the NanoSIMS maps with blue and red arrows pointing towards the locations of the prismatic oOSL (**a**) and crossed-acicular iOSL (**b**) sampling locations, respectively. Asterisk marks the inner shell surface. Scale bars are 10 µm.