

Supplement of Biogeosciences, 16, 3941–3958, 2019
<https://doi.org/10.5194/bg-16-3941-2019-supplement>
© Author(s) 2019. This work is distributed under
the Creative Commons Attribution 4.0 License.



Supplement of

Microbial community composition and abundance after millennia of submarine permafrost warming

Julia Mitzscherling et al.

Correspondence to: Susanne Liebner (susanne.liebner@gfz-potsdam.de)

The copyright of individual parts of the supplement might differ from the CC BY 4.0 License.

Supplementary Online Material:

Contents:

Figures S1-S5: including DNA concentrations, DOC contents and additional information to Figure 2 (PCA) as well as a schematic description of the formation of the sediments units.

Tables S1-S14: including a site description, an overview of selected physicochemical parameters and of the microbial abundances, detailed information about the molecular samples, sequencing primers and barcode sequences, sequencing statistics as well as the results of all statistical tests




Figure S1: Quality control of the extracted genomic DNA, exemplarily shown for two samples from core C2 (CK1210 SDS: 0.05 m bsf, 265 ng/g and CK1247 SDS: 52.7 m bsf, 33.4 ng/g). The examples show that there is not much fragmentation likely due to constantly freeze-locked conditions. Hence, we used the DNA extracts without gel purification for downstream analyses.




Figure S2: Boxplot of DOC concentrations of Unit II in mg C L⁻¹. Median lines are indicated within the boxes of which the size corresponds to $\pm 25\%$ of the data, whereas the whiskers show the minimum and maximum of all data. C1: n=74, C4: n=5, C3: n=3, C2: n=12.




Figure S3: DNA concentrations in ng g⁻¹ sediment wet weight of the cores C1, C4, C3 and C2. Box plots contain the mean values of all samples, obtained from two technical replicates each. Median lines are indicated within the boxes of which the size corresponds to $\pm 25\%$ of

the data, whereas the whiskers show the minimum and maximum of all data. C1, C4 and C3: n=6, C2: n=17.









Figure S4: Loadings plots belonging to Figure 3: PCA of environmental, sedimentological and pore water data from Unit II. Shown are the correlations to a) PC1 and b) PC2 which were used to choose the physicochemical factors that are mainly responsible for the variance between samples in the PCA.





D. modern coastal lowland of the western Laptev Sea



C. supposed Late Weichselian flood plain area of Ice Complex accumulation



B. supposed Early to Middle Weichselian fluvial dominated terrestrial shelf



A. supposed Eemian coastal lowland with brackish thermokarst lagoons and terrestrial thermokarst basins




Figure S5: Schematic representation of the late Quaternary landscape dynamics in the western Laptev Sea coastal region and formation of the sediments units (modified after Winterfeld et al. (2011)).

Table S1: Site description of each borehole location. Mbsf stands for meters below sea floor and indicates the depth in the submarine cores (C4, C3, C2), while mbs stands for meters below surface and is used for depth indication in the terrestrial core C1.

	C1	C4	C3	C2
Distance to coast [km]	- 0.1	1.0	3.0	11.5
Water depth [m]	-	2.2	4.4	6.0
Frost table depth [mbsf]	0.0	1.7	7.6	29.0
Upper boundary of Unit II [mbs/mbsf]	22.0	13.3	8.6	35.0
Lower boundary of Unit II	61.0	29.8	27.1	58.5
Uppermost sample depth [mbs/mbsf]	27.3	13.3	8.6	38.5
Lowermost sample depth	44.3	29.8	24.9	58.4

Table S2: Minimum, maximum and mean values of environmental factors in Unit II significantly contributing to the bacterial community composition and microbial abundance.

	Core	Minimum	Maximum	Mean	Std.Dev.	n
Temperature [°C]	C1	-12.5	-12.4	-12.4	0.0	8
	C4	-7.1	-5.8	-6.4	0.5	4
	C3	-1.8	-1.2	-1.4	0.2	4
	C2	-1.6	-1.5	-1.5	0.0	4
Salinity [PSU]	C1	0.0	1.6	0.5	0.4	184
	C4	0.9	17.6	5.6	4.8	10
	C3	0.5	3.7	1.0	0.6	38
	C2	0.0	12.5	0.8	1.7	67
$\delta^{18}\text{O}$ [‰] vs. SMOW]	C1	-30.8	-14.9	-22.3	4.0	184
	C4	-27.7	-18.8	-22.8	3.0	10
	C3	-20.6	-19.1	-20.1	0.3	38
	C2	-30.0	20.2	-27.6	1.5	67
δD [‰] vs. SMOW]	C1	-241.8	-115.7	-177.2	32.9	184
	C4	-219.1	-144.0	-178.8	25.7	10
	C3	-162.9	-149.4	-158.4	2.9	38
	C2	-232.7	-156.8	-213.3	11.3	67

Table S3: Geochemical, pore water and environmental data of all samples at each drill site.

See “SuppInfo_Table_3”

Table S 4: Sample names of the molecular samples, their depth relative to sea level (meters below sea level, m bsl), relative to surface (meters below surface (m bs) in the terrestrial core and meters below sea floor (m bsf) in the submarine cores), and their corresponding lithology.

Sample Name	Depth [m bsl]	Depth [m bs/ m bsf]	Lithology
C1-1	1.3	27.3	sandy
C1-2	9.9	35.9	sandy
C1-3	12.1	38.1	sandy
C1-4	17.2	43.2	plant/wood detritus
C1-5	17.4	43.4	plant/wood detritus
C1-6	18.4	44.4	plant/wood detritus
C4-1	15.5	13.3	plant/wood detritus
C4-2	22.0	19.8	small peat inclusions
C4-3	25.0	22.8	small peat inclusions
C4-4	28.5	26.3	sandy with quartz gravel
C4-5	30.0	27.8	sandy with quartz gravel
C4-6	32.0	29.8	sandy with quartz gravel
C3-1	13.0	8.6	small peat inclusions
C3-2	16.0	11.6	small peat inclusions
C3-3	19.0	14.6	small peat inclusions
C3-4	21.5	17.1	sandy
C3-5	24.6	20.2	sandy
C3-6	29.2	24.8	sandy
CK1232	44.5	38.5	
C2-1	46.0	40.0	sandy
C2-2	48.1	42.1	sandy
CK1235	49.3	43.3	
CK1236	50.7	44.7	
CK1237	51.8	45.8	
C2-4	52.2	46.2	sandy
C2-5	54.6	48.6	plant/wood detritus
CK 1241	54.7	48.7	
C2-7	55.0	49.0	plant/wood detritus
C2-8/1244	56.1	50.1	plant/wood detritus
CK1245	57.8	51.8	
CK1246	58.2	52.2	
CK1247	58.7	52.7	
CK1248	61.6	55.6	
C2-9	62.9	56.9	plant/wood detritus
C2-10	64.4	58.4	sandy silt

Table S5: Oligonucleotide primers for Illumina MiSeq sequencing and quantitative PCR.

Target	Primer Sets	Primer Sequence 5'-3'	Size bp	T (°C)	No. of PCR Cycles	References
Illumina MiSeq sequencing						
Bacterial 16S rRNA	S-D-Bact-0341-b-S-17	CCT ACG GGA GGC AGC AG	464	55	35	(Muyzer et al., 1993) (Herlemann et al., 2011)
	S-D-Bact-0785-a-A-21	GAC TAC HVG GGT ATC TAA TCC				
Quantitative PCR						
Bacterial 16S rRNA	S-D-Bact-0341-b-S-17	CCT ACG GGA GGC AGC AG	193	55.7	40	(Muyzer et al., 1993) (Muyzer et al., 1993)
	S-D-Bact-0517-a-A-18	ATT ACC GCG GCT GCT GG				

Table S6: Barcode sequences for Illumina MiSeq sequencing.

Barcode ID Forward Primer	Barcode Sequence	Barcode ID Reverse Primer	Barcode Sequence
Bac-01-For	ACGAGTGCGT	Bac-01-Rev	ACGAGTGCGT
Bac-02-For	ACGCTCGACA	Bac-02-Rev	ACGCTCGACA
Bac-03-For	AGACGCACT	Bac-04-Rev	AGCACTGTAG
Bac-06-For	ATATCGCGAG	Bac-05-Rev	ATCAGACACG
Bac-07-For	CGTGTCTCTA	Bac-06-Rev	ATATCGCGAG
Bac-08-For	CTCGCGTGT	Bac-07-Rev	CGTGTCTCTA
Bac-11-For	TGATACGTCT	Bac-08-Rev	CTCGCGTGTCT
Bac-13-For	CATAGTAGTG	Bac-11-Rev	TGATACGTCT
Bac-15-For	ATACGACGTA	Bac-13-Rev	CATAGTAGTG
Bac-16-For	TCACGTAATA	Bac-14-Rev	CGAGAGATAC
Bac-17-For	CGTCTAGTA	Bac-17-Rev	CGTCTAGTAC
Bac-19-For	TGTACTACT	Bac-18-Rev	TCTACGTAGC
Bac-23-For	TACTCTCGTG	Bac-19-Rev	TGTACTACTC
Bac-24-For	TAGAGACGAG	Bac-22-Rev	TACGAGTATG
Bac-25-For	TCGTGCTCG	Bac-23-Rev	TACTCTCGTG
Bac-26-For	ACATACGCGT	Bac-24-Rev	TAGAGACGAG
Bac-27-For	ACGCGAGTAT	Bac-25-Rev	TCGTGCTCG
Bac-28-For	ACTACTATGT	Bac-26-Rev	ACATACGCGT
Bac-31-For	AGCGTCGTCT	Bac-28-Rev	ACTACTATGT
Bac-33-For	ATAGAGTACT	Bac-30-Rev	AGACTATACT
Bac-34-For	CACGCTACGT	Bac-31-Rev	AGCGTCGTCT
Bac-35-For	CAGTAGACGT	Bac-33-Rev	ATAGAGTACT
Bac-36-For	CGACGTGACT	Bac-34-Rev	CACGCTACGT
Bac-38-For	TACACGTGAT	Bac-35-Rev	CAGTAGACGT
Bac-39-For	TACAGATCGT	Bac-36-Rev	CGACGTGACT
Bac-40-For	TACGCTGTCT	Bac-37-Rev	TACACACACT
Bac-41-For	TAGTGTAGAT	Bac-38-Rev	TACACGTGAT
Bac-42-For	TCGATCACGT	Bac-39-Rev	TACAGATCGT

Bac-44-For	TCTAGCGACT	Bac-40-Rev	TACGCTGTCT
Bac-45-For	TCTATACTAT	Bac-41-Rev	TAGTGATAGAT
Bac-49-For	ACGCGATCGA	Bac-44-Rev	TCTAGCGACT
Bac-50-For	ACTAGCAGTA	Bac-45-Rev	TCTATACTAT
		Bac-46-Rev	TGACGTATGT
		Bac-49-Rev	ACGCGATCGA
SfiA-MW00	ACACGT	SfiB-MW10	CAGTCA
SfiA-MW01	ACGTAC	SfiB-MW11	CATGAC
SfiA-MW02	ACTGCA	SfiB-MW12	GACTAG
SfiA-MW02	ACTGCA	SfiB-MW13	GAGATC
SfiA-MW03	AGAGTC	SfiB-MW14	GATCGA
SfiA-MW04	AGCTGA	SfiB-MW14	GATCGA
SfiA-MW05	AGTCAG	SfiB-MW15	GTACAC
SfiA-MW06	ATATCG	SfiB-MW15	GTACAC
SfiA-MW07	ATCGAT	SfiB-MW16	GTCACA
		SfiB-MW17	GTGTGT
		SfiB-MW18	TCAGAG
		SfiB-MW19	TCGAGA

Table S 7: Overview of sequencing reads: number of reads after the removal of singletons, number of reads that were removed when the background filter of 0.5% was applied, number of reads representing chloroplast, mitochondrial and archaeal taxa and finally the number of quality reads after the application of all filters. Critical samples with less than 15.000 raw reads are shaded red. Critical samples where the relative abundances within duplicates are comparable are colored light red. The dark red colored sample was not used for the calculation of the mean relative abundance as the relative abundances within duplicates differed.

	Raw reads	Background Reads (0.5%)	Chloroplast	Mitochondrial Taxa	Archaeal taxa	Quality reads
C1-1a	74760	20214	0	0	0	54546
C1-1b	89992	22879	0	0	0	67113
C1-2a	23789	15150	0	0	0	8639
C1-2b	25100	16330	0	0	0	8770
C1-3a	41727	24707	0	0	0	17020
C1-3b	8666	5185	0	0	0	3481
C1-4a	31071	22620	0	0	0	8451
C1-4b	5142	3761	0	0	0	1381
C1-5a	208578	100128	0	0	0	108450
C1-5b	147753	69180	0	0	0	78573
C1-6a	244866	100302	0	0	1790	142774
C1-6b	255535	113256	0	0	6331	135948
C4-1a	231425	70205	0	0	0	161220
C4-1b	103692	30996	0	0	0	72696
C4-2a	312930	64767	0	0	0	248163
C4-2b	18603	2840	0	0	0	15763
C4-3a	269853	99103	1765	0	0	168985
C4-3b	94463	33930	0	0	0	60533

C4-4a	170018	49851	0	0	1050	119117
C4-4b	180556	52147	0	0	0	128409
C4-5a	9823	4402	0	0	0	5421
C4-5b	17374	7566	0	0	0	9808
C4-6a	56201	23871	0	0	0	32330
C4-6b	20949	8951	0	0	0	11998
C3-1a	52885	27176	0	0	588	25121
C3-1b	180772	94154	1165	0	2402	83051
C3-2a	35528	13955	0	0	1511	20062
C3-2b	102122	39090	652	0	4780	57600
C3-3a	73935	22562	0	0	0	51373
C3-3b	25589	7940	0	0	0	17649
C3-4a	53553	16543	0	0	1899	35111
C3-4b	22552	6552	0	0	398	15602
C3-5a	128366	34045	1091	0	0	93230
C3-5b	16643	4179	152	0	0	12312
C3-6a	80004	20997	0	0	1349	57658
C3-6b	89902	21079	867	0	0	67956
CK1232-1	127284	59052	0	0	0	68232
CK1232-2	161752	70043	0	0	0	91709
C2-1a	53571	16678	0	0	0	36893
C2-1b	25615	7060	0	0	0	18555
C2-2a	55698	20602	0	0	0	35096
C2-2b	84301	25809	0	0	0	58492
CK1235-1	206215	122462	11152	0	0	72601
CK1235-2	74429	43777	4130	0	0	26522
CK1236-1	20564	11265	0	0	0	9299
CK1236-2	55725	30832	0	0	0	24893
CK1237-1	102376	62617	0	0	0	39759
CK1237-2	139700	86444	0	0	0	53256
C2-4a	136761	74460	0	0	0	62301
C2-4b	216318	118201	0	0	0	98117
C2-5a	48354	27604	0	0	0	20750
C2-5b	92506	55950	0	0	0	36556
CK1241-1	177526	115666	0	0	0	61860
CK1241-2	142667	88091	0	0	0	54576
C2-7a	63745	36419	0	0	0	27326
C2-7b	159960	88818	0	0	0	71142
C2-8a	22420	14376	0	0	0	8044
C2-8b	130842	85938	0	0	0	44904
CK1244-1	99934	54354	0	0	0	45580
CK1244-2	15808	9077	0	0	0	6731
CK1245-1	81822	42330	0	0	0	39492
CK1245-2	49130	24254	0	0	0	24876
CK1246-1	52169	30142	0	0	0	22027
CK1246-2	70027	43178	0	0	0	26849
CK1247-1	32592	14991	0	0	0	17601

CK1247-2	21821	9398	0	0	0	12423
CK1248-1	25455	16365	0	0	0	9090
CK1248-2	48980	32070	0	0	0	16910
C2-9a	37303	24313	0	0	0	12990
C2-9b	43272	26410	0	0	0	16862
C2-10a	1889	1288	0	0	0	601
C2-10b	213822	155149	1128	0	0	57545

Table S 8: Spearman correlation of DNA concentration, bacterial 16S rRNA gene abundance and total cell counts. P-values are shown above the diagonal and the correlation coefficient r_s below.

	DNA	16S rRNA gene copies	TCC
DNA		>0.0001	>0.0001
16S rRNA gene copies	0.87		0.0001
TCC	0.68	0.61	

Table S9: Minimum. maximum. mean values and standard deviation of microbial and bacterial abundance. n indicates the number of samples.

	Core	Min	Max	Mean	Std. dev.	n
DNA concentration [ng g ⁻¹]	C1	28.6	331.3	141.6	105.6	6
	C4	6.2	277.5	88.5	102.6	6
	C3	5.6	51.9	19.8	17.8	6
	C2	8.7	341.5	106.9	94.0	17
16S rRNA gene copies [g ⁻¹ sediment]	C1	2.4E+07	4.3E+08	1.6E+08	1.4E+08	6
	C4	1.7E+06	1.6E+08	3.6E+07	5.8E+07	6
	C3	4.1E+06	6.2E+07	1.7E+07	2.1E+07	6
	C2	5.4E+06	1.5E+09	2.9E+08	4.0E+08	17
16S rRNA gene copies [ng ⁻¹ DNA]	C1	7.6E+05	1.4E+06	1.0E+06	2.6E+05	6
	C4	5.8E+04	5.8E+05	2.7E+05	1.6E+05	6
	C3	4.9E+05	1.2E+06	7.6E+05	2.2E+05	6
	C2	2.6E+05	1.7E+07	2.7E+06	4.2E+06	17
TCC [g ⁻¹ sediment]	C1	6.8E+06	8.2E+07	5.0E+07	2.8E+07	6
	C4	1.6E+06	4.4E+07	1.3E+07	1.5E+07	6
	C3	3.4E+05	3.4E+06	1.5E+06	1.0E+06	6
	C2	1.4E+06	4.9E+07	1.5E+07	1.3E+07	17

Table S10: Rank-based Spearman correlation of DNA concentration, bacterial 16S rRNA gene abundance and total cell counts with environmental factors and pore water data. Values in bold are significant (< 0.05) when omitting a p-value correction. R_s - values highlighted red show a negative correlation, whereas r_s -values highlighted green show a positive correlation.

	16S Bacteria	16S/DNA	TCC	Temp	Salinity	Depth [mbsl]	Depth [mbs/mbsf]	Ba ²⁺	Ca ²⁺	K ⁺	Mg ²⁺	Na ⁺	Si _{aq}	Cl ⁻	SO ₄ ²⁻	Br ⁻	NO ₃ ⁻	δ18O	δD	pH	TC	TN	TS	TOC	Clay	Silt	Sand	Grav. Water Content
p-value																												
DNA	>0.001	>0.001	>0.001	0.030	0.039	0.813	0.076	0.658	0.604	0.020	0.061	0.021	0.872	0.011	0.410	0.015	0.593	0.027	0.055	0.008	0.017	0.329	0.175	0.045	0.307	0.111	0.130	0.006
16S Bact.		>0.001	>0.001	0.173	0.003	0.164	0.002	0.860	0.248	0.003	0.005	>0.001	0.475	>0.001	0.128	0.001	0.587	0.023	0.054	0.002	0.009	0.175	0.056	0.021	0.821	0.886	0.926	0.007
16S / DNA			0.03	0.503	>0.001	0.216	0.004	0.799	0.005	0.001	>0.001	>0.001	0.268	>0.001	0.016	>0.001	0.425	0.369	0.528	0.001	0.218	0.171	0.135	0.284	0.055	0.102	0.084	0.153
TCC				>0.001	0.008	0.465	0.138	0.024	0.193	0.012	0.029	0.002	0.262	0.001	0.593	0.002	0.890	0.028	0.027	0.097	0.749	0.759	0.233	0.429	0.184	0.572	0.524	0.369
correlation coefficient r_s																												
DNA	0.87	0.47	0.68	-0.37	-0.35	0.04	0.30	-0.08	-0.09	-0.39	-0.32	-0.39	-0.03	-0.43	-0.14	-0.41	0.09	-0.37	-0.33	-0.44	0.40	0.17	-0.23	0.34	0.18	0.27	-0.26	0.47
16S Bact.		0.79	0.61	-0.24	-0.48	0.24	0.51	0.03	-0.20	-0.49	-0.46	-0.57	-0.12	-0.56	-0.26	-0.54	0.09	-0.38	-0.33	-0.52	0.44	0.23	-0.33	0.39	-0.04	0.03	-0.02	0.47
16S / DNA			0.36	-0.12	-0.63	0.21	0.47	0.04	-0.47	-0.55	-0.60	-0.71	-0.19	-0.67	-0.40	-0.66	-0.14	-0.16	-0.11	-0.54	0.21	0.24	-0.26	0.19	-0.33	-0.28	0.30	0.26
TCC				-0.64	-0.44	-0.13	0.26	-0.38	-0.23	-0.42	-0.37	-0.50	-0.19	-0.52	-0.09	-0.50	0.02	-0.37	-0.37	-0.28	0.06	0.05	-0.21	0.14	-0.23	-0.10	0.11	0.16

Table S11: Significance of the variance introduced by environmental factors into the microbial community tested by Permutational MANOVA (PerMANOVA)

	Dim1	Dim2	r ²	p-value
Depth [mbs/msbf]	-0.53174	-0.84691	0.3322	0.006
Temperature	0.13632	0.99067	0.2487	0.015
Ba	0.94807	0.31805	0.1859	0.031
Si	0.90304	-0.42956	0.1541	0.056
Ca	0.50032	0.86584	0.0100	0.835
K	0.81761	0.57578	0.0612	0.341
Mg	0.80879	0.58809	0.0684	0.297
Na	0.99177	0.12804	0.0813	0.241
Nitrate	-0.81210	0.58351	0.0280	0.637
Chloride	0.98966	0.14344	0.0527	0.391
Sulfate	-0.28689	0.95796	0.1014	0.161
Bromide	0.92727	0.37439	0.0629	0.326
Salinity	0.99532	0.09660	0.0459	0.443
δ18O	0.99329	-0.11569	0.3753	0.001
δD	0.98430	-0.17648	0.3914	0.001
pH	-0.42785	0.90385	0.6412	0.001
TC	0.41379	-0.91037	0.1053	0.149
TN	-0.38942	-0.92106	0.0268	0.640
TS	0.03653	0.99933	0.2694	0.004
TOC	0.40692	-0.91346	0.0974	0.170
Clay	0.47503	0.87997	0.1123	0.132
Silt	0.76336	0.64597	0.0532	0.405
Sand	-0.70792	-0.70629	0.0630	0.330
Conductivity	0.98987	0.14199	0.0419	0.478

Table S12: One-way PerMANOVA of OTU data from each drill site. Summary presents the overall test statistics. Pairwise analysis shows Bonferroni corrected p-values above the diagonal and F-values below.

Summary		Pairwise				
			C1	C4	C3	C2
Permutation N:	9999					
Total sum of squares:	26.49	C1		0.0012	0.0006	0.0006
Within-group sum of squares:	19.41	C4	4.014		0.0006	0.0006
F:	8.276	C3	12.400	7.368		0.0006
p (same):	0.0001	C2	5.833	5.156	16.350	

Table S13: Analysis of variance (ANOVA) of DOC concentrations between all four cores and Tukey's pairwise post-hoc test with p-values adjusted according to Copenhaver-Holland above and the Tukey's Q below the diagonal.

	Sum of squares	df	Mean square	F	p (same)
Between groups:	24714.2	3	8238.06	4.814	0.003712
Within groups:	155731	91	1711.34		Permutation p (n=99999)
Total:	180446	94			0.02357

	C1	C4	C3	C2
C1		0.066	0.996	0.052
C4	3,540		0.299	0.002
C3	0.310	2.490		0.739
C2	3.676	5.209	1.441	

Table S14: Fossil bioindicators according to Schirrmeister et al. (2008); Winterfeld et al. (2011); Müller et al. (2009) and their stratigraphical and paleoenvironmental interpretation.

Units	Bioindicator	Stratigraphy	Landscape, facies	Vegetation	Climate
II d	<ul style="list-style-type: none"> • Pollen: <i>Cyperaceae</i>, <i>Poaceae</i>, <i>Artemisia</i>, <i>Salix</i> • Spores: <i>Encalypta</i>, <i>Glomus</i> • Green algae: <i>Botryococcus</i>, <i>Pediastrum</i> • Ostracodes • Plant macro remains: <i>Carex</i>, <i>Salix</i> sp., <i>Saxifraga hirculus</i>, <i>Dryas Kobresia myosuroides</i>, <i>Thlaspitea rotundifolii</i> • Testacea: hygrophillic (<i>Diffugia</i>), sphagnobiotic (<i>Heleopera</i>, <i>Nebela</i>, <i>Argynnia</i> sp.) • Mammals: <i>Equus caballus</i>, <i>Mammuthus primigenius</i> 	<p>Middle</p> <p>Weichselian</p> <p>Interstadial</p>	<p>Floodplain,</p> <p>alluvial,</p> <p>boggy,</p> <p>periodically</p> <p>flooded</p>	<p>Grass-sedge</p> <p>tundra</p>	<p>Moderate</p> <p>, humid</p>
II c	No determinable fossil records found	<p>Early</p> <p>Weichselian</p> <p>Stadial</p>	fluvial		
II b	<p>Pollen: <i>Larix</i>, <i>Alnus fruticosa</i>, <i>Betula nana</i>,</p> <p>Ericales</p>	<p>Eemian</p> <p>Interglacial</p>	<p>Thermokarst</p> <p>lake</p>	Shrub tundra	Temperate
II a	No determinable fossil records found	<p>Early</p> <p>Weichselian</p> <p>Stadial</p>	fluvial		
I	<ul style="list-style-type: none"> • Marine diatoms: <i>Hyalodiscus</i> sp., <i>Paralia sulcata</i>, <i>Porosira glacialis</i>, <i>Thalassiosira</i> sp., <i>Thalassiothrix longissima</i>, <i>Centralea</i> 	Eemian	Thermokarst	Shrub tundra	temperate

ind.	Interglacial	lagoon	e
<ul style="list-style-type: none"> • Fresh water diatoms: <i>Naicula radiosa</i>, <i>Eunotia praerupta</i>, <i>Pinnularia gibba</i>, <i>tetracyclus lacustris</i> • Sponge spicula • Pollen: <i>Larix</i>, <i>Alnus fruticosa</i>, <i>Betula nana</i>, Ericales • Spores: <i>Sphagnum</i> 			

Herlemann, D. P., Labrenz, M., Jürgens, K., Bertilsson, S., Waniek, J. J. and Andersson, A. F.: Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic Sea., *ISME J.*, 5(10), 1571–9, doi:10.1038/ismej.2011.41, 2011.

Müller, S., Bobrov, A. A., Schirrmeister, L., Andreev, A. A. and Tarasov, P. E.: Testate amoebae record from the Laptev Sea coast and its implication for the reconstruction of Late Pleistocene and Holocene environments in the Arctic Siberia, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 271(3–4), 301–315, doi:10.1016/J.PALAEO.2008.11.003, 2009.

Muyzer, G., De Waal, E. C. and Uitterlinden, A. G.: Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes coding for 16S rRNA., *Appl. Environ. Microbiol.*, 59(3), 695–700, 1993.

Schirrmeister, L., Grosse, G., Kunitsky, V., Magens, D., Meyer, H., Dereviagin, A., Kuznetsova, T., Andreev, A., Babiy, O., Kienast, F., Grigoriev, M., Overduin, P. P. and Preusser, F.: Periglacial landscape evolution and environmental changes of Arctic lowland areas for the last 60000 years (western Laptev Sea coast, Cape Mamontov Klyk), *Polar Res.*, 27(2), 249–272, doi:10.1111/j.1751-8369.2008.00067.x, 2008.

Winterfeld, M., Schirrmeister, L., Grigoriev, M. N., Kunitsky, V. V., Andreev, A., Murray, A. and Overduin, P. P.: Coastal permafrost landscape development since the Late Pleistocene in the western Laptev Sea, Siberia, *Boreas*, 40(4), 697–713, doi:10.1111/j.1502-3885.2011.00203.x, 2011.