



Organic-carbon-rich sediments: benthic foraminifera as bio-indicators of depositional environments

Elena Lo Giudice Cappelli¹, Jessica Louise Clarke^{1,2}, Craig Smeaton¹, Keith Davidson³, and William Edward Newns Austin^{1,3}

¹School of Geography and Sustainable Development, University of St Andrews, St Andrews, KY16 9AL, Scotland, UK

²School of Natural and Environmental Sciences, Newcastle University, Newcastle, NE1 7RU, UK

³Scottish Association for Marine Science, Scottish Marine Institute, Oban, PA37 1QA, Scotland, UK

Correspondence: Elena Lo Giudice Cappelli (elena.logiudice.cappelli@gmail.com)

Received: 5 April 2019 – Discussion started: 20 May 2019

Revised: 18 September 2019 – Accepted: 25 September 2019 – Published: 6 November 2019

Abstract. Fjords have been described as hotspots for carbon burial, potentially playing a key role within the carbon cycle as climate regulators over multiple timescales. Nevertheless, little is known about the long-term fate of the carbon that may become stored in fjordic sediments. One of the main reasons for this knowledge gap is that carbon arriving on the seafloor is prone to post-depositional degradation, posing a great challenge when trying to discriminate between an actual change in the carbon deposition rate and post-depositional carbon loss. In this study, we evaluate the use of modern benthic foraminifera as bio-indicators of organic carbon content in six voes (fjords) on the west coast of Shetland. Benthic foraminifera are known to be sensitive to changes in organic carbon content in the sediments, and changes in their assemblage composition therefore reflect synchronous variations in the quantity and quality of carbon reaching the seafloor. We identified four environments based on the relationship between benthic foraminiferal assemblages and organic carbon content in the sediments: (1) land-locked regions influenced by riverine and/or freshwater inputs of organic matter, namely the head of fjords with a restricted geomorphology; (2) stressed environments with a heavily stratified water column and sediments rich in organic matter of low nutritional value; (3) depositional environments with moderate organic content and mild or episodic current activity; and (4) marginal to coastal settings with low organic content, such as fjords with an unrestricted geomorphology. We conclude that foraminifera potentially provide a tool to disentangle primary organic carbon signals from post-depositional degradation and loss of organic carbon because of their en-

vironmental sensitivity and high preservation potential in the sedimentary record.

1 Introduction

Globally, fjords sequester about 18 Mt of organic carbon (OC) annually in their sediments, partly acting to reduce the remineralisation of OC into the environment and thus buffering the release of carbon dioxide (CO₂) to the atmosphere (e.g. Smith et al., 2015; Cui et al., 2016). A recent study by Smeaton et al. (2017) estimated that the sediments of Scotland's 111 major fjords hold 295 ± 52 Mt OC. In Shetland, these fjordic systems are different from their mainland counterparts, notably in that they are shallower and less glaciated (Edwards and Sharples, 1986) and peat or peaty soils dominate their catchments (Soil Survey of Scotland, 1981). The erosion of terrestrial organic matter (OM) is known to be a major source of OC in fjordic sediments (Cui et al., 2017), and Scottish peatlands hold 1620 Mt of OC (Chapman et al., 2009). Shetland peatlands cover over one-third of the archipelago's total area, and their erosion accounts for a loss of 1 to 4 cm yr⁻¹ from peat surfaces (Birnie, 1993). Eroding peatlands may therefore hold the potential to contribute significantly to OC burial and accumulation in Shetland's voes and to the overall coastal OC store of this region. Nevertheless, little is known about the long-term fate of the OC that may accumulate and become stored in these sediments. One of the main reasons for this knowledge gap is that OC is prone to post-depositional modification and degradation. It is

therefore difficult to discern whether a down-core decrease in OC content is due to an actual change in carbon deposition over time or simply to this post-depositional loss (Hülse et al., 2017; Stolpovsky et al., 2017).

Benthic foraminiferal assemblages have long been used as indicators of environmental change and to reconstruct the intrinsic characteristics of marine ecosystems. A recent study (Duffield et al., 2017) explored the relationship between foraminiferal assemblages and variations in OC content in the surficial and recent sediments of Norwegian fjords, highlighting temporal trends in OC fluxes and ecological quality status. In this study, we evaluate the use of modern benthic foraminifera as bio-indicators of OC content in six voes (fjords) on the west coast of Shetland whose sedimentation dynamics are likely influenced by the erosion of OC-rich peat from the surrounding catchments (Fig. 1). As benthic foraminifera are sensitive to changes in OC content in the sediments (Alve et al., 2016, and references therein), changes in their assemblage composition should reflect synchronous variations in the amount of OC reaching the seafloor. Given the high preservation potential of their tests in marine sediments (e.g. Murray, 2003; Thomas et al., 1995), foraminifera potentially offer a means to discriminate primary OC signals from post-depositional OC degradation and may provide a tool to validate studies and models reconstructing past changes in the carbon cycle (Hülse et al., 2017; Stolpovsky et al., 2017). Nevertheless, the relationship between OC and foraminiferal assemblages is not straightforward and warrants further investigation.

Early studies developed conceptual models explaining benthic foraminiferal microhabitat preferences in deep sea settings based on changes in the organic flux reaching the seafloor and bottom water oxygenation (Jorissen et al., 1995; Van Der Zwaan et al., 1999). Researchers have found that in well-oxygenated bottom waters, food is usually the limiting factor controlling the distribution of benthic foraminifera, whereas under eutrophic conditions, bottom water oxygenation may become the dominant control on the composition of foraminiferal assemblages (Jorissen et al., 1995; Van Der Zwaan et al., 1999). Typically, food availability is strongly connected to OC fluxes and the transport of organic matter to the seafloor. In this context, it becomes crucial to understand not only food availability in terms of the quantity of OM reaching the seafloor, but also to understand the quality of this food source. For example, refractory organic matter (ROM), often derived from reworked terrestrial sources, is known to be of low nutritional value for benthic foraminifera compared to the more bio-available form of labile organic matter (LOM) (e.g. Fontanier et al., 2008; Jorissen et al., 1998). It should also be noted that the OC accumulation rate and bottom water oxygen concentrations are often related, with low oxygen concentrations reducing the reworking or remineralisation of OM and thus favouring its preservation in marine sediments. In certain environments, this implies that similar OC fluxes could result in different OC accumu-




Figure 1. Location map of Shetland and the sampling stations.

lation rates depending on bottom water oxygenation, the latter defining the composition of benthic foraminiferal assemblages (Van Der Zwaan et al., 1999).

To investigate the relationships between sedimentary OC in six west Shetland voes and the associated changes in benthic foraminiferal assemblages, this study combines geochemical and micropalaeontological techniques to do the following: (1) fingerprint the source (terrestrial vs. marine) and quality (refractory vs. labile) of organic matter and the form (organic vs. inorganic) of sedimentary carbon; (2) establish the benthic foraminiferal distribution in Shetland's voes from recent surficial sediments; and (3) investigate the use of benthic foraminifera as bio-indicators of OC content in coastal sediments and their potential for palaeo-OC reconstruction purposes.

Table 1. Station data.

Location	Sample ID	Longitude	Latitude
Clift Sound	MD15-01	−1.277267	60.11943
Clift Sound	MD15-02	−1.284533	60.09987
Clift Sound	MD15-03	−1.2975	60.07995
Clift Sound	MD15-04	−1.307783	60.06635
Clift Sound	MD15-05	−1.315883	60.04765
Sand Sound	MD15-06	−1.40295	60.24485
Sand Sound	MD15-07	−1.404183	60.24477
Sand Sound	MD15-08	−1.379517	60.24237
Sand Sound	MD15-09	−1.35835	60.24178
Sand Sound	MD15-10	−1.370517	60.22822
Olna Firth	MD15-11	−1.295883	60.36277
Olna Firth	MD15-12	−1.32315	60.36425
Aith Voe	MD15-13	−1.374883	60.28945
Aith Voe	MD15-14	−1.373333	60.3055
Busta Voe	MD15-15	−1.361217	60.39075
Busta Voe	MD15-16	−1.358883	60.38063
Busta Voe	MD15-17	−1.370517	60.3694
Vaila Sound	MD15-18	−1.564083	60.21943
Vaila Sound	MD15-19	−1.5624	60.213
Vaila Sound	MD15-20	−1.582167	60.21428
Vaila Sound	MD15-21	−1.567467	60.20567

2 Material and methods

2.1 Area of study and sampling strategy

Shetland is a subarctic archipelago covering a total area of 1466 km², lying about 170 km north of mainland Scotland, UK (Fig. 1). Several voes dissect Shetland's coastline, providing an extended route for the transport of OC-rich material from land to sea (Cui et al., 2016). The current study focuses on six voes on Shetland's west coast: Clift Sound, Sand Sound, Olna Firth, Aith Voe, Busta Voe and Vaila Sound (Fig. 1, Table 1). These six voes are characterised by different geomorphologies, freshwater inputs, proximity to the open sea and intensity of local currents; together these characteristics determine the amount of carbon deposited in the sediments. Clift Sound lies south of Shetland, with the islands of Trondra and East Burra to the west, and has steep sides and a relatively unrestricted geomorphology (Fig. 1). Two main types of soil are present in the catchment of Clift Sound: humus–iron podzols cover the islands of East Burra and Trondra, while peaty soils dominate the eastern coast of Clift Sound (Fig. S1 in the Supplement). Winds are generally from the southwest and can be channelled, resulting in intensified currents due to the geometry of the sound and the surrounding highlands. Tidal energy can also be channelled into more vigorous currents (Cefas, 2007a).

Sand Sound is somehow T-shaped, located in the southwest of Shetland and comprises three areas: the head of the voe, the inner basin and the outer basin (Fig. 1). The head of the voe, almost perpendicular to the inner and outer basins,

is a sheltered area that receives freshwater from a significant number of rivers draining the surrounding land. Three main types of soil characterise this area: peaty gleys, peaty podzols and peaty rankers (Fig. S1). Here, the voe is very shallow and drying can occur in its eastern and western periphery (Cefas, 2007b, 2008b). In contrast, the rest of the voe is much deeper. The inner basin gently slopes away, reaching more than 20 m at its maximum depth; drying can still occur where this basin meets the head of the voe (Cefas, 2007b). The outer basin has steep sides; it is 42 m deep at its deepest point and completely open to sea. A shallow sill divides the inner from the outer basins (Cefas, 2007b).

Busta Voe, Olna Firth and Aith Voe are part of a major inlet on the southern coastline of St Magnus Bay on the west coast of Shetland (Fig. 1). Olna Firth represents the eastern branch of the inlet and is roughly oriented east–west; it is the furthest away from the open sea (Fig. 1). A large area of this voe exceeds 30 m of water depth; the northern coastline gradually slopes into the voe, whereas the south side shows a steeper gradient. Olna Firth is classified as micro-tidal, and due to the generally low level of energy in this system, stratification may occur, especially during summertime (Cefas, 2013).

Aith Voe is the southern component of the inlet, with a north–south orientation (Fig. 1). Due to the hilly landscape (up to 100 m) surrounding the voe, Aith Voe is exposed to winds from the north, which can significantly alter local circulation and prevent stratification of the water column. Two types of soil surround Aith voe: peaty and organic soils (Fig. S1); few streams drain these soils, with the largest of them discharging on the east coast of Aith Voe (Cefas, 2010).

Busta Voe lies in the northern part of the inlet and is oriented north–south, sheltered and with a maximum water depth of 39 m (Fig. 1). Three main types of soil surround this area: peaty gleys, peaty podzols and peaty rankers (Fig. S1); two rivers drain a relatively small catchment area into Busta Voe. Tidal flow is weak in the voe and wind-generated currents are predominant; stratification of the water column, especially during warm periods, can occur (Cefas, 2008a).

Vaila Sound lies on the western coast of Shetland, with the island of Vaila to the southwest and the isle of Linga in the middle (Fig. 1). Three types of soil dominate the catchment of this voe: peaty gleys, peaty podzols and peaty rankers (Fig. S1). Most streams discharge in the northern part of the sound; hence, locations in this area might become more affected by terrigenous inputs. Additionally, Linga offers protection from wind and currents, which could facilitate the localised accumulation of terrigenous material. Shelter from strong winds is also provided by the island of Vaila to the south. Here the tidal range is small and the associated energy weak; however, Vaila Sound is connected to the Atlantic Ocean west and east of the island of Vaila (Cefas, 2009).

The MV *Moder Dy* 2015 cruise in west Shetland surveyed the six voes in August 2015 (Table 1). Marine surface sediments were sampled using a Duncan and Associates Van Veen grab with a sampling area of 0.1 m². A total of 23 sur-

face sediment samples were obtained by scraping the top layer (~ 1 cm thick) of each grab with a domestic spoon; samples were then stored in a cold box in sealed plastic bottles.

An earlier field survey of Shetland voes carried out in August 2009 measured bottom water temperature (BWT), salinity (BWS) and oxygen (O_2) at the same locations as this study (Fig. 2); however, no data were collected in Vaaila Sound. Oceanographic measurements were made using a Sea-Bird SBE 19 plus conductivity–temperature–depth (CTD) profiler with a dissolved oxygen probe. Bottom water temperature varies between 14.14 and 12.20 °C, being on average 12.74 °C (Fig. 2a). Bottom water salinity remains almost constant across these voes, ranging between 35 and 35.2 psu with the exception of MD15-06, where BWS reached a minimum of 34.7 psu (Fig. 2b). Similarly, bottom waters are mostly well-oxygenated in all the voes ($O_2 > 5 \text{ mg L}^{-1}$) with the exception of Olua Firth where lower O_2 was observed (Fig. 2c).

2.2 Particle size measurement

Particle size analyses were performed on a subset of surface sediments using a Coulter LS230 particle size analyser to quantify the volume (%) of the various grain sizes in each sample. We identified three categories based on the International Organisation for Standardisation (ISO) scale: “clay” includes particles $< 2 \mu\text{m}$, “silt” particles between 2 and $63 \mu\text{m}$, and “sand” particles $> 63 \mu\text{m}$ (upper limit of $2000 \mu\text{m}$). Before analysis, sediment samples were digested using 30 % hydrogen peroxide to remove organic matter and 10 % hydrochloric acid to remove carbonate, following a similar methodology to that outlined by Austin and Evans (2010). Digestions were carried out to obtain an insoluble residue useful for comparing sediments having very different carbonate and organic matter content, which otherwise might skew the measurements because of in situ processes.

2.3 Loss on ignition (LOI) analysis

Loss on ignition measurements were carried out to quantify the percentage of total organic matter (TOM) and the respective amounts of labile (LOM) and refractory organic matter (ROM). The initial combustion temperature was set to 250 °C, as significant mass loss of LOM has been recorded at this temperature (Mook and Hoskin, 1982). Sediments were successively heated to 550 °C to quantify ROM, as refractory terrestrial and aquatic OM will likely burn off at this temperature (Kristensen, 1990). About 1 g of dried sediment was precisely weighted into crucibles of known weight for each surface sample (M_0). These were ashed in a muffle furnace at 250 °C for 4 h, cooled and weighted (M_{250}). Samples were then returned to the furnace at 550 °C for 4 h, cooled and reweighted (M_{550}). TOM, LOM and ROM were calculated

as follows:

$$\text{LOM} = [(M_0 - M_{250})/M_S] \cdot 100$$

$$\text{ROM} = [(M_{250} - M_{550})/M_S] \cdot 100$$

$$\text{TOM} = \text{LOM} + \text{ROM},$$

where M_S is the sample weight minus the crucible weight. Replicate samples were measured at stations MD15-01A, MD15-01B, MD15-05A and MD15-05B, which resulted in a mean relative error of $\pm 0.07 \%$ for LOM, $\pm 0.06 \%$ for ROM and $\pm 0.03 \%$ for TOM, pointing to very good data reproducibility and representation of local conditions.

2.4 Bulk carbon elemental and stable isotope analyses

Surface sediment samples were analysed to determine bulk elemental C and stable isotope ($\delta^{13}\text{C}$) values. Each sediment sample was freeze-dried and homogenised, and approximately 12 mg of milled sediment was placed into silver capsules; a further 10 mg was placed into tin capsules.

The samples encapsulated in silver underwent an acid fumigation step (Harris et al., 2001) to remove inorganic carbon (IC). After drying for 24 h at 40 °C, both OC and $\delta^{13}\text{C}$ were measured using an elemental analyser coupled to an isotope ratio mass spectrometer (IRMS) at the NERC Life Science Mass Spectrometer Facility (Lancaster, UK). The standard deviation of $\delta^{13}\text{C}$ triplicate measurements was 0.07 ‰, and $\delta^{13}\text{C}$ values are reported in standard delta notation relative to Vienna Pee Dee Belemnite (VPDB).

The samples in tin capsules were analysed for total carbon (TC) using an Elementar elemental analyser (EA) at the School of Geography and Sustainable Development, University of St Andrews (Verardo et al., 1990). The precision of the analysis is calculated based on repeat measurements of standard reference material B2178 (medium organic content standard from Elemental Microanalysis, UK) with $C = 0.08 \%$.

These results were combined together to calculate the quantity of IC in each sample; the percentage of OC was subtracted from the percentage of TC to obtain the percentage of IC.

2.5 Binary mixing model

To discriminate between marine-sourced and terrestrially sourced OC, we used a two-endmember (binary) mixing model assuming a two-point source of OC, marine and terrestrial, following Thornton and McManus (1994). We used $\delta^{13}\text{C}$ measurements in the sediment samples as a tracer of the source of OC in west Shetland voes and calculated the fraction of terrestrially sourced OC (OC_{terr}) as follows:

$$\text{OC}_{\text{terr}} = (\delta^{13}\text{C}_{\text{mar}} - \delta^{13}\text{C}_{\text{terr}})/(\delta^{13}\text{C}_{\text{mar}} - \delta^{13}\text{C}_{\text{sample}})$$

$$\text{OC}_{\text{terr}} + \text{OC}_{\text{mar}} = 1,$$

where the endmember value for marine-sourced OC ($\delta^{13}\text{C}_{\text{mar}}$) was taken from Smeaton and Austin (2017) based




Figure 2. CTD profiles from the 2009 survey. (a) Temperature profiles and bottom water temperature (BWT) at each sampling station. (b) Salinity profiles and bottom water salinity (BWS) at each sampling station. (c) Dissolved oxygen profiles and oxygen concentration in bottom waters at each sampling station. The dashed line at 5 mg L^{-1} indicates the possible onset of hypoxic conditions.

on phytoplankton, zooplankton, macroalgae and benthic microalgae carbon stable isotope values, while the endmember for terrestrially sourced OC ($\delta^{13}\text{C}_{\text{terr}}$) was based upon the work of Thornton et al. (2015), who studied the distribution of carbon stable isotope in Scotland's topsoil; OC_{mar} indicates the fraction of marine-derived OC. We assumed no isotope discrimination between the source and the surface sediment samples.

2.6 Benthic foraminiferal counts and statistics

Aliquots of surface sediments were stained with rose bengal to allow for the identification of living foraminifera, as detailed in Schönfeld et al. (2012). However, it should be noted that foraminiferal counts are “total” (live + dead) because the main objective of this study is to provide a tool

for the interpretation of fossil foraminiferal assemblages and their relationship with changes in OM and OC content in sediments over time (e.g. Conradsen, 1993). Our priority is to understand how benthic foraminifera respond to prevailing (long-term) environmental conditions rather than to seasonal variability. Additionally, scraping the top layer ($\sim 1 \text{ cm}$ thick) of each grab with a domestic spoon may lead to a potential underestimation of living fauna if mixing of the top layer occurred during sampling. In this scenario, the number of living foraminifera at the sediment surface will be “diluted” due to the presence of dead and/or fossil specimens from deeper sediments. Having said this, a recent study by Rillo et al. (2019) reported that historical sediment samples collected in a way that could have caused disturbance of the sediment surface (sounding and dredge) are still represen-

tative of surface conditions, and their foraminiferal assemblages can be used to reconstruct environmental changes reliably. At two sites, MD15-01 and MD15-05, replicate samples were taken (MD15-01A and MD15-01B; MD15-05A and MD15-05B, respectively) to check measurement reproducibility and seafloor heterogeneity.

As part of an unpublished work, rose-bengal-stained surface sediments were wet-sieved over a 63 µm mesh; the residues were oven-dried at < 60 °C, then weighed and dry-sieved into 63–150 µm and > 150 µm size fractions. As part of an associated work aiming to understand the influence of different size fractions on the composition of benthic foraminiferal assemblages (Lo Giudice Cappelli and Austin, 2019), both size fractions were analysed independently, and the results demonstrated that a more holistic picture of environmental change is obtained when benthic foraminiferal assemblages of both size fractions are compiled together. This improves the representation of the entire assemblage and results in more statistically robust environmental reconstructions. Similar findings were also reached by Weinkauff and Milker (2018, and references therein). Depending on sample volume, each sample was divided into a number of splits using a standard dry splitter and, when possible, at least 300 specimens were dry-picked to ensure statistical significance when discussing benthic foraminiferal assemblage diversity and composition (Schönfeld et al., 2012). Species were identified following Austin (1991) and counted using a tally sheet; notes on tests preservations were made when signs of etching were observed.

Relative benthic foraminiferal abundances were calculated for each sample. It should be noted that we grouped under the name *E. excavatum* both forma *selseyense* and forma *clavata* despite them being recognised as genetically different species (Darling et al., 2016). It was not possible to consistently identify and discriminate between these two species; hence, we used a common identifier.

The Palaeontological Statistics software package PAST (version 3.16; Hammer et al., 2001) was used to analyse relationships between benthic foraminiferal assemblages through cluster analysis, canonical component analysis (CCA) and non-metric multidimensional scaling (MDS) based on the Bray–Curtis similarity index and relative abundance data. Cluster analysis showed good reproducibility of benthic foraminiferal assemblages (Fig. 4b) with the two pairs of replicate stations exhibiting assemblage reproducibility of 90 % (MD15-01A and MD15-01B) and 94 % (MD15-05A and MD15-05B).

3 Results

3.1 Sediment analyses

3.1.1 Particle size distribution in west Shetland voes

A highly variable pattern is evident in the particle size distribution of the sediments of west Shetland voes (Fig. 3a). Clift Sound, Olva Firth, stations MD15-08 and MD15-10 in Sand Sound, and station MD15-13 in Aith Voe have silt-dominated sediments (> 65 %; Fig. 3a). Conversely, stations MD15-06, MD15-07 and MD15-09 in Sand Sound, MD15-14 in Aith Voe, MD15-15 and MD15-17 in Busta Voe, and MD15-20 and MD15-21 in Vaila Sound are dominated by sandy sediments (> 50 %) (Fig. 3a).

3.1.2 Organic matter content in west Shetland voes

Total OM is generally higher in proximity to land than at sites more exposed to the open sea (Fig. 3c). Site MD15-20 (Vaila Sound) has the lowest percentage of total OM (3.05 %), ROM (2.24 %) and LOM (0.81 %), whereas site MD15-08 (Sand Sound) has the highest percentage of total OM (20.85 %) and ROM (12.37 %). LOM is the highest (8.81 %) at site MD15-11 (Olva Firth) (Table 3; Fig. 3c).

3.1.3 Carbon content and source in west Shetland bulk sediments

The percentage of total carbon (TC) in west Shetland voes varies between 4.31 % and 14.32 % (Fig. 3d), with the highest percentage found at station MD15-05 (Clift Sound) and the lowest at station MD15-09 (Sand Sound). Organic carbon (OC) content ranges between 2.0 % and 8.51 %, peaking at station MD15-11 (Olva Firth) and reaching a minimum at site MD15-17 (Busta Voe) (Fig. 3d; Table 3). The percentage of inorganic carbon (IC) varies between 0.35 % and 7.51 % and is the highest at station MD15-03 (Clift Sound) and the lowest at station MD15-14 (Aith Voe) (Fig. 3d). We excluded from the dataset site MD15-15, as this sample contains coal, which is believed to have been shed accidentally from a commercial vessel. In general, there is a tendency for high OC and low IC at the head of the voes, and the reverse occurs where the voes connect to the sea (Fig. 3d).

In terms of a carbon source, OC_{terr} is generally higher at sites close to land than in proximity to the open sea (Fig. 3b). It reaches a peak of 72 % at station MD1-06 in Sand Sound and a minimum of 21 % at station MD15-03 (Fig. 3b).

3.1.4 Carbon stable isotopes ($\delta^{13}\text{C}$)

Bulk sediment $\delta^{13}\text{C}$ varies between -25.41‰ and -21.26‰ with an average value of -22.67‰ (Table 3). The lowest value (-25.41‰) is recorded at site MD15-06 (Sand Sound) and the highest (-21.26‰) at station MD15-




Figure 3. Sediment physical properties. **(a)** Particle size analysis. **(b)** Fraction of terrestrially sourced OC (OC_{terr}) and marine-sourced OC (OC_{mar}) at each site. **(c)** Refractory and labile organic matter content at each site and carbon stable isotope. **(d)** Inorganic and organic carbon percentage at each site and carbon stable isotope.

03 (Clift Sound); high $\delta^{13}\text{C}$ values characterise marine-dominated locations within each voe (Fig. 3c and d).

3.2 Benthic foraminiferal assemblages in west Shetland voes

We observed a total of 64 species of benthic foraminifera at the 21 stations analysed in this study, including the two replicates samples (Fig. S2). Nine of the observed species are agglutinated: *Ammoscalaria runiana*, *E. scaber*, *Connemarella rudis*, *Haplophragmoides* sp., *Reophax fusiformis*, *Reophax scotii*, *Spiroplectinella wrightii*, *Textularia earlandii* and *Trochammina* sp. Of these, only *E. scaber* and *Trochammina* sp. were observed at all 21 stations. However, the *E. scaber* relative abundance varies from 0.4 % in Vaila Sound (MD15-20) to 43.0 % in Sand Sound (MD15-10) (Fig. 5), whereas the *Trochammina* sp. relative abundance reaches a maximum of 2.3 % in Clift Sound. Other common agglutinated species are *Haplophragmoides* sp. and *S. wrightii*, with a relative abundance ranging from 0.1 % (Vaila Sound) to 2.9 % (Busta Voe) and 0.1 % (Olna Firth) to 3.9 % (Clift Sound), respectively. The distribution of the remaining agglutinated species is patchy, and their relative abundance varies around the average value of 1 %.

A total of 6 of the 64 observed species belong to the porcelaneous group Miliolina: *Cornuspira* sp., *Miliolinella subrotunda*, *Quinqueloculina bicornis*, *Quinqueloculina seminula*,

Quinqueloculina sp. and *Spiroloculina rotunda*. Of these, the most abundant species is *Q. seminula*, which is present in low numbers in five of the six studied fjords; it does not occur in Aith Voe. The general distribution of Miliolina is patchy, and their relative abundance varies around the average value of 0.5 %, with the highest value of 2.4 % observed at the mouth of Clift Sound (MD15-05B).

The remaining 49 species are hyaline foraminifera of which 9 species have relative abundances of > 10 % for at least one of the 21 stations studied: *Ammonia* spp., *Bulimina marginata*, *Buliminella elegantissima*, *Cibicides* spp., *Elphidium gerthi*, *Elphidium margaritaceum*, *Elphidium excavatum*, *Rosalina* spp. and *Stainforthia fusiformis* (Table 2). Additionally, of all the hyaline species, 20 are rare, having relative abundances < 1 %, while 18 species are frequent, having a relative abundance between 1 % and 5 %, and only 2 species are common, having relative abundances ranging between 5 % and 10 % (Fig. S2).

Overall, 10 taxa of benthic foraminifera were observed in each of the six voes and had relative abundances of more than 10 % for at least one of the 21 stations studied (Table 2): *Ammonia* spp., *Bulimina marginata*, *Buliminella elegantissima*, *Cibicides* spp., *Eggerelloides scaber*, *Elphidium gerthi*, *Elphidium margaritaceum*, *Elphidium excavatum*, *Rosalina* spp. and *Stainforthia fusiformis*. These 10 taxa were categorised as dominant and imaged with a scanning electron mi-




Figure 4. Statistical analyses. **(a)** Cluster analysis of sediment and seawater physical properties. **(b)** Cluster analysis of benthic foraminiferal assemblages. **(c)** Canonical component analysis of the combined datasets (sediment and seawater + benthic foraminifera relative abundance).

croscopie (SEM) at the Scottish Oceans Institute, University of St Andrews (Fig. 5). Altogether, these 10 dominant taxa account for more than 70 % of the total assemblage at each station (Table 2). In general, the most abundant taxa observed in west Shetland's six voes is the hyaline *Cibicides* spp., followed by the agglutinated *E. scaber* (Table 2 and Fig. 5).

3.2.1 Statistical analyses of benthic foraminifera data and environmental parameters

We used a cluster analysis of environmental parameters (BWT, BWS, O₂, % Clay, OC, IC, ROM, LOM, δ¹³C and WD) based on Euclidean distance and of benthic foraminifera relative abundance based on the Bray–Curtis index to illustrate similarities between locations (Fig. 4a and b). We performed CCA on the combined dataset (benthic foraminifera + environmental parameters) to examine the response of benthic foraminiferal assemblages to environmental gradients (Fig. 4c). Additionally, we run non-metric MDS based on the Bray–Curtis similarity index and foraminifera relative abundance data to analyse similarities between the populations of the 21 studied stations, independently from environmental forcing (Fig. 5). Note that temperature, salinity and oxygen measurements are from 2009 and may not fully represent the conditions in 2015 when the sediment samples were collected.

Based on both CCA and non-metric MDS, we defined the same four groups with 75 % confidence; in the CCA plot, axis 1 explains 57 % of the total variability and axis 2 explains 26 % based on the 10 environmental parameters (Figs. 4c and 5). Group 1 includes assemblages dominated by the genus *Elphidium* and by *E. scaber*; the head of Clift Sound falls into this cluster (MD15-06 to MD15-09). Group 2 contains assemblages dominated by *E. scaber* with *S. fusiformis* and *B. marginata* as associated species; the inner basin of Sand Sound (MD15-10), Olua Firth (MD15-11 and MD15-12) and MD15-16 in Busta Voe are part of this cluster. Group 3 consists of assemblages characterised by relatively high diversity and no obvious dominant species; MD15-01A and MD15-01B (head of Clift Sound), MD15-13 and MD15-14 (Aith Voe), MD15-15 (Busta Voe), and MD15-18 and MD15-19 (head of Vaila Sound) belong to this group. Group 4 comprises assemblages dominated by the genus *Cibicides* with *Rosalina* as associated taxa and includes stations MD15-02 to MD15-05 in Clift Sound and stations MD15-20 and MD15-21 in Vaila Sound (Figs. 4c and 5). Station MD15-17 in Busta Voe falls just outside the 75 % confidence interval defining Group 4; however, cluster analysis (Fig. 4b) shows ~ 82 % similarity between the assemblage of this site and station MD15-20 and more than the 70 % similarity with the rest of the locations belonging to




Table 2. Relative abundance of the 10 dominant taxa.

Location	Sample ID	<i>Ammonia</i> spp.	<i>B. marginata</i>	<i>B. elegantissima</i>	<i>Cibicoides</i> spp.	<i>E. scaber</i>	<i>E. gerthi</i>	<i>E. margaritaceum</i>	<i>E. excavatum</i>	<i>Rosalina</i> spp.	<i>S. fusiformis</i>	Total %
Clift Sound	MD15-01A	0.2	0.5	2.1	14.5	22.0	10.7	20.1	7.8	3.8	3.0	84.6
Clift Sound	MD15-01B	0.2	0.6	2.1	15.0	19.9	10.7	24.3	2.1	3.2	4.4	82.5
Clift Sound	MD15-02		0.7	0.3	47.6	10.6	5.4	3.1	1.9	6.6	3.8	80.3
Clift Sound	MD15-03	0.2	1.2	0.7	53.7	3.3	10.5	0.8	1.2	5.4	4.6	81.5
Clift Sound	MD15-04	0.3	0.9	0.3	48.9	5.2	13.9	0.2		4.7	2.9	77.3
Clift Sound	MD15-05A		0.3		57.6	2.1	0.4	0.1		9.7	0.7	71.0
Clift Sound	MD15-05B		0.6		56.7	2.1	0.2	0.0	0.5	10.4	1.5	72.0
Sand Sound	MD15-06	4.8	0.5	4.6	2.0	25.5	15.6	6.1	15.3		1.3	75.7
Sand Sound	MD15-07	6.5	0.8	4.8	5.5	27.7	11.7	7.6	21.5	0.7		86.7
Sand Sound	MD15-08	3.0	1.7	11.4	2.0	29.7	7.5	1.4	22.4	0.2	6.3	85.6
Sand Sound	MD15-09	5.4	0.9	8.8	1.9	19.7	9.8	10.1	25.6	0.2	1.6	83.9
Sand Sound	MD15-10	0.9	3.8	1.9	8.3	43.0	4.7	1.3	5.2	1.4	11.3	81.8
Olna Firth	MD15-11		14.8	0.8	10.9	31.9	4.4	1.8	0.8	0.5	23.1	88.8
Olna Firth	MD15-12	8.5	8.6	0.7	13.7	22.9	2.2	0.9	2.4	0.3	26.0	86.2
Aith Voe	MD15-13	3.0	5.6	12.6	14.2	11.7	9.3	4.0	10.0	0.3	7.3	78.1
Aith Voe	MD15-14	12.6	4.1	4.6	26.1	10.0	8.2	0.9	4.3	1.8	8.2	80.8
Busta Voe	MD15-15	1.0	2.8	2.0	30.1	12.9	7.0	5.1	5.6	0.8	8.2	75.5
Busta Voe	MD15-16	5.7	8.5	0.9	7.6	26.4	5.0	1.2	5.1	0.2	23.1	83.7
Busta Voe	MD15-17	2.2	1.2	0.1	70.1	2.8	1.4	0.8	1.5	1.1	4.8	86.1
Vaila Sound	MD15-18	6.1	0.9	11.5	24.4	9.5	7.6	9.0	9.2	2.3	4.0	84.5
Vaila Sound	MD15-19	10.2	2.1	1.8	29.6	8.8	8.3	3.4	9.2	4.5	4.3	82.3
Vaila Sound	MD15-20		0.4	0.1	68.7	0.4	1.2	0.9	1.2	14.7	0.5	88.2
Vaila Sound	MD15-21	0.3	1.4	0.3	56.8	1.9	1.9	1.0	2.9	11.3	1.4	78.9

Group 4. Therefore, for the purposes of our discussion, we include station MD15-17 in Group 4.

4 Discussion

4.1 Particle size distribution in west Shetland voes

The particle size data from the voes of Shetland exhibit a large range with the silt and sand fractions but little variation in the proportion of clay (Fig. 3a). These samples suggest a significant size sorting, which is consistent with the tidal and estuarine dynamics of these voes. In Vaila Sound, intensified bottom currents and winnowing of finer sediments owing to the influx of Atlantic Ocean waters (Cefas, 2009) appear to control local grain size distribution (Fig. 3a). Here, stations exposed to the open sea and intensified bottom currents are characterised by coarser sediments (MD15-20 and MD15-21), whereas more sheltered and calm locations (MD15-18 and MD15-19) are characterised by finer sediments (Fig. 3a). In contrast, sediments in Clift Sound are silt-dominated (Fig. 3a) despite the unrestricted geomorphology (Fig. 1) and vigorous bottom currents (Cefas, 2007a). It appears that a mechanism other than winnowing of finer

sediments by intensified bottom currents drives particle size distribution here. In Clift Sound, the overall low percentage of OM, together with the relatively high IC content and $\delta^{13}\text{C}$ values, points to a reduction in terrigenous input as the main driver of particle size distribution (Fig. 3). Conversely, in Sand Sound, an increased terrigenous input appears to be responsible for the coarser sediments found at stations MD15-06, MD15-07 and MD15-09 (Fig. 3). The head of the voe is a sheltered area that receives freshwater from a significant number of rivers draining the surrounding land (Cefas, 2007b, 2008b), resulting in an overall high percentage of OM and OC, low $\delta^{13}\text{C}$ values (Fig. 3c and d), and coarser sediments (Fig. 3a). In contrast, station MD15-08 at the head of Sand Sound is characterised by finer sediments (Fig. 3a) despite the high terrigenous supply at this location (Fig. 3c and d). It is possible that, due to the restricted geomorphology of Sand Sound (Fig. 1), weak currents are unable to transport the coarse material to station MD15-08, the deepest station at the head of the voe (11 m vs. 6 m). In the Sand Sound inner basin (MD15-10), silty sediments dominate (Fig. 3a). Here, a combination of reduced terrigenous input and/or weaker currents could be driving particle size distribution. Nørgaard-Pedersen et al. (2006) found similar

Table 3. Sediment physical properties.

Location	Sample ID	Water depth (m)	$\delta^{13}\text{C}$	%IC	%OC	%LOM	%ROM	%OC _{terr}
Clift Sound	MD15-01A	13	−23.16	3.55	5.44	4.35	6.99	45
Clift Sound	MD15-01B	13	−23.16	3.55	5.44	3.94	7.29	45
Clift Sound	MD15-02	15	−21.92	7.07	2.80	1.59	2.95	30
Clift Sound	MD15-03	23	−21.26	7.51	3.48	1.60	3.38	21
Clift Sound	MD15-04	21	−21.51	4.72	3.58	1.78	3.79	25
Clift Sound	MD15-05A	20	−21.35	7.12	7.20	1.32	4.47	23
Clift Sound	MD15-05B	20	−21.35	7.12	7.20	1.40	4.09	23
Sand Sound	MD15-06	6	−25.41	2.41	4.33	5.32	8.50	72
Sand Sound	MD15-07	6	−24.07	0.53	4.51	2.63	4.26	56
Sand Sound	MD15-08	11	−23.84	1.67	8.28	8.48	12.37	53
Sand Sound	MD15-09	6	−23.64	0.53	3.78	3.07	4.24	50
Sand Sound	MD15-10	21	−22.87	5.57	4.80	4.92	6.68	41
Olna Firth	MD15-11	31	−22.36	3.02	8.51	8.81	10.89	35
Olna Firth	MD15-12	35	−22.65	2.15	7.54	8.07	11.83	38
Aith Voe	MD15-13	12	−23.55	4.70	5.44	5.54	7.07	49
Aith Voe	MD15-14	36	−23.38	0.35	4.48	2.06	3.92	47
Busta Voe	MD15-15	18	–	–	–	2.99	8.27	–
Busta Voe	MD15-16	26	−23.06	7.40	3.27	5.48	9.19	43
Busta Voe	MD15-17	24	−22.59	5.47	2.06	2.54	3.93	38
Vaila Sound	MD15-18	14	−21.82	3.70	4.79	3.03	6.00	28
Vaila Sound	MD15-19	21	−21.54	3.91	3.51	2.86	6.38	25
Vaila Sound	MD15-20	19	−21.6	4.07	4.07	0.81	2.24	26
Vaila Sound	MD15-21	21	−21.75	5.52	2.89	1.16	3.03	27

drivers of particle size distribution in Loch Etive, a sea loch in mainland western Scotland. Vigorous bottom currents and/or enhanced terrigenous supply would result in locally coarse sediments, whereas weaker currents and/or limited terrigenous supply would result in finer sediments. Due to the duality of drivers determining particle size distribution in west Shetland voes (bottom current intensity vs. terrigenous supply), it is challenging to consistently group locations based on granulometry data (Fig. 3a) or identify a relationship between grain size distribution and OM and OC content in the sediments (Fig. S3).

4.2 Spatial distribution, quantity, quality, and source of organic matter and carbon in west Shetland voes

A clear pattern is evident with regard to the quantity, quality and source of organic matter and carbon measured in the six west Shetland voes (Fig. 3). Typically, carbon is present in the sediments as IC and OC, and the distributions of these two forms of carbon tend to mirror one another, with low IC and high OC percentages towards land, while the reverse occurs close to the open sea (Fig. 3d). Organic matter follows the same pattern of OC distribution regardless of its quality (refractory vs. labile); however, ROM content in the sediments is always higher than LOM (Fig. 3c) owing to its resilience to degradation.

About the source of OM and OC, low $\delta^{13}\text{C}$ and high OC_{terr} values characterise the head of the voes where high ROM

and OC are accumulated, indicating that the source of this material is terrestrial (Fig. 3). Terrestrial OM is more stable and resistant to decay than marine OM (Batten, 1996) and it is enriched in ^{12}C from soils, resulting in lower $\delta^{13}\text{C}$ values than marine OM derived from algae (e.g. Balasse et al., 2005; Ficken et al., 1998; Marconi et al., 2011; Schiener et al., 2014; Schmidt and Gleixner, 2005). Sand Sound is dominated by organic-rich and terrestrially sourced material as it receives freshwater and terrestrial OM from the numerous streams draining the surrounding land (Fig. 3). In contrast, Clift and Vaila Sound are characterised by marine-sourced material and high IC content in their sediments (Fig. 3) as the inflow of Atlantic Ocean waters improves the voes' potential for IC storage (Burrows et al., 2017; Smeaton et al., 2016, 2017). Olna Firth has the highest LOM percentage as this relatively deep and micro-tidal basin (Cefas, 2013) is prone to water mass stratification, which can lower bottom water oxygenation and slow down the reworking or remineralisation of OM, favouring the preservation of the more labile material (Fig. 3).

A slight deviation from this overall trend was recorded at station MD15-05. We speculate that the unusually high TC content at station MD15-05 at the mouth of Clift Sound (Fig. 3d) could relate to the MV *Braer* oil spill of 1993. It appears that after the spill, due to the adverse weather, the oil became deposited in the sediment of Burra Haaf, just west of Clift Sound (Fisheries Research Services). Despite the fact that oil levels in the contaminated sediments have been de-

creasing steadily over the intervening years, it is plausible that the OC measurements in this study detected remnants of this event. Nevertheless, the benthic foraminiferal assemblage of this location is dominated by *Cibicides* spp. (see Sect. 4.3.4 for more details), which does not reflect the oil spill or the high OC content, leaving an open question regarding what drives OC deposition and accumulation at the mouth of Clift Sound.

4.3 Modern benthic foraminiferal distribution in Shetland

4.3.1 Group 1 – *Elphidium* spp. and *E. scaber*

The genus *Elphidium* becomes the dominant taxa (relative abundance $\sim 38\%$) at the head of Sand Sound (Fig. 5), a very shallow (max depth 11 m) and organic-rich basin (OC 5.2 %, ROM 7.3 % and LOM 4.9 %; Fig. 4c). Specifically, *E. excavatum* is the most abundant *Elphidium* ($\sim 21\%$; Fig. 5), as it can live close to freshwater outflows in low-energy environments and tolerates high OC and OM concentrations (Fig. 4c) (Alve and Murray, 1999; Jennings, 2004; Korsun et al., 2014; Mendes et al., 2012). This is consistent with streams draining into the head of Sand Sound transporting terrestrial and organic-rich material into the voe (Figs. 3 and 4c). Additionally, the high relative abundance of *E. scaber* (26 %) at the same locations (Fig. 5) purports a low-energy environment, possible water mass stratification and high ROM (Fig. 4c) (Evans et al., 2002; Fontanier et al., 2002; Murray, 1992; Nørgaard-Pedersen et al., 2006). Thus, assemblages dominated by *Elphidium* spp. and *E. scaber* are likely to be found in land-locked regions influenced by riverine and/or freshwater inputs like the head of fjords with a restricted geomorphology.

4.3.2 Group 2 – *E. scaber*, *B. marginata* and *S. fusiformis*

Eggerelloides scaber, *B. marginata* and *S. fusiformis* are species typical of organic-rich sediments, stratified waters and possibly low oxygen concentrations (Fig. 4c), often suggesting stressed environments (Alve, 1994, 2003; Evans et al., 2002; Fontanier et al., 2002; Murray, 1992; Nørgaard-Pedersen et al., 2006). *Bulimina marginata* ($\sim 9\%$) seems to thrive in regions with high ROM (9.6 %) due to its ability to use OM of low nutritional value as food (Klitgaard-Kristensen and Buhl-Mortensen, 1999) (Figs. 4c and 5). *Stainforthia fusiformis* ($\sim 21\%$) is a known opportunistic species which follows high OM gradients at stratification fronts (Alve, 2003; Alve and Murray, 1997; Scott et al., 2003). In general, high organics in the sediments can result in low oxygen concentrations at the seafloor (Fig. 4c) due to eutrophication and increased respiration (Bianchi et al., 2016). Additionally, the production of humic acids as a by-product of organic matter biodegradation may lower the pH

(Bauer and Bianchi, 2012), making the environment overall more hostile for foraminifera to calcify, which could result in the dominance of agglutinated species like *E. scaber* ($\sim 31\%$) (Fig. 5). Etching of *Ammonia* spp. shells (Fig. S2) in Olva Firth supports the local acidification of bottom waters, which is substantiated by the low IC content (2.6 %) of stations MD15-11 and MD15-12 (Fig. 3). Therefore, assemblages characterised by *E. scaber*, *B. marginata* and *S. fusiformis* reflect a stressed environment with a heavily stratified water column, possibly low oxygen levels and organic-rich sediment, especially in ROM of low nutritional value. Despite the fact that seawater properties (T , S and O_2) were measured in 2009, they seem to be representative of seawater characteristics in 2015 when the sediment samples were collected, as the two sets of measurements (seawater and sediments) appear to be in agreement, reflecting long-term environmental dynamics.

4.3.3 Group 3 – no obvious dominant taxa

The assemblage of Group 3 is characterised by a high relative abundance of taxa with affinities for OC (4.9 %) and OM (6.3 %) like *Elphidium* (8 %), *E. scaber* (13 %) and *B. elegantissima* (5 %) (e.g. Mendes et al., 2012; Alve and Murray, 1999), as well as high-energy taxa like *Cibicides* (22 %) and *Ammonia* (5 %) (Scott et al., 2003) (Figs. 4c and 5). The low relative abundance of *B. marginata* (2 %) and *S. fusiformis* (5 %) additionally suggests that OM at these locations has good nutritional value, and current activity is present at these sites (reduced stratification, possibly vertical mixing). Therefore, assemblages showing this composition are likely to reflect depositional environments with mild or episodic current activity and moderate organic content.

4.3.4 Group 4 – *Cibicides* spp. and *Rosalina* spp.

In Group 4, the dominant taxa is *Cibicides* ($\sim 57\%$) followed by *Rosalina* (8 %) (Fig. 5), which reflect marginal to open-sea conditions and vigorous bottom currents (Fig. 4c) (Alve and Murray 1999; Jennings, 2004; Klitgaard-Kristensen and Sejrup, 2002). A clear relationship is evident between low organics (OC 4.2 %, ROM 3.2 % LOM 1.5 %) (Fig. 4c) and the dominance of *Cibicides* (Figs. 4c and 5). All locations falling in Group 4 are on the path of Atlantic Ocean waters entering the fjords, meaning that this assemblage composition is most likely found in areas dominated by sea processes.

4.3.5 Other drivers influencing modern benthic foraminifera distribution

Seawater properties can compete with OC and OM in influencing the distribution of benthic foraminifera across the six voes; BWS and the O_2 concentration in particular may exert a 2nd-order control on the composition of foraminiferal assemblages (Jorissen et al., 1995; Van Der Zwaan et al., 1999). A highly variable BWS generally influences the distribution of

stenohaline (strictly marine species) and euryhaline species (able to adapt to a wide range of salinity) along BWS gradients; however, in Shetland, BWS is almost constant across the voes (Fig. 2), having a very minimal influence on benthic foraminiferal assemblages (Fig. 4c). Similarly, bottom waters are mostly well-oxygenated ($[O_2] > 5 \text{ mg L}^{-1}$) in all the voes (Fig. 2), meaning that food availability is the main driver of changes in the composition of benthic assemblages (Jorissen et al., 1995; Van Der Zwaan et al., 1999). It should be noted that the seawater properties reported in Fig. 2 were measured in 2009 and may differ from the actual conditions in 2015 when sediment samples were collected; nevertheless, our results show that similar dynamics were most likely in place in 2015, as the two sets of measurements compare very well in depicting local environmental dynamics.

Food availability is deeply connected to organic fluxes and the transport of organic matter to the seafloor; hence, relationships are found between benthic foraminiferal assemblages and depositional environments (Figs. 4c and 5). Only one station in Olna Firth (MD15-11) has poorly oxygenated bottom waters ($O_2 < 5 \text{ mg L}^{-1}$), which could represent the onset of hypoxic conditions (Vaquer-Sunyer and Duarte, 2008) potentially affecting the distribution of benthic foraminifera (Fig. 2). The benthic foraminiferal assemblage of this location reflects not only organic-rich sediments, but also stratified waters and low oxygen concentrations, pointing to a stressed environment (Fig. 4c). However, this is far from the dysoxic–anoxic conditions required for a switch from food-limited to oxygen-limited conditions (Fig. 2). Therefore, the quantity and quality (labile vs. refractory) of organic matter reaching the seafloor is still the major control on the benthic foraminiferal assemblage found at station MD15-11.

4.4 Comparison with other fjords

4.4.1 Spatial distribution and sources of OM and C

The processes governing OM and C inputs in west Shetland voes are comparable with those reported for other Scottish fjords by Smeaton et al. (2017), who found a relationship between the OC content in Scottish fjords and their physical characteristics such as tidal range, precipitation, catchment area and runoff. However, as catchments in west Shetland transport sediment from the adjacent C-rich peatlands into the marine system, west Shetland voes overall store more TC (Fig. S1) compared to other Scottish fjords (Smeaton et al., 2016). On average, the TC content of the six west Shetland voes is 8.8 %, whereas TC in other Scottish fjords ranges between 0 % and 8 % (Smeaton et al., 2017 in Fig. 5). Nevertheless, as stated in Smeaton et al. (2017), there are similarities between Scotland's mainland fjords and these six voes in west Shetland. For example, fjords like Loch Broom and Little Loch Broom in Scotland have relatively unrestricted geomorphologies, which makes them highly efficient in cap-

turing IC, similar to Clift Sound and Vaila Sound in Shetland (Fig. 3). Transient low oxygen concentrations in Scotland's mainland fjords seem to be an important factor in the burial and preservation of OC (Gillibrand et al., 2006). Similarly, low oxygen levels in Olna Firth (Fig. 2), also indicated by benthic foraminiferal assemblages (Fig. 4c), are associated with the preservation of labile organic matter and OC (Figs. 3 and 4c). Additionally, carbon stable isotope values in Shetland are in range with carbon stable isotope measurements of other Scottish sea lochs: Sunart (average -22.37 ‰ , ranging from -24.42 ‰ to -21.37 ‰), Teacius (average -24.79 ‰ , $n = 15$) (Smeaton and Austin, 2017), Creran (average -21.98 ‰ varying between -24.7 ‰ and -17.3 ‰) and Etive (average -25.8 ‰ , $n = 3$) (Loh et al., 2008).

4.4.2 Benthic foraminiferal assemblages

Our knowledge of benthic foraminiferal assemblages in Scottish fjords is limited to studies based on fauna from Loch Etive (Murray et al., 2003; Nørgaard-Pedersen et al., 2006) and their response to changes in sea level and terrigenous input entering the catchment. Species in common with west Shetland voes include *Cibicides lobatulus*, *Ammonia batavus* and *E. scaber*. Assemblages dominated by *C. lobatulus* and *A. batavus* in Loch Etive indicate marine conditions and intensified bottom currents possibly associated with deepwater renewal events (Nørgaard-Pedersen et al., 2006). This agrees well with the assemblages dominated by *Cibicides* spp. and *Rosalina* spp. observed in Clift Sound and Vaila Sound suggestive of high-energy environments (Fig. 5). The dominance of agglutinated species such as *E. scaber* occurs in Loch Etive when OM deposition increases, eventually resulting in low oxygen levels in bottom waters and carbonate dissolution (Murray et al., 2003; Nørgaard-Pedersen et al., 2006). We observed similar environmental conditions and assemblages in Olna Firth, the middle of Busta Voe and the inner basin of Sand Sound (Figs. 4c and 5).

Similarities can also be observed between assemblages from the Scottish coast and shelf and west Shetland voes. Klitgaard-Kristensen and Sejrup (2002) and Murray (2003) found a high abundance of *C. lobatulus* and *Rosalina* sp. in areas with strong currents, low organics and coarse-grained sediments. We do not see a correlation between grain size and *Cibicides* or *Rosalina* in our samples due to the multiple forcings driving particle size distribution in Shetland (bottom current intensity vs. terrigenous supply); however, we do observe a clear correlation between a high relative abundance of *Cibicides* and *Rosalina* and low organics and high-energy environments (Figs. 4c and 5). Klitgaard-Kristensen and Sejrup (2002) and Murray (2003) also found a relationship between the dominance of *B. marginata* and low oxygen levels, high organics and fine sediments. Additionally, Murray (2003) also observed *S. fusiformis*-dominated assemblages in association with a high availability of LOM and low

oxygen levels. Overall, this compares really well with the assemblage we described in Group 2 (Figs. 4c and 5).

A recent study by Murray and Alve (2016) synthesises benthic foraminiferal biogeography in NW European fjords, mostly focusing on Norwegian fauna. Similarities with west Shetland voes include *C. lobatulus* as an “Atlantic species”, pointing to the influence of Atlantic Ocean waters on local estuarine and fjordic circulation. Assemblages dominated by *E. scaber* have been observed in Oslofjord and Drammensfjord in Norway (Alve, 1995; Alve and Nagy, 1986) and Gullmar Fjord in Sweden (Qvale et al., 1984), generally associated with organic-rich sediments and shallow waters. Additionally, Klitgaard-Kristensen and Buhl-Mortensen (1999) found a high relative abundance of the opportunistic species *S. fusiformis* in Norwegian fjords with a high percentage of sand and organic carbon, as well as assemblages characterised by *B. marginata* in environments rich in ROM.

In our sample set, we had no means to distinguish in situ tests from potentially advected tests when picking and counting unstained specimens, and the close match between our benthic foraminiferal assemblages and published data suggests that advection of material from other locations, especially in the more energetic environments, is negligible. Overall, benthic foraminiferal assemblages do respond to variations in the amount of OM and OC becoming deposited in fjord sediments (Fig. 4c), and comparisons can be drawn between fjordic systems. We recognise that this is a qualitative approach and we are not able to quantify changes in OM or OC content reaching the seafloor based on foraminiferal assemblages. However, this study provides an initial insight into the use of foraminifera to discriminate between post-depositional OC degradation and actual OC burial and accumulation in fjords, as foraminifera would only preserve the latter information in their assemblage composition. We reckon, however, that extreme conditions at the seafloor may result in the post-depositional dissolution of benthic foraminiferal tests, which would obviously compromise the assemblage composition. For this reason, we recommend carefully considering the level of preservation of each assemblage and, when necessary, making appropriate remarks on etching tests and other possible signs of test dissolution (e.g. Khanna et al., 2013). Additionally, assemblages dominated by agglutinated species like in Group 2 might have a lower diagnostic potential in their application down-core compared to their hyaline and porcelainous counterpart due to a tendency to be poorly preserved in the fossil record. Nevertheless, foraminifera are generally abundant in the fossil record; this approach shows down-core potential to support reconstructions of changes in OC burial and accumulation over time and could be paired with statistical and spatial model studies to tease out post-depositional OC degradation.

5 Conclusions

Our records from west Shetland voes provide new insights into sediment provenance and type, modern benthic foraminiferal spatial distribution across environmental gradients, and their use as indicators for OC enrichment in fjordic systems. Our results support a terrestrial source of both OM and OC in land-locked areas and calm depositional settings in west Shetland voes. Additionally, in locations characterised by low bottom water oxygenation, LOM is present in higher percentages than at well-ventilated sites. Conversely, marine processes govern IC storage, with higher concentrations in marginal to open marine settings. Benthic foraminiferal assemblages reliably reflect changes in OC content reaching the seafloor and show great potential as bio-indicators to trace these changes over time. We were able to identify four groups (75 % confidence) based on foraminiferal relative abundance data, which reflect four types of depositional setting: (1) land-locked with high organics and very weak currents; (2) stressed environments with high organics, a heavily stratified water column and low oxygen concentrations; (3) transitional environments characterised by moderate concentrations of organics and mild or intermittent currents; and (4) marginal to coastal settings with low organics and a high-energy environment.

Therefore, studies aiming to reconstruct past variations in the amount of OC reaching the seafloor and becoming stored therein could benefit from the use of benthic foraminifera assemblages as a qualitative approach to understand OC preservation in the sediment column and, for example, ground truth model predictions.

Data availability. Foraminiferal data are archived and available at <https://doi.org/10.1594/PANGAEA.907351> (Cappelli et al., 2019).

Supplement. The supplement related to this article is available online at: <https://doi.org/10.5194/bg-16-4183-2019-supplement>.

Author contributions. WENA and ELGC designed the study. ELGC, JC and CS analysed sediment physical properties, while ELGC and WENA worked on benthic foraminiferal assemblages. KD provided hydrographic data. ELGC prepared the paper with contributions from all co-authors.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. The crew of the MV *Moder Dy* are gratefully acknowledge for support of the work during the cruise in Shetland. We are thankful to undergraduate students Laura Smith, Leonie Strobl and Bethan Hudd for their help in the microscopy lab

assisting with benthic foraminifera picking. We thank two anonymous referees for valuable and constructive comments that improved the paper.

Financial support. This project was supported by BB-SRC/NERC (ref. BB/M026620/01). Isotope analysis was supported by the NERC Life Science Mass Spectrometry Facility (CEH_L_098_11_2015). KD was funded by the EU Interreg IV Northern Periphery project WATER (Warning of Algal Toxin Events to Support Aquaculture in the NPP Coastal Zone Region) and by the northwest Europe Interreg project PRIMROSE.

Review statement. This paper was edited by Hiroshi Kitazato and reviewed by three anonymous referees.

References

- Alve, E.: Opportunistic features of the foraminifer *Stainforthia fusiformis* (Williamson): evidence from Frierfjord, Norway, *J. Micropalaeontol.*, 13, 24–24, <https://doi.org/10.1144/jm.13.1.24>, 1994.
- Alve, E.: Benthic foraminiferal responses to estuarine pollution; a review, *J. Foramin. Res.*, 25, 190–203, <https://doi.org/10.2113/gsjfr.25.3.190>, 1995.
- Alve, E.: A common opportunistic foraminiferal species as an indicator of rapidly changing conditions in a range of environments, *Estuar. Coast. Shelf S.*, 57, 501–514, [https://doi.org/10.1016/S0272-7714\(02\)00383-9](https://doi.org/10.1016/S0272-7714(02)00383-9), 2003.
- Alve, E. and Murray, J. W.: High benthic fertility and taphonomy of foraminifera: A case study of the Skagerrak, North Sea, *Mar. Micropaleontol.*, 31, 157–175, [https://doi.org/10.1016/S0377-8398\(97\)00005-4](https://doi.org/10.1016/S0377-8398(97)00005-4), 1997.
- Alve, E. and Murray, J. W.: Marginal marine environments of the Skagerrak and Kattegat: A baseline study of living (stained) benthic foraminiferal ecology, *Palaeogeogr. Palaeoclimatol.*, 146, 171–193, [https://doi.org/10.1016/S0031-0182\(98\)00131-X](https://doi.org/10.1016/S0031-0182(98)00131-X), 1999.
- Alve, E. and Nagy, J.: Estuarine foraminiferal distribution in Sandebukta, a branch of Oslo Fjord, *J. Foramin. Res.*, 16, 261–284, <https://doi.org/10.2113/gsjfr.16.4.261>, 1986.
- Alve, E., Korsun, S., Schönfeld, J., Dijkstra, N., Golikova, E., Hess, S., Husum, K., and Panieri, G.: Foraminiferal AMBI: A sensitivity index based on benthic foraminiferal faunas from North-East Atlantic and Arctic fjords, continental shelves and slopes, *Mar. Micropaleontol.*, 122, 1–12, <https://doi.org/10.1016/j.marmicro.2015.11.001>, 2016.
- Austin, W. E. N. and Evans, J. R.: NE Atlantic benthic foraminifera: modern distribution patterns and palaeoecological significance, *J. Geol. Soc. London*, 157, 679–691, <https://doi.org/10.1144/jgs.157.3.679>, 2010.
- Austin, W. E. N.: Late Quaternary benthic foraminiferal stratigraphy of the western U.K. continental shelf, University College of North Wales, Bangor, 1991.
- Balasse, M., Tresset, A., Dobney, K., and Ambrose, S. H.: The use of isotope ratios to test for seaweed eating in sheep, *J. Zool.*, 266, 283–291, <https://doi.org/10.1017/S0952836905006916>, 2005.
- Batten, D. J.: Sedimentary organic matter. Organic facies and palynofacies, *Mar. Petrol. Geol.*, 13, 595–596, [https://doi.org/10.1016/0264-8172\(96\)88335-8](https://doi.org/10.1016/0264-8172(96)88335-8), 1996.
- Bauer, J. E. and Bianchi, T. S.: Dissolved Organic Carbon Cycling and Transformation, in: *Treatise on Estuarine and Coastal Science*, Vol. 5, 7–67, Elsevier Inc., <https://doi.org/10.1016/B978-0-12-374711-2.00502-7>, 2012.
- Bianchi, T. S., Schreiner, K. M., Smith, R. W., Burdige, D. J., Woodard, S., and Conley, D. J.: Redox Effects on Organic Matter Storage in Coastal Sediments During the Holocene: A Biomarker/Proxy Perspective, *Annu. Rev. Earth Planet. Sci.*, 44, 295–319, <https://doi.org/10.1146/annurev-earth-060614-105417>, 2016.
- Birnie, R. V.: Erosion rates on bare peat surfaces in Shetland, *Scot. Geogr. Mag.*, 109, 12–17, <https://doi.org/10.1080/00369229318736871>, 1993.
- Burrows, M., Hughes, D., Austin, W. E. N., Smeaton, C., Hicks, N., Howe, J., Allen, C., Taylor, P., and Vare, L.: Assessment of Blue Carbon Resources in Scotland's Inshore Marine Protected Area Network, Vol. 957, Scottish Natural Heritage, 2017.
- Cappelli, E. L. G., Clarke, J. L., Smeaton, C., Davidson, K., Austin, W. E. N.: Benthic foraminifera relative abundance in surface sediments of voes on Shetland's west coast, *PANGAEA*, <https://doi.org/10.1594/PANGAEA.907351>, 2019.
- Cefas: Scottish Sanitary Survey Project-Sanitary Survey Report Clift Sound and Stream Sound, available at: https://www.cefas.co.uk/media/52667/20090218_sanitarysrappendices_10_clift-sound.pdf (last access: 5 November 2019), 2007a.
- Cefas: Scottish Sanitary Survey Project-Sanitary Survey Report Sandsound Voe, available at: <https://www.cefas.co.uk/media/41384/sandsound-voe-sanitary-survey.pdf> (last access: 5 November 2019), 2007b.
- Cefas: Sanitary Survey Report, Busta Voe Lee North, SI 327, available at: https://www.cefas.co.uk/media/52671/20090221_sanitarysr_20_busta-voe-lee-north.pdf (last access: 5 November 2019), 2008a.
- Cefas: Scottish Sanitary Survey Project-Sanitary Survey Report Report East Firth and Firth, available at: https://www.cefas.co.uk/media/53114/20090220_sanitarysr_10_the-firth_redacted.pdf (last access: 5 November 2019), 2008b.
- Cefas: Scottish Sanitary Survey Project-Sanitary Survey Report Vaila Sound and Vaila Sound: Riskness, available at: https://www.cefas.co.uk/media/52690/vaila-sound_sanitary-survey-v10_redacted.pdf (last access: 5 November 2019), 2009.
- Cefas: Scottish Sanitary Survey Project-Sanitary Survey Report Report East Burra Firth and Aith Voe, available at: https://www.cefas.co.uk/media/53148/20100512_sanitarysr_10_east-burra-firth_redacted.pdf (last access: 5 November 2019), 2010.
- Cefas: Scottish Sanitary Survey Report-Sanitary Survey Report Olna Firth, available at: <https://www.cefas.co.uk/media/41375/olna-firth-sanitary-survey-report-v10-final.pdf> (last access: 5 November 2019), 2013.
- Chapman, S. J., Bell, J., Donnelly, D., and Lilly, A.: Carbon stocks in Scottish peatlands, *Soil Use Manage.*, 25, 105–112, <https://doi.org/10.1111/j.1475-2743.2009.00219.x>, 2009.

- Conradsen, K.: Recent benthic foraminifera in the southern Kattegat, Scandinavia: distributional pattern and controlling parameters, *Boreas*, 22, 367–382, <https://doi.org/10.1111/j.1502-3885.1993.tb00200.x>, 1993.
- Cui, X., Bianchi, T. S., Savage, C., and Smith, R. W.: Organic carbon burial in fjords: Terrestrial versus marine inputs, *Earth Planet. Sc. Lett.*, 451, 41–50, <https://doi.org/10.1016/j.epsl.2016.07.003>, 2016.
- Cui, X., Bianchi, T. S., and Savage, C.: Erosion of modern terrestrial organic matter as a major component of sediments in fjords, *Geophys. Res. Lett.*, 44, 1457–1465, <https://doi.org/10.1002/2016GL072260>, 2017.
- Darling, K. F., Schweizer, M., Knudsen, K. L., Evans, K. M., Bird, C., Roberts, A., Filipsson, H. L., Kim, J. H., Gudmundsson, G., Wade, C. M., Sayer, M. D. J., and Austin, W. E. N.: The genetic diversity, phylogeography and morphology of Elphidiidae (Foraminifera) in the Northeast Atlantic, *Mar. Micropaleontol.*, 129, 1–23, <https://doi.org/10.1016/j.marmicro.2016.09.001>, 2016.
- Duffield, C., Alve, E., Andersen, N., Andersen, T., Hess, S., and Strohmeier, T.: Spatial and temporal organic carbon burial along a fjord to coast transect: A case study from Western Norway, *Holocene*, 27, 1325–1339, <https://doi.org/10.1177/0959683617690588>, 2017.
- Edwards, A. and Sharples, F.: Scottish sea Lochs: a catalogue, Scottish Marine Biological Association Nature Conservancy Council, Oban, 1986.
- Evans, J. R., Austin, W. E. N., Brew, D. S., Wilkinson, I. P., and Kennedy, H. A.: Holocene shelf sea evolution offshore northeast England, *Mar. Geol.*, 191, 147–164, [https://doi.org/10.1016/S0025-3227\(02\)00529-7](https://doi.org/10.1016/S0025-3227(02)00529-7), 2002.
- Ficken, K. J., Barber, K. E., and Eglinton, G.: Lipid biomarker, $\delta^{13}\text{C}$ and plant macrofossil stratigraphy of a Scottish montane peat bog over the last two millennia, *Org. Geochem.*, 28, 217–237, [https://doi.org/10.1016/S0146-6380\(97\)00126-5](https://doi.org/10.1016/S0146-6380(97)00126-5), 1998.
- Fontanier, C., Jorissen, F. J., Licari, L., Alexandre, A., Anschutz, P., and Carbonel, P.: Live benthic foraminiferal faunas from the Bay of Biscay: Faunal density, composition, and microhabitats, *Deep-Sea Res. Pt. I*, 49, 751–785, [https://doi.org/10.1016/S0967-0637\(01\)00078-4](https://doi.org/10.1016/S0967-0637(01)00078-4), 2002.
- Fontanier, C., Jorissen, F. J., Lansard, B., Mouret, A., Buscail, R., Schmidt, S., Kerhervé, P., Buron, F., Zaragosi, S., Hunault, G., Ernoult, E., Artero, C., Anschutz, P., and Rabouille, C.: Live foraminifera from the open slope between Grand Rhône and Petit Rhône Canyons (Gulf of Lions, NW Mediterranean), *Deep-Sea Res. Pt. I*, 55, 1532–1553, <https://doi.org/10.1016/j.dsr.2008.07.003>, 2008.
- Gillibrand, P., Cromey, C., and Black, K.: Identifying the Risk of Deoxygenation in Scottish Sea Lochs with Isolated Deep Water, Scottish Association for Marine Science, Dunstaffnage Marine Laboratory, Oban, Argyll, 1–45, 2006.
- Hammer, Ø., Harper, D. A. T., and Ryan, P. D.: PAST: Paleontological Statistics Software Package for Education and Data Analysis, *Palaeontol. Electron.*, 4, 1–9, <https://doi.org/10.1016/j.bcp.2008.05.025>, 2001.
- Harris, D., Horwath, W. R., and van Kessel, C.: Acid fumigation of soils to remove carbonates prior to total organic carbon or CARBON-13 isotopic analysis, *Soil Sci. Soc. Am. J.*, 65, 1853, <https://doi.org/10.2136/sssaj2001.1853>, 2001.
- Hülse, D., Arndt, S., Wilson, J. D., Munhoven, G., and Ridgwell, A.: Understanding the causes and consequences of past marine carbon cycling variability through models, *Earth-Sci. Rev.*, 171, 349–382, <https://doi.org/10.1016/j.earscirev.2017.06.004>, 2017.
- Jennings, A. E.: Modern Foraminiferal Faunas of the South-western To Northern Iceland Shelf: Oceanographic and Environmental Controls, *J. Foramin. Res.*, 34, 180–207, <https://doi.org/10.2113/34.3.180>, 2004.
- Jorissen, F. J., de Stigter, H. C., and Widmark, J. G. V.: A conceptual model explaining benthic foraminiferal microhabitats, *Mar. Micropaleontol.*, 26, 3–15, [https://doi.org/10.1016/0377-8398\(95\)00047-X](https://doi.org/10.1016/0377-8398(95)00047-X), 1995.
- Jorissen, F. J., Wittling, I., Peypouquet, J. P., Rabouille, C., and Relexans, J. C.: Live benthic foraminiferal faunas off Cape Blanc, NW-Africa: Community structure and microhabitats, *Deep-Sea Res. Pt. I*, 45, 2157–2188, [https://doi.org/10.1016/S0967-0637\(98\)00056-9](https://doi.org/10.1016/S0967-0637(98)00056-9), 1998.
- Khanna, N., Godbold, J. A., Austin, W. E. N., and Pateron, D. M.: The impact of ocean acidification on the functional morphology of foraminifera, *PLoS One*, 8, 10–13, <https://doi.org/10.1371/journal.pone.0083118>, 2013.
- Klitgaard-Kristensen, D. and Buhl-Mortensen, L.: Benthic foraminifera along an offshore-fjord gradient: A comparison with amphipods and molluscs, *J. Nat. Hist.*, 33, 317–350, <https://doi.org/10.1080/002229399300281>, 1999.
- Klitgaard-Kristensen, D., Fetter Sejrup, H., and Hafidason, H.: Distribution of recent calcareous benthic foraminifera in the northern North Sea and relation to the environment, *Polar Res.*, 21, 275–282, <https://doi.org/10.3402/polar.v21i2.6488>, 2002.
- Korsun, S., Hald, M., Golikova, E., Yudina, A., Kuznetsov, I., Mikhailov, D., and Knyazeva, O.: Intertidal foraminiferal fauna and the distribution of Elphidiidae at Chupa Inlet, western White Sea, *Mar. Biol. Res.*, 10, 153–166, <https://doi.org/10.1080/17451000.2013.814786>, 2014.
- Kristensen, E.: Characterization of biogenic organic matter by step-wise thermogravimetry (STG), *Biogeochemistry*, 9, 135–159, <https://doi.org/10.1007/BF00692169>, 1990.
- Lo Giudice Cappelli, E. and Austin, W. E. N.: Size matters: analyses of benthic foraminiferal assemblages across differing size fractions, *Front. Mar. Sci.*, in review, 2019.
- Loh, P. S., Miller, A. E. J., Reeves, A. D., Harvey, S. M., and Overnell, J.: Assessing the biodegradability of terrestrially-derived organic matter in Scottish sea loch sediments, *Hydrol. Earth Syst. Sci.*, 12, 811–823, <https://doi.org/10.5194/hess-12-811-2008>, 2008.
- Marconi, M., Giordano, M., and Raven, J. A.: Impact of taxonomy, geography, and depth on $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ Variation in a large collection of macroalgae, *J. Phycol.*, 47, 1023–1035, <https://doi.org/10.1111/j.1529-8817.2011.01045.x>, 2011.
- Mendes, I., Dias, J. A., Schonfeld, J., and Ferreira, O.: Distribution of living benthic foraminifera on the Northern Gulf of Cadiz continental shelf, *J. Foramin. Res.*, 42, 18–38, <https://doi.org/10.2113/gsjfr.42.1.18>, 2012.
- Mook, D. H. and Hoskin, C. M.: Organic determinations by ignition: Caution advised, *Estuar. Coast. Shelf S.*, 15, 697–699, [https://doi.org/10.1016/0272-7714\(82\)90080-4](https://doi.org/10.1016/0272-7714(82)90080-4), 1982.
- Murray, J. W.: Distribution and population dynamics of benthic foraminifera from the southern North Sea, *J. Foramin. Res.*, 22, 114–128, <https://doi.org/10.2113/gsjfr.22.2.114>, 1992.

- Murray, J. W.: Foraminiferal assemblage formation in depositional sinks on the continental shelf west of Scotland, *J. Foramin. Res.*, 33, 101–121, 2003.
- Murray, J. W. and Alve, E.: Benthic foraminiferal biogeography in NW European fjords: A baseline for assessing future change, *Estuar. Coast. Shelf S.*, 181, 218–230, <https://doi.org/10.1016/j.ecss.2016.08.014>, 2016.
- Murray, J. W., Alve, E., and Cundy, A.: The origin of modern agglutinated foraminiferal assemblages: Evidence from a stratified fjord, *Estuar. Coast. Shelf S.*, 58, 677–697, [https://doi.org/10.1016/S0272-7714\(03\)00179-3](https://doi.org/10.1016/S0272-7714(03)00179-3), 2003.
- Nørgaard-Pedersen, N., Austin, W. E. N., Howe, J. A., and Shimmiel, T.: The Holocene record of Loch Etive, western Scotland: Influence of catchment and relative sea level changes, *Mar. Geol.*, 228, 55–71, <https://doi.org/10.1016/j.margeo.2006.01.001>, 2006.
- Qvale, G., Markussen, B., and Thiede, J.: Benthic foraminifers in fjords: response to water masses, *Nor. Geol. Tidsskr.*, 64, 235–249, 1984.
- Rillo, M. C., Kucera, M., Ezard, T. H. G., and Miller, C. G.: Surface Sediment Samples From Early Age of Seafloor Exploration Can Provide a Late 19th Century Baseline of the Marine Environment, *Front. Mar. Sci.*, 5, 1–6, <https://doi.org/10.3389/fmars.2018.00517>, 2019.
- Schiener, P., Black, K. D., Stanley, M. S., and Green, D. H.: The seasonal variation in the chemical composition of the kelp species *Laminaria digitata*, *Laminaria hyperborea*, *Saccharina latissima* and *Alaria esculenta*, *J. Appl. Phycol.*, 27, 363–373, <https://doi.org/10.1007/s10811-014-0327-1>, 2014.
- Schmidt, M. W. I. and Gleixner, G.: Carbon and nitrogen isotope composition of bulk soils, particle-size fractions and organic material after treatment with hydrofluoric acid, *Eur. J. Soil Sci.*, 56, 407–416, <https://doi.org/10.1111/j.1365-2389.2004.00673.x>, 2005.
- Schönfeld, J., Alve, E., Geslin, E., Jorissen, F., Korsun, S., Spezzaferri, S., Abramovich, S., Almogi-Labin, A., du Chatelet, E. A., Barras, C., Bergamin, L., Bicchi, E., Bouchet, V., Cearreta, A., Di Bella, L., Dijkstra, N., Disaro, S. T., Ferraro, L., Frontalini, F., Gennari, G., Golikova, E., Haynert, K., Hess, S., Husum, K., Martins, V., McGann, M., Oron, S., Romano, E., Sousa, S. M., and Tsujimoto, A.: The FOBIMO (FOraminiferal BIo-MONitoring) initiative: Towards a standardised protocol for soft-bottom benthic foraminiferal monitoring studies, *Mar. Micropaleontol.*, 94–95, 1–13, <https://doi.org/10.1016/j.marmicro.2012.06.001>, 2012.
- Scott, G. A., Scourse, J. D., and Austin, W. E.: The distribution of benthic foraminifera in the Celtic Sea: The significance of seasonal stratification, *J. Foramin. Res.*, 33, 32–61, <https://doi.org/10.2113/0330032>, 2003.
- Smeaton, C. and Austin, W. E. N.: Sources, Sinks and Subsidies: Terrestrial Carbon Storage in Mid-Latitude Fjords, *J. Geophys. Res.-Biogeo.*, 122, 2754–2768, <https://doi.org/10.1002/2017JG003952>, 2017.
- Smeaton, C., Austin, W. E. N., Davies, A. L., Baltzer, A., Abell, R. E., and Howe, J. A.: Substantial stores of sedimentary carbon held in mid-latitude fjords, *Biogeosciences*, 13, 5771–5787, <https://doi.org/10.5194/bg-13-5771-2016>, 2016.
- Smeaton, C., Austin, W. E. N., Davies, A. L., Baltzer, A., Howe, J. A., and Baxter, J. M.: Scotland's forgotten carbon: a national assessment of mid-latitude fjord sedimentary carbon stocks, *Biogeosciences*, 14, 5663–5674, <https://doi.org/10.5194/bg-14-5663-2017>, 2017.
- Smith, R. W., Bianchi, T. S., Allison, M., Savage, C., and Galy, V.: High rates of organic carbon burial in fjord sediments globally, *Nat. Geosci.*, 8, 450–453, <https://doi.org/10.1038/NGEO2421>, 2015.
- Soil Survey of Scotland: Soil maps of Scotland (partial coverage) at a scale of 1 V 25 000, Soil Survey of Scotland, Aberdeen, 1981.
- Stolpovsky, K., Dale, A. W., and Wallmann, K.: A new look at the multi-G model for organic carbon degradation in surface marine sediments for coupled benthic–pelagic simulations of the global ocean, *Biogeosciences*, 15, 3391–3407, <https://doi.org/10.5194/bg-15-3391-2018>, 2018.
- Thomas, E., Booth, L., Maslin, M., and Shackleton, N. J.: North-eastern Atlantic benthic foraminifera and implications of productivity during the last 40,000 years, *Paleoceanography*, 10, 545–562, <https://doi.org/10.1029/94PA03056>, 1995.
- Thornton, B., Martin, G., Procee, M., Miller, D. R., Coull, M., Yao, H., Chapman, S. J., Hudson, G., and Midwood, A. J.: Distributions of carbon and nitrogen isotopes in Scotland's topsoil: A national-scale study, *Eur. J. Soil Sci.*, 66, 1002–1011, <https://doi.org/10.1111/ejss.12289>, 2015.
- Thornton, S. F. and McManus, J.: Application of organic carbon and nitrogen stable isotope and C/N ratios as source indicators of organic matter provenance in estuarine systems: Evidence from the tay estuary, Scotland, *Estuar. Coast. Shelf S.*, 38, 219–233, <https://doi.org/10.1006/ecss.1994.1015>, 1994.
- Van Der Zwaan, G. J., Duijnste, I. A. P., Den Dulk, M., Ernst, S. R., Jannink, N. T., and Kouwenhoven, T. J.: Benthic foraminifers: Proxies or problems? A review of paleocological concepts, *Earth Sci. Rev.*, 46, 213–236, [https://doi.org/10.1016/S0012-8252\(99\)00011-2](https://doi.org/10.1016/S0012-8252(99)00011-2), 1999.
- Vaquier-Sunyer, R. and Duarte, C. M.: Thresholds of hypoxia for marine biodiversity, *P. Natl. Acad. Sci. USA*, 105, 15452–15457, <https://doi.org/10.1073/pnas.0803833105>, 2008.
- Verardo, D. J., Froelich, P. N., and McIntyre, A.: Determination of organic carbon and nitrogen in marine sediments using the Carlo Erba NA-1500 analyzer, *Deep-Sea Res. Pt. A*, 37, 157–165, [https://doi.org/10.1016/0198-0149\(90\)90034-S](https://doi.org/10.1016/0198-0149(90)90034-S), 1990.
- Weinkauff, M. F. G. and Milker, Y.: The Effect of Size Fraction in Analyses of Benthic Foraminiferal Assemblages: A Case Study Comparing Assemblages From the > 125 and > 150 μm Size Fractions, *Front. Earth Sci.*, 6, 1–10, <https://doi.org/10.3389/feart.2018.00037>, 2018.