



Supplement of

Understanding the uncertainty in global forest carbon turnover

Thomas A. M. Pugh et al.

Correspondence to: Thomas A. M. Pugh (t.a.m.pugh@bham.ac.uk)

The copyright of individual parts of the supplement might differ from the CC BY 4.0 License.

Forest-type	TBM PFTs (forest-type classification in parentheses)									
classifications	CABLE-POP	JULES	LPJ-GUESS	LPJmL	ORCHIDEE	SEIB-DGVM				
Tropical broadleaved	Deciduous broadleaf	Broadleaf deciduous tree	Tropical broadleaf	Tropical broad-leaved	Tropical broad-leaved	Tropical broad-leaved				
deciduous (TrBD)	forest (TrBD, TeBD,	(TrBD, TeBD, BBD ^a)	raingreen tree (TrBD)	raingreen tree (TrBD)	raingreen tree (TrBD)	raingreen tree (TrBD)				
	BBD ^a)									
Tropical broadleaved	Evergreen broadleaf	Tropical broadleaved	Tropical broadleaf	Tropical broad-leaved	Tropical broad-leaved	Tropical broad-leaved				
evergreen (TrBE)	forest (TrBE, TeBE ^a)	evergreen tree (TrBE)	evergreen tree (TrBE) ^b	evergreen tree (TrBE)	evergreen tree (TrBE)	evergreen tree (TrBE)				
Temperate	Deciduous broadleaf	Broadleaf deciduous tree	Temperate broadleaf	Temperate broad-leaved	Temperate broad-leaved	Temperate broad-leaved				
broadleaved	forest (TrBD, TeBD,	(TrBD, TeBD, BBD ^a)	summergreen tree (TeBD,	summergreen tree (TeBD)	summergreen tree	summergreen tree (TeBD)				
deciduous (TeBD)	BBD ^a)		BBD) ^b		(TeBD)					
Temperate	Evergreen broadleaf	Temperate broadleaved	Temperate broadleaf	Temperate broad-leaved	Temperate broad-leaved	Temperate broad-leaved				
broadleaved evergreen	forest (TrBE, TeBE ^a)	evergreen tree (TeBE)	evergreen tree (TeBE)	evergreen tree (TeBE)	evergreen tree (TeBE)	evergreen tree (TeBE)				
(TeBE)										
Boreal broadleaved	Deciduous broadleaf	Broadleaf deciduous tree	NA ^c	Boreal broad-leaved	Boreal broad-leaved	Boreal broad-leaved				
deciduous (BBD)	forest (TrBD, TeBD,	(TrBD, TeBD, BBD ^a)		summergreen tree (BBD)	summergreen tree (BBD)	summergreen tree (BBD)				
	BBD ^a)									
Needleleaved	Deciduous needleleaf	Needleleaf deciduous tree	Boreal needleleaf	NA ^c	Boreal needleleaf	Boreal needle-leaved				
deciduous (ND)	forest (ND)	(ND)	summergreen tree (ND)		summergreen tree (ND)	summergreen tree (ND)				
Needleleaved	Evergreen needleleaf	Needleleaf evergreen tree	Boreal needleleaf	Temperate needle-leaved	Temperate needleleaf	Temperate needle-leaved				
evergreen (NE)	forest (NE)	(NE)	evergreen tree (NE) ^b	evergreen tree (NE)	evergreen tree (NE)	evergreen tree (NE)				
				Boreal needle-leaved	Boreal needleleaf	Boreal needle-leaved				
				evergreen tree (NE)	evergreeen tree (NE)	evergreen tree (NE)				
Non-forest (NF),	Shrub (NF)	Deciduous shrub (NF)	NA ^c	NA ^c	NA ^c	NA ^c				
shrub		Evergreen shrub (NF)								
Non-forest (NF),	C3 grass (NF)	C3 grass (NF)	C3 grasses (NF)	Tropical herbaceous (NF)	C3 grass (NF)	Tropical herbaceous (NF)				
herbaceous	C4 grass (NF)	C4 grass (NF)	C4 grasses (NF)	Temperate herbaceous (NF)	C4 grass (NF)	Temperate herbaceous (NF)				

Table S1. Plant functional types (PFTs) simulated by each terrestrial biosphere model (TBM) and their grouping into forest-type classifications.

^a PFT assigned to more than one forest type: tropical forest when latitude is between 23° N and 23° S, temperate forest when latitude is >23° N to \leq 55° N or >23° S, and boreal forest when latitude is >55° N. ^b Both shade-tolerant and shade intolerant PFTs simulated.

^c PFT not simulated for this study.

Forest type ^c						Mo	del						
	CABLE-POP		JULES		LPJ-C	LPJ-GUESS		LPJmL		ORCHIDEE		SEIB-DGVM	
	Leaf	Root ^d	Leaf	Root ^d	Leaf	Root ^d	Leaf	Root ^d	Leaf	Root ^d	Leaf	Root ^d	
Boreal needleleaved	0.52	0.7	1.0ª	6.67	1.0 ^b	1.4	1.0	1.0	1.0 ^b	1.0 ^b	1.0 ^b	5.0	
deciduous													
Boreal needleleaved	3.5	0.7	4.0 ^a	6.67	3.0	1.4	2.0	2.0	3.0	3.0	4.55	2.38	
evergreen													
Boreal broadleaved	n/a	n/a	n/a	n/a	n/a	n/a	1.0	1.0	1.0 ^b	1.0 ^b	1.0 ^b	2.13	
deciduous													
Temperate needleleaved	3.5	0.7	n/a	n/a	n/a	n/a	2.0	2.0	3.0	3.0	4.55	1.56	
evergreen													
Temperate broadleaved	3.5	0.7	2.0 ^a	4.0	3.0	1.4	1.0	1.0	1.0	1.0	2.63	1.56	
evergreen													
Temperate broadleaved	2	0.7	1.0 ^a	4.0	1.0 ^b	1.4	1.0	1.0	1.0^{b}	1.0 ^b	1.0^{b}	1.56	
deciduous													
Tropical broadleaved	3.5	0.7	4.0 ^a	4.0	2.0	1.4	2.0	2.0	2.5	2.5	1.69	1.32	
evergreen													
Tropical broadleaved	n/a	n/a	1.0 ^a	4.0	1.0 ^b	1.4	1.0	1.0	1.0 ^b	1.0 ^b	1.0 ^b	0.63	
deciduous													
C3 herb	0.5	0.29	0.33	4.0	1.0	1.4	1.0	2.0	0.5	0.5	2.0	2.50	
C4 herb	0.5	0.21	0.33	4.0	1.0	1.4	1.0	2.0	0.5	0.5	2.0	1.3	

Table S2. Phenological longevity parameters (years) in the terrestrial biosphere model (TBM) ensemble.

^a JULES modifies phenology on a baseline turnover rate according to if the ambient temperature dips below a defined leaf-off temperature. The values in this table assume that the leaf-off temperature is never achieved for every recent trees, but is achieved every year for deciduous trees.

^b Reported here as one for deciduous phenologies on the assumption that only one leaf flush per year will be permitted. Actual leaf longevity is shorter.

^c Where forest-type classifications cover multiple PFTs, the most representative PFT is listed.

^d All root values are reported for fine roots only.

Table S3. Mappi	g of European	Space Agency	(ESA) landc	over classes to f	forest types used	in this analysis.
- mart at the pro-	8	~ r	(,

Code	Forest-type classification	ESA landcover	Additional conditions	
		classes ¹		
TrBE	Tropical broadleaved evergreen	50	latitude $\leq 23^{\circ}$	
TrBD	Tropical broadleaved deciduous	60, 61, 62	latitude \leq 23°	
OTr	Other tropical forest	100, 160, 170	latitude $\leq 23^{\circ}$	
TeBE	Temperate broadleaved evergreen	50	latitude > 23°	
TeBD	Temperate broadleaved deciduous	60, 61, 62	latitude > 23°	
NE	Needleleaved evergreen	70, 71, 72	n/a	
ND	Needleleaved deciduous	80, 81, 82	n/a	
MX	Broadleaved-needleleaved mixed forest	90	n/a	
Other	Other forest	100, 160, 170	latitude > 23°	

¹ESA (2017), Pugh et al. (2019b)

PFT type ^a	LA:SA (m ² cm ⁻²)							
	CABLE-POP	JULES	LPJ-GUESS	LPJmL	ORCHIDEE	SEIB-DGVM		
Boreal needleleaved deciduous	4000	b	5000	8000	no constraint	6000		
Boreal needleleaved evergreen	4000	b	5000	8000	no constraint	6000		
Boreal broadleaved deciduous	4000	n/a	n/a	n/a	n/a	8500		
Temperate needleleaved evergreen	4000	n/a	n/a	8000	no constraint	4800		
Temperate broadleaved evergreen	4000	b	6000	8000	no constraint	4800		
Temperate broadleaved deciduous	4000	b	6000	8000	no constraint	14500		
Tropical broadleaved evergreen	4000	b	6000	8000	no constraint	no constraint		
Tropical broadleaved deciduous	4000	b	6000	8000	no constraint	no constraint		

Table S4. Leaf area to sapwood area ratios used in the terrestrial biosphere model (TBM) ensemble

^a Where forest-type classifications cover multiple PFTs, the value for the most representative PFT is listed. ^b JULES does not define an area ratio, but instead relates leaf area to mass of sapwood for the purpose of calculating respiration (Eq. 46 in Clark et al., 2011).

Model	N^1	Fraction of observed drought-induced events with increased mortality							
		All mechanisms		Most drought-sensitive relevant process					
		Drought	Random ²	Drought	Random	Process-type			
		period + 5		period + 5		(Table 3)			
		years		years					
CABLE-	25	0.08	0.03	0.12	0.02	Growth efficiency			
POP									
JULES	25	0.16	0.04	0.16	0.12	Self-thinning			
LPJ-GUESS	26	0.04	0.02	0.12	0.04	Growth efficiency			
LPJmL	19	0.11	0.05	0.11	0.03	Self-thinning			
ORCHIDEE	22	0.05	0.04	n/a	n/a	n/a			
SEIB-	25	0.04	0.04	0.27	0.04	Bioclimatic limits			
DGVM									

Table S5. Performance of terrestrial biosphere models (TBMs) in capturing observed drought-induced mortality events.

¹ Total number of events for which the model simulated forest and a mortality flux was recorded in at least one year during the period 1901-2015.
² Mean of 10 randomly selected year ranges.



Figure S1. Evaluation of grid-cells where TBMs simulate forest for the period 1985-2014 (see Methods for definition) against satellite observations for the year 2000 (Hansen et al., 2013). Blue shading indicates where the TBM simulates forest and satellite observations find at least 10% of the grid-cell to be covered by forest with a 50% canopy-cover threshold (Pugh et al., 2019a). Red shading indicates where forest is observed in the satellite data but not simulated by the TBM. Grey shading indicates where forest is simulated by the TBM but is not observed in the satellite data.



Figure S2. Forest type regions based on the European Space Agency (ESA) land-cover map (ESA 2017). Reproduced from Pugh et al. (2019b), Fig. S9. See Table S3 for forest type codes.



Figure S3. NPP mean for the period 1985-2014 as forced by the CRU-NCEP climate (units of kg C m⁻² y⁻¹). Colour scale is capped at 2.5 kg C m⁻² y⁻¹. Maps show areas which are simulated as forest for each model and have at least 10% of the grid-cell covered by closed-canopy forest (see Methods).



Figure S4. Leaf C cost as a function of longevity for three of the TBMs. Leaf C cost is defined as the reciprocal of specific leaf area divided by leaf longevity.



Figure S5. Annual mean maximum value of the monthly LAI (m² m⁻²) over the period 1985-2014 in the CRU-NCEP-forced simulation.



Figure S6. Fraction of F_{turn} which results from mortality (F_{mort} / F_{turn}), for the period 1985-2014 as forced by the CRU-NCEP climate. The fraction of F_{turn} resulting from phenological processes is the remainder. Masking as for Fig. 1.



Figure S7. Distribution of the model simulated forest types during 1985-2014 based on the CRU-NCEP simulation.



Figure S8. Standard deviation of turnover fluxes in space for F_{mort} and F_{phen} , calculated by model forest types over the period 1985-2014 from the CRU-NCEP-forced simulation. For "within forest type" variance was calculated for each forest type, before taking the square root of the mean variance across all forest types. For "across forest type" the mean of fluxes across each forest type was taken, before calculating the standard deviation of these forest-type means. Calculations were only made across grid-cells with at least 10% forest cover. Comparisons of absolute numbers from JULES with those of other TBMs should be avoided because of the different spatial resolution.



Figure S9. Percentage change in τ_{mort} (C_{veg}/F_{mort}) mean between the periods 1985-2014 and 2070-2099 as forced by the IPSL-CM5A-LR climate (units of years). Masking as for Fig. 1.



Figure S10. Distribution of the model simulated forest types during 1985-2014 (a-f) and 2070-2099 (g-l) in the TBM simulations forced by IPSL-CM5A-LR RCP 8.5 bias-corrected climate data.



Figure S11. Mortality rate (F_{mort}/C_{veg}) for CABLE-POP split by conceptual process grouping (Table 3) and observational forest type for 1985-2099 in the simulation forced by IPSL-CM5A-LR RCP 8.5 bias-corrected climate data. 31-year running means are plotted for clarity and thus only 2000-2085 is shown. Rates are calculated based on grid-cell total C_{veg} and mortality fluxes.



Figure S12. Mortality rate (F_{mort}/C_{veg}) for JULES split by conceptual process grouping (Table 3) and observational forest type for 1985-2099 in the simulation forced by IPSL-CM5A-LR RCP 8.5 bias-corrected climate data. 31-year running means are plotted for clarity and thus only 2000-2085 is shown. Rates are calculated based on grid-cell total C_{veg} and mortality fluxes.



Figure S13. Mortality rate (F_{mort}/C_{veg}) for LPJ-GUESS split by conceptual process grouping (Table 3) and observational forest type for 1985-2099 in the simulation forced by IPSL-CM5A-LR RCP 8.5 bias-corrected climate data. 31-year running means are plotted for clarity and thus only 2000-2085 is shown. Rates are calculated based on grid-cell total C_{veg} and mortality fluxes.



Figure S14. Mortality rate (F_{mort}/C_{veg}) for LPJmL split by conceptual process grouping (Table 3) and observational forest type for 1985-2099 in the simulation forced by IPSL-CM5A-LR RCP 8.5 bias-corrected climate data. 31-year running means are plotted for clarity and thus only 2000-2085 is shown. Rates are calculated based on grid-cell total C_{veg} and mortality fluxes.



Figure S15. Mortality rate (F_{mort}/C_{veg}) for ORCHIDEE split by observational forest type for 1985-2099 in the simulation forced by IPSL-CM5A-LR RCP 8.5 bias-corrected climate data. No mechanism breakdown was available. 31-year running means are plotted for clarity and thus only 2000-2085 is shown. Rates are calculated based on grid-cell total C_{veg} and mortality fluxes.



Figure S16. Mortality rate (F_{mort}/C_{veg}) for SEIB-DGVM split by conceptual process grouping (Table 3) and observational forest type for 1985-2099 in the simulation forced by IPSL-CM5A-LR RCP 8.5 bias-corrected climate data. 31-year running means are plotted for clarity and thus only 2000-2085 is shown. Rates are calculated based on grid-cell total C_{veg} and mortality fluxes.



Figure S17. Change in NPP sums by observational forest type for 1985-2099 in the simulation forced by IPSL-CM5A-LR RCP 8.5 bias-corrected climate data. 31-year running means are plotted for clarity and thus only 2000-2085 is shown.

References

- Clark, D. B., Mercado, L. M., Sitch, S., Jones, C. D., Gedney, N., Best, M. J., Pryor, M., Rooney, G. G., Essery, R. L. H., Blyth, E., Boucher, O., Harding, R. J., Huntingford, C. and Cox, P. M.: The Joint UK Land Environment Simulator (JULES), model description – Part 2: Carbon fluxes and vegetation dynamics, Geosci. Model Dev., 4(3), 701–722, doi:10.5194/gmd-4-701-2011, 2011.
- ESA: ESA CCI Land Cover dataset (v 2.0.7), [online] Available from: http://maps.elie.ucl.ac.be/CCI/viewer/ (Accessed 29 June 2017), 2017.
- Hansen, M. C., Potapov, P. V, Moore, R., Hancher, M., Turubanova, S. A., Tyukavina, A., Thau, D., Stehman, S. V, Goetz, S. J., Loveland, T. R., Kommareddy, A., Egorov, A., Chini, L., Justice, C. O., and Townshend, J. R. G.: High-resolution global maps of 21st-century forest cover change., Science, 342, 850–853, https://doi.org/10.1126/science.1244693, 2013.
- Pugh, T. A. M., Arneth, A., Kautz, M., Poulter, B., and Smith, B.: Important role of forest disturbances in the global biomass turnover and carbon sinks, Nat. Geosci., 12, 730–735, https://doi.org/10.1038/s41561-019-0427-2, 2019a.
- Pugh, T. A. M., Lindeskog, M., Smith, B., Poulter, B., Arneth, A., Haverd, V., and Calle, L.: The role of forest regrowth in global carbon sink dynamics, Proc. Natl. Acad. Sci. U. S. A., 116(10), 4382–4387, https://doi.org/10.1073/pnas.1810512116, 2019b.