



Corrigendum to

“Using ^{226}Ra and ^{228}Ra isotopes to distinguish water mass distribution in the Canadian Arctic Archipelago” published in Biogeosciences, 17, 4937–4959, 2020

Chantal Mears^{1,2}, Helmuth Thomas^{2,1}, Paul B. Henderson³, Matthew A. Charette³, Hugh MacIntyre¹, Frank Dehairs⁴, Christophe Monnin⁵, and Alfonso Mucci⁶

¹Department of Oceanography, Dalhousie University, Halifax, NS, Canada

²Institute for Coastal Research, Helmholtz Centre Geesthacht, Geesthacht, Germany

³Department of Marine Chemistry and Geochemistry, Woods Hole Oceanographic Institution, Woods Hole, MA, USA

⁴Analytical, Environmental and Geo-Chemistry, Vrije Universiteit Brussel, Brussels, Belgium

⁵CNRS – Université Paul Sabatier-IRD-OMP, Geosciences Environnement Toulouse (GET), Toulouse, France

⁶Geotop and Department of Earth and Planetary Sciences, McGill University, Montréal, QC, Canada

Correspondence: Helmuth Thomas (helmuth.thomas@hzg.de)

Published: 23 November 2020

During figure production, an error was introduced in Figs. 2d–h and 3 of the abovementioned paper: the color-coding of stations CAA4 and CAA6 in the original publication was incorrect. The corrected figures can be found below.




Figure 2. Vertical distributions of (a) practical salinity, (b) temperature, (c) density, (d) AT, (e) Ba, (f) $\delta^{18}\text{O}$, (g) DIC, and (h) normalized DIC (DIC_{norm}) observed at stations CAA1 to CAA7 throughout the CAA.




Figure 3. (a) ^{226}Ra , (b) ^{228}Ra , and (c) $^{228}\text{Ra}/^{226}\text{Ra}$ profiles ranging from the surface to depth for stations CAA1 to CAA7 (where pink dots indicate station CAA2, which was only sampled at the surface and bottom) throughout the Canadian Arctic Archipelago.