




*Supplement of*

**Assessing the impacts of agricultural managements on soil carbon stocks, nitrogen loss, and crop production – a modelling study in eastern Africa**


**Jianyong Ma et al.**

*Correspondence to:* Jianyong Ma ([jianyong.ma@kit.edu](mailto:jianyong.ma@kit.edu))

The copyright of individual parts of the supplement might differ from the article licence.



**Figure S1.** Comparison of daily climate between GSWP3-W5E5 data set ( $34.25^{\circ}E$ ,  $0.25^{\circ}N$ ) and field-based weather records at Kenyan INM3 site ( $34.40^{\circ}E$ ,  $0.14^{\circ}N$ ) from 2004-2014 as time series (a), and scatter plots shown in (b). AB and RB are mean absolute bias and relative bias, respectively, presented in percent (%).



**Figure S2.** The modelled SOC stocks (0-150cm) by LPJ-GUESS at the CT1 and INM3 sites between 1901 and 2002. The shaded areas represent the period of cropland systems. All the above-ground biomass under grassland systems was returned to the soils in the simulations (see Sect. 2.3.1).

Following Olin et al. (2015), relationships between allocation to sorghum leaves ( $P_{leaf}$ ), stem ( $P_{stem}$ ), root ( $P_{root}$ ), and grains ( $P_{yield}$ ) based on the allocation model of Penning de Vries et al. (1989) are established using a logistic growth function, a Richards curve (Richards, 1959) (Eq. S1):

$$f_i = a_i + \frac{b_i - a_i}{1 + e^{-c_i \times (DS - d_i)}} \quad (S1)$$

where  $DS$  is crop development stage, ranging from 0 to 2 ( $DS=0$ , sowing;  $DS=1$ , flowering;  $DS=2$ , harvest); and  $a_i$ ,  $b_i$ ,  $c_i$ ,  $d_i$  are fitting coefficients for the three functions (specific values given in Table S1 below).

The relative relationships of daily assimilate allocation to the organs are described by Eqs. S2:


$$f_1 = \frac{P_{root}}{P_{veg}}, f_2 = \frac{P_{leaf}}{P_{veg} - P_{root}}, f_3 = \frac{P_{yield}}{P_{veg} + P_{yield}}, P_{veg} + P_{yield} = 1 \quad (S2)$$

where  $P_{veg}$  are the fraction of carbon allocated to vegetative organs (i.e.,  $P_{veg} = P_{leaf} + P_{stem} + P_{root}$ ), ranging from 0 to 1.

We can achieve dynamic carbon allocation to the plant organs over the growing season by combining Eqs. S2:

$$\begin{cases} P_{root} = f_1 \times (1 - f_3) \\ P_{leaf} = f_2 \times (1 - f_1) \times (1 - f_3) \\ P_{stem} = P_{veg} - P_{root} - P_{leaf} = (1 - f_1) \times (1 - f_2) \times (1 - f_3) \\ P_{yield} = f_3 \end{cases} \quad (S3)$$

Partitioning functions are plotted for sorghum in Fig. S3. More details on how these relationships were derived can be found in Olin et al. (2015).




**Figure S3.** The organ's assimilate partitioning to roots, leaves, stem, and yields for sorghum. Solid lines represent the fitted Richards functions used in this study, and dashed lines are the allocation scheme from Penning de Vries et al. (1989).


**Table S1.** CFT-specific parameters for the daily carbon allocation (Eqs. S1-S3), specific leaf area (SLA), minimum C:N value of the leaves (leaf C:N<sub>min</sub>), and the fraction of the total mineral N applied at the three crop development stages (DS, in which DS=0, sowing; DS=0.5, halfway through the vegetative stage; DS=1.0, flowering).

Parameter	Maize <sup>I</sup>	Sorghum <sup>II</sup>	Spring wheat <sup>I</sup>	Rice <sup>I</sup>	Pulses <sup>III</sup>	Soybean <sup>III</sup>	Unit
f <sub>1</sub>	a <sub>1</sub>	0.24	0.49	0.62	0.14	0.59	0.56
	b <sub>1</sub>	1.22	0	-0.02	-2.45	0	0
	c <sub>1</sub>	18.1	6.74	5.80	2.79	5.53	3.74
	d <sub>1</sub>	1.12	0.54	0.55	2.19	0.51	0.53
f <sub>2</sub>	a <sub>2</sub>	0.68	0.56	0.86	0.52	0.75	0.67
	b <sub>2</sub>	-0.06	0.09	0.19	-0.04	0	0
	c <sub>2</sub>	12.48	25.75	28.65	10.16	7.69	30.78
	d <sub>2</sub>	0.81	0.90	0.55	0.79	1.38	1.73
f <sub>3</sub>	a <sub>3</sub>	0	0	0	0	0	0
	b <sub>3</sub>	1	1	1	1	1	1
	c <sub>3</sub>	28.52	10.54	8.27	121.21	9.59	8.93
	d <sub>3</sub>	1.03	1.12	1.10	1.09	1.46	1.41
SLA	45	45	35	45	45	40	m <sup>2</sup> kg <sup>-1</sup> C <sup>-1</sup>
Leaf C:N <sub>min</sub>	15	15	15	10	10	8	kg C kg N <sup>-1</sup>
N <sub>app</sub> , DS=0	11%	11%	11%	3%	50%	50%	fraction
N <sub>app</sub> , DS=0.5	50%	50%	50%	46%	0%	0%	fraction
N <sub>app</sub> , DS=1.0	39%	39%	39%	51%	50%	50%	fraction

<sup>I</sup> Olin et al. (2015); <sup>II</sup> This study; <sup>III</sup> Ma et al. (2022)




**Figure S4.** Inputs used for simulations in Eastern Africa: (a) simulated crop-specific areas (rain-fed and irrigated) over the historic period compared to statistics from FAO (ha); (b) mean rates of mineral N fertilizer and manure applied to each crop type ( $\text{kg N ha}^{-1}$ ) from 1901-2014; (c) annual  $\text{CO}_2$  concentration (ppm, right scale), and mean temperature ( $^{\circ}\text{C}$ , left scale) from CRUJRA (B1, Table 1) and five GCMs (C1 and C2, Table 1). The faba bean and common bean are simulated as pulses in (a) and (b). The black, blue, green, and red thick solid lines in (c) denote the temperature averaged by five GCMs for historical, SSP1-26, 3-70 and 5-85 scenarios, respectively; thin lines in light color represent the temperature from individual GCM. Dashed lines show  $\text{CO}_2$  concentrations.




**Figure S5.** Modelled and observed maize yields (long rainy season, i.e. the main growing period) for the evaluation treatments at INM3 (a) and CT1 (b) sites, with two levels of mineral N fertilizer input (N30 and N90). The dashed line is the modelled yields, and the closed circle represents the observed value averaged by the four replicates in the trials, with standard deviation as given in the vertical bar.

**Table S2.** Evaluation of the model performance to simulate maize yields in long rainy season (n=12) using multi-years average, absolute bias (AB), root mean square error (RMSE), and Person correlation coefficient (*r*) at INM3 and CT1 sites.


<b>Treatment</b>	<b>Observed mean (t ha<sup>-1</sup> yr<sup>-1</sup>)</b>	<b>Simulated mean (t ha<sup>-1</sup> yr<sup>-1</sup>)</b>	<b>AB (%)</b>	<b>RMSE (t ha<sup>-1</sup> yr<sup>-1</sup>)</b>	<b><i>r</i></b>
<b>INM3</b>					
N30_NoMan_RR	4.00	3.95	26	1.08	0.02 ( <i>p</i> >0.05)
N90_NoMan_RR	4.25	4.95	32	1.31	0.07 ( <i>p</i> >0.05)
N30_Man_NoRR	4.56	4.98	21	1.01	0.27 ( <i>p</i> >0.05)
N90_Man_NoRR	5.19	5.16	28	1.41	-0.06 ( <i>p</i> >0.05)
N30_Man_RR	4.58	4.96	33	1.37	0.14 ( <i>p</i> >0.05)
N90_Man_RR	4.95	5.17	27	1.32	-0.03 ( <i>p</i> >0.05)
N30_NoMan_NoRR	3.56	3.91	26	1.04	0.05 ( <i>p</i> >0.05)
N90_NoMan_NoRR	3.65	4.92	46	1.59	0.23 ( <i>p</i> >0.05)
<b>CT1</b>					
N30_NoTill_RR	2.83	3.27	34	0.88	0.30 ( <i>p</i> >0.05)
N90_NoTill_RR	4.10	4.59	39	1.28	0.13 ( <i>p</i> >0.05)
N30_Till_NoRR	3.31	3.26	17	0.61	0.59 ( <i>p</i> <0.05)
N90_Till_NoRR	4.41	4.53	24	1.04	0.17 ( <i>p</i> >0.05)
N30_Till_RR	3.28	3.31	17	0.62	0.53 ( <i>p</i> <0.05)
N90_Till_RR	4.37	4.58	32	1.26	0.18 ( <i>p</i> >0.05)
N30_NoTill_NoRR	2.81	3.22	43	1.07	-0.12 ( <i>p</i> >0.05)
N90_NoTill_NoRR	3.78	4.54	58	1.66	-0.15 ( <i>p</i> >0.05)



**Figure S6.** The modelled relative response (%) of cropland SOC, N loss, and yield to alternative managements (see Table 3 for abbreviations) compared to the conventional management ( $F_{std}$ ) in maize (a), sorghum (b), wheat (c), and pulses (d) cropping systems over Eastern Africa. Box plots denote the 5th and 95th percentiles with whiskers, median and interquartile range with box lines, and mean with asterisks across all the simulated crop grid cells (maize-403; sorghum-398; wheat-382; pulses-425).




**Figure S7.** The optimal soil C sequestration practice ( $F_{opt}$ , Eq.4) simulated by LPJ-GUESS in maize (a), sorghum (b), wheat (c), and pulses (d) cropping systems in Kenya and Ethiopia (B2, Table 1). Numbers in plots represent the grid cell proportion of each optimal management to the total crop grid cells (maize-403; sorghum-398; wheat-382; pulses-425). The conventional management ( $F_{std}$ , black in plots) was chosen when none of other alternative managements yielded a net increase in SOC. See table 3 for management abbreviations.



**Figure S8.** The simulated cropland soil C stocks from the conventional management ( $F_{std}$ ) for GCM-based historical period (C1, Table 1) (a), and SOC potential response to alternative management options with future SSP1-26, 3-70 and 5-85 scenarios (C2, Table 1) in Ethiopia (b) and Kenya (c). Blue triangle for Ethiopia and sun cross for Kenya denote the seamless connection between historical period and future simulations. Filled circle represents the averaged result between 2091 and 2100, with a range from five GCMs as given in vertical bar. See Table 3 for management abbreviations.

**Table S3.** The relative (%) number of cropland grid cell and areas regarding the potential transition of optimal soil C sequestration practice ( $F_{opt}$ , Eq. 4), comparing the historical period (GCM-based climate; C3, Table 1) with three future SSP scenarios from 2091-2100 (C2, Table 1). See Table 3 for management abbreviations.

From (GCMs historical)	To (future)	Amount of cropland grid cells (%)			Amount of cropland areas (%)		
		SSP1-26	SSP3-70	SSP5-85	SSP1-26	SSP3-70	SSP5-85
$F_{std}$	$F_{RR}$	0.70	1.17	1.17	0.00	0.01	0.01
	$F_{Man}$	2.80	3.04	2.57	0.03	0.03	0.02
	$F_{CC-BNF}$	1.87	1.17	1.17	0.03	0.03	0.03
$F_{RR}$	$F_{std}$	0.23	0.47	0.23	0.00	0.49	0.00
	$F_{Man}$	0.47	0.00	0.23	0.04	0.00	0.01
	$F_{CC-BNF}$	12.85	14.02	16.12	17.74	17.73	19.40
$F_{Man}$	$F_{std}$	1.64	1.87	2.34	0.01	0.01	0.02
	$F_{RR}$	3.27	4.21	4.21	1.12	0.38	0.50
	$F_{CC-BNF}$	14.02	15.65	15.89	9.88	11.10	11.12
$F_{CC-BNF}$	$F_{std}$	0.00	0.00	0.00	0.00	0.00	0.00
	$F_{Man}$	0.47	0.00	0.00	0.01	0.00	0.00
	$F_{RR}$	1.17	2.10	1.87	0.84	1.52	1.46
Total change		39.5	43.7	45.8	29.7	31.3	32.6



**Figure S9.** The N fixation rate ( $\text{kg N ha}^{-1} \text{ yr}^{-1}$ ) of herbaceous legumes modelled by LPJ-GUESS in the FCC-BNF simulation under the present-day climate, averaged over the last ten years of simulation (B2, Table 1).

## Reference

- Ma, J., Olin, S., Anthoni, P., Rabin, S. S., Bayer, A. D., Nyawira, S. S. and Arneth, A.: Modeling symbiotic biological nitrogen fixation in grain legumes globally with LPJ-GUESS ( v4.0 , r10285 ), *Geosci. Model Dev.*, 15(2), 815–839, 2022.
- Olin, S., Schurgers, G., Lindeskog, M., Wärlind, D., Smith, B., Bodin, P., Holmér, J. and Arneth, A.: Modelling the response of yields and tissue C : N to changes in atmospheric CO<sub>2</sub> and N management in the main wheat regions of western Europe, *Biogeosciences*, 12(8), 2489–2515, doi:10.5194/bg-12-2489-2015, 2015.
- Penning de Vries, F. W. T., Jansen, D. M., Berge, H. F. M. ten and Bakema, A.: Simulation of ecophysiological processes of growth in several annual crops, Centre for Agricultural Publishing and Documentation, Wageningen, Wageningen., 1989.
- Richards, F. J.: A flexible growth function for empirical use, *J. Exp. Bot.*, 10(2), 290–301, doi:10.1093/jxb/10.2.290, 1959.