



Origin of secondary fatty alcohols in atmospheric aerosols in a cool–temperate forest based on their mass size distributions

Yuhao Cui^{1,2}, Eri Tachibana², Kimitaka Kawamura³, and Yuzo Miyazaki²

¹Division of Earth System Science, Graduate School of Environmental Science, Hokkaido University, Sapporo, 060-0810, Japan

²Institute of Low Temperature Science, Hokkaido University, Sapporo, 060-0819, Japan

³Chubu Institute for Advanced Studies, Chubu University, Kasugai, 487-8501, Japan

Correspondence: Yuzo Miyazaki (yuzom@lowtem.hokudai.ac.jp)

Received: 24 April 2023 – Discussion started: 23 May 2023

Revised: 18 October 2023 – Accepted: 31 October 2023 – Published: 16 December 2023

Abstract. Fatty alcohols (FAs) are major components of surface lipids in plant leaves and serve as surface-active organic aerosols (OAs), which can act as primary biological aerosol particles (PBAPs). To elucidate the origin and formation of secondary fatty alcohols (SFAs) in atmospheric aerosols, their mass size distribution in aerosol samples obtained from a deciduous forest canopy was measured in spring, summer and autumn. The SFAs showed the highest concentration in spring (growing season), with *n*-nonacosan-10-ol being the most abundant. In spring and summer, the size peak of *n*-nonacosan-10-ol was in the particle size range $> 10.0\ \mu\text{m}$, whereas it was in the $1.9\text{--}3.0\ \mu\text{m}$ range in autumn. The size distribution of *n*-nonacosan-10-ol did not show any significant correlation with that of the known biogenic tracers of pollen, soil and fungal spores in spring and summer. The overall results, together with SFAs measured in plant leaves, as well as the literature, suggest that SFAs originate mostly from plant waxes and that leaf senescence status is likely an important factor controlling the size distribution of SFAs. This study provides new insights into the possible sources of PBAPs and their effects on the ice nucleation activity of aerosols based on seasonal changes in particle size.

Kawamura et al., 2003). Once emitted into the atmosphere as primary biological aerosol particles (PBAPs), FAs can act as surface-active organic aerosols (OAs). Among various effects, surface-active OAs can change the surface tension at air–water interfaces (Donaldson et al., 2006; You et al., 2020) and affect the atmospheric lifetime of particles (Gilman et al., 2004). In addition, FAs can nucleate ice (Perkins et al., 2020), which is important for cloud formation. Qiu et al. (2017) suggested that monolayers of *n*-alkyl alcohols, with carbon numbers up to 30, can act as efficient ice nucleants – a common ability of long-chain FAs (Vazquez de Vasquez et al., 2020). The ice nucleation properties of PBAPs (e.g. bacteria, fungal spores and pollen) have been widely studied (e.g. Lukas et al., 2020; O’Sullivan et al., 2015; Hader et al., 2013). However, data on the surface lipids or plant waxes from terrestrial higher plants emitted as PBAPs are limited.

Monolayers of long-chain (carbon number ≥ 16) linear primary alcohols (PAs) have historically been used to initiate ice nucleation (Huang et al., 2018). PAs can be oxidised to carboxylic acids, with aldehydes as intermediate components, in the aqueous phase (Yu et al., 2005). When emitted into the atmosphere, FA can be a component of the surfactant film coating on aqueous aerosols, which affects droplet growth interactions (Li et al., 2019). Primary FAs (PFAs) are less stable in the atmosphere than secondary FAs (SFAs) because H abstraction from secondary alcohols occurs less frequently than that from primary alcohols (Koivisto et al., 2015). SFAs are less affected by atmospheric photochemical reactions than PFAs and may, therefore, serve as tracers of specific sources. Previous studies have shown that SFAs

1 Introduction

Terrestrial lipids are key tracers for investigating the origins of atmospheric aerosols (Chen et al., 2021). Fatty alcohols (FAs), with chains varying between C_{20} and C_{34} , are major components of surface lipids (Jetter and Kunst, 2008;

predominate in more than 50 % of the epicuticular waxes in some plants, such as *Tropaeolum majus*, *Pinus pinaster*, *Pinus halepensis* and most conifers (e.g. Koch et al., 2006; Matas et al., 2003; Nikolić et al., 2020).

Among SFAs, *n*-nonacosan-10-ol, forming tubular aggregates (Jetter et al., 1994), is a common (in some plants, the most abundant) component of epicuticular leaf waxes (Spangenberg et al., 2010) and one of the most abundant substances in plant tubules (Wang et al., 2015). For atmospheric aerosols, Oros and Simoneit (2001) detected *n*-nonacosan-10-ol in smoke samples from conifers subjected to controlled burning. Another SFA compound, *n*-nonacosan-5,10-ol, was identified in ambient aerosol samples collected in the western North Pacific during the Asian Pacific Regional Aerosol Characterization Experiment (ACE-Asia) campaign (Simoneit et al., 2004). Gagosian et al. (1987) emphasised that aerosol lipids can be transported from the land surface to the oceans in a relatively shorter timescale of a few days, indicating the possible impact of SFAs on the ice nucleation (IN) activity in aerosol particles, not only on local scales but also on regional scales. Miyazaki et al. (2019) identified five SFAs in aerosols in a forest atmosphere for the first time and showed distinct seasonal variations in their mass concentrations. The above-mentioned studies suggest that SFAs in plant leaves can potentially be emitted into the atmosphere in natural terrestrial environments and can subsequently act as PBAPs.

Despite the importance of SFAs in atmospheric aerosols in terms of their possible effects on atmospheric chemistry and climate, studies are limited, particularly on the origins of SFAs in atmospheric aerosols, owing to a lack of aerosol samples obtained in the vicinity of possible source regions. This study was aimed at elucidating the origin and formation of SFAs based on size-segregated aerosol samples collected at a deciduous forest site during different seasons.

2 Experimental

2.1 Size-segregated aerosol sampling

Aerosol samplings were conducted at the Sapporo Forest meteorology research site (42°59' N, 141°23' E; 182 m a.s.l.) in the western part of Hokkaido, the northernmost major island of Japan (Miyazaki et al., 2012). The sampling site (Fig. 1) is situated in a cool temperate zone with an annual mean temperature of 6.5 °C and an annual precipitation of 1100 mm (Nakai et al., 2003). This mixed cool-temperate forest consists of a mature and secondary deciduous forest and a human-made coniferous forest with varied forest floor cover. The experimental site was covered with a broad-leaf forest consisting of white birch (*Betula platyphylla*), Sasa bamboo (*Sasa kurilensis*) and other species (Nakai et al., 2003). The canopy has a mean height of 20 m, and the forest floor is 0.5–2 m high. The evolution pattern of leaf area



Figure 1. Location of the aerosol sampling site (© GeoTechnologies).

index (LAI) at the sampling site was similar to that at other temperate deciduous forests. The LAI determined by Nakai et al. (2003), showing seasonal variations after the initial foliating period, remained almost constant at approximately 4.0 from the end of June to mid-September. The predominant local wind direction and wind speed from May to October at this forest site indicated that the majority of aerosols sampled were likely influenced by emissions from forested areas (Miyazaki et al., 2012). Indeed, the source apportionment study suggested that the majority of aerosols originated from the local forest in spring, summer and autumn, whereas the observed aerosols in winter were significantly affected by marine and anthropogenic sources (Miyazaki et al., 2012). Because we focus on the effects of forest vegetation on SFAs in aerosols, the data obtained in spring, summer and autumn are shown and discussed in this study.

To obtain the mass size distributions of the aerosol SFAs and related compounds, size-segregated aerosol sampling was performed using an Andersen-type cascade impactor (AN-2100, Tokyo Dylec. Corp.). The nine-stage impactor collected supermicrometre particles (with diameters (D_p) of 1.0–1.9, 1.9–3.0, 3.0–4.3, 4.3–6.4, 6.4–10.0 and $> 10.0 \mu\text{m}$) in six stages and submicrometre particles (with $D_p < 0.39$, 0.39–0.58 and 0.58–1.0 μm) in three stages. The size-segregated aerosol samples were collected on prebaked quartz-fibre filters (i.d. 8 cm) at a flow rate of 120 L min^{-1} without temperature and humidity control. The impactor was placed 2 m above the forest floor. The duration of each sam-




Figure 2. (a) Molecular structures of secondary fatty alcohols (SFAs) identified in this study; *n*-heptacosan-10-ol, *n*-heptacosan-5,10-diol, *n*-nonacosan-10-ol, *n*-nonacosan-10,13-diol and *n*-nonacosan-5,10-diol are defined as SFA1, SFA2, SFA3, SFA4 and SFA5, respectively. (b) Mass size distributions of SFAs during the period between 21 and 28 May 2010.

pling was approximately 1 week. Size-segregated aerosol data obtained from 26 April to 28 May in 2010, 22 June to 13 August in 2010 and 6 to 28 October in 2009, are shown; these periods are defined as spring, summer and autumn, respectively. The collected filter samples were stored in glass jars at -20°C until analysis. In total, nine sets of samples were used to measure SFAs and related compounds.

2.2 Chemical analysis

To measure SFAs and related compounds, a quarter of each sample filter with an area of 10.18 cm^2 was extracted using a mixture of dichloromethane and methanol (2 : 1, *v/v*). The hydroxyl functional group ($-\text{OH}$) in the extracted sample was derivatised with a mixture of $50\ \mu\text{L}$ *N*, *O*-bis-(trimethylsilyl) trifluoroacetamide (BSTFA) and $10\ \mu\text{L}$ pyridine to form trimethylsilyl (TMS) ether ($-\text{OTMS}$) (Fu et al., 2009). The TMS derivatives were injected into a capillary gas chromatograph (GC8890, Agilent), equipped with a fused silica capillary column (DB-5MS) and coupled to a mass spectrometer (MSD5977B, Agilent), for analysis. Total ion chromatograms (TICs) of the five derivatised SFAs were identified. The identification of each peak was based on a detailed interpretation of the electron ionization (EI) mass spectral data (mass spectrometric fragmentation patterns) together with their comparison to data in the literature as well as exact mass measurements using a high-resolution gas chromatograph–time-of-flight mass spectrometer (GC–TOF-MS; JMS-T100GCV, JEOL) at the Creative Research Institute of Hokkaido University, as described by Miyazaki et al. (2019). Figure 2a shows molecular structures of five SFAs, namely *n*-heptacosan-10-ol (SFA1), *n*-heptacosan-5,10-diol (SFA2), *n*-nonacosan-10-ol (SFA3), *n*-nonacosan-10,13-diol (SFA4) and *n*-nonacosan-5,10-diol (SFA5), identified in this study. TMS derivatives of these compounds were reported previously (Miyazaki et al., 2019). Besides SFAs,

sugar compounds (sucrose, trehalose, arabinol and mannitol) were measured using the method described above.

In this study, water-insoluble organic carbon (WIOC) is defined as a parameter calculated from the difference between the measured mass of organic carbon (OC) and water-soluble organic carbon (WSOC). A total organic carbon (TOC) analyser (Model TOC-L_{CHP}, Shimadzu) was used to determine the WSOC mass concentration. A filter cut of 19.63 cm^2 was extracted and then filtered with a disc filter (Millex-GV, $0.22\ \mu\text{m}$, Millipore, Billerica, MA, USA), followed by injection of dissolved OC in the extracts into the analyser. The measured carbon is defined as the WSOC. The data presented here were corrected using blanks. The mass concentration of OC was measured using a Sunset Laboratory OC–EC analyser. A filter punch of 1.54 cm^2 was used for OC analysis. After measuring the OC and WSOC, the mass concentrations of WIOC were defined as the difference between those of OC and WSOC ($\text{WIOC} = \text{OC} - \text{WSOC}$).

3 Results and discussion

3.1 Mass size distributions of secondary fatty alcohols in the aerosol samples

Figure 2b shows the typical mass size distributions of the five identified SFAs obtained from samples collected in spring. SFA3 was the dominant SFA identified in all samples, followed by SFA5. Table 1 summarises the mass concentrations and fractions of SFAs for each size range of the sample. The predominance of SFA3 among SFAs identified in aerosol samples was also reported previously at other forest sites (Miyazaki et al., 2019). Overall, the concentrations of SFAs were the highest in spring, which is the growing season (Table 1). The average concentrations of SFA3 ($10.2 \pm 10.3\ \text{ng m}^{-3}$) and SFA5 ($0.52 \pm 0.69\ \text{ng m}^{-3}$) in the bulk aerosol (the sum of submicrometre and supermicrometre aerosol mass) during spring are smaller than those (~ 100 and $\sim 8\ \text{ng m}^{-3}$, respectively) in total suspended particulate (TSP) matter obtained at the Tomakomai (TMK) experimental forest ($42^{\circ}43'\text{ N}$, $141^{\circ}36'\text{ E}$) (Miyazaki et al., 2019). If most of these SFAs originated from forest vegetation, the difference in the concentration levels is attributable to the larger area (2715 ha) and leaf area index (LAI; $\sim 4\text{--}6$) with many more plant species in the TMK experimental forest (Hiura, 2001) compared to those (150 ha and < 4 , respectively) of the current research forest site (Nakai et al., 2003).

In all seasons, the majority of the SFA mass resided in the supermicrometre size range. Specifically, the mass of *n*-nonacosan-10-ol in the supermicrometre size accounted for $93.4 \pm 7.1\%$, $84.2 \pm 17.8\%$ and $88.2 \pm 2.3\%$ of the total SFA mass in spring, summer and autumn, respectively. In the following sections, the mass size distributions of *n*-nonacosan-10-ol are discussed as a representative SFAs.

Table 1. Concentrations of secondary fatty alcohols (SFAs) in different size ranges of the aerosol sample. ND indicates that the compound was not detected (i.e. below the lower detection limit).

Season	Sampling period	Concentration (ng m ⁻³)	
		Submicrometre ($D_p < 1.0 \mu\text{m}$)	Supermicrometre ($D_p > 1.0 \mu\text{m}$)
(a) <i>n</i>-nonacosan-10-ol (SFA3)			
Spring	26 April–7 May 2010	1.38	7.96
	7–14 May 2010	0.01	0.48
	21–28 May 2010	0.57	20.52
Summer	22–29 June 2009	0.82	14.91
	8–16 July 2010	0.11	1.73
	6–13 August 2010	0.90	1.57
Autumn	6–13 October 2009	0.68	5.24
	13–21 October 2009	1.35	8.17
	21–28 October 2009	0.45	4.27
(b) <i>n</i>-nonacosan-5,10-diol (SFA5)			
Spring	26 April–7 May 2010	0.04	0.29
	7–14 May 2010	ND	0.01
	21–28 May 2010	0.03	1.30
Summer	22–29 June 2009	0.03	0.78
	8–16 July 2010	ND	0.07
	6–13 August 2010	0.05	0.10
Autumn	6–13 October 2009	0.02	0.27
	13–21 October 2009	0.03	0.32
	21–28 October 2009	0.01	0.17
(c) <i>n</i>-heptacosan-10-diol (SFA1)			
Spring	26 April–7 May 2010	0.03	0.18
	7–14 May 2010	ND	0.06
	21–28 May 2010	0.02	0.49
Summer	22–29 June 2009	ND	0.33
	8–16 July 2010	ND	0.05
	6–13 August 2010	0.04	0.04
Autumn	6–13 October 2009	ND	0.24
	13–21 October 2009	ND	0.45
	21–28 October 2009	ND	0.25

3.2 Seasonal changes in the mass size distributions of the secondary fatty alcohols

Figure 3 shows the average mass size distributions of SFA3 and SFA5 for each season. As described in the previous section, SFAs exhibited the highest concentrations in spring (the growing season). It is apparent that in spring, the peak diameter was greater than $10 \mu\text{m}$. Interestingly, the peak diameter shifted to smaller sizes from spring to autumn, with the relative amount of mass in the $1.9\text{--}3.0 \mu\text{m}$ diameter range becoming larger as the season progressed. The large standard deviation shown in Fig. 3 can be partly explained by the effect of precipitation that occurred in spring and summer (Miyazaki

et al., 2012), which might cause a decrease in the concentrations of SFAs in aerosols.

Because SFAs are compound groups of water-insoluble organic carbon (WIOC), their contribution to WIOC in terms of mass should be examined. The average mass concentration ratio of SFA3 : WIOC in spring was 0.79 %, whereas this period had the most abundant SFA mass concentration among all the samples. Indeed, the R^2 value between *n*-nonacosan-10-ol and bulk WIOC concentrations was as high as 0.78 for that season (Fig. 4b). This result suggests that, although the mass fraction of SFAs identified was small, a possible source of SFA is an important factor in controlling the WIOC

Table 1. Continued.

Season	Sampling period	Concentration (ng m ⁻³)	
		Submicrometre ($D_p < 1.0 \mu\text{m}$)	Supermicrometre ($D_p > 1.0 \mu\text{m}$)
(d) <i>n</i>-nonacosan-10,13-ol (SFA4)			
Spring	26 April–7 May 2010	ND	0.10
	7–14 May 2010	ND	ND
	21–28 May 2010	ND	0.44
Summer	22–29 June 2009	ND	0.18
	8–16 July 2010	ND	0.01
	6–13 August 2010	0.02	ND
Autumn	6–13 October 2009	ND	0.06
	13–21 October 2009	ND	0.09
	21–28 October 2009	ND	0.04
(e) <i>n</i>-heptacosan-5,10-diol (SFA2)			
Spring	26 April–7 May 2010	ND	0.02
	7–14 May 2010	ND	ND
	21–28 May 2010	ND	0.12
Summer	22–29 June 2009	ND	0.04
	8–16 July 2010	ND	ND
	6–13 August 2010	ND	ND
Autumn	6–13 October 2009	ND	ND
	13–21 October 2009	ND	0.00
	21–28 October 2009	ND	0.00




Figure 3. The average mass size distributions of (a, b, c) *n*-nonacosan-10-ol (SFA3) and (d, e, f) *n*-nonacosan-5,10-diol (SFA5) in spring (a, d), summer (b, e) and autumn (c, f). Spring: 26 April to 7 May 2010, 7 to 14 May 2010 and 21 to 28 May 2010; summer: 22 to 29 June 2009, 8 to 16 July 2010 and 6 to 13 August 2010; autumn: 6 to 13 October 2009, 13 to 21 October 2009 and 21 to 28 October 2009.




Figure 4. (a) Mass size distribution of bulk water-insoluble organic carbon (WIOC) compared with that of *n*-nonacosan-10-ol during the period between 21 and 28 May 2010. (b) A scatterplot of the mass concentrations between *n*-nonacosan-10-ol and bulk WIOC in spring (21–28 May 2010).

mass in spring. The increase in the concentrations of WIOC with diameters greater than 10 μm was likely due to the influence of larger plant debris in spring (Matsumoto et al., 2022), which has been regarded as a possible source of SFAs. However, this was not evident in other seasons, when the mass concentrations were lower than those in spring.

3.3 Possible origin and formation of secondary fatty alcohols in the aerosol

In this section, the origin and processes of formation of SFAs in the observed aerosols are described from the viewpoint of mass size distributions and seasonal differences in abundance. As indicated by the dominant mass fraction in the supermicrometre size range, SFAs in the observed aerosols were most likely PBAPs that are emitted into the atmosphere by the local surface wind (Tegen et al., 2018). To support the discussion on the origin of PBAPs in relation to other PBAPs, correlations between the size-segregated mass concentrations of SFAs and several biogenic tracers were examined for each season.

3.3.1 Spring and summer

Figure 5 shows the mass size distributions of sucrose and trehalose compared with those of *n*-nonacosan-10-ol in each season. Sucrose is the most abundant component of airborne pollen grains (Zhu et al., 2015) and was used as a pollen tracer compound. As expected, sucrose showed the largest mass concentration in spring, with the peak diameter (D_p) in the range of 4.3 to 6.4 μm (Fig. 5a), whereas peak diameters (i.e. the shape of size distributions) were different between *n*-nonacosan-10-ol and sucrose in spring as well as in summer. Correlations between *n*-nonacosan-10-ol and sucrose concentrations were not apparent during these two seasons (Fig. 6a). In contrast to *n*-nonacosan-10-ol and sucrose, trehalose had the lowest mass concentration in spring (Fig. 5d–f). Trehalose is the most abundant sugar in soil (Jia

and Fraser, 2011), and its contribution to soil dust has been previously reported (e.g. Jia et al., 2010; Medeiros et al., 2006). These studies suggest that trehalose can be used as a tracer for surface soil in atmospheric aerosols. The concentrations of *n*-nonacosan-10-ol did not show any significant correlation with trehalose concentrations in spring and summer (Fig. 6b).

Figure 7 shows a comparison of the mass size distributions of *n*-nonacosan-10-ol with those of arabinol and mannitol during each season. Arabinol and mannitol have been used as tracers for airborne fungal spores (Fu et al., 2012). The size distributions for these two tracers were similar to each other but dissimilar to that of SFAs in each season; the mass concentrations showed insignificant correlations with those of *n*-nonacosan-10-ol in spring and summer ($R^2 < 0.25$) (Fig. 8). These insignificant correlations suggest that fungal spores are not the dominant source of *n*-nonacosan-10-ol during the spring and summer. In summary, the size distribution of *n*-nonacosan-10-ol did not show any significant correlation with those of known biogenic tracers in spring, when SFAs had maximum concentrations. Insignificant correlations between *n*-nonacosan-10-ol and these tracers were also observed in the summer. Overall, the results indicate that the majority of SFAs did not originate from pollen, soil or fungal spores during spring and summer.

As the most probable source of SFAs in the observed aerosols, *n*-nonacosan-10-ol is recognised as a major compound in epicuticular waxes found in gymnosperm species (e.g. Tulloch, 1976; Schulten et al., 1986; Sharma et al., 2018), such as *Sequoiadendron giganteum* (Yamamoto et al., 2008). Vioque and Kolattukudy (1997) measured secondary fatty diols in other plant species, such as *Pisum sativum*. The main production of SFAs begins with the reduction in fatty acids through the formation of aldehydes as intermediates to produce FAs, which can form waxes in plants (Mudge, 2005). The wax-homologous series of long-chain lipids, including nonacosan-10-ol, are predominant (Gülz et




Figure 5. The average mass size distributions of (a, b, c) sucrose and (d, e, f) trehalose in spring (26 April to 28 May 2010) (a, d), summer (22 June to 13 August 2010) (b, e) and autumn (6 to 28 October 2009) (c, f). Each size distribution was compared with that of *n*-nonacosan-10-ol during each corresponding period (grey).




Figure 6. Scatterplots of the mass concentrations between (a) *n*-nonacosan-10-ol and sucrose and (b) *n*-nonacosan-10-ol and trehalose in spring and summer.

al., 1994). In the aerosol samples, the carbon numbers of the identified SFAs were 27 and 29 (odd numbers). Indeed, SFAs with odd carbon numbers can be produced by decarbonylation of FAs in plant leaves (Gülz et al., 1994). In order to confirm SFAs in a plant leaf which is representative of the sampling region, leaf samples of a Norway spruce (*Picea abies*) were collected at the nearby sampling site on the campus of Hokkaido University. Figure 9 presents an example GC–MS total ion chromatogram (TIC) of the TMS extract of the leaf sample. The most abundant SFA, *n*-nonacosan-10-ol, found in the aerosol samples was indeed detected in the leaf, at an average mass concentration of $46.4 \pm 11.3 \text{ mg g}^{-1}$. This re-

sult, together with the literature, supports the idea that the majority of the SFAs in the aerosols measured in this study originated from plant wax.

The strength of atmospheric emissions of SFAs may partly depend on the degree of epicuticular wax accumulation, which depends on the age of the leaf linked with phenology. The generation and regeneration of wax in plants occurs mostly in spring (Mohammadian et al., 2007). This explains the current results of the maximum concentrations of SFAs in the growing season, which is attributable to a possible increase in the generation of plant wax. Waxes can be aerosolised from living plants and can be transported by wind (Nelson et al., 2018); thus, the most plausible process of formation of SFA aerosol particles is wind-driven emission from the leaf surface.

3.3.2 Autumn

The average mass concentrations of *n*-nonacosan-10-ol in autumn and summer were similar (Table 1). In contrast, trehalose and mannitol had the highest concentrations in autumn (Figs. 5 and 7). The masses of these tracer compounds were mostly in the supermicrometre range. Similarly, the mass of arabitol was mostly in the supermicrometre size range, with a concentration greater than that in summer and the R^2 between *n*-nonacosan-10-ol and arabitol concentrations being 0.24. The R^2 values of concentrations of *n*-nonacosan-10-ol with those of trehalose and mannitol were




Figure 7. The average mass size distributions of (a, b, c) arabitol and (d, e, f) mannitol in spring (26 April to 28 May 2010) (a, d), summer (22 June to 13 August 2010) (b, e) and autumn (6 to 28 October 2009) (c, f). Each size distribution was compared with that of *n*-nonacosan-10-ol during each corresponding period (grey).




Figure 8. Scatterplots of the mass concentrations between (a) *n*-nonacosan-10-ol and arabitol and (b) *n*-nonacosan-10-ol and mannitol in spring and summer.

0.64 and 0.50 ($p < 0.01$ for both), respectively. However, these relationships are probably a coincidence because of the different seasonal changes in the mass concentrations of these tracer compounds and *n*-nonacosan-10-ol, as well as the differences in the possible sources discussed in the previous section.

The concentration of *n*-nonacosan-10-ol positively correlated with that of sucrose in autumn (Fig. S1 in the Supplement). The flowering periods of the vegetation at the study site differed, and the possible sources of pollen were diverse. The two major types of vegetation at the study site are *Betula platyphylla* and *Quercus crispula* Blume (Nakai




Figure 9. GC-MS total ion chromatogram obtained for the trimethylsilyl (TMS) derivatives from leaf samples.

et al., 2003), the flowering periods of which are from April to May and June to August, respectively. On the forest floor, *Sasa kurilensis* flowers were present at the study site in autumn. Particularly during the flowering period, when large atmospheric emissions of pollen are expected, pollen is often ejected in clumps that stick to nearby vegetation and is blown away after drying (Jones and Harrison, 2004). The positive correlation between *n*-nonacosan-10-ol and sucrose in autumn can be partly explained by the adhesion of pollen to plants (Visez et al., 2020), followed by atmospheric emissions together with *n*-nonacosan-10-ol from plant wax.

The predominance of the smaller peaks ($D_p = 1.9\text{--}3.0\ \mu\text{m}$) in the size distributions of *n*-nonacosan-10-ol in au-




Figure 10. Seasonal changes in the average mass size distribution of *n*-nonacosan-10-ol.

tumn compared with that in the other seasons (Fig. 10) can be partly explained by leaf senescence. The crystal transformation and degradation of epicuticular wax depend on the age of the leaf, which is linked to plant phenology (Mohammadian et al., 2007). Modification of epicuticular structures occurs during the ageing of leaf surfaces, when epicuticular wax starts to degrade in autumn (Reicosky and Hanover, 1978). As a result, senescing leaves have less epicuticular wax than mature leaves (Cao et al., 2013). The relatively large contribution of the smaller peak to the mass of *n*-nonacosan-10-ol is likely due to wax degradation. It is noted that relative humidity can influence particle size distribution. However, SFAs are water-insoluble fractions of organic aerosol (Fig. 4) which might be less hygroscopic, and their size distributions may be less affected by relative humidity. Indeed, the size distributions of SFAs observed in this study exhibited peaks in the larger size ranges in spring, despite the highest relative humidity being observed in summer at the site, suggesting insignificant effects of relative humidity on the size distributions in this study.

In summary, the major source of SFAs in the atmospheric aerosol samples collected in this study was probably plant wax; this notion is supported by the data for SFAs measured in the leaf samples and surrounding vegetation at the sampling site and those in the literature. Further studies are required to measure SFAs in other types of plant leaves simultaneously with aerosol samples in a forest environment. The current findings on SFAs in aerosols provide new insights into the origins and possible effects of PBAPs. Specifically, this study suggests that the ageing processes of plants can affect the mass size distributions of SFAs and might thus alter the IN activity (Zhang et al., 2021).

4 Conclusions

In this study, we investigated the origin and formation of SFAs in atmospheric aerosols based on their mass size distribution in aerosol samples obtained from a deciduous forest site. Five SFAs were identified, and their mass size distributions in the forest environment were measured for the first time. The mass concentrations of SFAs in the aerosols were the highest in the growing season (spring), which supports the results of a previous study at other forest sites; *n*-nonacosan-10-ol (SFA3) was the most abundant compound among the SFAs. On average, the mass of *n*-nonacosan-10-ol was mostly (> 85 %) in the supermicrometre size range, whereas the peak of the size range shifted to smaller sizes as the season progressed.

In spring, the mass size distributions were highly correlated with those of bulk WIOC, indicating that *n*-nonacosan-10-ol is an important factor controlling the WIOC mass. In spring and summer, the size distribution of *n*-nonacosan-10-ol was not significantly correlated with that of known biogenic tracers, such as pollen (sucrose), soil (trehalose) and fungal spores (arabitol and mannitol). Furthermore, *n*-nonacosan-10-ol was detected in Norway spruce leaf samples. The results of the aerosol and leaf samples, together with the literature on SFAs in plant leaves, suggest that SFAs mainly originated from plant wax. Moreover, the current results indicate that *n*-nonacosan-10-ol identified in this study can act as a possible new tracer for PBAPs.

In autumn, the mass peak of SFAs in particle sizes greater than 10 μm, as determined in spring, was not apparent, whereas the mass in the smaller size range of 1.9–3.0 μm was relatively greater compared with the larger size range. The relatively large contribution of the smaller peaks to the size distribution of *n*-nonacosan-10-ol in autumn can be partly explained by leaf senescence, followed by the degradation of plant wax and subsequent atmospheric emissions. This study provides new insights into possible sources of PBAPs, namely SFAs. Furthermore, our results suggest that the different growth stages of plants can result in differences in the size distributions of PBAPs emitted into the atmosphere, which may affect the IN properties of aerosol particles.

Data availability. The measurement data for the aerosol samples are provided in the Supplement. All other data are available upon request.

Supplement. The supplement related to this article is available online at: <https://doi.org/10.5194/bg-20-4969-2023-supplement>.

Author contributions. YC and YM designed the study and wrote the manuscript. YC performed the GC–MS experiments. YM performed the aerosol sampling. KK managed the sampler. YC, YM and ET analysed the data.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

Special issue statement. This article is part of the special issue "Land surface–atmosphere interactions – from the microbial to the global scale". It is a result of the iLEAPS – OzFlux joint 2023 Conference, Auckland, New Zealand, 31 January–3 February 2023.

Acknowledgements. We thank Katsumi Yamanoi for managing the sampling site and for his help with the aerosol sampling.

Financial support. This research was supported by Grants-in-Aid for Scientific Research (B) (16H02931) from the Ministry of Education, Culture, Sports, Science and Technology (MEXT), Japan, and by JST SPRING (grant number JPMJSP2119) via the DX Doctoral Fellowship of Hokkaido University.

Review statement. This paper was edited by Semeena Valiyaveetil Shamsudheen and reviewed by two anonymous referees.

References

- Cao, J., Song, Y., Wu, H., Qin, L., Hu, L., and Hao, R.: Ultrastructural Studies on the Natural Leaf Senescence of *Cinnamomum Camphora*, *Scanning*, 35, 336–343, <https://doi.org/10.1002/sca.21065>, 2013.
- Chen, J., Kawamura, K., Hu, W., Liu, C. Q., Zhang, Q., and Fu, P.: Terrestrial lipid biomarkers in marine aerosols over the western North Pacific during 1990–1993 and 2006–2009, *Sci. Total Environ.*, 797, 149115–149115, <https://doi.org/10.1016/j.scitotenv.2021.149115>, 2021.
- Donaldson, D. J. and Vaida, V.: The Influence of Organic Films at the Air–Aqueous Boundary on Atmospheric Processes, *Chem. Rev.*, 106, 1445–1461, <https://doi.org/10.1021/cr040367c>, 2006.
- Fu, P., Kawamura, K., Chen, J., and Barrie, L.: Isoprene, monoterpene, and sesquiterpene oxidation products in the high Arctic aerosols during late winter to early summer, *Environ. Sci. Technol.*, 43, 4022–4028, 2009.
- Fu, P., Kawamura, K., Kobayashi, M., and Simoneit, B. R. T.: Seasonal variations of sugars in atmospheric particulate matter from Gosan, Jeju Island: Significant contributions of airborne pollen and Asian dust in spring, *Atmos. Environ.*, 55, 234–239, <https://doi.org/10.1016/j.atmosenv.2012.02.061>, 2012.
- Gagosian, R. B., Peltzer, E. T., and Merrill, J. T.: Long-range transport of terrestrially derived lipids in aerosols from the South Pacific, *Nature*, 325, 800–803, 1987.
- Gilman, J. B., Eliason, T. L., Fast, A., and Vaida, V.: Selectivity and stability of organic films at the air–aqueous interface, *J. Colloid Interface Sci.*, 280, 234–243, <https://doi.org/10.1016/j.jcis.2004.07.019>, 2004.
- Gülz, P. G.: Epicuticular leaf waxes in the evolution of the plant kingdom, *J. Plant Physiol.*, 143, 453–464, [https://doi.org/10.1016/S0176-1617\(11\)81807-9](https://doi.org/10.1016/S0176-1617(11)81807-9), 1994.
- Hader, J. D., Wright, T. P., and Petters, M. D.: Contribution of pollen to atmospheric ice nuclei concentrations, *Atmos. Chem. Phys.*, 14, 5433–5449, <https://doi.org/10.5194/acp-14-5433-2014>, 2014.
- Hiura, T.: Stochasticity of species assemblage of canopy trees and understory plants in a temperate secondary forest created by major disturbances, *Ecol. Res.*, 16, 887–893, 2001.
- Huang, H., Yarmush, M. L., and Usta, O. B.: Long-term deep-supercooling of large-volume water and red cell suspensions via surface sealing with immiscible liquids, *Nat. Commun.*, 9, 3201, <https://doi.org/10.1038/s41467-018-05636-0>, 2018.
- Jetter, R. and Kunst, L.: Plant surface lipid biosynthetic pathways and their utility for metabolic engineering of waxes and hydrocarbon biofuels, *Plant J.*, 54, 670–683, <https://doi.org/10.1111/j.1365-313X.2008.03467.x>, 2008.
- Jetter, R. and Riederer, M.: Epicuticular crystals of nonacosan-10-ol: in-vitro reconstitution and factors influencing crystal habits, *Planta*, 195, 257–270, 1994.
- Jia, Y. and Fraser, M. P.: Characterization of saccharides in size-fractionated ambient particulate matter and aerosol sources: the contribution of primary biological aerosol particles (PBAPs) and soil to ambient particulate matter, *Environ. Sci. Technol.*, 45, 930–936, <https://doi.org/10.1021/es103104e>, 2011.
- Jia, Y. L., Clements, A. L., and Fraser, M. P.: Saccharide composition in atmospheric particulate matter in the southwest US and estimates of source contributions, *J. Aerosol Sci.*, 41, 62–73, <https://doi.org/10.1016/j.jaerosci.2009.08.005>, 2010.
- Jones, A. M. and Harrison, R. M.: The effects of meteorological factors on atmospheric bioaerosol concentrations: a review, *Sci. Total Environ.*, 326, 151–180, <https://doi.org/10.1016/j.scitotenv.2003.11.021>, 2004.
- Kawamura, K., Ishimura, Y., and Yamazaki, K.: Four years' observations of terrestrial lipid class compounds in marine aerosols from the western North Pacific, *Global Biogeochem. Cy.*, 17, 1003, <https://doi.org/10.1029/2001GB001810>, 2003.
- Koch, K., Dommissé, A., and Barthlott, W.: Chemistry and crystal growth of plant wax tubules of Lotus (*Nelumbo nucifera*) and Nasturtium (*Tropaeolum majus*) leaves on technical substrates, *Crystal Growth Des.*, 6, 2571–2578, <https://doi.org/10.1021/cg060035w>, 2006.
- Koivisto, E., Ladommatos, N., and Gold, M.: Systematic study of the effect of the hydroxyl functional group in alcohol molecules on compression ignition and exhaust gas emissions, *Fuel*, 153, 650–663, <https://doi.org/10.1016/j.fuel.2015.03.042>, 2015.

- Li, S., Cheng, S., Du, L., and Wang, W.: Establishing a model organic film of low volatile compound mixture on aqueous aerosol surface, *Atmos. Environ.*, 200, 15–23, <https://doi.org/10.1016/j.atmosenv.2018.11.052>, 2019.
- Lukas, M., Schwidetzky, R., Kunert, A.T., Pöschl, U., Fröhlich-Nowoisky, J., Bonn M., and Meister, K.: Electrostatic Interactions Control the Functionality of Bacterial Ice Nucleators, *J. Am. Chem. Soc.*, 142, 15, 6842–6846, <https://doi.org/10.1021/jacs.9b13069>, 2020.
- Matas, A. J., Sanz, M. J., and Heredia, A.: Studies on the structure of the plant wax nonacosan-10-ol, the main component of epicuticular wax conifers, *Int. J. Biol. Macromol.*, 33, 31–35, [https://doi.org/10.1016/S0141-8130\(03\)00061-8](https://doi.org/10.1016/S0141-8130(03)00061-8), 2003.
- Matsumoto, K., Kodama, S., Sakata, K., and Watanabe, Y.: Atmospheric deposition fluxes and processes of the water-soluble and water-insoluble organic carbon in central Japan, *Atmos. Environ.*, 271, 118913, <https://doi.org/10.1016/j.atmosenv.2021.118913>, 2022.
- Medeiros, P. M., Conte, M. H., Weber, J. C., and Simoneit, B. R. T.: Sugars as source indicators of biogenic organic carbon in aerosols collected above the Howland Experimental Forest, Maine, *Atmos. Environ.*, 40, 1694–1705, <https://doi.org/10.1016/j.atmosenv.2005.11.001>, 2006.
- Miyazaki, Y., Fu, P. Q., Kawamura, K., Mizoguchi, Y., and Yamanoi, K.: Seasonal variations of stable carbon isotopic composition and biogenic tracer compounds of water-soluble organic aerosols in a deciduous forest, *Atmos. Chem. Phys.*, 12, 1367–1376, <https://doi.org/10.5194/acp-12-1367-2012>, 2012.
- Miyazaki, Y., Gowda, D., Tachibana, E., Takahashi, Y., and Hiura, T.: Identification of secondary fatty alcohols in atmospheric aerosols in temperate forests, *Biogeosciences*, 16, 2181–2188, <https://doi.org/10.5194/bg-16-2181-2019>, 2019.
- Mohammadian, M. A., Watling, J. R., and Hill, R. S.: The impact of epicuticular wax on gas-exchange and photoinhibition in *Leucadendron lanigerum* (Proteaceae), *Acta Oecolog.*, 31, 93–101, <https://doi.org/10.1016/j.actao.2006.10.005>, 2007.
- Mudge, S. M.: Fatty alcohols – A review of their natural synthesis and environmental distribution, *Soap Deterg. Assoc.*, 132, 1–141, <https://doi.org/10.1016/j.febslet.2011.10.016>, 2005.
- Müller, A., Miyazaki, Y., Tachibana, E., Kawamura, K., and Hiura, T.: Evidence of a reduction in cloud condensation nuclei activity of water-soluble aerosols caused by biogenic emissions in a cool-temperate forest, *Sci. Rep.-UK*, 7, 8452, <https://doi.org/10.1038/s41598-017-08112-9>, 2017.
- Nakai, Y., Kitamura, K., Suzuki, S., and Abe, S.: Year-long carbon dioxide exchange above a broadleaf deciduous forest in Sapporo, Northern Japan, *Tellus*, B, 55, 305–312, <https://doi.org/10.3402/tellusb.v55i2.16755>, 2003.
- Nelson, D. B., Ladd, S. N., Schubert, and C. J., and Kahmen, A.: Rapid atmospheric transport and large-scale deposition of recently synthesized plant waxes, *Geochim. Cosmochim. Ac.*, 222, 599–617, <https://doi.org/10.1016/j.gca.2017.11.018>, 2018.
- Nikolić, B., Todosijević, M., Đorđević, I., Stanković, J., Mitić, Z., Tešević, V., and Marin, P.: Nonacosan-10-ol and *n*-Alkanes in Leaves of *Pinus pinaster*, *Nat. Prod. Commun.*, 15, 1–4, <https://doi.org/10.1177/1934578X20926073>, 2020.
- O’Sullivan, D., Murray, B. J., Ross, J. F., Whale, T. F., Price, H. C., Atkinson, J. D., Umo, N. S., and Webb, M. E.: The relevance of nanoscale biological fragments for ice nucleation in clouds, *Sci. Rep.-UK*, 5, 8082, <https://doi.org/10.1038/srep08082>, 2015.
- Oros, D. R. and Simoneit, B. R. T.: Identification and emission factors of molecular tracers in organic aerosols from biomass burning Part 1. Temperate climate conifers, *Appl. Geochem.*, 16, 1513–1544, [https://doi.org/10.1016/S0883-2927\(01\)00021-X](https://doi.org/10.1016/S0883-2927(01)00021-X), 2001.
- Perkins, R. J., Vazquez de Vasquez, M. G., Beasley, E. E., Hill, T. C. J., Stone, E. A., Allen, H. C., and DeMott, P. J.: Relating structure and ice nucleation of mixed surfactant systems relevant to sea spray aerosol, *J. Phys. Chem. A*, 124, 8806–8821, <https://doi.org/10.1021/acs.jpca.0c05849>, 2020.
- Qiu, Y., Odendahl, N., Hudait, A., Mason, R., Bertram, A. K., Paesani, F., DeMott, P. J., and Molinero, V.: Ice nucleation efficiency of hydroxylated organic surfaces is controlled by their structural fluctuations and mismatch to ice, *J. Am. Chem. Soc.*, 139, 3052–3064, <https://doi.org/10.1021/jacs.6b12210>, 2017.
- Reicosky, D. A. and Hanover, J. W.: Physiological effects of surface waxes I. Light reflectance for glaucous and nonglucous *Picea pungens*, *Plant Physiol.*, 62, 101–104, <https://doi.org/10.1104/pp.62.1.101>, 1978.
- Schulten, H.-R., Simmleit, N., and Rump, H. H.: Soft ionization mass spectrometry of epicuticular waxes isolated from coniferous needles, *Chem. Phys. Lipids*, 41, 209–224, [https://doi.org/10.1016/0009-3084\(86\)90023-X](https://doi.org/10.1016/0009-3084(86)90023-X), 1986.
- Sharma, P., Kothari, S. L., Rathore, M. S., and Gour, V. S.: Properties, variations, roles and potential applications of epicuticular wax: A review, *Turk. J. Bot.*, 42, 135–149, <https://doi.org/10.3906/bot-1702-25>, 2018.
- Simoneit, B. R. T., Kobayashi, M., Mochida, M., Kawamura, K., Lee, M., Lim, H. J., Turpin, B. J., and Komazaki, Y.: Composition and major sources of organic compounds of aerosol particulate matter sampled during the ACE-Asia campaign, *J. Geophys. Res.*, 109, D19S10, <https://doi.org/10.1029/2004JD004598>, 2004.
- Spangenberg, J. E., Ferrer, M., Tschudin, P., Volken, M., and Hafner, A.: Microstructural, chemical and isotopic evidence for the origin of late neolithic leather recovered from an ice field in the Swiss Alps, *J. Archaeol. Sci.*, 37, 1851–1865, <https://doi.org/10.1016/j.jas.2010.02.003>, 2010.
- Tegen, I. and Schepanski, K.: Climate Feedback on Aerosol Emission and Atmospheric Concentrations, *Curr. Clim. Change Rep.*, 4, 1–10, <https://doi.org/10.1007/s40641-018-0086-1>, 2018.
- Tulloch, A. P.: Chemistry of waxes of higher plants, chemistry and biochemistry of natural waxes, edited by: Kolattukudy, P. E., Amsterdam, Elsevier, 235–287, 1976.
- Vazquez de Vasquez, M., Carter-Fenk, K. A., Beasley, E. E., Simpson, J. B., Allen, H. C., and McCaslin, L.: Ice nucleation insights: interfacial electric fields and fatty alcohol and acid hydration, *AGU Fall Meeting Abstracts*, 2020, A035-0009, <https://ui.adsabs.harvard.edu/abs/2020AGUFMA035.0009V> (last access: 23 April 2023), 2020.
- Vioque, J. and Kolattukudy, P. E.: Resolution and purification of an aldehydegenerating and an alcohol-generating fatty acyl-CoA reductase from pea leaves (*Pisum sativum* L), *Arch. Biochem. Biophys.*, 340, 64–72, 1997.
- Visez, N., Ivanovsky, A., Roose, A., Gosselin, S., Sénéchal, H., Poncet, P., and Choël, M.: Atmospheric particulate mat-

- ter adhesion onto pollen: A review, *Aerobiologia*, 36, 49–62, <https://doi.org/10.1007/s10453-019-09616-9>, 2020.
- Wang, Y., Wang, J., Chai, G., Li, C., Hu, Y., Chen, X., and Wang, Z.: Developmental changes in composition and morphology of cuticular waxes on leaves and spikes of glossy and glaucous wheat (*Triticum aestivum* L.), *PLOS ONE*, 10, e0143671, <https://doi.org/10.1371/journal.pone.0141239>, 2015.
- Yamamoto, S., Otto, A., and Simoneit, B. R. T.: GC-MS analysis of wax in leaf of *Sequoiadendron giganteum*, *Sequoioideae*, *Cupressaceae*, *Res. Org. Geochem.*, 224, 159–171, 2008.
- You, B., Li, S., Tsona, N. T., Li, J., Xu, L., Yang, Z., Cheng, S., Chen, Q., George, C., Ge, M., and Du, L.: Environmental Processing of Short-Chain Fatty Alcohols Induced by Photosensitized Chemistry of Brown Carbons, *ACS Earth Space Chem.*, 4, 631–640, <https://doi.org/10.1021/acsearthspacechem.0c00023>, 2020.
- Yu, L. E., Shulman, M. L., Kopperud, R., and Hildemann, L. M.: Characterization of organic compounds collected during the southeastern aerosol and visibility study: Water-soluble organic compounds, *Environ. Sci. Technol.*, 39, 707–715, <https://doi.org/10.1021/es0489700>, 2005.
- Zhang, M., Khaled, A., Amato, P., Delort, A.-M., and Ervens, B.: Sensitivities to biological aerosol particle properties and ageing processes: potential implications for aerosol–cloud interactions and optical properties, *Atmos. Chem. Phys.*, 21, 3699–3724, <https://doi.org/10.5194/acp-21-3699-2021>, 2021.
- Zhu, C., Kawamura, K., and Kunwar, B.: Organic tracers of primary biological aerosol particles at subtropical Okinawa Island in the western North Pacific Rim, *J. Geophys. Res.-Atmos.*, 120, 5504–5523, <https://doi.org/10.1002/2015jd023611>, 2015.