



Ancient clays support contemporary biogeochemical activity in the Critical Zone

Vanessa M. Alfonso¹, Peter M. Groffman^{1,2}, Zhongqi Cheng¹, and David E. Seidemann¹

¹Department of Earth and Environmental Sciences, Brooklyn College of the City University of New York, 2900 Bedford Avenue, Brooklyn, NY 11210, USA

²Advanced Science Research Center at the Graduate Center, City University of New York, 85 St. Nicholas Terrace, New York, NY 10031, USA

Correspondence: Vanessa M. Alfonso (vanessa.alfonso23@bcmail.cuny.edu)

Received: 18 April 2024 – Discussion started: 18 July 2024

Revised: 28 March 2025 – Accepted: 1 April 2025 – Published: 10 July 2025

Abstract. Late Cretaceous clays exposed at sites located on the north shore of Long Island, New York, USA, were sampled to explore questions about how contemporary factors and processes interact with ancient geological materials that are often assumed to not be biologically active. Chemically and biologically catalyzed weathering processes have produced multi-colored clays belonging to the kaolin group with inclusions of hematite, limonite, and pyrite nodules. We sampled exposed clays at three sites to address three questions: (1) do these exposed clays support significant amounts of microbial biomass and activity, i.e., are they alive? (2) Do these clays support significant nitrogen (N) cycle activity? (3) Are these clays a potential non-anthropogenic source of reactive N in the contemporary landscape? Samples were analyzed for total carbon (C) and N content, microbial biomass C and N content, microbial respiration, organic matter (OM) content, potential net N mineralization and nitrification, soil nitrate (NO_3^-) and ammonium (NH_4^+) content, and denitrification potential. Results strongly support the idea that ancient geologic materials play a role in contemporary N and C cycling in the Critical Zone. Respiration (average $4.098 \mu\text{g C g}^{-1} \text{d}^{-1}$) was detectable in all samples and was strongly correlated to OM, indicating a living microbial community on the clays. There was evidence of an active N cycle. Higher levels of denitrification potential (average $1.376 \mu\text{g N g}^{-1} \text{d}^{-1}$) compared to both potential net nitrification (average $0.061 \mu\text{g N g}^{-1} \text{d}^{-1}$) and potential net N mineralization (average $0.144 \mu\text{g N g}^{-1} \text{d}^{-1}$) indicate that these clays act more as a sink rather than as a source of reactive N in the landscape.

1 Introduction

The Critical Zone is Earth's constantly evolving boundary layer where rock, soil, water, air, and living organisms interact (Schroeder, 2018). It is comprised of the solid phase matter in which water circulates and is stored, expanding from the top of the vegetation canopy down into the water bearing bedrock. Critical Zone processes are key drivers of chemical transfers between biota and geological materials (Brantley et al., 2006).

A major question in Critical Zone science is how contemporary factors and processes interact with ancient geological materials. These materials are often assumed to have little or no biogeochemical activity as they have not been involved in active biological activity for long periods of time and may lack active microbial communities and labile pools of carbon (C) and nitrogen (N) that drive this activity. These interactions are particularly obvious and important in the lower boundary of the Critical Zone, and in places where ancient geologic materials become exposed to contemporary environments (Dass et al., 2021). Major questions center on the ability of ancient materials to support biogeochemical processes related to the cycling of C and N that underlie plant and microbial activity, which underlies environmental and ecosystem “services” of interest to society.

The discovery of significant amounts of N in sedimentary deposits increased interest in the role of geological materials in contemporary N cycling (Morford et al., 2011). Up to 17 % of the currently cycling N in some ecosystems may originate from rock materials deep in the Critical Zone (Houlton et al.,

2018). Such observations motivate the analysis of surface-exposed ancient rocks in this study. Both NH_4^+ and NO_3^- are highly soluble and the negative charge of NO_3^- makes it highly mobile, thus driving hydrologic losses of N. In excessive amounts, these nutrients lead to ecological stresses such as eutrophication. NH_4^+ is converted to NO_3^- by nitrifying bacteria, and NO_3^- is converted to N_2 by denitrifying bacteria leading to gaseous losses of N back into the atmosphere (Seitzinger et al., 2006).

In this study, we sampled surface exposures of unconsolidated bulk clay units exposed in outcrops that showed no evidence of modern soil development and measured microbial biomass and activity to address three questions: (1) do these exposed clays support significant amounts of microbial biomass and activity, i.e., are they alive? To address this question we measured microbial biomass C, an index of the living microbial biomass in soil, and microbial respiration, a direct measure of microbial activity (Paul, 2014). Total C and organic matter (OM) content were measured as energy sources for microbial biomass and activity. (2) Do these clays support significant amounts of N cycle activity? N cycle activity was assessed with measurements of microbial biomass N content, potential net N mineralization, and total N content. Microbial biomass N provides an index of the net flux of N through microbial pools. Mineralization results from microbial degradation of N compounds resulting in the production of inorganic, plant-available forms of N. Total N content was measured to quantify the total amount of N potentially available for active cycling. (3) Are these clays a potential non-anthropogenic source of reactive N in the contemporary landscape? The potential for the clays' microbial activity to be a source or "sink" for reactive N was evaluated by measuring potential net nitrification, denitrification potential, and pools of ammonium (NH_4^+) and nitrate (NO_3^-).

2 Background

2.1 Clay minerals and their role in ecosystem function and nutrient cycling

Clay minerals are hydrous alumino-silicates, more specifically hydrous phyllosilicates, and are some of the most stable products of chemical weathering at surface conditions. These minerals produce a specialized microhabitat and their ability to store and release nutrients make them ecologically important (Kleber et al., 2021). Geochemical composition and pore space of these materials, affect microbial activity and therefore rates of biogeochemical processes in C and N cycles (Li et al., 2023). Clay minerals have high surface adhesion capabilities and sorption and desorption of OM in soils varies with mineral assemblage. Sorption of organic carbon (OC) onto phyllosilicates and hydrous iron (Fe) oxides affects accumulation and stabilization of OC in soils (Saidy et al., 2013). The 1 : 1 layers of hydrated kaolinite clays and 2 : 1

layers of mixed layer clays, along with OM, serve as nutrient exchange sites between biomass and subsurface weathering horizons. Therefore, the layered structures of clays are key facilitators of seasonal cycling of nutrients including NH_4^+ among others (Eby, 2016; Halama and Bebout, 2021). Chemical weathering of silicate minerals is a significant mechanism for the availability, uptake, storage, and transport of key nutrients in ecosystems. Variations in mineral weathering and nutrient availability occur due to microorganism and mineral speciation, while intensity of mineral weathering is influenced by a mineral's potential to provide nutrients (Bennett et al., 2001). The pathways of C and N from the atmosphere to incorporation into the lithosphere progress from fixation into OM to storage in low temperature silicate phases such as clay minerals (Busigny and Bebout, 2013; Halama and Bebout, 2021). This is facilitated by the aqueous solutions and living microbes characteristic of the Critical Zone.

2.2 Geological background

Late Cretaceous clays on the north shore of Long Island, New York, USA, offer an opportunity to study the effects of contemporary factors and processes on ancient Critical Zone materials. Long Island is composed of Pleistocene sediments deposited on top of Late Cretaceous formations (Sirkin, 1991) and includes aquifers, confining units, and clay-rich deposits, such as the Raritan, Magothy, Gardiners, and Wantagh Clay formations, that overlie gneissic bedrock. Coastal exposures of the Raritan and Magothy formations present opportunities for biogeochemical analysis of these materials. At these locations, clays are exposed in outcrops, offering the chance to investigate their contemporary biogeochemical activity at surface conditions (Fig. 1). Long Island's clay strata have been mainly accessed through core drilling for hydrological studies; in most instances the clays are a secondary detail rather than the main subject of study. The same is true for palynology studies in which the clay is the assemblage containing the ancient pollen being studied. In contrast, this study focuses on biogeochemical activity and how the clay strata affect and interact with surrounding environments in the Critical Zone. The kaolinitic materials at these sites and their abundance in oxides contribute to long lasting micro-environments (Six et al., 2000).

All three study sites are located on the north shore of Long Island, NY. Two of the sites are located along the shoreline – Garvies Point Preserve (GP) on Hempstead Bay and Caumsett State Historic Park Preserve (CSP) along Long Island Sound; Hempstead Harbor Woods (HHW) is located inland from the western side of Hempstead Bay. Long Island contains numerous clay beds such as Gardiner's Clay, Raritan Confining Unit, Wantagh Clay, and Smithtown Clay, as well as clay lenses in the Magothy Formation (Mills and Wells, 1974). The study area was shaped by the Wisconsin glaciation approximately 75 000 to 11 000 years ago. During this period, Cretaceous strata were sheared off, transported, and



Figure 1. Map of study sites on Long Island, New York (maps constructed using map data from Esri and © Google Earth).

re-deposited. Glacially induced thrusting of the strata facilitated bulging of the clay (Mills and Wells, 1974), thus leading to eventual exposure. Evidence of this can be observed in the outcrop sampled for this study at GP, where the clay layers are currently oriented vertically rather than in a near horizontal depositional position. Although there is consensus that the Late Cretaceous deposits of the western north shore of Long Island were formed in a shallow delta or estuary (Fuller, 1914; Swarzenski, 1963) and that they are composed of sand, silt, gravel, and clay, further augmented by eolian action, it is not completely clear to which formation the exposed North Shore clays belong. It is generally accepted that the Late Cretaceous formations exposed on the North Shore tentatively belong to the Magothy formation but may include some younger formations (Isbister, 1966). It is presumed that Pleistocene deep permafrost formation must have preceded glacial thrusting (Mills and Wells, 1974), which provided the sedimentary cohesion needed to produce the tilted clay strata and layered shale (not sampled for this study) observed today. The three sites of this study also differ in their position on the Manhasset Plateau. The HHW and GP localities are situated on the Upper Manhasset Plateau while CSP is situated on the Lower Manhasset Plateau. The Lower Manhasset Plateau is thought to have sustained a more prolonged grinding by the ice sheet (Fuller, 1914). The most abundant clay species identified from core drills in northwestern Long Island is kaolinite of light gray, brown-yellow, and tan colors, along with sparse chlorite, vermiculite, and montmorillonite, with the kaolinite ratio to other clays decreasing with depth (Liebling, 1973).

Fe content, among other factors, has contributed to variations in coloring and produced distinctive clay strata, allowing for color based grouping of the samples. The color variations are the result of differences in constituents and geo-

chemical processes that evolved the clays into their modern state. The Fe oxides in hematite are likely sources of the red and brownish coloring and the Fe oxides in goethite are likely sources of the yellow coloring (Davey et al., 1975). This is further supported by portable x-ray fluorescence (pXRF) scan results (Sects. 4 and 5) which show red, yellow, and brown samples to have some of the highest Fe contents. Mn also contributes to brown coloring (Jakobsson et al., 2000) which is further supported by pXRF scans that show brown samples to have the highest Mn content. Recent conditions have added sand and water to some clays to yield different textures, providing another observational classification criterion.

3 Methods

3.1 Study sites

Garvies Point Preserve (GP) is located in Glen Cove, NY, on the eastern shoreline of Hempstead Bay at $40^{\circ}51'35''$ N $73^{\circ}39'07''$ W. The exposed clays were accessed via trails leading to the beach and samples were collected along approximately 450 m of shoreline, with special focus on the main outcrop which features five types of clays exposed at this site. The most prominent outcrop at GP is approximately 4 m high and yielded samples from five differently colored clays (light gray, dark charcoal gray, white, yellow, and dark red/purplish). The adjacent outcrop is approximately 2 m high containing light and dark gray clay.

Hempstead Harbor Woods (HHW) is located in North Hempstead, NY, at $40^{\circ}50'11''$ N $73^{\circ}39'57''$ W, on the inland western side of Hempstead Bay, approximately 300 m from the shoreline. Samples were collected from 0–1 m from the ground level throughout the wooded area. Although this location does not have direct exposures to the bay, some of the sampled clay materials have indicators of being part of the same formation as those exposed across the bay at GP. Specifically, the red and light gray packed clays collected at HHW share the color and texture attributes of those at GP.

At Caumsett State Historic Park Preserve (CSP) samples were collected from exposures along approximately 1000 m of shoreline. A prominent exposure at CSP located at $40^{\circ}56'21''$ N $73^{\circ}28'13''$ W has an elevation of approximately 40 m. Samples were collected starting from the bottom of the cliff, just above the beach floor, and as high as 6 m above the beach floor. Additional samples were collected from exposures at $40^{\circ}56'08''$ N $73^{\circ}28'14''$ W approximately 4 m above the beach floor, and at $40^{\circ}56'11''$ N $73^{\circ}28'43''$ W approximately 6 m above the beach floor.

3.2 Sampling

At each site samples were collected from locations that indicated the highest likelihood of yielding multiple clay types. Large outcrops containing multiple variations of clay at GP

and CSP were sampled, yielding 14 and 5 samples respectively. The prominent outcrops at these locations feature variety in both color and texture. While our sampling facilitated comparison of these obvious differences, we did not have any *a priori* hypotheses about their causes and biogeochemical effects. HHW featured scattered locations, each featuring clay of a single texture and color, yielding 9 samples. Samples were collected from unconsolidated bulk clay units that showed no evidence of modern soil development, i.e. accumulation of recent organic matter or development of horizons. One sample from HHW (sample 3N-HHW) was collected from a naturally exposed soil horizon (B horizon) underneath topsoil (O horizon).

Samples were collected from the surface of the exposures. Locations where clay had at least a few inches of depth were chosen. The uppermost layer (~ 1 cm) was scraped off to ensure only clay was collected and to remove any field debris such as loose soil, rocks, sand, and loose plant material. Samples were collected from approximately 5–10 cm of depth. The samples were collected using a trowel, placed in labeled Ziplock bags, and packed loosely with some air remaining in the bag. Rocks, roots, and leaves were removed by hand right after collection. The samples were then refrigerated until laboratory processing. Samples were classified into groups of color (brown, dark gray, light gray, red, white, yellow) and texture (packed clay, sandy clay, watery clay) by visual observation. The packed clay was homogeneous, contained some moisture, and had a putty like texture that easily formed into a ribbon several inches long. The sandy clay was drier and had fine sand mixed in. The watery clay was collected from a small basin of waterlogged clay that was homogeneous, had very fine particles, and a thick viscous texture.

3.3 Laboratory analysis

Samples were analyzed for moisture content by oven drying for 24 h at 105 °C. Moisture content was used to calculate values for all variables on a per g of dry soil basis.

OM content was determined by loss on ignition (LOI). Oven-dried samples were combusted at 450 °C in a muffle furnace and % OM was calculated from weight loss after 8 h of heating.

For elemental analysis by pXRF, samples were oven-dried for 24 h at 80 °C, ground with a mortar and pestle, sieved through a #230 (63 μ m) sieve, tightly packed into 2.75 cm diameter holders, and scanned with a portable Olympus DC-4000 XRF scanner.

Total C, total N, and the C/N ratio were measured using flash combustion/oxidation. Oven-dried and ground samples were pressed into 1 g pellets. The pellets were analyzed in an Elementar vario MAX cube elemental analyzer.

For analysis of exchangeable NO_3^- and NH_4^+ , samples were blended with 2 M KCl on an orbital shaker at 125 rpm for 1 h, followed by filtration (Whatman #42 filter) into scintillation vials that were immediately refrigerated until analy-

sis. The samples were pipetted into microplates and analyzed on a SpectraMax M2 Multi-Mode Microplate Reader from Molecular Devices using wavelengths of 450 nm for NO_3^- , and 650 nm for NH_4^+ (Doane and Horwath, 2003; Sims et al., 1995). Total inorganic N (TIN) was calculated as the sum of NH_4^+ and NO_3^- .

The chloroform fumigation and incubation method (CFIM) (Jenkinson and Powlson, 1976) was used to determine the C and N content of microbial biomass. Samples (10 g) were fumigated with chloroform for 24 h, inoculated with 0.2 g of unfumigated clay and incubated for 10 d in 946 mL Mason jars with lids fitted with septa. At the end of the incubation, gas samples were taken by syringe and analyzed for carbon dioxide (CO_2) with a Shimadzu GC-2014 gas chromatograph. After gas sampling, fumigated soils were extracted using KCl as described above after 10 d and the NO_3^- and NH_4^+ produced over the 10 d incubation was taken as an estimate of microbial biomass N. A proportionality constant of 0.41 was used to calculate microbial biomass C from the CO_2 produced over the 10 d incubation (Robertson, 1999).

Unfumigated samples were also incubated for 10 d and provided estimates of microbial respiration and potential net N mineralization and nitrification. These samples were incubated and sampled as described above and production of CO_2 during the 10 d incubation was taken as an estimate of microbial respiration. Production of NO_3^- and NH_4^+ over the 10 d incubation was taken as an estimate of potential net N mineralization and production of NO_3^- was taken as an estimate of potential net nitrification.

A denitrification enzyme assay (DEA) was used to measure the rate of potential denitrification (Groffman et al., 1999; Smith and Tiedje, 1979). A quantity of 5 g of each sample was placed in 125 mL Erlenmeyer flasks and amended with 10 mL of DEA media. DEA media was prepared by adding 0.72 g KNO_3^- as a substrate for existing enzymes, 0.5 g glucose as a source of C/energy, and 0.125 g chloramphenicol to block production of new enzymes during incubation to 1 L of nanopure water. The flasks were sealed with rubber stoppers, flushed repeatedly with N_2 gas to create anaerobic conditions, amended with acetylene (C_2H_2) gas, and placed on a shaker at 125 rpm. The headspace of the flasks was sampled (8 mL) by syringe after 30 and 90 min of incubation. Samples were analyzed for nitrous oxide (N_2O) with a Shimadzu GC-2014 gas chromatograph.

3.4 Statistical analysis

SPSS version 28 was used for all analyses. One-way analysis of variance (ANOVA) with post hoc multiple comparison tests were run on all response variables using location (GP, HHW, CSP), texture (packed, sandy, watery), and color (yellow, white, red, light gray, dark gray, brown) as grouping factors. A Sidak adjustment was applied to the post hoc multiple comparison tests because the number of samples was

not equal across groups. Spearman's rho was used to evaluate linear correlations because the data were not normally distributed and had a heavy positive skew. ANOVA pretest results for homogeneity of variance were reinforced with Welch and Brown–Forsythe tests. Kruskal–Wallis analysis was run to increase confidence in results for groupings by color, texture, and location. Mann–Whitney analysis with the Monte Carlo option was used to further reinforce ANOVA results grouped by location. Significance levels were evaluated based on $\alpha < 0.01$ indicating a strong statistically different significance, $\alpha < 0.05$ indicating a standard statistically different significance, and $\alpha < 0.1$ indicating a marginal statistically different significance. Since ANOVA results yielded the same significance parameters for the majority of variables as Kruskal–Wallis and Mann–Whitney tests, ANOVA values are reported in the results section.

4 Results

4.1 Basic characteristics of clays

All samples had detectable amounts of C, N, and OM (Fig. 2). The dark gray samples had significantly higher OM, total C, total N, and C/N ratio than several of the other clay color types. The packed clay had the highest values of all these variables among the texture groups and significant differences were found for OM between packed and sandy clays. ANOVA post hoc analysis identified further significant differences in the C/N ratio when comparing dark gray clay to the Fe bearing red, yellow, and brown clays. A number of packed, dark gray and light gray samples contained concentrations of N exceeding $1000 \text{ mg N kg}^{-1}$ which is considered ecologically significant for geologic N (Holloway and Dahlgren, 2002).

Semi-quantitative results from pXRF scans provide estimates of selected elements of interest (Table 1a). Phosphorus (P), a critical nutrient, was highest in brown clay and in watery clay with marginal differences between GP and HHW. Chlorine (Cl), which plays a role in nutrient transport, was highest in yellow, brown, and red clays, and in watery clays with significant differences among color groups and locations. Titanium (Ti), which plays a role in nutrient cycling and enhances clay adsorption properties, was highest in yellow, red, and brown clays, and in packed clays with significant differences among packed and sandy clay textures and marginal differences between GP and CSP locations. Manganese (Mn), which plays a role in nitrogen metabolism, was highest in brown clays and watery clays, with significant differences among color groups, textures, and locations. Fe, a critical nutrient, was highest in brown, red, yellow clays, and in watery clays, with significant and marginal differences among color groups and locations. Rubidium (Rb) can shed light on clay speciation and therefore nutrient cycling capacity and was highest in dark gray, and in packed clays,




Figure 2. Total C, N, OM content, and C/N ratio of different colored clays. Values are mean $\pm$ SD. Values with different superscripts are significantly different ($p < 0.05$) except: in total C dark gray clay was marginally different ($p < 0.10$) from yellow clay; in total N dark gray clay was marginally different ($p < 0.10$) from brown and white clays; C/N ratio in dark gray clay was marginally different ($p < 0.10$) from brown clay.

with significant differences among color groups and locations. Strontium (Sr) can interact with other elements such as Calcium (Ca) and had similar estimated abundances across color groups with the highest concentrations in the light gray, dark gray, brown, and yellow samples. Zirconium (Zr), which can enhance the adsorption properties of clays, was highest in brown and white samples, and in sandy and watery clays, with significant and marginal differences among color groups, textures, and locations. Zinc (Zn), an essential nutrient, was highest in brown and dark gray, and in watery clays, with significant and marginal differences among color groups and locations. The Rb/Sr ratio can serve as a proxy for environmental changes that affected the clays. It was highest in dark gray clays, was nearly the same across textures and had significant differences among color groups and locations (locations had additional marginal differences). There were significant correlations between P and Cl, P and Mn, P and Fe, Cl and Mn, Cl and Fe, Ti and Zr, Mn and Fe, Mn and Zn, Zn and Rb, Zn and Sr, Rb and Sr, Sr and Zr, and marginal correlations between P and Ti, and Ti and Fe.

4.2 Do these exposed clays support significant amounts of microbial biomass and activity, i.e., are they alive?

Most clay types exhibited detectable amounts of microbial biomass and activity (Fig. 3). Microbial biomass C was significantly higher (by 135 %) in the watery clay than in the packed or sandy clay. A significant difference between watery and sandy clays was identified by ANOVA post hoc

Table 1. (a) Estimated amounts of selected elements of interest detected by pXRF analysis; values are means in ppm. (b) Standard deviations of estimated elemental content for groups with more than one sample.

(a)	P	Cl	Ti	Mn	Fe	Rb	Sr	Zr	Zn	Rb/Sr ratio
Garvies Point										
dark gray	10 991.38	140 312.87	4171.20	67.73	6736.80	139.20	73.93	288.47	65.47	1.89
light gray	3372.50	133 413.83	3441.33	49.33	4820.00	111.33	70.17	284.33	45.17	1.56
red	30 710.67	161 559.78	4884.67	52.33	27106.89	98.67	67.67	294.67	48.56	1.46
white	9194.00	126 827.67	4351.00	85.67	6296.33	137.00	75.67	335.33	48.33	1.81
yellow	25 721.33	160 147.78	4566.11	54.00	16 648.44	110.33	72.22	340.44	34.00	1.52
Hempstead Harbor Woods										
brown	14 576.00	152 885.00	3806.00	204.67	14 309.00	93.67	80.33	330.00	73.00	1.17
light gray	8068.67	123 039.50	3832.17	40.17	6617.50	96.61	76.56	376.56	34.06	1.37
red	5371.50	139 207.00	4649.67	23.33	7586.33	42.67	41.00	448.33	33.67	1.04
white	442.00	135 305.00	3644.33	32.33	4786.33	79.67	51.00	804.67	23.00	1.56
Caumsett State Park										
brown	31 116.83	161 317.92	3584.17	352.92	25 723.83	98.08	72.08	676.67	74.75	1.39
red	3942.00	142 768.00	3741.00	81.00	18 065.67	67.33	62.33	387.00	59.67	1.08
(b)	SD of P	SD of Cl	SD of Ti	SD of Mn	SD of Fe	SD of Rb	SD of Sr	SD of Zr	SD of Zn	SD of Rb/Sr
Garvies Point										
dark gray	5524	18 068	210	18	1633	8	6	32	21	0.1
light gray	380	1430	647	5	1257	41	9	16	30	0.4
red	23 507	12 890	906	14	20 172	13	6	34	30	0.1
yellow	19 354	10 021	318	3	11 599	18	8	60	2	0.1
Hempstead Harbor Woods										
light gray	5363	7575	560	11	1325	9	28	130	10	0.4
Caumsett State Park										
brown	15 314	20 994	528	100	4136	9	15	194	18	0.2

analysis. There were no significant differences in microbial biomass C with clay color. Soil respiration was detectable in all materials but there were no significant differences with clay color or texture. There were significant correlations between microbial biomass C and total N, total C (Fig. 4), OM, and respiration. Respiration was significantly correlated with total N (Fig. 4) and OM, and was also marginally correlated with total C and C/N ratio (Fig. 4).

4.3 Do these clays support significant amounts of N cycle activity?

There were significant differences in microbial biomass N among textures, and amounts were highest in dark gray and packed clays. We detected both potential net N mineralization and N immobilization (negative net N mineralization) in our samples (Fig. 3). Potential net N mineralization was highest in brown and in packed clays, with significant differences among samples from all locations. Net N mineralization was marginally different between the GP and HHW locations.

Microbial biomass N was significantly correlated with total C, total N, C/N ratio (Fig. 6), potential net nitrification, OM, and TIN. Total N was significantly correlated with microbial biomass C, respiration (Fig. 4), OM, TIN (Fig. 5), microbial biomass N, potential net N mineralization (Fig. 6), and total C. Potential net N mineralization was significantly correlated with total C, total N, C/N ratio (Fig. 6), TIN, microbial biomass N, and potential net nitrification. Total N and the C/N ratio were marginally correlated.

4.4 Are these clays a potential non-anthropogenic source of reactive N in the contemporary landscape?

The highest amounts of potential net nitrification were found in light gray (Fig. 3) and packed clays, with significant differences between the GP and HHW locations identified by ANOVA post hoc analysis. Light gray clays at HHW had 100 % higher values than the same color group at GP. The highest amounts of NO_3^- were found in light gray (Fig. 7) and packed clays (light gray clays at HHW had 6447 % higher values than the same color group at GP), with significant




Figure 3. Potential net N mineralization, potential net nitrification, denitrification potential, microbial biomass N content, microbial biomass C, and respiration. Values with different superscripts are significantly different at $p < 0.05$ except that dark gray clay was marginally different ($p < 0.10$) from brown and white clays. All values are mean $\pm$ SD.

differences between GP and HHW, and between HHW and CSP identified by ANOVA post hoc analysis. The highest amounts of NH_4^+ were found in dark gray (Fig. 7), packed clays at GP, with significant differences among color groups (763 % higher amount in dark gray than red clays at GP). The highest amounts of TIN were found in dark gray and packed clays (280 % higher than light gray, and 1057 % higher than sandy clay, respectively), with significant differences among color and texture groups. Kruskal–Wallis analysis identified a marginally significant difference for TIN by location. There were no significant differences in denitrification potential.

Potential net nitrification was significantly correlated with total C, C/N ratio (Fig. 8), microbial biomass N, potential net N mineralization, and marginally correlated with TIN. Denitrification potential was significantly correlated with the C/N ratio (Fig. 8), suggesting joint control by C and N availability. There were significant correlations between NH_4^+ pools and the following variables: total C, total N, C/N ratio (Fig. 9), microbial biomass N, potential net N mineralization, TIN, and OM. NO_3^- pools were significantly correlated with total N (Fig. 9), microbial biomass N, and potential net N mineral-

ization. There was a marginal correlation between NO_3^- and total C (Fig. 9).

5 Discussion

Our study sought to explore organic-mineral interactions by investigating the functional performance of clays through measurements of biogeochemical processes occurring within their structures. This research aimed to expand knowledge beyond the traditional approach of investigating C and N cycle dynamics based on single predictor values such as phyllosilicate clay, abundance of a certain mineral species, and specific surface area and adsorption values (Kleber et al., 2021) by making a suite of measurements of microbial biomass and activity relevant to C and N cycles. This approach allowed us to determine that ancient clays are contributing to contemporary Critical Zone biogeochemical processes at ecosystem and landscape scales.

5.1 N, C, and OM in geological materials

Over geological time scales, N and C cycle from the Earth's surface to depth through subduction zones and are returned to the surface through arc magmatism. Geological cycling of N is greatly facilitated by storage of NH_4^+ in silicate minerals. NH_4^+ in minerals is a major influence on fluxes between reservoirs (Halama and Bebout, 2021), thus giving geological materials such as the clays we have analyzed a potential role in ecosystem function.

For contemporary ecosystem processes, it is generally assumed that geologic materials are not an important source of N (Schlesinger, 2013), as the dominant global pool of N is in the atmosphere. The focus of much N cycle research is on the energetically expensive movement of atmospheric N into biological pools (Galloway et al., 2004). The largest pools of N in ecosystems are in particulate and dissolved OM pools in soils, sediments, and the ocean (Groffman et al., 2021). While it is reasonable to assume that some of the N in our samples could be of modern biological origin and is adsorbed onto the surface of the clay particles, our sampling approach that targeted areas with no evidence of modern OM or soil development suggests that our analysis was of older N associated with geologic materials. Increasingly, recognition of the fact that abundant organic N becomes incorporated into geological materials has fueled interest in the role of these materials in contemporary ecosystem processes. Relevant analyses show that 10^{21} g of global fixed N were incorporated into meta-sedimentary mica-rich rocks by the burial of OM in marine and freshwater sediments during the Cretaceous (Morford et al., 2011). These analyses have stimulated studies of the movement of deeply buried organic N into actively cycling pools of N in soils and vegetation (Houlton et al., 2018) and interest in the exposure of ancient materials at the soil surface, such as the clays collected for this study.




Figure 4. Regression of microbial biomass C and respiration against total C, total N, and C/N ratio. Values are represented for each location; trend line is indicated among all locations. Significant differences are indicated by p value and R^2 .

At our study sites, burial of OM likely occurred in a Late Cretaceous shallow delta (Fuller, 1914). The N content of these samples is within the range of previous reports of sedimentary and metasedimentary rock N content of 200–1200 mg N kg^{-1} (Holloway and Dahlgren, 2002; Morford et al., 2011). Organic-rich marine sediments commonly exceed 1000 mg N kg^{-1} and some of our samples fit this criterion (Li, 1991; Morford et al., 2011). As these bedrock materials weather, N is released in plant available forms that stimulate ecosystem productivity and C storage (Dahlgren, 1994; Morford et al., 2011). Bedrock is also a source of N to aquifers, which is relevant for our samples which have a hydrogeologic origin as part of the clay confining units of Long Island's aquifer system. The samples were collected at ex-

posed aquifer margins in this region with great concern about groundwater reactive N (Karamouz et al., 2020). The limited literature on the N content of geological materials does not include any studies of clay materials similar to those studied here.

The surface exposure of the clays in our study allowed us to directly measure microbial biomass and activities that are central to biogeochemical cycling of C and N. These measurements shed light on the role that these secondary minerals, that were produced by the weathering of primary silicate minerals which originated at depth, may be playing in the contemporary N cycle on Long Island and in the Critical Zone elsewhere on Earth. These measurements also allowed for comparison of these geological materials with surface and




Figure 5. Regression of OM and TIN against total C, total N, and C/N ratio. Values are represented for each location; trend line is indicated among all locations. Significant differences are indicated by p value and R^2 .

subsurface soils in the region that have been assayed with the same methods (Groffman et al., 2009; Morse et al., 2014). Our analysis showed significant microbial biomass and activity in many samples, with much of the variation in activity driven by the total C and N content of the samples. The results strongly support the idea that ancient geologic materials play a role in contemporary N and C cycling in the Critical Zone.

Biogeochemical processes are influenced by microbial-mineral associations that influence the rates and magnitudes at which biogeochemical reactions occur. The structure of the mineral space influences the extent of biologically mediated reactions occurring within this habitat. Clays are ex-

pected to support less of these processes due to their tightly bound layered structure and fine particle size (Kleber et al., 2021). This is consistent with the lower amounts of activity detected in our samples compared to other soils, discussed in Sect. 5.2. The tight structure of the clay offers a micro-environment that is more constant over time, as compared to the more dynamic conditions found within soil (Kleber et al., 2021). The longevity of a micro-environment is affected by both OM and mineralogy, with kaolinitic 1 : 1 clays and oxides producing longer lasting environments (Six et al., 2000). The smaller pore size of clay facilitates anaerobic conditions, leading to higher denitrification rates in clay rich materials than in sandy ones (Li et al., 2023; Pihlatie et al., 2004). This




Figure 6. Regression of microbial biomass N and net N mineralization against total C, total N, and C/N ratio. Values are represented for each location; trend line is indicated among all locations. Significant differences are indicated by p value and R^2 .




Figure 7. Exchangeable NO_3^- , NH_4^+ , and TIN in clays of different colors. Values are means $\pm$ SD. Values with different superscripts are significantly different at $p < 0.05$.




Figure 8. Regression of net nitrification and denitrification potential against total C, total N, and C/N ratio. Values are represented for each location; trend line is indicated among all locations. Significant differences are indicated by p value and R^2 .

relationship between particle size and microbial activity was evident in the analysis results obtained from our samples.

What then are the sources of the OC supporting microbial biomass and activity in the samples in this study, which varied with clay color and texture? Plant litter and soil debris from the overlying deposits are certain contributors of OC to the exposed clays at these locations. Ample C in coarse clay ($0.2\text{--}2.0\mu\text{m}$) has been documented to be in the form of charcoal or black carbon (BC) (Laird et al., 2008). We have observed BC within the clay layers at the GP site, both distinctly layered with white clay and as small inclusions in some of the gray clay, which contains the highest amount of OM at that site (up to 15.8 %). This presence indicates that BC is one of the contributing sources of OC and has become incorporated into silt and clay fraction minerals in the geo-

logical unit that is exposed at the GP site through processes such as adsorption of dissolved biogenic compounds onto the clay particle surfaces (Laird et al., 2008).

Clay color is also affected by Fe, which likely contributed to the pigmentation of the red, yellow, and brown clays in our study. Our pXRF scans detected Fe content as high as 4.55 % in some samples. Possible sources of Fe include oxidative reactions and organic ligand bonding, both of which can be catalyzed by bacteria. The Fe content of Long Island's aquifers is driven by processes that include oxidative dissolution of minerals such as pyrite (FeS_2) (Brown and Schoonen, 2004). Pyrite inclusions occur in the clays at these study sites and our pXRF scans detected the presence of sulfur (S) in addition to the Fe in our samples. Chelation of ions and organic acids is common in the Critical Zone. In the pres-




Figure 9. Regression of exchangeable NO_3^- and NH_4^+ against total C, total N, and C/N ratio. Values are represented for each location; trend line is indicated among all locations. Significant differences are indicated by p value and R^2 .

ence of water, Fe is one of the metal ions that can chelate with organic acids and become mobilized through the subsurface (Schroeder, 2018). Furthermore, Fe, Ti, and Mn, all of which were detected in our samples, play a key role in mineral and organic oxidation reactions. Fe(II), FeO_2 , and TiO_2 bearing clays produce the highest amounts of reactive oxygen species, which in turn react with organic C to transform soil and sediment OM, and to produce CO_2 as the end-product of organic C oxidation (Kleber et al., 2021). Mn oxides are the strongest naturally occurring oxidants and play an important role in organic C transformation (Remucal and Ginder-Vogel, 2014). The clay color variations, indicating

variation in oxidative or reducing conditions, may reflect biogeochemical conditions and activity during their formation that have legacy effects on contemporary activity.

The much higher occurrence of significant differences among the clays when grouped by color and texture, as opposed to when grouped by location, indicates that land use history and other local characteristics have less or no effect, and that the main differentiator for microbial activity is the clay material itself and variations therein, such as Fe oxides, trace elements, and variations in clay speciation. It also indicates that the clays from the different locations are likely

part of the same larger formation (presumably Raritan or Magothy).

5.2 Are these ancient clays contributing to contemporary biogeochemical processes?

Our study was driven by three questions, the first of which was the most fundamental: do these materials support living microbial biomass? Our analysis detected a living microbial community on most of the clay samples. There were strong positive relationships between total C content, total N content, and microbial biomass and activity. The dark gray clays had the highest total N, total C, and microbial biomass N, and microbial biomass C was highest in both dark gray and brown clays. White clays had the lowest contents of these variables, indicating that inclusions of Fe and C (discussed in Sect. 5.1) in the clays play a role in their capacity to sequester C and N and support microbial biomass and activity.

While our clay samples had significant amounts of microbial biomass and activity, it is important to compare our results with other materials and soils in our region (the northeastern USA) to evaluate just how important this activity might be. Both surface organic and subsurface mineral horizons of fully developed soils often have substantial C and N pools (Bohlen et al., 2001), with variations in distribution. We compared the data of our clay samples to two studies that used the same analysis methods (Groffman et al., 2009; Morse et al., 2014) (Table 2) to elucidate the differences in microbial biomass and activity between different soils and the clays. In a comparison with forest, agricultural, and grassland soils in the Baltimore, Maryland metropolitan area, respiration $4.1 \mu\text{g C g}^{-1} \text{d}^{-1}$; microbial biomass C $114 \mu\text{g C g}^{-1}$; ratio 28) was closest to the agricultural soil (respiration $6.7 \text{ mg C kg}^{-1} \text{d}^{-1}$; microbial biomass C 224 mg C kg^{-1} ; ratio 33) (Groffman et al., 2009) and lower than the forest or grassland soils.

We further compared our results to soils in various spodic hydropedologic settings in a northern hardwood forest at the Hubbard Brook Experimental Forest, New Hampshire (Morse et al., 2014). These settings included typical Podzols (T), bimodal Podzols (Bi), Bh Podzols (Bh), and seeps. Three depths (Oi/Oe, Oa/A, B horizons) were sampled. Common features of Spodosols are complexation with aluminum (Al) and Fe (often organic complexation) and enrichment with Fe and Mn (Van Ranst et al., 2018), thus sharing some elemental features with our clay samples since Al is a major constituent of clay and pXRF scans confirmed the presence of Mn and Fe in our samples.

Our samples, all of which were collected from the surface, were most comparable to the B horizons and to the Bi setting from Morse et al. (2014) (Table 2). B horizons in the Bi setting soils had the lowest total C and microbial biomass C and N contents of the Hubbard Brook soils but were still much higher than our clay values. The clay had a higher microbial

biomass C/total C ratio (0.014) than the surface (0.012) and B (0.005) horizons at Hubbard Brook suggesting relatively high OM quality in the clays.

Morse et al. (2014), found that differences in total C and N content were more notable across soil horizons than across hydropedologic setting, where the B horizon, which accumulated Al and Fe, had less C, N, microbial biomass, and microbial activity. The dominance of chemical (OM quality) versus location controls is consistent with the stronger differences that we observed with color than with location. The composition of OM in clay fractions differs from that in sand and silt fractions, and there is usually a decrease in C/N ratio as particle size decreases from coarse silt to fine clay (Laird et al., 2008). In clay, the packed textures dominated by smaller particle size contain more OM than larger particle sandy samples, but the C/N ratio increases with smaller particle size. In our samples, dark gray clays had the highest total N, total C, and microbial biomass N; microbial biomass C was highest in both dark gray and brown clays. White clays had the lowest contents of these variables, indicating that inclusions (such as Fe, BC, and others) in the clays play a role in increasing their capacity to accommodate rich C and N contents.

The second question our study addressed is whether the clays support an active N cycle. We measured several indices of N cycling activity that showed that the exposed clays do indeed support an active N cycle and have potential to supply N to support plant growth. Microbial biomass N is an index of the size of the actively cycling labile N pool in soil. Mineralization, the production of simple, soluble, inorganic forms of N that are a dominant source of N for plant growth, is strongly tied to the C cycle (Hart et al., 1994). When microbes degrade N containing compounds during mineralization, their N is converted to proteins which release ammonia (NH_3) that converts to NH_4^+ , and their C is converted to biomass or CO_2 (Groffman et al., 2021).

Our findings have general similarities to those of Morse et al. (2014), who found that clay rich B horizons have lower rates of biogeochemical activity (lower net N mineralization and net nitrification potentials) as well as smaller C and N pools than surface soils with less clay and Al. Clay minerals play a role in the C and N adsorption and stabilization in soil. The storage potential of C is influenced by the size of the silicate mineral's surface area and the amount of cations adhering to these minerals (Kahle et al., 2002). Adsorption rates and amounts of OM on mineral surfaces are influenced by variations in aqueous solution dynamics, mineralogy, and OM chemistry, while OM affects mineral growth, transformation, and dissolution (Kleber et al., 2021).

The third question we asked is whether the clays are a non-anthropogenic source of reactive N. There is great concern about N pollution of groundwater and coastal waters in our region of study (Karamouz et al., 2020). There is particular concern about NO_3^- as it is the most highly mobile form of reactive N that is a drinking water pollutant and a prime cause

Table 2. Comparison of response variables from this study to data from Groffman et al. (2009) and Morse et al. (2014).

Groffman et al. (2009) (Table 3)		Total C	Respiration	MBC	MBN	NH ₄ ⁺	NH ₄ ⁺	NO ₃ ⁻	NO ₃ ⁻	TIN	TIN	Ratio of	Ratio of
		mg C kg ⁻¹	mg C kg ⁻¹ d ⁻¹	mg C kg ⁻¹	mg N g ⁻¹	mg N kg ⁻¹	mg N g ⁻¹	mg N kg ⁻¹	mg N g ⁻¹	mg N g ⁻¹	mg N g ⁻¹	MBC/total C	MBC/respiration
Forest mean Agriculture mean Grassland mean			10.3	346		2.1	0.9	0.4	8.7	2.5			33.592
			6.7	224		0.9	0.5	1.2		9.6			35.433
			8.2	306						1.7			37.317
Morse et al. (2014) (Table 1)													
Hydropedologic setting	Horizon	Total C mg C kg ⁻¹		MBC µg C g ⁻¹	MBN µg N g ⁻¹	NH ₄ ⁺ µg N g ⁻¹	NO ₃ ⁻ µg N g ⁻¹	TIN µg N g ⁻¹	ratio of MBC/Total C				
T: typical Podzols	O ₁ /O _e	506 000		4780	774	152	16.8	168.8	0.009				
	O _a /A	329 000		2320	320	30.4	18.3	48.7	0.007				
	mean of surface O ₁ /O _e , O _a /A	417 500		3550	547	91.2	17.55	108.75	0.008				
Bt: bimodal Podzols	B > 10 cm	74 000		481	32.4	2.4	3.33	5.73	0.007				
	O ₁ /O _e	378 000		6210	511	83.7	9.6	93.3	0.016				
	O _a /A	158 000		1300	226	6.7	9.97	16.67	0.008				
Bh: Bh Podzols	mean of surface O ₁ /O _e , O _a /A	268 000		3755	368.5	45.2	9.785	54.985	0.012				
	B > 10 cm	56 000		275	21.6	2.12	0.53	2.65	0.005				
Scepp	O ₁ /O _e	443 000		8230	710	144	17.8	161.8	0.019				
	O _a /A	234 000		3310	333	4.93	15.9	20.83	0.014				
	mean of surface O ₁ /O _e , O _a /A	338 500		5770	521.5	74.465	16.85	91.315	0.016				
This study	B > 10 cm	60 000		569	36	1.46	2.67	4.13	0.009				
	O _a /A	232 000		5060	238	9.18	1.02	10.2	0.022				
	Total C mg C kg ⁻¹		µg C g ⁻¹ d ⁻¹	MBC µg C g ⁻¹	MBN µg N g ⁻¹	NH ₄ ⁺ µg N g ⁻¹	NO ₃ ⁻ µg N g ⁻¹	TIN µg N g ⁻¹	ratio of MBC/Total C	ratio of MBC/respiration			
All locations, surface, mean		8220.18	4.1	113.53	7.64	4.63	1.47	6.11	0.014	27.690			

Abbreviations – HPS: hydropedologic setting, T: typical Podzols, Bt: bimodal Podzols, Bh: Bh Podzols; MBC: microbial biomass C, MBN: microbial biomass N.

of eutrophication in coastal waters (Conley et al., 2009). We therefore assessed the potential of these clays to contribute to high levels of NO_3^- in the environment by measuring both NO_3^- pools as well as processes that produce (nitrification) and consume (denitrification) NO_3^- .

Nitrification is carried out by chemoautotrophic bacteria that oxidize NH_4^+ into NO_2^- which is further oxidized into NO_3^- . When this process is stimulated by the addition of N through application of fertilizers, atmospheric deposition, groundwater and runoff sources, it can lead to excessive production of NO_3^- (Groffman et al., 2021). Denitrification is an anaerobic process that converts NO_3^- to gaseous forms NO , N_2O , and N_2 (Robertson and Groffman, 2015), removing reactive N from the soil and facilitating the cycling of N between the biosphere and lithosphere to the atmosphere.

5.3 N pollution due to reactive N species

N pollution occurs in the air, soil, and water. In the air it is the addition of volatile forms of N gases, such as NO_y , caused by industrial and commercial activity; in soil there is often an overload of nutrients from overuse of fertilizers and deposition from the atmosphere; in water a major cause is sewage system drainage and agricultural runoff, causing eutrophication which leads to dead zones in the oceans in the most extreme scenario. The main source for N pollution (in the form of NO_3^-) to Long Island's aquifers is from septic systems, with additional inputs from agriculture, lawn care, and atmospheric deposition (Szymczycha et al., 2017). Nitrate fluxes in Long Island's aquifers can negatively impact coastal ecosystems (Karamouz et al., 2020) and drinking water quality. The clay units included in Long Island's aquifer and aquitard strata can act as confining units, and as sinks or non-anthropogenic sources of reactive N.

The clays examined in this study could act as a minor source or a potentially significant sink for reactive N in an aquifer system. Clays with significant rates of potential net nitrification (especially the light gray clay) could be a source of NO_3^- if they were in contact with the aquifer. However, rates of denitrification potential were generally much higher than potential net nitrification, suggesting that these clay materials are more likely to act as NO_3^- sinks. The light gray clays have the potential to act as both NO_3^- sources and sinks depending on environmental conditions, while the white and Fe-bearing (red, yellow) clays have the potential to act as NO_3^- sinks.

6 Conclusions

This investigation shed light on the amount and type of microbial activity that occurs in geological microhabitats in the Critical Zone at a coastal exposure of temperate northern latitude and allowed us to evaluate their potential local and regional impact.

Our analyses have taken a step towards better understanding the nature and extent of the biogeochemical activity that these types of microhabitats support. The approach of using laboratory measurements of microbial biomass and activity in ancient materials was successful in characterizing the biogeochemical potential of these materials, even at low levels, and could be applied in other Critical Zone studies.

The results from this study provide quantitative data showing microbial activity of silt and clay fraction materials of hydrogeologic origin, and confirm that these materials contain ecologically significant concentrations of geologic N exceeding $1000 \text{ mg N kg}^{-1}$ (Holloway and Dahlgren, 2002). Their surface exposure allowed us to explore the interaction between ancient geological stratigraphic components and modern day environmental conditions. This type of interaction is illustrative of the conceptual reach of Critical Zone science and the cohesive understanding of multiple differing factors that it provides. The unity of various disciplines and their individual approaches to investigation allow for greater understanding of the ensuing conditions that occur in the Critical Zone, where the resulting processes work together to support all living organisms.

Our results advance the emerging science of the geological N cycle and clearly show that ancient geological materials are contributing to contemporary biogeochemical processes in the Critical Zone of our study region. Further, we have shown that these materials support a wide range of N cycle processes encompassing mineralization, immobilization, nitrification, and denitrification. There is a clear need for further research on how clay physical and chemical characteristics influence the flows of N to and from clays and biogeochemical processes in the Critical Zone, and also to see how these materials and processes are contributing to the growth of vegetation and the dynamics of pollutants in the Critical Zone of this dynamic, densely populated, and environmentally sensitive region.

Data availability. All the data referred to in the text are included in the figures, tables, and figures and tables in the Supplement.

Supplement. The supplement related to this article is available online at <https://doi.org/10.5194/bg-22-3357-2025-supplement>.

Author contributions. VMA conceptualized the study; performed fieldwork, sample collection, laboratory analyses, and data analysis; and wrote and revised the manuscript. PMG designed the study; provided laboratory methodology, access, and supplies; interpreted results; and co-wrote and revised the manuscript. ZC provided laboratory equipment and access, interpreted results, and revised and edited the manuscript. DES interpreted results and revised and edited the manuscript.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

Acknowledgements. The authors thank Clare Kohler and Kaitlin McLaughlin for help with laboratory analyses and thank Veronica Natale and Herbert Mills for help obtaining samples.

Review statement. This paper was edited by Paul Stoy and reviewed by three anonymous referees.

References

- Bennett, P. C., Rogers, J. R., Chol, W. J., and Hiebert, F. K.: Silicates, Silicate Weathering, and Microbial Ecology, *Geomicrobiol. J.*, 18, 3–19, <https://doi.org/10.1080/01490450151079734>, 2001.
- Bohlen, P. J., Groffman, P. M., Driscoll, C. T., Fahey, T. J., and Siccama, T. G.: PLANT–SOIL–MICROBIAL INTERACTIONS IN A NORTHERN HARDWOOD FOREST, *Ecology*, 82, 965–978, [https://doi.org/10.1890/0012-9658\(2001\)082\[0965:PSMIIA\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2001)082[0965:PSMIIA]2.0.CO;2), 2001.
- Brantley, S. L., White, T. S., White, A. F., Sparks, D. L., Richter, D., Pregitzer, K. S., Derry, L. A., Chorover, J., Chadwick, O. A., April, R., Anderson, S. P., and Amundson, R. C.: Frontiers in exploration of the critical zone, Re-port of a Workshop Sponsored by the National Science Foundation(NSF), Newark, DE, 30, 2006.
- Brown, C. J. and Schoonen, M. A. A.: The origin of high sulfate concentrations in a coastal plain aquifer, Long Island, New York, *Appl. Geochem.*, 19, 343–358, [https://doi.org/10.1016/S0883-2927\(03\)00154-9](https://doi.org/10.1016/S0883-2927(03)00154-9), 2004.
- Busigny, V. and Bebout, G. E.: Nitrogen in the Silicate Earth: Speciation and Isotopic Behavior during Mineral-Fluid Interactions, *Elements*, 9, 353–358, <https://doi.org/10.2113/gselements.9.5.353>, 2013.
- Conley, D. J., Paerl, H. W., Howarth, R. W., Boesch, D. F., Seitzinger, S. P., Havens, K. E., Lancelot, C., and Likens, G. E.: Controlling Eutrophication: Nitrogen and Phosphorus, *Science*, 323, 1014–1015, <https://doi.org/10.1126/science.1167755>, 2009.
- Dahlgren, R. A.: Soil acidification and nitrogen saturation from weathering of ammonium-bearing rock, *Nature*, 368, 838–841, <https://doi.org/10.1038/368838a0>, 1994.
- Dass, P., Houlton, B. Z., Wang, Y., Wårlind, D., and Morford, S.: Bedrock Weathering Controls on Terrestrial Carbon-Nitrogen-Climate Interactions, *Glob. Biogeochem. Cycles*, 35, e2020GB006933, <https://doi.org/10.1029/2020GB006933>, 2021.
- Davey, B. G., Russell, J. D., and Wilson, M. J.: Iron oxide and clay minerals and their relation to colours of red and yellow podzolic soils near Sydney, Australia, *Geoderma*, 14, 125–138, [https://doi.org/10.1016/0016-7061\(75\)90071-3](https://doi.org/10.1016/0016-7061(75)90071-3), 1975.
- Doane, T. A. and Horwath, W. R.: Spectrophotometric Determination of Nitrate with a Single Reagent, *Anal. Lett.*, 36, 2713–2722, <https://doi.org/10.1081/AL-120024647>, 2003.
- Eby, N.: Principles of environmental geochemistry, Waveland Press, Inc, Long Grove, Illinois, 514 pp., ISBN 10: 1-4786-3164-3, ISBN 13: 978-1-4786-3164-4, 2016.
- Fuller, M.: The Geology of Long Island, New York, *J. Geol.*, 24, 303–304, <https://doi.org/10.1086/622335>, 1914.
- Galloway, J. N., Dentener, F. J., Capone, D. G., Boyer, E. W., Howarth, R. W., Seitzinger, S. P., Asner, G. P., Cleveland, C. C., Green, P. A., Holland, E. A., Karl, D. M., Michaels, A. F., Porter, J. H., Townsend, A. R., and Vorismarty, C. J.: Nitrogen Cycles: Past, Present, and Future, *Biogeochemistry*, 70, 153–226, <https://doi.org/10.1007/s10533-004-0370-0>, 2004.
- Groffman, P. M., Holland, E. A., Myrold, D. D., Robertson, G. P., and Zou, X.: Denitrification, in: Standard soil methods for long-term ecological research, edited by: Robertson, G.P., Coleman, D.C., Bledsoe, C. S., and Sollins, P., Oxford University Press, New York, 272–288, <https://doi.org/10.1093/oso/9780195120837.003.0014>, 1999.
- Groffman, P. M., Williams, C. O., Pouyat, R. V., Band, L. E., and Yesilonis, I. D.: Nitrate Leaching and Nitrous Oxide Flux in Urban Forests and Grasslands, *J. Environ. Qual.*, 38, 1848–1860, <https://doi.org/10.2134/jeq2008.0521>, 2009.
- Groffman, P. M., Rosi, E. J., and Fulweiler, R. W.: The nitrogen cycle, in: Fundamentals of Ecosystem Science, edited by: Weathers, K. C., Strayer, D. L., and Likens, G. E., Academic Press, 161–188, ISBN 978-0-12-088774-3, 2021.
- Halama, R. and Bebout, G.: Earth's Nitrogen and Carbon Cycles, *Space Sci. Rev.*, 217, 45, <https://doi.org/10.1007/s11214-021-00826-7>, 2021.
- Hart, S. C., Nason, G. E., Myrold, D. D., and Perry, D. A.: Dynamics of Gross Nitrogen Transformations in an Old-Growth Forest: The Carbon Connection, *Ecology*, 75, 880–891, <https://doi.org/10.2307/1939413>, 1994.
- Holloway, J. M. and Dahlgren, R. A.: Nitrogen in rock: Occurrences and biogeochemical implications: BIOGEOCHEMICAL IMPLICATIONS OF N IN ROCK, *Glob. Biogeochem. Cycles*, 16, 1118, <https://doi.org/10.1029/2002GB001862>, 2002.
- Houlton, B. Z., Morford, S. L., and Dahlgren, R. A.: Convergent evidence for widespread rock nitrogen sources in Earth's surface environment, *Science*, 360, 58–62, <https://doi.org/10.1126/science.aan4399>, 2018.
- Isbister, J.: Geology and hydrology of northeastern Nassau County, Long Island, New York, US Government Printing Office, <https://doi.org/10.3133/wsp1825>, 1966.
- Jakobsson, M., Løvlie, R., Al-Hanbali, H., Arnold, E., Backman, J., and Möhrth, M.: Manganese and color cycles in Arctic Ocean sediments constrain Pleistocene chronology, *Geology*, 28, 23, [https://doi.org/10.1130/0091-7613\(2000\)28<23:MACCIA>2.0.CO;2](https://doi.org/10.1130/0091-7613(2000)28<23:MACCIA>2.0.CO;2), 2000.
- Jenkinson, D. S. and Powlson, D. S.: The effects of biocidal treatments on metabolism in soil – V, *Soil Biol. Biochem.*, 8, 209–213, [https://doi.org/10.1016/0038-0717\(76\)90005-5](https://doi.org/10.1016/0038-0717(76)90005-5), 1976.

- Kahle, M., Kleber, M., and Jahn, R.: Review of XRD-based quantitative analyses of clay minerals in soils: the suitability of mineral intensity factors, *Geoderma*, 109, 191–205, [https://doi.org/10.1016/S0016-7061\(02\)00175-1](https://doi.org/10.1016/S0016-7061(02)00175-1), 2002.
- Karamouz, M., Mahmoodzadeh, D., and Oude Essink, G. H. P.: A risk-based groundwater modeling framework in coastal aquifers: a case study on Long Island, New York, USA, *Hydrogeol. J.*, 28, 2519–2541, <https://doi.org/10.1007/s10040-020-02197-9>, 2020.
- Kleber, M., Bourg, I. C., Coward, E. K., Hansel, C. M., Myneni, S. C. B., and Nunan, N.: Dynamic interactions at the mineral–organic matter interface, *Nat. Rev. Earth Environ.*, 2, 402–421, <https://doi.org/10.1038/s43017-021-00162-y>, 2021.
- Laird, D. A., Chappell, M. A., Martens, D. A., Wershaw, R. L., and Thompson, M.: Distinguishing black carbon from biogenic humic substances in soil clay fractions, *Geoderma*, 143, 115–122, <https://doi.org/10.1016/j.geoderma.2007.10.025>, 2008.
- Li, L., Shields, J., Snow, D. D., Kaiser, M., and Malakar, A.: Labile carbon and soil texture control nitrogen transformation in deep vadose zone, *Sci. Total Environ.*, 878, 163075, <https://doi.org/10.1016/j.scitotenv.2023.163075>, 2023.
- Li, Y. H.: Distribution patterns of the elements in the ocean: A synthesis, *Geochim. Cosmochim. Acta*, 55, 3223–3240, [https://doi.org/10.1016/0016-7037\(91\)90485-N](https://doi.org/10.1016/0016-7037(91)90485-N), 1991.
- Liebling, R. S.: Clay Minerals of the Weathered Bedrock Underlying Coastal New York, *Geol. Soc. Am. Bull.*, 84, 1813, [https://doi.org/10.1130/0016-7606\(1973\)84<1813:CMOTWB>2.0.CO;2](https://doi.org/10.1130/0016-7606(1973)84<1813:CMOTWB>2.0.CO;2), 1973.
- Mills, H. C. and Wells, P. D.: Ice-Shove Deformation and Glacial Stratigraphy of Port Washington, Long Island, New York, *Geol. Soc. Am. Bull.*, 85, 357, [https://doi.org/10.1130/0016-7606\(1974\)85<357:IDAGSO>2.0.CO;2](https://doi.org/10.1130/0016-7606(1974)85<357:IDAGSO>2.0.CO;2), 1974.
- Morford, S. L., Houlton, B. Z., and Dahlgren, R. A.: Increased forest ecosystem carbon and nitrogen storage from nitrogen rich bedrock, *Nature*, 477, 78–81, <https://doi.org/10.1038/nature10415>, 2011.
- Morse, J. L., Werner, S. F., Gillin, C. P., Goodale, C. L., Bailey, S. W., McGuire, K. J., and Groffman, P. M.: Searching for biogeochemical hot spots in three dimensions: Soil C and N cycling in hydropedologic settings in a northern hardwood forest: Biogeochemical hotspots in soil profiles, *J. Geophys. Res.-Biogeosci.*, 119, 1596–1607, <https://doi.org/10.1002/2013JG002589>, 2014.
- Paul, E. A.: Soil microbiology, ecology, and biochemistry, Fourth Edn., Academic Press, Amsterdam, ISBN 9780123914118, 2014.
- Pihlatie, M., Syväsalö, E., Simojoki, A., Esala, M., and Regina, K.: Contribution of nitrification and denitrification to N₂O production in peat, clay and loamy sand soils under different soil moisture conditions, *Nutr. Cycl. Agroecosystems*, 70, 135–141, <https://doi.org/10.1023/B:FRES.0000048475.81211.3c>, 2004.
- Remucal, C. K. and Ginder-Vogel, M.: A critical review of the reactivity of manganese oxides with organic contaminants, *Environ. Sci. Process. Impacts*, 16, 1247, <https://doi.org/10.1039/c3em00703k>, 2014.
- Robertson, G. P. (Ed.): Standard soil methods for long-term ecological research, Oxford University Press, New York, 462 pp., <https://doi.org/10.1093/oso/9780195120837.001.0001>, 1999.
- Robertson, G. P. and Groffman, P. M.: Nitrogen transformations., in: *Soil Microbiology, Ecology, and Biogeochemistry*, Academic Press, Burlington, MA, 421–446, <https://doi.org/10.1016/B978-0-12-415955-6.00014-1>, 2015.
- Saidy, A. R., Smernik, R. J., Baldock, J. A., Kaiser, K., and Sanderman, J.: The sorption of organic carbon onto differing clay minerals in the presence and absence of hydrous iron oxide, *Geoderma*, 209–210, 15–21, <https://doi.org/10.1016/j.geoderma.2013.05.026>, 2013.
- Schlesinger, W. H.: An estimate of the global sink for nitrous oxide in soils, *Glob. Change Biol.*, 19, 2929–2931, <https://doi.org/10.1111/gcb.12239>, 2013.
- Schroeder, P. A.: Clays in the Critical Zone, Cambridge University Press, Cambridge New York, 178–179, <https://doi.org/10.1017/9781316480083>, 2018.
- Seitzinger, S., Harrison, J. A., Böhlke, J. K., Bouwman, A. F., Lowrance, R., Peterson, B., Tobias, C., and Drecht, G. V.: DENITRIFICATION ACROSS LANDSCAPES AND WATERSCAPES: A SYNTHESIS, *Ecol. Appl.*, 16, 2064–2090, [https://doi.org/10.1890/1051-0761\(2006\)016\[2064:DALAWA\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2006)016[2064:DALAWA]2.0.CO;2), 2006.
- Sims, G. K., Ellsworth, T. R., and Mulvaney, R. L.: Microscale determination of inorganic nitrogen in water and soil extracts, *Commun. Soil Sci. Plant Anal.*, 26, 303–316, <https://doi.org/10.1080/00103629509369298>, 1995.
- Sirkin, L.: Stratigraphy of the Long Island Platform, *J. Coast. Res.*, 217–227, <http://www.jstor.org/stable/25735581> (last access: 12 April 2023), 1991.
- Six, J., Elliott, E. T., and Paustian, K.: Soil Structure and Soil Organic Matter II. A Normalized Stability Index and the Effect of Mineralogy, *Soil Sci. Soc. Am. J.*, 64, 1042–1049, <https://doi.org/10.2136/sssaj2000.6431042x>, 2000.
- Smith, M. S. and Tiedje, J. M.: Phases of denitrification following oxygen depletion in soil, *Soil Biol. Biochem.*, 11, 261–267, [https://doi.org/10.1016/0038-0717\(79\)90071-3](https://doi.org/10.1016/0038-0717(79)90071-3), 1979.
- Swarzenski, W. V.: Hydrogeology of Northwestern Nassau and Northeastern Queens counties, Long Island, New York, USGS Geological Survey Water Supply Paper 1657, 88 pp., <https://doi.org/10.3133/wsp1657>, 1963.
- Szymczycha, B., Kroeger, K. D., Crusius, J., and Bratton, J. F.: Depth of the vadose zone controls aquifer biogeochemical conditions and extent of anthropogenic nitrogen removal, *Water Res.*, 123, 794–801, <https://doi.org/10.1016/j.watres.2017.06.048>, 2017.
- Van Ranst, E., Wilson, M. A., and Righi, D.: Spodic Materials, in: *Interpretation of Micromorphological Features of Soils and Regoliths*, Elsevier, 633–662, <https://doi.org/10.1016/B978-0-444-63522-8.00022-X>, 2018.