



# Extraordinary bloom of toxin-producing phytoplankton enhanced by strong retention on the offshore Patagonian shelf

Valeria Ana Guinder<sup>1</sup>, Urban Tillmann<sup>2</sup>, Martin Rivarossa<sup>3</sup>, Carola Ferronato<sup>1</sup>, Fernando J. Ramírez<sup>1</sup>, Bernd Krock<sup>2</sup>, Haifeng Gu<sup>4</sup>, and Martin Saraceno<sup>5,6,7</sup>

<sup>1</sup>Instituto Argentino de Oceanografía (IADO), Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Universidad Nacional del Sur (UNS), Camino La Carrindanga km 7.5, 8000 Bahía Blanca, Argentina

<sup>2</sup>Alfred-Wegener-Institut (AMI), Helmholtz Zentrum für Polar- und Meeresforschung, Am Handelshafen 12, 27570 Bremerhaven, Germany

<sup>3</sup>Instituto Nacional de Investigación y Desarrollo Pesquero (INIDEP), and CONICET, Paseo Victoria Ocampo, Escollera Norte 1, B7602HSA Mar del Plata, Buenos Aires, Argentina

<sup>4</sup>Third Institute of Oceanography (TIO), Ministry of Natural Resources, Xiamen, 361005, China

<sup>5</sup>Centro de Investigaciones del Mar y de la Atmósfera (CIMA), CONICET, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires (UBA), C1063ACV, Buenos Aires, Argentina

<sup>6</sup>Instituto Franco-Argentino de Estudio sobre el Clima y sus Impactos (IFAECI) CNRS-IRD-UBA-CONICET, Buenos Aires, Argentina

<sup>7</sup>Departamento de Ciencias del Océano y de la Atmósfera (DCAO), Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires (UBA), C1063ACV, Buenos Aires, Argentina

**Correspondence:** Valeria Ana Guinder (vguinder@iado-conicet.gob.ar)

Received: 11 October 2024 – Discussion started: 23 October 2024

Revised: 12 April 2025 – Accepted: 15 April 2025 – Published: 11 July 2025

**Abstract.** The extensive Patagonian continental shelf in the Atlantic Ocean is renowned for its high productivity associated with nutrient-rich waters that fertilize massive phytoplankton blooms, especially along the shelf-break frontal system. Growing evidence reflects this ecosystem as a hotspot for harmful algal blooms (HABs). Whether these HABs reach coastal areas or are exported to the adjacent ocean basin by energetic edge currents remains unexplored. During two oceanographic cruises in spring 2021, a bloom of dinoflagellates of the Amphidomataceae family was sampled over the outer shelf with a 10 d interval, at stations 40 km apart. The bloom was first sampled on 16 November, with  $32 \times 10^6$  cells  $L^{-1}$ , and was still persistent on 25 November, with  $14 \times 10^6$  cells  $L^{-1}$ . The magnitude of this bloom is a global record for this group so far reported in the literature. The toxin azaspiracid-2 (AZA-2) was detected in both stages of the bloom, with values up to 2122 pg  $L^{-1}$ . The most likely source of AZA-2 was *Azadinium spinosum* ribotype B. The bloom developed in vertically stable waters (60 m mixed layer depth) with elevated chlorophyll concentration. Water

retention and the presence of fronts induced by horizontal stirring controlled the persistence and trajectory of the bloom in a localized area over the continental shelf, as evidenced by analysis of geostrophic surface currents, Lyapunov coefficients, and particle advection modelling. These findings underscore the importance of monitoring HABs in offshore environments and the need to understand biophysical interactions that govern bloom taxa assemblages and transport pathways.

## 1 Introduction

In marine environments, dinoflagellates are the primary toxin-producing group of protistan plankton and key causative agents of harmful algal blooms (HABs). As the most diverse group of toxic microorganisms, e.g. *Alexandrium* spp., *Karenia* spp., *Dinophysis* spp., *Azadinium* spp., and *Amphidoma* spp., dinoflagellates produce a wide range of toxins. Phycotoxins are natural intracellular metabolites

synthesized by certain microalgae that can be transferred through the food web, having severe impacts on marine biota, ecosystems, and human health (Anderson et al., 2015; Sunesen et al., 2021). Broadly, HABs were long thought to occur exclusively in coastal regions due to their visible impacts on water quality and human-related activities, as documented for instance in the Argentine Patagonian gulfs (Wilson et al., 2015; D'Agostino et al., 2019) and the Beagle Channel (Almandoz et al., 2019; Cadaillon et al., 2024). However, the perception of HABs as solely coastal events was biased, primarily due to greater monitoring efforts in coastal areas compared to the fewer studies conducted offshore (Hallegraeff et al., 2021; Sunesen et al., 2021; Anderson et al., 2021). In line with this trend, the expansion of the monitored area in the Argentine Sea from the coast to the outer continental shelf over recent decades has confirmed that toxic species are indeed common in offshore waters (Ramírez et al., 2022), especially associated with the shelf-break front (reviewed in Guinder et al., 2024). It is well known that dinoflagellates possess advantageous strategies for thriving in frontal systems, such as effective swimming, mucus and cyst formation, mixotrophy, and toxin production (Smayda, 2002; Glibert, 2016). However, the biophysical mechanisms explaining the development of large harmful blooms on hydrographically complex shelves are still not fully understood, mainly due to the lack of simultaneous taxonomic data and velocity fields at synoptic scales.

In oceanic waters, the permanence and spatial extent of discrete plankton blooms are influenced by dispersal mechanisms that rely on diffusion and horizontal advection (Abraham et al., 2000; Mahadevan, 2016; Lehahn et al., 2017). Typically, the dispersion and stirring of phytoplankton blooms in the ocean have been studied using the remote sensing of chlorophyll *a* and models (Lehahn et al., 2007; Lévy et al., 2018; Ser-Giacomi et al., 2023), with few studies considering in situ sampling (Abraham et al., 2000; Giddings et al., 2014; Hernández-Carrasco et al., 2020) to assess the biophysical couplings of bloom development and persistence. In addition to the considerable complexity of studying the dynamics of phytoplankton blooms in the ocean, which involve multiple physical (temperature, salinity, and mixing of water masses), biological (species competition, grazing), and chemical (nutrients) factors, monitoring HABs presents an even greater challenge (Pitcher et al., 2010; Wells et al., 2020; Iriarte et al., 2023). This includes the identification of the species involved in these blooms. It is precisely in this area where detailed studies, employing multiple approaches such as microscopy, molecular techniques, and toxin profiling (Smayda, 2002; Wells et al., 2020), make a significant contribution to understanding HABs' development and potential impacts.

The extensive Patagonian shelf-break front (35–55° S) in the SW Atlantic Ocean is a high-productivity ecosystem, located ~200 to ~900 km offshore (Martinetto et al., 2019; Guinder et al., 2024). This permanent thermohaline front is

associated with the upwelling of nutrient-rich waters of the westerly edge of the Malvinas Current, which fertilizes the surface waters (Palma et al., 2008; Matano et al., 2010). Massive phytoplankton proliferations occur over the outer shelf in spring and summer (García et al., 2008; Carreto et al., 2016; Ferronato et al., 2023), including a variety of HAB-forming taxa and associated phycotoxins (Ramírez et al., 2022; Guinder et al., 2024). In fact, the increase in oceanographic studies focused on the detection of HABs along the outer Patagonian shelf has led to several new records of toxin-producing species and phycotoxins in the South Atlantic (Akselman et al., 2015; Guinder et al., 2018; Tillmann et al., 2019). The most conspicuous HABs are those formed by the nano-dinoflagellates of the Amphidomataceae family (Akselman and Negri, 2012; Tillmann et al., 2018; 2019; Fabro et al., 2019; Guinder et al., 2020), producers of toxic azaspiracids (AZAs). These large HABs have emerged as important hazards in the productive Patagonian shelf-break frontal ecosystem. Amphidomataceans include four AZA-producing species, i.e. *Azadinium dexteroporum*, *Az. poporum*, *Az. spinosum*, and *Amphidoma languida* (Krock et al., 2019) and have been reported from different marine regions globally (Tillmann, 2018a; Salas et al., 2021; Liu et al., 2023). So far, the maximum bloom abundances reported in the literature are from the Argentine Sea (Akselman and Negri, 2012). During the springs of 1991 and 1992, these dinoflagellates reached between 3 and  $9 \times 10^6$  cells L<sup>-1</sup> and caused water discolouration in the northern area (38–42° S; 58–56° W) of the Patagonian shelf (Akselman and Negri, 2012). No toxin screening was performed at that time, but in spring 2015, an AZA-producing *Azadinium* strain was isolated from another large bloom in the area (Tillmann et al., 2019). Moreover, AZAs have been detected in the tissue of the scallop *Zygochlamys patagonica* since the early 1990s (Turner and Goya, 2015). These scallops form large seabed banks along the 100 m isobath between 38 and 48° S (Alemany et al., 2024), associated with the high phytoplankton productivity over the outer Patagonian shelf.

Despite the limited synoptic sampling in offshore waters, the prevalence of HABs in the Patagonian front highlights this ecosystem as a hotspot that requires further monitoring. The notably high abundance of Amphidomataceans over the outer shelf holds greater significance when assessing the potential risks that a bloom dissemination may pose to both regional and global ecosystems. The development and transport of HABs remain poorly understood, as does the question of whether they may reach coastal areas or be exported offshore into the stirring Atlantic Ocean. The accumulation and dispersion of phytoplankton patches in complex shelf waters need to be addressed from an interdisciplinary perspective, combining remote sensing, modelling, and in situ sampling (Ferronato et al., 2025). In a recent study conducted on the outer Patagonian shelf – the core of the Malvinas Current and adjacent open-ocean waters – we emphasized the key role of mesoscale energetic variability in modulating phytoplankton



spring blooms (Ferronato et al., 2025). However, these observational studies combining multiple approaches at synoptic scale remain scarce in the region.

In this study, we characterized the biophysical aspects of a large multispecific spring bloom of Amphidomataceans, detected through an unusual sampling effort that involved two research expeditions in November 2021. This HAB was observed at two sampling sites 40 km apart within a span of 10 d. We combined field observations of protistan plankton species composition and associated toxins with remotely sensed ocean colour images of chlorophyll *a* and geostrophic surface currents, particle tracking experiments, and Lyapunov coefficient analysis to assess the horizontal displacement and retention of the Amphidomataceae bloom within a mesoscale eddy. Furthermore, we aim to explore whether this HAB that developed in offshore shelf waters might reach coastal areas or be advected by the Malvinas Current, facilitating the dispersal of toxic species to other shelves and ocean basins.

## 2 Materials and methods

### 2.1 Hydrography and productivity in the Patagonian continental shelf

Along the external margin of the Patagonian continental shelf, between 35 and 55° S, a thermohaline front develops throughout the year, characterized by high biological productivity. The development of this front and the associated upwelling is due to the interaction of the energetic western boundary current system with the steep slope and the waters of the shelf, as well as the effect of winds and tides (reviewed in Piola et al., 2024). The Malvinas Current originates at ~55° S as a branch of the Antarctic Circumpolar Current (Fig. 1) and runs northwards at high velocity (mean surface velocities of 45 cm s<sup>-1</sup>, Piola et al., 2024) along the shelf break in two jets that meet at ~44° S (Frey et al., 2023). Then, at ~38° S, the Malvinas Current meets the warm and oligotrophic Brazil Current, which runs southwards in the so-called Brazil–Malvinas Confluence, and waters are exported eastwards into the South Atlantic Ocean basin (Fig. 1). In addition, another branch of the Antarctic Circumpolar Current gives origin to the Patagonian Current which runs northwards over the continental shelf, carrying diluted subantarctic waters (Fig. 1).

The interaction of the Malvinas Current with the irregular bottom topography generates the upwelling of cold, nutrient-rich waters that fertilize phytoplankton over the shelf, together with the Patagonian Current (reviewed in Guinder et al., 2024). Phytoplankton blooms expand over the mid and outer shelf as reflected by a persistent satellite chlorophyll *a* band, wider in spring and narrower in summer along the shelf-beak front (Guinder et al., 2024). The magnitude of the upwelling has low seasonal variability and is heterogeneous


along the extensive latitudinal range of the slope (Combes and Matano, 2018). Hence, productivity over the shelf varies spatially and temporally, and in consequence, multiple bioregions emerge, each characterized by unique phytoplankton phenological patterns, as revealed by climatological analysis of satellite-derived chlorophyll *a* (Delgado et al., 2023).

### 2.2 Research cruises

Two oceanographic expeditions were carried out during late spring (November 2021) to study microbial plankton communities on the Argentine continental shelf and adjacent ocean basin (Fig. 1). The first research cruise, Ana María Gayoso (hereafter Gayoso, GA), took place aboard the R/V *Bernardo Houssay* (operated by PNA and CONICET, Argentina). This cruise was planned so as to analyse the pre-bloom conditions of the coccolithophore *Gephyrocapsa huxleyi*, which blooms in early summer (December) on the Patagonian shelf. The location of the sampling stations was delineated by tracking daily satellite-derived signals of chlorophyll *a* and particulate inorganic carbon (see Ferronato et al., 2025). Sampling occurred between 16 and 22 November 2021 at 10 stations distributed along the outer Patagonian shelf – the core of the Malvinas Current and the adjacent open-ocean waters (Fig. 1). The second cruise, Agujero Azul (AA), took place aboard the R/V *Victor Angelescu* (INIDEP and CONICET, Argentina). This multidisciplinary expedition was part of a national project aimed at assessing pelagic and benthic biodiversity in a highly productive area known as the Agujero Azul (44–47° S, 62–57.5° W) (Fig. 1), with particular emphasis on commercially important fish species (Alemany et al., 2024). Sampling took place between 25 November and 3 December 2021 at 23 stations arranged along two cross-shelf transects in the Agujero Azul area (Fig. 1). We first present an overview of the in situ measurements (CTD) and water sampling conducted during both cruises for the analysis of plankton and nutrients. We then focus on the Amphidomataceae bloom observed over the shelf at station GA01 on 16 November and station AA09 on 25 November, located approximately 40 km apart (Fig. 1).

### 2.3 Remote sensing of surface Chl *a* and SST

In order to contextualize the discrete observations of plankton within broader spatio-temporal dynamics, we explored satellite-derived surface chlorophyll *a* concentration (Chl *a*) data from the days of the cruises. In this sense, a daily merged product with a 4 km resolution provided by the GlobColour project (distributed by ACRI ST, France: <https://hermes.acri.fr/>, last access: September 2024) were downloaded. This ocean colour product is generated from the fusion of the SeaWiFS, MERIS, MODIS-Aqua, and OLCI sensors and estimates the average Chl *a* concentration in the surface layer (Maritorena et al., 2010). The fusion of data from different satellite sensors, combined with the quality control



**Figure 1.** Sampling stations of the research cruises Gayoso (pink dots, on board R/V *Houssay*) and Agujero Azul (blue dots, on board R/V *Angelescu*) and the main circulation pattern in the Argentine continental shelf and shelf break. The stations where the bloom of Amphydomataceae was observed are indicated with a red square: GA01 and AA09, sampled on 16 and 25 November, respectively. Micrographs show the Amphydomatacean bloom and a cell of *Azadinium spinosum* ribotype B, producer of azaspiracid-2 (AZA-2). ACC: Antarctic Circumpolar Current, MC: Malvinas Current, BC: Brazil Current, BMC: Brazil–Malvinas Confluence, and PC: Patagonian Current. Bathymetry from GEBCO (2021). Isobaths of 200, 1000, and 1800 m are displayed.

criteria used by GlobColour, enables enhanced spatial and temporal coverage. Eight-day temporal averages were calculated for the periods 12–19 and 20–27 November 2021. Additionally, daily Chl *a* images were assessed from 14 to 27 November, focused on the bloom area to track its short-term evolution. Due to high cloud covering, data are missing from 22–24 November. To analyse the sea surface temperature (SST) during the sampling periods, daily NSST MODIS-Aqua level L3 images with a 4 km resolution were downloaded from <https://oceancolor.gsfc.nasa.gov/>, last access: May 2023. Eight-day temporal averages were also constructed for the periods of 12–19 and 20–27 November 2021. All images were processed using SeaDAS (version 8.3) and QGIS (version 3.38), mapped to the WGS84 reference system (datum WGS84, ellipsoid WGS84) and restricted to the study area. The images were smoothed using a “non-linear mean  $3 \times 3$ ” filter.

## 2.4 In situ measurements and sample collection

At each sampling station, continuous vertical profiles of temperature, salinity, and fluorescence were measured. In the Gayoso cruise, a Sea-Bird 9 plus CTD and a fluorometer sensor Wet Labs FLRTD-5105 were used. During the Agujero Azul cruise, a Sea-Bird SBE 9 Digiquartz CTD coupled with an ancillary SeaPoint SCF chlorophyll fluorometer was used. To assess the vertical stability of the water column, the Brunt–Väisälä buoyancy frequency ( $\text{cyc h}^{-1}$ ) was computed using the function *swN2* of the package *oce* (Kelley et al., 2022) in R statistical software. Thereafter, the mixed layer depth (MLD, in metres) was defined at the depth where the maximum value of the Brunt–Väisälä frequency was detected.

Niskin bottles attached to the CTD rosette were used to collect water samples at the surface (5 m depth) for the analysis of chlorophyll *a*, dissolved inorganic macronutrients, pro-

tistan plankton by microscopy, genetic analysis of the species diversity, and phycotoxins in field samples.

For chlorophyll *a*, a volume of 400 mL was filtered through GF/F fibre-glass filters pre-combusted at 450 °C for 4 h. A volume of 10 mL of 90 % acetone was used for pigment extraction during 24 h (4 °C) and thereafter quantified using an Agilent Cary 60 UV-Vis spectrophotometer. Concentration was estimated using the equations developed by Jeffrey and Humphrey (1975).

For inorganic nutrients, the water samples filtered through GF/F fibre-glass filters pre-combusted at 450 °C for 4 h were stored at −20 °C in alkali-rinsed (NaOH, 0.1 M) polyethylene bottles. Nitrite, nitrate, ammonium, silicate, and phosphate were measured using a spectrophotometer Agilent Cary 60 UV-Vis following standard seawater methods outlined by Hansen and Koroleff (1999). These protocols for chlorophyll and nutrient determinations have been applied and intercalibrated for other areas of the Patagonian shelf (e.g. Ferronato et al., 2025; Gilabert et al., 2025).

Duplicate samples for plankton counts collected with Niskin bottles were preserved with Lugol (1 % final concentration) and formaldehyde (1 % *f/c*) in glass bottles (250 mL) and kept in the dark and at 4 °C for their analyses under microscopy. Similarly, duplicate water samples were collected by three vertical net tows (20 µm mesh size) integrating the first 30 m depth for the identification of protists' taxa.

For the quantification of azaspiracids (AZAs) as well as for genetic analysis of species diversity, the same sampling protocol was applied. A volume of 4–5 L of seawater from the Niskin bottles was pre-screened through a 20 µm mesh-size net and subsequently filtered through 5 µm pore-size polycarbonate filters (Millipore, Eschborn, Germany) under gentle vacuum (< 200 mbar). Filters were placed in 50 mL centrifuge tubes and preserved at −80 °C for further analyses in the laboratory.

## 2.5 Plankton diversity analyses

Morphological aspects of plankton cells were carefully observed under different light microscopes all equipped with epifluorescence and differential interference contrast optics: a Nikon Eclipse E-400 microscope, a ZEISS Axioskop 2 microscope, and an inverted Axiovert 200 M. In order to measure the length and width of cells, micrographs were taken at 1000 magnification under a ZEISS Axio Vert.A1 equipped with a digital camera Axiocam 208 Color and under an Axioskop 2 equipped with an Axiovision digital camera. Thereafter, micrographs were processed with the software ZEN (version 2.7, ZEISS) and Axiovision (version 4.8, ZEISS). Further, scanning electron microscopy (SEM, FEI Quanta FEG 200) was used to assess detailed taxonomic features of the dinoflagellate species (e.g. arrangement of thecal plates, presence of pores and spines). SEM samples were treated following the protocol described in Tillmann et al. (2017a).

For the estimation of total protist abundance (in cells L<sup>−1</sup>), seawater samples collected with Niskin bottles and fixed with Lugol were settled in sedimentation chambers, and single cells were counted under inverted microscope using a magnification of 400 following traditional techniques (Hasle, 1978). All protists larger than 5 µm in cell size were counted and classified into species or genera taxonomic levels or merged into taxonomic/functional groups organized in size ranges (e.g. ciliates 10–20 µm, cryptophytes < 10 µm, *Gymnodinium*-type cell, Kareniaceae-type cell). In addition, to assess the relative abundance of the Amphidomataceae species responsible for the multispecific bloom of this clade, subsamples (10 mL) were carefully counted with high taxa resolution.

As an additional and complementary mean for analysing species diversity in the field samples, metabarcoding was performed, specifically targeting the internal transcribed spacer (ITS1) region, following Liu et al. (2023). This information was used as a complement to the exhaustive morphological taxonomy performed under light microscopy and SEM. In particular, metabarcoding also allows the specific identification of ribotypes of Amphidomatacean species. For example, this is important because for *Az. spinosum*, different ribotypes have been shown to differ in toxin production and toxin profile (Tillmann et al., 2019). A detailed rationale for the Amphidomatacean species and ribotype designation is given in Appendix B.

## 2.6 Toxin identification and quantification

Filters were repeatedly rinsed with 500 µL methanol until complete discolouration of the filters. The methanolic extracts were transferred to a spin filter (0.45 µm pore size, Millipore) and centrifuged at 800 × *g* for 30 s, followed by transfer to autosampler vials and stored at −20 °C until analysis. Toxin analyses were performed using high-performance liquid chromatography coupled with tandem mass spectrometry HPLC-MS/MS in the selected reaction monitoring (SRM) mode for the detection of known AZA variants. In addition, precursor experiments of the ions *m/z* 348, 350, 360, 362, and 378 were carried out to find potentially new AZA variants. Screened mass transitions and instrument parameters are detailed in Tillmann et al. (2021).

## 2.7 Lagrangian simulations and finite-size Lyapunov exponent analysis

To explore the physical mechanisms that might explain the concentration of Amphidomataceae measured in the two locations sampled, we used two complementary analyses: Lagrangian advection of virtual particles and finite-size Lyapunov exponents (FSLEs).

The first technique consists of the analysis of trajectories of virtual neutrally buoyant particles that were obtained with an algorithm that represents the advection process caused by

surface currents. The advection equation is as follows:

$$X(t + \Delta t) = X(t) + \int_t^{t+\Delta t} v(x, \tau) d\tau, \quad (1)$$

where  $X$  is the three-dimensional position of a particle and  $v(x, \tau)$  is the three-dimensional velocity field, integrated using a fourth-order Runge–Kutta scheme. Particles were released at the surface along 46° S and every 0.05 grades in the four regions, as indicated in Appendix D. The algorithm computes the particle positions based on initial location and knowledge of the velocity field. A time step of 1 h was considered. The accuracy of the trajectories obtained relies on the accuracy of the velocity field used. For this experiment we considered geostrophic velocities obtained from satellite altimetry. In the northern portion of the Argentine continental shelf, such surface velocities were shown to be well correlated with in situ current measurements (Lago et al., 2021). We therefore assume that the surface dynamics can be represented by satellite-altimetry-derived data and use this as the input velocity field for the algorithm to advect the virtual particles. Geostrophic velocities derived from gridded absolute dynamic topography (ADT) of daily temporal resolution and 1/4° spatial resolution maps were downloaded from CMEMS (<https://marine.copernicus.eu/>, last access: September 2023).

The second technique consists of the analysis of FSLE images with a spatial resolution of 1/25° grid that were obtained from AVISO (<https://www.aviso.altimetry.fr>, last access: September 2023). FSLE ridges approximate the so-called Lagrangian coherent structures, which are a generalization of stable hyperbolic trajectories of time-independent flow. They are defined as the larger eigenvalues of the Cauchy–Green strain tensor of the flow map. FSLEs are defined by the exponential rate  $\lambda$  (d<sup>−1</sup>) of separation of two neighbouring particles during a time advection  $t$ :

$$\lambda = t^{-1} \log(\delta f / \delta 0), \quad (2)$$

where  $\delta 0$  and  $\delta f$  are the initial and final separation distance that are fixed before computation. FSLEs are commonly used as an indicator of frontal activity and stirring intensity (d'Ovidio et al., 2004). Relatively large FSLE values are associated with formerly distant water masses, whose confluence creates a transport front (d'Ovidio et al., 2004, 2009). Fronts identified as positive maxima (ridges) of FSLEs exhibit a convergent dynamics transverse to them, whereby passive particles in their vicinity are attracted to the front and then advected along it (Della Penna et al., 2015). Conversely, negative values of FSLEs indicate divergent dynamics, leading to the dispersion of particles away from the front. In order to examine meso- and submesoscale frontal structures during phytoplankton blooms, daily FSLEs images from 10 to 25 November were analysed. The daily images were used to create a video (see the Video supplement) to illustrate the daily evolution of the FSLEs in the area where the phytoplankton bloom developed.

### 3 Results

#### 3.1 Satellite-derived chlorophyll *a* during the sampling period


During the Gayoso cruise (12–19 November, Fig. 2a), a uniform, large band of high-surface Chl *a* concentration (between 4.0 and 32 mg m<sup>−3</sup>) expanded over the mid and outer shelf. During the Agujero Azul cruise (20–27 November, Fig. 2b), the band of Chl *a* disaggregated and showed lower intensity (1.5 and 34 mg m<sup>−3</sup>), but the Chl *a* concentration was still high in the area of the sampling stations (Fig. 2b). The SST showed warming of the inner mid-shelf waters north of 44° S over the 8 d average periods (around 14.5 °C in mid-shelf) (Fig. 2c, d), but the SST remained constant at the sampling area, at around 11 °C (see Fig. 4 and Table 1). The daily images of Chl *a* from 14 to 25 November (Fig. 3) showed an abrupt proliferation of phytoplankton on 15 November, which notably intensified on 16 November when it reached the maximum concentrations during the studied period (up to 35 mg m<sup>−3</sup>). On this date, the extraordinary bloom of Amphidomataceae was sampled at GA01. During the following days, the chlorophyll levels remained high in the area but became more disaggregated into variable patches. On 25 November, the sampling day at station AA09, Chl *a* had decreased at station GA01 but remained intense in the area (Fig. 3) with still extraordinary densities of Amphidomataceans at station AA09.

#### 3.2 In situ biogeochemical properties and water column structure

A reddish water discolouration was observed in the bloom area (45.5–46° S, 62–61° W) during the cruises. Surface water temperature and salinity remained similar at both stations GA01 and AA09, sampled with a 10 d interval (Fig. 4, Table 1). Moreover, both stations displayed the same vertical structure in terms of temperature and salinity (Fig. 4), the mixed layer depth (MLD), and the subsurface chlorophyll maximum (SCM) (Table 1). In situ chlorophyll *a* concentration at the surface was 20 µg L<sup>−1</sup> at GA01 and 4.5 µg L<sup>−1</sup> at AA09 (Table 1). Dissolved inorganic macronutrients in surface waters were similar in both stages of the bloom, except for nitrate and silicate, which were higher at AA09. In particular, the silicate recovered by ~ 5 times towards the advanced stage of the bloom.

#### 3.3 Multispecificity of the Amphidomatacean bloom and azaspiracids

Total protistan plankton of cell size larger than 5 µm reached up to 31.68 × 10<sup>6</sup> cells L<sup>−1</sup> at station GA01; and 10 d later at station AA09, the abundance was 13.69 × 10<sup>6</sup> cells L<sup>−1</sup>. Of all this total protist abundance, the Amphidomataceae clade represented up to 99 % and 98 %, respectively (Fig. 5), mostly dominated by the non-toxicogenic *Azadinium spinosum*




**Figure 2.** Eight-day time mean of (a, b) satellite chlorophyll *a* (Chl *a* in  $\text{mg m}^{-3}$ ) and (c, d) sea surface temperature (SST in  $^{\circ}\text{C}$ ) during the sampling period: (a, c) 12–19 November 2021 and (b, d) 20–27 November 2021. From left to right, isobaths of 200, 1000, and 1800 m are shown. The numbers in the column on the left side of each panel correspond to the percentage of cloud-free pixels in each daily satellite image. The sampling stations GA01 (pink dot) and AA09 (blue dot) are shown.

**Table 1.** Surface values of physical and chemical variables measured at stations GA01 and AA09: sea surface temperature (SST), sea surface salinity (SSS), concentration of in situ chlorophyll *a*, and macronutrients. The depth of the subsurface chlorophyll maximum (SCM) and the mixed layer depth (MLD) is also displayed.

	GA01	AA09
Date	16 Nov	25 Nov
SST ( $^{\circ}\text{C}$ )	10.5	10.9
SSS	33.5	33.5
SCM (m)	15	10–25
MLD (m)	60	55
Chlorophyll <i>a</i> ( $\mu\text{g L}^{-1}$ )	20.0	4.5
Nitrite ( $\mu\text{M}$ )	0.48	0.33
Nitrate ( $\mu\text{M}$ )	1.77	3.57
Ammonium ( $\mu\text{M}$ )	1.32	1.32
Silicate ( $\mu\text{M}$ )	4.54	22.35
Phosphate ( $\mu\text{M}$ )	0.33	0.44

ribotype C and *Az. dalianense* (Figs. 5 and 6), representing together > 95 % of total Amphidomataceae (Fig. 5). Taxonomic identification up to species level was possible after an exhaustive morphological examination of cells under light microscopy (Fig. 5) and scanning electron microscopy (Fig. 6). In Appendix A, more micrographs of Amphidomataceans are shown, taken under light microscopy and scanning electron microscopy (Figs. A1 to A10), along with the rationale for the Amphidomatacean species designations. The ITS-based metabarcoding of species diversity detected in the field samples at GA01 and AA09 (shown in Appendix B) also supports the dominance of *Azadinium spinosum* ribotype C and *Az. dalianense*. In the detailed counting of protistan species in 10 mL subsamples, the well-recognized Amphidomataceae species under inverted light microscopy for their individual quantification were *Az. spinosum* ribotype B and ribotype C, *Az. dalianense*, *Az. obesum*, and a group of smaller taxa including *Az. dexteroporum*, *Amphidoma parvula*, and *Am. languida* (Fig. 5). This distinction was based on morphological aspects combining the cell size and shape, such as the length/width relation and other taxonomic





**Figure 3.** Daily satellite-derived surface chlorophyll *a* in the area of the sampling stations: GA01 (pink dot, sampled on 16 November 2021) and AA09 (blue dot, sampled on 25 November 2021).


aspects, for instance, a slender shape: *Az. spinosum* ribotype B; round: *Az. obesum*; short, tiny: *Az. dexterporum*, *Am. parvula*, and *Am. languida*; with a bump in the hypotheca and a pyrenoid in the hyposome: *Az. dalianense*; and with a spine in the hypotheca: *Az. spinosum* ribotype C, *Az. dalianense*, and *Az. dexterporum*. These and other taxonomic features were further examined by SEM (Fig. 6), for example the number and arrangement of the thecal plates, the presence of thecal pores (see Appendix A). Finally, protists other than Amphidomataceans (Fig. 7) contributed up to 1.0 % and 2.2 % of the total abundance at stations GA01 and AA09, respectively. Most of the other protists were heterotrophic and mixotrophic dinoflagellates and ciliates. No diatoms were observed in the field samples. For an overview of the pure Amphidomataceae bloom in the field samples (e.g. no mucus formation, no aggregates, cells undergoing cell division), low-magnification micrographs obtained through light mi-

croscopy are presented in Appendix C. The screen of all known and potentially novel variants of azaspiracids, which are produced by Amphidomataceae, revealed the presence of solely azaspiracid-2 (AZA-2) in both stages of the bloom, with field values of 2122 pg L<sup>-1</sup> at GA01 and 620 pg L<sup>-1</sup> at AA09.

### 3.4 Surface currents, particle advection model, and description of the frontal systems

The mean and standard deviation of the ADT during the sampling period (16 to 28 November 2021) evidenced a mesoscale anticyclonic eddy of about 100 km in diameter in the area where the Amphidomataceae bloom was observed at the two sampling locations (Fig. 8). In addition, the modelled trajectory of the particles released along the zonal transect at 46° S on 10 November showed high retention at the





**Figure 4.** Vertical profiles of temperature, salinity, and stability measured at the two blooming stations, GA01 (sampled on 16 November) and AA09 (sampled on 25 November). Strong stratification is denoted by the water column stability (Brunt–Väisälä buoyancy frequency), which indicates a mix layer depth (MLD) of around 60 m.

blooming area over the continental shelf after running for 20 d (Fig. 9). On the contrary, high advection within the flow of the Malvinas Current was evidenced (Fig. 9). Notably, in the eddy area, particles slightly displaced southwards, remaining trapped in the area after 20 d since their release. The particles advected by geostrophic velocities suggest that the anticyclonic eddy acted as a potential mechanism to retain the Amphidomataceae bloom within the same location during the two synoptic samplings (Fig. 9). All the other particles released east and west of the eddy displayed a different behaviour (Fig. 9), also shown in Appendix D where four parcels of particles were advected from 16 to 25 November. West of the eddy they were advected northwards at a rather slow speed (average  $5 \text{ cm s}^{-1}$ ), while east of the eddy they increased their speed towards the north as they approached the continental slope. Within the core of the Malvinas Current, speeds were as high as  $0.8 \text{ m s}^{-1}$  (Fig. 9).


Moreover, the finite-size Lyapunov exponent (FSLE) ridges highlighted the stirring and hydrologically complex nature of the southwestern Atlantic Ocean, associated with the high hydrographical heterogeneity of the oceanic waters (Fig. 10). Although FSLEs were less intense over the shelf than in the adjacent oceanic waters, in the area where the Amphidomataceae bloom was sampled (dashed pink square in Fig. 10), two relatively strong FSLE ridges consistently kept both bloom stations within the same water mass during the period between the two synoptic samplings (Fig. 11).

## 4 Discussion

This study is unique from both biological and physical perspectives due to the following factors: (i) the Amphidomataceae bloom observed in spring 2021 in the Argentine Sea, with up to  $32 \times 10^6 \text{ cells L}^{-1}$ , represents the largest bloom of this clade ever recorded globally; (ii) unusual sampling in offshore shelf waters with two vessels over a 10 d interval allowed for synoptic observations of the bloom at two active developmental stages; (iii) simultaneous ecological characterization of the bloom, surface currents, and fronts provided insights into patch stirring and the short-term evolution of the bloom; (iv) abundance data of Amphidomataceae are rare in the field, and to our knowledge, this is the first detailed description of species abundance in field samples, combining light microscopy, electron microscopy, and metabarcoding; (v) the relatively low abundance of *Azadinium spinosum* ribotype B indicated high AZA-2 cell quotas; (vi) the fine-taxon assessment of the Amphidomataceae bloom revealed biogeographical patterns and strain-specific toxic potential; and (vii) the use of interdisciplinary approaches sheds light on the biophysical coupling underlying the persistence and horizontal transport of this extraordinary bloom in offshore shelf waters.

### 4.1 Amphidomataceae blooms over the Patagonian shelf

The phytoplankton spring and summer blooms in the Patagonian shelf display a southward progression related to the seasonal thermal cycle. In early spring (September–October), the water column stratifies north of  $\sim 45^\circ \text{S}$ , favouring the proliferation of opportunistic micro-diatoms in the nutrient-rich, well-lit surface layers (Ferronato et al., 2023). South of  $\sim 45^\circ \text{S}$ , the bloom initiates later in spring/early summer (December–January) and continues until autumn (March) (revised in Guinder et al., 2024). The distribution of surface Chl *a* during the time of the cruises was indeed indicative of a mid–late-spring phytoplankton bloom over the Patagonian shelf, following the thermal stratification. Here, blooms of nanoflagellates and dinoflagellates are triggered by combined vertical stability and nutrient-depleted surface waters (especially silicates) after the early-spring blooms of large diatoms (Balch et al., 2014; Carreto et al., 2016). During the Gayoso cruise, detailed analyses of plankton communities and microbial carbon fluxes at all sampling stations revealed a latitudinal shift from more heterotrophic to more phototrophic activity towards the south (Ferronato et al., 2025; Gilabert et al., 2025). The massive proliferation of Amphidomataceans in mid-November 2021 was in line with this successional pattern. These nano-dinoflagellates bloomed in a stratified water mass with a deep mixed layer depth ( $\sim 60 \text{ m}$ ). While dinoflagellates are less effective at nutrient resorption compared to diatoms, they can move throughout the stable water column to find light and nutrients (Glibert, 2016), espe-



**Figure 5.** Relative abundance (in %) of the Amphidomataceae species identified under light microscopy at stations GA01 and AA09. The total abundance of protists at each station was 32 and 14  $\times 10^6$  cells  $L^{-1}$ , respectively. From the total cells counted, Amphidomataceae represented 99.0 % and 97.8 %, respectively. The colours in the pie charts correspond to the same species at both stations. Scale bar: 5  $\mu m$ .


cially at low phosphate levels (Lin et al., 2016) as observed at the stations GA01 and AA09 with high nitrate-to-phosphate ratios.

The success of the multispecific bloom of Amphidomataceae may be attributed to a combination of multiple intrinsic and extrinsic factors. For instance, their small cell size, unique swimming modes, and the production of azaspiracids may have alleviated grazing pressure (Tillmann et al., 2019). In fact, in the fixed samples, it was observed that many cells were obviously active and undergoing cell division and that these nanoflagellates were overwhelmingly the predominant photosynthetic protists responsible for the high Chl *a* levels, with negligible abundance of micro-grazers accompanying the bloom development. The presence of less than 2 % of other protists (mixotrophs and heterotrophs) could be also related to a delay in the recovery of predators following the early blooms of microdiatoms, as well as to an abrupt development of the Amphidomataceae bloom, which may have prevented micrograzers from taking advantage of the available food. Another observation supporting the active persistence of the bloom was the pristine condition of the microenvironment surrounding the dense populations of Amphidomataceae (see Appendix C), with no aggregates or mucus formation as is typically observed in the late stages of blooms (Genitsaris et al., 2019). Furthermore, no competitors for light and nutrients were detected in the samples; specif-

ically, no diatoms were present. This may explain the rapid recovery of silicates (from 5 to 22  $\mu M$ ) over the 10 d persistence of the Amphidomataceae bloom, as these silicates were not being utilized by silicate-requiring species. A similar observation was noticed during a bloom induced by iron fertilization in the Southern Ocean, where diatoms predominated and silicate levels decreased from 10 to 6  $\mu M$  within the bloom patch over the course of 12 d (Abraham et al., 2000).

#### 4.2 Highest abundance ever recorded for a bloom of Amphidomataceans

With 17 described species of *Azadinium* and 14 of *Amphidoma*, the Amphidomataceae represent a small but diverse group of dinoflagellates. Most of these species are very similar in size and shape, which makes the qualitative identification and, in particular, the species-specific quantification in field samples difficult. Hence, characterizing the cryptic species of the multispecific bloom of Amphidomataceae in this study represented a major challenge, where a reliable species identification was achieved by the combination of several diagnostic details using electron microscopy. Counting of fixed samples by light microscopy provided high quantitative accuracy, but reliable species identification was only possible in a few cases. Complementing the identification by microscopy with metabarcoding, specifically targeting the internal transcribed spacer (ITS1) region (Liu et al., 2023),



**Figure 6.** Scanning electron microscopy of Amphidomataceae species (a–f) and other dinoflagellates (g–j) at stations GA01 and AA09. (a) *Azadinium spinosum*, (b) *Az. dalianense*, (c) *Az. obesum*, (d) *Az. dexteroporum*, (e) *Amphidoma parvula*, (f) *Am. languida*, (g) *Oxytoxum gracile*, (h) *O. laticeps*, (i) *Gyrodinium* sp., and (j) unidentified gymnodinoid species. Scale bars = 2  $\mu\text{m}$ . See Appendix A for more micrographs of Amphidomataceans and for evidence and rationale for the Amphidomatacean species designations.

allowed for the detailed characterization of Amphidomatacean species diversity in the field samples. By combining the three approaches, we were able to identify ribotypes and the toxic species, including previously described Amphidomataceae species for the Argentine Sea (Fabro et al., 2019; Tillmann et al., 2019, 2021) or species still undescribed in the global seas (see Appendix A).

In the North Atlantic, AZA-1 (and its producing species, *Az. spinosum* ribotype A) has been identified as one of the most prevalent toxins among a wide range of AZA variants (Tillmann et al., 2021). The bloom density of Am-

phidomataceae around Ireland has been reported as  $8.3 \times 10^4$  cells  $\text{L}^{-1}$  for *Az. spinosum* (Wietkamp et al., 2020) and  $47 \times 10^5$  cells  $\text{L}^{-1}$  for *Amphidoma languida* (McGirr et al., 2022) and a small bloom of *Am. languida* in the North Sea with  $1.2 \times 10^5$  cells  $\text{L}^{-1}$  has also been described (Wietkamp et al., 2020). In this region, cases of human intoxication with AZAs have been linked to the consumption of contaminated mussels from the Irish west coast, where blooms in the shelf-break area can reach coastal shellfish beds through wind-driven advection (Raine, 2014). Notably, in the Argentine Sea, only AZA-2 has been detected in field and culture sam-






**Figure 7.** Micrographs of other protists (1.0–2.2 % of total abundance) present in the bloom: (a) *Oxytoxum laticeps*, (b) *Gyrodinium spiralis*, (c) *O. gracile*, (d) *Karenia* sp., (e) unidentified dinoflagellate, (f) *Gyrodinium* sp., (g, h, j) unidentified gymnodinoid cells, (i) *Katodinium* sp., (k–l) naked ciliates, (m) *Peridiniella* sp., (n) Kareniaceae-type cell, (o) Euglenophyte, (p) *Lessardia elongata*, (q) *Torodinium robustum*, (r) ciliate with an Amphidomataceae cell (arrow), (s) ciliate, and (t) *Dinophysis* sp. Scale bars = 10 µm, except in photos (g) and (h) where scale bars = 5 µm.


ples (Turner and Goya, 2015; Fabro et al., 2019; Tillmann et al., 2019; Guinder et al., 2020), and so far, no poisoning events have been attributed to AZAs.

In this study, relatively high levels of solely AZA-2 were detected in bloom samples. A toxin profile of solely AZA-2 is up to now only known for the Argentine strain H-1-D11 of *Azadinium spinosum* ribotype B (Tillmann et al., 2019), and this ribotype was also identified in the present bloom. Relat-

ing AZA quantities to the relatively low abundance (0.1 % to 0.4 % of total Amphidomataceae) of *Az. spinosum* ribotype B revealed AZA-2 cell quotas of 17–42 fg per cell, which is an order of magnitude higher than the cell quota of 2 to 9 fg per cell for strain H-1-D11 grown under laboratory conditions (Tillmann et al., 2019). In fact, *Az. spinosum* of ribotype A producing AZA-1 and AZA-2 is the primary causative agent of AZA poisoning in Europe (Tillmann, 2018). How-



**Figure 8.** (a) Mean and (b) standard deviation (SD) of the absolute dynamic topography (ADT, in metres), displayed in colour scale during the period 16 to 28 November, embracing the sampling period on both bloom stations: GA01 (pink dot, Gayoso cruise) and AA09 (blue dot, Agujero Azul cruise).



**Figure 9.** Particle advection after 20 d since particle release on 10 November 2021. Initial points: one particle every 0.05 grades along 46° S. Note the high retention and the distinct behaviour of particles in the area of the sampling stations: GA01 (pink dot) and AA09 (blue dot). Satellite altimetry geostrophic velocities averaged for the same 20 d are also displayed with black vectors; the magnitude of the averaged speeds is represented with colours in the background ( $\text{m s}^{-1}$ ).


ever, the large majority of cells of *Az. spinosum* in the present bloom sample is of ribotype C, which is, based on analyses of several strains from Argentina, non-toxicogenic (Tillmann et al., 2019). All globally available strains (including

strains from Argentina) of the other co-dominant species in the bloom, *Az. dalianense* (Tillmann et al., 2019), also do not produce AZAs. In the Chilean continental shelf in the SE Pacific, AZAs have been detected in scallops and mussels (López-Rivera et al., 2010), but no intoxication events or large blooms of this clade have been documented (Iriarte et al., 2023). Other *Azadinium* cells have been observed widely distributed in inner seas along the coast of Chile but with no further characterization of their potential toxicity (Iriarte et al., 2023; Corredor-Acosta et al., 2025). Only *Az. poporum* has been described as an AZA producer in Chilean waters (Tillmann et al., 2017a). Likewise in Peru, a relatively dense bloom (up to  $1.0 \times 10^6$  cells  $\text{L}^{-1}$ ) of *Az. polongum* was detected in the summer of 2014, with no AZA production (Tillmann et al., 2017b). Although the continental shelves of Chile-Peru and Argentina have different hydrology, both span similar latitudinal gradients along the South American coasts and are influenced by the Humboldt and Malvinas currents, which share a common origin in the Circumpolar Antarctic Current. Strikingly, both shelves exhibit different populations of Amphidomataceae, despite the expectation that ocean currents could serve as transport pathways for HAB species, promoting their dispersion (Giddings et al., 2014). This suggests that additional factors, such as coastline configuration and bottom topography, play a role in water mixing and retention processes and ultimately the development of HABs (Pitcher et al., 2010).

#### 4.3 Spatio-temporal evolution of the bloom: retention and stirring

Phytoplankton bloom initiation, magnitude, and persistence rely on a host of biogeochemical and physical processes. As






**Figure 10.** FSLE ridges computed as finite-size Lyapunov exponent (FSLE) for 16 November, shown against a greyscale background. All sampling stations from the Gayoso cruise (pink dots) and Agujero Azul cruise (blue dots) are indicated. The date 16 November corresponds to the sampling of the Amphidomataceae bloom at station GA01 (pink dot within the square marked by a dashed pink line). The square highlights the area shown in Fig. 11.

discussed in previous sections, the explosive onset of the bloom of multiple species of Amphidomataceae was associated with a combination of water column stability, the negligible presence of micrograzers, and the ecological traits of this group that facilitate massive proliferation. However, these conditions alone do not fully explain the persistence of this bloom, which was sampled 10 d later at a location 40 km away in the offshore waters of the Patagonian shelf, where strong surface currents were expected to disperse plankton blooms. The persistence of this extraordinary bloom, characterized by its remarkable magnitude and consistent species composition, indicates the retention and accumulation of the bloom patch within the same water mass. In addition to the biological evidence confirming the presence of the same Amphidomataceae bloom at both sampling stages, analyses of circulation through satellite altimetry showed that an anticyclone of about 100 km in diameter was the key feature responsible for retaining the bloom within the two sampling stations. The retention that the eddy caused was evidenced by two Lagrangian experiments. In the first one, the trajectories of virtual particles released within the eddy show that almost none of the particles escaped from the eddy during the 10 d that separated the two sampling stations. In the second one, FSLE maps showed that no mesoscale fronts separated the two sampling stations during those days. Both stations remained within the same water mass delimited by two FSLEs with negative values, which indicate a divergence of plankton transverse to the filaments, enhancing the reten-

tion of the bloom. Two potential scenarios could explain this: (1) the same patch remained in the area over the 10 d of sampling, occupying an area of 40 km or larger (Fig. 12a and b), or (2) a smaller bloom patch was initially detected at station GA01 and then transported by stirring towards AA09 (Fig. 12c). Sampling with higher spatial and temporal resolution of bloom initiation, development, species succession, and collapse would have been necessary to distinguish between scenarios (1) and (2). A less likely scenario is that (3) two Amphidomataceae blooms developed independently at both locations (Fig. 12d). This scenario is improbable given the complex interplay of physical and biological processes that govern bloom development. On the physical side, the advection, accumulation, and stirring of water masses act as selective forces (Abraham et al., 2000; Lehahn et al., 2007; Della Penna et al., 2015), favouring the proliferation of certain species or functional groups over others, depending on local and transient conditions (Lévy et al., 2018; Hernández-Carrasco et al., 2020; Mangolte et al., 2023). Biologically, additional layers of variability – including interspecific competition, grazing pressure, successional dynamics, toxin production, and cyst formation – further shape bloom composition and trajectory. Considering the influence of such dynamic and site-specific factors, the independent development of blooms with identical species composition and relative abundances at two separate locations is unlikely.

Additionally, the hydrographically complex southwestern Atlantic creates a variety of microhabitats at the meso-






**Figure 11.** Finite-size Lyapunov exponent (FSLE) ridges in the area of the two locations with the Amphidomataceae bloom: GA01 (pink dot) sampled on 16 November and AA09 (blue dot) sampled on 25 November. See the video of these daily images in the Video supplement. The 200 m isobath is indicated in yellow.

and submesoscale. These include areas of upwelling, downwelling, eddies, retention, and dispersion (Becker et al., 2023; Beron-Vera et al., 2020; Salyuk et al., 2022; Saraceno et al., 2024). This spatial heterogeneity enhances the development of variable nutrient patches and phytoplankton productivity (Lehahn et al., 2017; Lévy et al., 2018; Hernández-Carrasco et al., 2020; Ser-Giacomi et al., 2023). During the Gayoso cruise, contrasting plankton assemblages and bloom types were observed at all sampling locations, including distinct blooms of dinoflagellates, coccolithophores, diatoms,

and nanoflagellates. These variations were related to substantial heterogeneity in surface velocities and environmental conditions across the region (Feronato et al., 2025). It is important to highlight that the bloom of Amphidomataceae at GA01 and AA09 was not observed at any other station during the Agujero Azul cruise. During the Gayoso cruise, this clade was observed in relatively high abundance only at GA10, located in the northern area over the shelf, where species assemblages were significantly different (Feronato et al., 2025). In this study, while the estimation of the re-



**Figure 12.** Hypothetical scenarios of the spatio-temporal evolution of the Amphidomataceae bloom during the 10 d period between the synoptic sampling at the two stations 40 km apart: GA01 (pink circle) and AA09 (blue circle). The bloom's biomass was  $32 \times 10^6$  cells  $L^{-1}$  at GA01 and  $14 \times 10^6$  cells  $L^{-1}$  at AA09. In the “Same Bloom Patch” (a–a' and b–b'), the bloom covered an area encompassing both sampling stations. The bloom developed in such a way that the biomass was either distributed homogeneously across the patch (a–a') or heterogeneously (b–b'), resulting in variable biomass patterns over time and space. In the “Drifting Bloom Patch” (c–c'), the water mass with the Amphidomataceae bloom detected at station GA01 was transported by currents towards station AA09, where the bloom was detected with a lower biomass but still with high intensity. The “Independent Bloom Patch” (d–d') suggests that two discrete, autonomous Amphidomatacean blooms developed locally at each sampling station.

tention of the Amphidomataceae bloom is certainly limited by the accuracy of satellite altimetry maps, the documented 100 km diameter of the eddy is a reasonable size that can be distinguished using the gridded satellite altimetry maps produced by CMEMS. Although this retention is transient, this particular circulation facilitated the massive development of the Amphidomataceae bloom, with no evidence that this patch was advected to the Malvinas Current, as observed with drifters released east of the bloom area in spring 2021 (Saraceno et al., 2024). Our results highlight the importance of studying the development of toxic phytoplankton blooms on continental shelves, focusing on the biophysical coupling that drives their patchy nature, persistence, and transport, in order to capture short-lived blooms and their potential to cause toxic outbreaks.

## 5 Conclusions

We present an example where mesoscale (100 km) circulation on the Patagonian shelf acted as a crucial physical driver that prevented the dispersion of an Amphidomatacean bloom. As a result, the bloom persisted for 10 d in the same area. While our study provides valuable insights, it has limi-

tations, as it only explores physical processes at a mesoscale resolution, whereas sub-mesoscale processes are crucial for understanding divergence, convergence, and the mixing of plankton and nutrients within and between water masses. We also note that the collection of samples from only two synoptic stations, along with the incidental detection of the Amphidomatacean bloom during two cruises with different ecological objectives, limited our ability to track the detailed spatial and temporal evolution of the bloom. One strategy to overcome these limitations would be continuous sampling of the bloom, which could be achieved through a Lagrangian experiment. In this approach, water samples would be collected following drifters that track surface currents. At the same time, detailed biodiversity analyses of the bloom should be conducted, including the succession of species and functional groups in relation to their differential uptake of nutrients, grazing pressure, and the potential exchange of plankton and nutrients between neighbouring water masses.

## Appendix A: Estimation and categorization of Amphidomatacean species diversity

For the present study, we combined light microscopy (LM) quantification, scanning electron microscopy (SEM) examination, and metabarcoding to characterize the field samples as accurately as possible, both qualitatively and quantitatively. Generally, the following species of Amphidomataceae were identified as detailed here.

### A1 *Azadinium spinosum*

Specimens were identified with SEM as *Az. spinosum* based on the combination of (1) the presence of an antapical spine, and (2) the presence of a ventral pore located on the right side of the suture of plate 1' and 1'' (Fig. A1). The vast majority of cells thus identified as *Az. spinosum* had a somewhat broader cell shape. Generally, identification of *Az. spinosum* is complicated as there are several different ribotypes (Tillmann et al., 2021), which notably differ in azaspiracid toxin presence and profile. In a previous study from the Argentine shelf region, it was shown that 23 out of 24 isolated strains of *Az. spinosum* were assigned to the non-toxicogenic ribotype group C, and cells from these strains also had a somewhat broader cell shape (Tillmann et al., 2019). Metabarcoding of the present bloom sample revealed that the most common sequences showed a high match with sequences of these Argentine ribotype C strains. A dominance of non-AZA-producing *Az. spinosum* (ribotype C) also aligns with the finding that no AZA-1, the marker toxin of the toxicogenic *Az. spinosum* ribotype A strains (Tillmann et al., 2021), was detected in the field samples. In the quantitative light microscopy (LM) counts, all medium-sized cells (length > 12 µm) with a rounded hypotheca and an antapical spine were thus categorized as *Az. spinosum* ribotype C.

Recent studies have revealed the presence of a new, molecularly distinct species of *Azadinium* in the North Atlantic, which is morphologically indistinguishable from *Az. spinosum* and is currently provisionally referred to as *Azadinium* cf. *spinosum* (Tillmann et al., 2021). Therefore, it cannot be ruled out that this species was also present in the samples, but metabarcoding showed no evidence of the presence of *Az. cf. spinosum* in the bloom samples.

In addition to these broader cells of *Az. spinosum* ribotype C cells, LM analysis revealed a (much rarer) presence of distinctly slender cells with an antapical spine (Fig. A2a). Such cells perfectly correspond to the cell shape of a single strain H-1-D11 from Argentina, identified as a ribotype B strain of *Az. spinosum* (Tillmann et al., 2019), and this strain is depicted here for comparison (Fig. A2c, d). In SEM, specimens of slender shape lacked the rim around the pore plate (Fig. A2b), which is the morphological diagnostic feature differing in ribotype B strains from ribotype A and C strains, which all have a thick rim. Additionally, metabarcoding showed conformity of some sequences with other ri-


botype B *Az. spinosum* strains (e.g. 99 % similarity with the Argentinian strain H-1-D11). Consequently, all slender cells with an antapical spine quantified in LM were categorized here as *Az. spinosum* ribotype B. Strain H-1-D11 from Argentina was shown to produce solely AZA-2 (Tillmann et al., 2019). As this was the only AZA detected in our field sample, this is additional support for this *Az. spinosum* ribotype B designation.

### A2 *Azadinium obesum*


Cells of the non-toxicogenic species *Az. obesum* were identified in the SEM samples based on the combination of the following features: (1) no antapical spine, (2) ventral pore on the left side of plate 1', and (3) no contact between plates 1a and 1'' (Fig. A3). All such cells had a distinctly broad oval shape and were relatively large. In the light microscope, all relatively large oval Amphidomataceae cells with a rounded hypotheca and no visible spine were therefore categorized as *Az. obesum*.

### A3 *Azadinium dalianense*

With the rounded hypotheca, cells of *Az. obesum* and *Az. spinosum* ribotype C were clearly distinguishable in the light microscope from cells with a distinctly protruding bump, at the tip of which a small spine was present (Fig. A4a). Corresponding cells detected in SEM preparations (Fig. A4b–e) were identified as *Az. dalianense*, based on the combination of the following features: (1) a ventral pore on the left side of the pore plate, (2) asymmetrical hypotheca with a bump and a distinct antapical spine, and (3) the presence of only three apical plates and two anterior intercalary plates. Since all three features were only rarely visible simultaneously due to the cell's orientation, the presence of the somewhat similar species *Az. perfusorium* cannot be ruled out. *Az. perfusorium* also has a posterior small bump with a spine and a ventral pore located on the left side of the pore plate, but it possesses four apical plates and three intercalary plates (Salas et al., 2021). However, neither the SEM nor metabarcoding provided any indication of its presence in the samples. The occurrence of *Az. dalianense* in the region is well documented by a series of strains isolated from the Argentine shelf in 2015 (Tillmann et al., 2019), and *Az. dalianense* was also identified as part of *Azadinium* blooms in 1991 (Tillmann and Akselman, 2016). Metabarcoding additionally indicated the presence of two different ribotypes of *Az. dalianense*, namely E and B as defined in Tillmann et al. (2019), where all previous strains from Argentina belong to the ribotype E clade. Accordingly, the majority of reads from the bloom station *Az. dalianense* were from ribotype E (represented by strains H-4-E8 and N-12-04 in the reference dataset), whereas reads of ribotype B (represented by strain IFR-ADA-01C) made only ~ 0.02 % of all *Az. dalianense* reads. All strains of *Az.*




**Figure A1.** LM (a) and SEM (b–f) of cells of the Amphidomatacean bloom stations identified as *Azadinium spinosum* ribotype C. (b–d) Cells in ventral view. (e, f) Cells in dorsal view. Note the pyrenoid (black arrow in a), the position of the ventral pore (black arrows in b, c, d), the rim around the pore plate (white arrows in b, c), and the distinct antapical spine. Scale bars = 2  $\mu$ m.



**Figure A2.** LM (a) and SEM (b) of cells of the Amphidomatacean bloom stations identified as *Azadinium spinosum*. For comparison, the LM (c) and SEM (d) of cells of strain H-1-D11 of *Azadinium spinosum* ribotype B isolated from the Argentine shelf in 2015. Note the elongated cell shape, the distinct antapical spine, and the lack of a rim around the pore plate (white arrows in b and d). Scale bars = 2  $\mu$ m.





**Figure A3.** LM (a) and SEM (b–f) of cells of the Amphidomatacean bloom stations identified as *Azadinium obesum*. (b–d) Cells in ventral view. (e, f) Epitheca in apical view. Note the lack of an antapical spine, the position of the ventral pore (white arrows in b, e, f), and the lack of contact between plates 1a and 1'' (kofoidian plate label notation) visible in (e) and (f). Scale bars = 2  $\mu\text{m}$ .

*dalianense* representing different ribotypes collected from various regions analysed so far were non-toxicogenic.


The classification and quantification of *Az. obesum* and *Az. dalianense*, however, is complicated by the fact that a number of cells of an unclear assignment were found in the samples (Fig. A5). These cells, like *Az. obesum*, (1) had a ventral pore on the left side of plate 1', and (2) no contact between plates 1a and 1'', but unlike *Az. obesum*, they had a distinct, albeit small, antapical spine. This combination of features is not known from any described *Azadinium* species, suggesting that this may be a new species. However, for a complete description as a new *Azadinium* species, further investigations are necessary, ideally adding sequence data and analyses of toxin production. In any case, it is clear that cells of this type will have been included in the categories *Az. obesum* or *Az. spinosum* during the light microscope analyses and quantifications.

#### A4 Smaller Amphidomatacean species: *Azadinium dexteroporum*, *Amphidoma parvula*, and *Amphidoma languida*

While *Az. spinosum*, *Az. obesum*, and *Az. dalianense* fall into a slightly larger size class, a number of smaller *Azadinium* species were identified and categorized in the samples. One

of them was *Az. dexteroporum* (Fig. A6), which was identified in the SEM by the following combination of features: (1) relatively small size, (2) presence of a distinct antapical spine, (3) a slightly posteriorly positioned ventral pore on the right side of the pore plate, and (4) a distinctly concave central intercalary plate 2a. Metabarcoding revealed a number reads for an *Azadinium* sp. 1 with *Az. dexteroporum* as closest species suggestion (line 7 in Table B1), however only with rather low similarity (90–95 %) compared to the reference database. Globally, there are only three available strains of *Az. dexteroporum*, and of those only one strain from the Mediterranean was identified as a producer of AZAs (Rossi et al., 2017). In contrast, two additional strains from the North Atlantic, which also had marked sequence differences compared to the Mediterranean strain, did not produce AZAs (Tillmann et al., 2020). The low similarity of *Az. dexteroporum* reads from the present bloom samples thus indicates that the local population may represent a new ribotype quite distinct from the AZA-producing ribotype, and strain isolation of local *Az. dexteroporum* is needed to clarify its identity and toxin production potential.

In the same size class, cells were also observed that were identified in SEM as *Amphidoma parvula* (Fig. A7) by (1) their flat hypotheca and (2) a characteristically shaped 1' plate. This non-toxicogenic species was described in 2018



**Figure A4.** LM (a) and SEM (b–e) of cells of the Amphidomatacean bloom stations identified as *Azadinium dalianense*. (b) Cell in ventral view. (c) Cell in dorsal view. (d) Epitheca in apical view. (e) Epitheca in dorsal view. Note the distinct apical spine on a triangular, bumpy hypotheca (a–c), the position of the ventral pore on the left side of the pore plate (white arrow in d), the presence of only two large anterior intercalary plates 1a and 2a (c–e), and the presence of only three apical plates in (d) (kofoidian plate label notation). Scale bars = 2  $\mu\text{m}$ .

based on a culture isolated from the Argentine shelf (Tillmann et al., 2018b). In accordance, a low number of reads with high similarity to *Am. parvula* strain H-1E9 (> 98 %) were recorded by metabarcoding. With its relatively long apical plates, *Am. parvula* could also be easily distinguished in the SEM from the similarly small *Am. languida*, which was also identified in SEM (Fig. A8). In *Am. languida*, (1) the small apical plates and (2) the presence of a large characteristic antapical pore are distinguishing features.

However, in 2024, a new species of *Amphidoma*, *Am. fulgens*, was described, which is morphologically almost identical to *Am. languida* but shows significantly different sequence data and, different to *Am. languida*, does not produce azaspiracids (Kuwata et al., 2024). *Amphidoma fulgens* was found to be widely distributed in the Pacific, but there are no records yet from the Atlantic Ocean. Despite its presence in the bloom samples, there were no reads related with higher similarities to *Am. languida* in the ITS metabarcoding dataset but only very few reads with rather low similarity to *Amphidoma* entries in the database. This seems to be in line with the general notion that ITS sequencing of cultured strains of *Am. languida* in many cases failed


(Wietkamp et al., 2019). In fact, data of a previous metabarcoding study (Liu et al., 2023) showed that *Am. languida* hits were abundant in Chinese waters in the LSU dataset but absent in the ITS-1 dataset. While *Am. languida* and also *Am. fulgens*, according to their original descriptions, only have one small pore on each precingulate plate (Tillmann et al., 2012; Kuwata et al., 2024), there were also several cells of *Amphidoma* with three or more small pores on individual precingulate plates (Fig. A9a–d). To what extent these cells represent *Am. languida* or other yet undescribed and closely related species requires further clarification.

Due to the very similar size of *Az. dexteroporum*, *Am. parvula*, and *Am. languida*, all three species as identified with SEM are present in the one category of small amphidomatacean cells used for light microscopic analysis and quantification.

#### A5 Other Amphidomataceae

In the SEM, a few other cells were observed which can also be assigned to the genus *Amphidoma*. The epitheca found in dorsal view in Fig. A9e, with its characteristic pore ridges on the large apical plates, corresponds to *Am. alata*, a species






**Figure A5.** SEM of yet unidentified cells of *Azadinium* sp. 1 of the Amphidomatacean bloom stations. (a) Cell in left-lateral ventral view. (b) Cell in dorsal view. (c) Cell in apical view. (d–f) Cells in dorsal view. Note the small antapical spine (white arrow in a, b, d–f) and the lack of contact between plates 1a and 1'' (kofoidian plate label notation) visible in (b)–(d) and (f). Scale bars = 2  $\mu\text{m}$ .


which was described from the Argentina shelf (Tillmann, 2018b). The cell depicted in Fig. A9f in ventral view likely represents a new species of *Amphidoma*. The cells in Fig. A9g–i likely correspond to *Am. trioculata*, another species described from Argentina (Tillmann, 2018b), although assigning an isolated hypotheca (Fig. A9i) is difficult.

Moreover, for a more complete description of the diversity of Amphidomataceae in the bloom sample, the following individual findings (Fig. A10) should also be mentioned: a cell in dorsal view (Fig. A10a) had a very distinct antapical spine, relatively large apical plates, and a small six-sided central intercalary plate. This combination of features has not been described in any known *Azadinium* species, suggesting that this may represent a new species. The cell depicted in Fig. A10b had a distinct antapical spine, a ventral pore on the right side of the pore plate, and a very short first apical plate, with the anterior sulcal plate extending far into the epitheca. Both cells in Fig. A10b and c resemble *Az. spinosum* but differ in that the ventral pore is centrally located within plate 1' in the apical area. They likely correspond to the cells designated as *Azadinium* sp. 3 in Fig. 14b and c, in Tillmann (2018). The cell in Fig. A10e had a ventral pore on the left side of plate 1' (like *Az. spinosum* and *Az. obesum*), but here the lateral apical plates were significantly


larger than in these species. The cell in Fig. A10f in dorsal view in terms of size and shape might correspond to *Az. asperum* described from the Argentine shelf (Tillmann, 2018a), undoubtedly an *Azadinium* species due to the apical pore and intercalary plates, does not match any previously described species based on size ( $\sim 20 \mu\text{m}$  cell length) and shape. This, along with the other cells in Fig. A9, which likely represents previously undescribed species due to the unique combination of features, highlights the great diversity of Amphidomataceae in this 2021 bloom sample.




**Figure A6.** LM (a) and SEM (b–f) of cells of the Amphidomatacean bloom stations identified as *Azadinium dexteroporum*. (b, c) Cells in ventral view. (d–f) Cells in dorsal view. Note the small size, the distinct antapical spine, the position of the ventral pore on the right side of the pore plate (white arrows in b, c), and the concave central intercalary plate 2a visible in (d)–(f). Scale bars = 2  $\mu\text{m}$ .



**Figure A7.** LM (a) and SEM (b–f) of cells of the Amphidomatacean bloom stations identified as *Amphidoma parvula*. (b, c) Cells in ventral view. (d) Cell in dorsal view. (e) Cell in ventral antapical view. (f) Cell in apical view. Note the small size, the flat hypotheca, the shape of the 1' plate visible in (a) and (b), the group of pores in the second antapical plate (white arrow in e), and the relatively long apical plates visible in (d) and (f). Scale bars = 2  $\mu\text{m}$ .




**Figure A8.** LM (a) and SEM (b–f) of cells of the Amphidomatacean bloom stations identified as *Amphidoma languida*. (b, c) Cells in ventral view. (d) Cell in dorsal view. (e) Cell with epitheca in ventral view and hypotheca in antapical view. (f) Epitheca in lateral apical view. (g) Epitheca in apical view. (h) Hypotheca in antapical view. Note the shape of the 1' plate visible in (b), the distinct antapical pore in the second antapical plate (white arrows in d, e, h), and the relatively short apical plates visible in (d), (f), (g). Also note that there are only single pores on precingular plates (small white arrows in c, e, f, and g). Scale bars = 2  $\mu\text{m}$ .



**Figure A9.** SEM of unidentified cells of *Amphidoma* spp. of the Amphidomatacean bloom stations. (a–d) Cells in apical (a, c) or dorsal (b, d) view resembling *Amphidoma languida* but with multiple pores in precingular plates (small white arrows in a–d). (e) Cell of *Amphidoma* sp. in dorsal view. Note the very long apical plates. The row of pores with a distinct rim on the apical plates (white arrow in e) resemble *Amphidoma alata*. (f) Cell of an unidentified *Amphidoma* sp. 1 in ventral view. Note the long apical plates, the ventral depression (white arrow in f), and the row of pores on the posterior suture of apical plates. (g, h) Two cells of *Amphidoma* sp. resembling *Amphidoma trioculata*. (i) Hypotheca in antapical view of an unidentified *Amphidoma*. Note the multiple pores on the plates and the presence of a very small antapical pore on the second antapical plate (white arrow in i). Scale bars = 2  $\mu$ m.





**Figure A10.** SEM of unidentified cells of *Azadinium* spp. of the Amphidomatacean bloom stations. **(a)** Cell in dorsal view. Note the very small and five-sided intercalary plate (white arrow in **a**). **(b)** Cell of *Azadinium* sp. 1 in ventral view. Note the very short 1' plate and the position of the ventral pore inside the pore plate (white arrow in **b**). It may be assumed but is not clear if **(a)** is the dorsal view of such an *Azadinium* sp. 1. **(c, d)** Two cells in ventral view of *Azadinium* sp. 2. Note the position of the ventral pore in apical position inside of plate 1' (white arrows in **c, d**). **(e)** Cell of *Azadinium* sp. 3 in lateral ventral view. Note the position of the ventral pore (white arrow in **e**) and the rather long apical plates. **(f)** Dorsal view of *Azadinium* sp. 4 resembling in size and shape *Azadinium asperum*. Scale bars = 2  $\mu$ m.

## Appendix B


Species diversity based on ITS1-based metabarcoding.

**Table B1.** Species detected by amplicon sequence variant (ASV) reads at stations GA01 and AA09.

Nr	species	Genbank	Identity %	GA01	AA09
1	<i>Amphidoma parvula</i>	KY996792	> 98	19	0
2	<i>Amphidoma</i> sp. 1	OQ360107	90–95	158	200
3	<i>Amphidoma</i> sp. 2	LC788745	90–95	16	0
5	<i>Azadinium dalianese</i> ribotype E	LS974150	90–95	41	63
			95–98	33	58
			> 98	93.775	98.503
6	<i>Azadinium dalianense</i> ribotype B	MF033117	95–98	9	0
			> 98	0	19
7	<i>Azadinium</i> sp. 1	OQ360091	90–95	36	41
8	<i>Azadinium</i> sp. 2	OQ360094	> 98	6	5
9	<i>Azadinium spinosum</i> ribotype C	MK405512	> 98	17.006	7.589
10	<i>Azadinium spinosum</i> ribotype B	LS974169	> 98	0	37
11	<i>Ansanella</i> sp.	MN604385	90–95	4	13
12	<i>Bicheleria</i> sp.	KC895487	90–95	12	52
13	<i>Bicheleria cincta</i>	KC895487	> 98	6	17
14	<i>Blastodinium oviforme</i>	JX473680	95–98	2	0
15	<i>Karlodinium decipiens</i>	LC521288	95–98	0	7
			> 98	885	106
16	<i>Karlodinium digitatum</i>	MN133932	> 98	3	0
17	<i>Kirithra asteri</i>	MW267275	> 98	41	186
18	<i>Pelagodinium beii</i>	KP843723	> 98	48	77

Note that a total of 849 ASVs were identified; however, only 118 of these were successfully annotated at the species level (with over 97 % similarity). This limitation may be attributed to the inadequacy of the ITS region database used in our study, which may have affected the taxonomic resolution of certain sequences.






**Figure B1.** Relative abundance (in %) of the Amphidomataceae species detected with metabarcoding targeting ITS1 regions at stations GA01 (a) and AA09 (b).

## Appendix C


Overall appearance of the pure Amphidomataceae bloom in fixed field samples



**Figure C1.** Micrographs of the Amphidomataceae bloom taken under light microscopy at low magnification: 200 $\times$ . Only (e) was taken under 400 $\times$  and (p) was taken using phase contrast in 100 $\times$ . The toxic *Azadinium spinosum* ribotype B is indicated in a red circle. The arrows indicate other protists: *Og*: *Oxytoxum gracile*, *K-type*: *Karenia*-type cell, *K*: *Katodinium* sp., *T*: *Tripos* sp., *C*: ciliate with a cell of Amphidomataceae inside, *Mr*: *Mesodinium rubrum*.

## Appendix D

Retention of particles in the blooming area.



**Figure D1.** Background colours: absolute dynamic topography (m) averaged from 16 to 25 November. Black lines correspond to the advection of particles for the same period after release at the points indicated with a black dot on 16 November.

**Data availability.** The CTD data and the abundance of prokaryotic species counted under light microscopy at the sampling stations GA01 and AA09 are publicly available after request. Data from the Gayoso cruise are available at <https://doi.org/10.1594/PANGAEA.971564> (Ferronato et al., 2024). Sequences obtained in this study are available at the National Center for Biotechnology Information, Sequence Read Archive (<http://www.ncbi.nlm.nih.gov/sra>, last access: 9 July 2025, project number PRJNA1289085).

**Video supplement.** The video showing the daily evolution from 10 to 25 November 2021 of the finite-size Lyapunov exponent (FSLE) ridges in the area of the two locations with the Amphidomataceae bloom: GA01, pink dot, sampled on 16 November; and AA09, blue dot, sampled on 25 November. The two stations remained within the same water mass separated by two maxima FSLE (Guinder, 2025).

**Author contributions.** VAG conceptualized and designed the study, coordinated the planning and execution of fieldwork and laboratory work, and secured funding. VAG and UT analysed the plankton samples using LM and SEM. MR processed the satellite-derived chlorophyll data and the Lyapunov coefficients. CF and FR conducted field research and processed the CTD data. VAG and BK processed the toxin samples. HG processed the DNA samples. MS analysed the geostrophic currents and performed the particle tracking modelling. All co-authors contributed to the interpretation of the results. VAG prepared the paper, with contributions from all co-authors.

**Competing interests.** The contact author has declared that none of the authors has any competing interests.

**Disclaimer.** Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

**Acknowledgements.** Valeria Ana Guinder and Martin Saraceno acknowledge the BioMMar consortium and the common grant received from the Argentinean Initiative Pampa Azul (PIDT A6) to carry out the Gayoso cruise. Valeria Ana Guinder acknowledges the academic mobility projects FitoxNorPat, CONICET-DAAD call-2020, and CoastCarb EUH2020-Marie Skłodowska-Curie (grant agreement no. 872690). Valeria Ana Guinder especially thanks Marcelo Acha from INIDEP, Argentina, for his invitation to participate in the WG of the Marine Priority Area of Agujero Azul, from the Pampa Azul Initiative. Valeria Ana Guinder also thanks Annegret Müller and Thomas Max for their help and guidance in the laboratory work on processing toxin samples; and Valeria Ana Guinder, Urban Tillmann, and Haifeng Gu thank Annegret Müller for DNA extraction. The authors thank the Argentinean authorities, the captain and crew of the R/V *Bernardo Houssay* (Prefectura Naval Argentina), and the R/V *Victor Angelescu* (INIDEP).

**Review statement.** This paper was edited by Sebastian Naeher and reviewed by Leonardo Guzman and one anonymous referee.

## References

- Abraham, E. R., Law, C. S., Boyd, P. W., Lavender, S. J., Maldonado, M. T., and Bowie, A. R.: Importance of stirring in the development of an iron-fertilized phytoplankton bloom, *Nature*, 407, 727–730, <https://doi.org/10.1038/35037555>, 2000.
- Akselman, R. and Negri, R. M.: Blooms of *Azadinium* cf. *spinosum* Elbrächter et Tillmann (Dinophyceae) in northern shelf waters of Argentina, Southwestern Atlantic, *Harmful Algae*, 19, 30–38, <https://doi.org/10.1016/j.hal.2012.05.004>, 2012.
- Akselman, R., Krock, B., Alpermann, T. J., Tillmann, U., Borel, C. M., Almandoz, G. O., and Ferrario, M. E.: *Protoceratium reticulatum* (Dinophyceae) in the austral Southwestern Atlantic and the first report on YTX-production in shelf waters of Argentina, *Harmful Algae*, 45, 40–52, <https://doi.org/10.1016/j.hal.2015.03.001>, 2015.
- Aleman, D., Zavatteri, A., Prandoni, N., and Giussi, A. R.: Fisheries in the Patagonian shelf break Front, Chap. 7, in: *The Patagonian shelf-break Front. Ecology, Fisheries, Wildlife Conservation*, edited by: Acha, M., Iribarne, O., and Piola, A., Springer Nature, ISBN: 978-3-031-71189-3, [https://doi.org/10.1007/978-3-031-71190-9\\_7](https://doi.org/10.1007/978-3-031-71190-9_7), 2024.
- Almandoz, G. O., Cefarelli, A. O., Diodato, S., Montoya, N. G., Benavides, Carignan, M., Hernando, M., Fabro, E., Metfies, K., Lundholm, N., Schloss, I. R., Alvarez, M., and Ferrario, M. E.: Harmful phytoplankton in the Beagle Channel (South America) as a potential threat to aquaculture activities, *Mar. Poll. Bull.*, 145, 105–117, <https://doi.org/10.1016/j.marpolbul.2019.05.026>, 2019.
- Anderson, C. R., Moore, S. K., Tomlinson, M. C., Silke, J., and Cusack, C. K.: Living with harmful algal blooms in a changing world: strategies for modelling and mitigating their effects in coastal marine ecosystems, in: *Coastal and marine hazards, risks, and disasters*, Elsevier, 495–561, ISBN: 978-0-12-396483-0, 2015.
- Anderson, D. M., Fensin, E., Gobler, C. J., Hoeglund, A. E., Hubbard, K. A., Kulis, D. M., Landsberg, J. H., Lefebvre, K. A., Provoost, P., Richlen, M. L., Smith, J. L., Solow, A. R., and Trainer, V. L.: Marine harmful algal blooms (HABs) in the United States: history, current status and future trends, *Harmful Algae*, 102, 101975, <https://doi.org/10.1016/j.hal.2021.101975>, 2021.
- Balch, W. M., Drapeau, D. T., Bowler, B. C., Lyczkowski, E. R., Lubelczyk, L. C., Painter, S. C., and Poulton, A. J.: Surface biological, chemical, and optical properties of the Patagonian Shelf coccolithophore bloom, the brightest waters of the Great Calcite Belt, *Limnol. Oceanogr.*, 59, 1715–1732, <https://doi.org/10.4319/lo.2014.59.5.1715>, 2014.
- Becker, F., Romero, S. I., and Pisoni, J. P.: Detection and characterization of submesoscale eddies from optical images: a case study in the Argentine continental shelf, *Int. J. Remote Sens.*, 44, 3146–3159, <https://doi.org/10.1080/01431161.2023.2216853>, 2023.
- Beron-Vera, F. J., Bodnariuk, N., Saraceno, M., Olascoaga, M. J., and Simionato, C.: Stability of the Malvinas current, *Chaos*, 30, 13152, <https://doi.org/10.1063/1.5129441>, 2020.
- Cadaillon, A. M., Mattered, B., Albizzi, A., Montoya, N., Maldonado, S., Rey, A. R., Riccialdelli, L., Almandoz, G. O., and Schloss, I. R.: Multispecies mass mortality in the Beagle Channel associated with paralytic shellfish toxins, *Harmful Algae*, 132, 102581, <https://doi.org/10.1016/j.hal.2024.102581>, 2024.
- Carreto, J. I., Montoya, N. G., Carignan, M. O., Akselman, R., Acha, E. M., and Derisio, C.: Environmental and biological factors controlling the spring phytoplankton bloom at the Patagonian shelf break front—Degraded fucoxanthin pigments and the importance of microzooplankton grazing, *Prog. Oceanogr.*, 146, 1–21, <https://doi.org/10.1016/j.pocean.2016.05.002>, 2016.
- Combes, V. and Matano, R. P.: The Patagonian shelf circulation: Drivers and variability, *Prog. Oceanogr.*, 167, 24–43, <https://doi.org/10.1016/j.pocean.2018.07.003>, 2018.
- Corredor-Acosta, A., Galán, A., Saldías, G. S., Mardones, J. I., Medellín-Mora, J., Frangopulos, M., Shiozaki, T., Harada, N., González, H. E., and Iriarte, J. L.: Oceanic phytoplankton structure off western Patagonia during the austral summer: Implications for harmful algal blooms, *Prog. Oceanogr.*, 231, 103409, <https://doi.org/10.1016/j.pocean.2024.103409>, 2025.
- D’Agostino, V. C., Krock, B., Degradi, M., Sastre, V., Santinelli, N., Krohn, T., and Hoffmeyer, M. S.: Occurrence of toxigenic microalgal species and phycotoxin accumulation in mesozooplankton in northern Patagonian gulfs, Argentina, *Environ. Toxicol. Chem.*, 38, 2209–2223, <https://doi.org/10.1002/etc.4538>, 2019.
- Delgado, A. L., Hernández-Carrasco, I., Combes, V., Font-Muñoz, J., Pratolongo, P. D., and Basterretxea, G.: Patterns and trends in chlorophyll *a* concentration and phytoplankton phenology in the biogeographical regions of Southwestern Atlantic, *J. Geophys. Res.-Ocean.*, 128, e2023JC019865, <https://doi.org/10.1029/2023JC019865>, 2023.
- Della Penna, A., De Monte, S., Kestenare, E., Guinet, C., and d’Ovidio, F.: Quasi-planktonic behavior of foraging top marine predators, *Sci. Rep.*, 5, 18063, <https://doi.org/10.1038/srep18063>, 2015.
- d’Ovidio, F., Fernández, V., Hernández-García, E., and López, C.: Mixing structures in the mediterranean sea from finite-size lyapunov exponents, *Geophys. Res. Lett.*, 31, 17203, <https://doi.org/10.1029/2004GL020328>, 2004.
- d’Ovidio, F., Isern-Fontanet, J., López, C., Hernández-García, E., and García-Ladona, E.: Comparison between Eulerian diagnostics and finite-size Lyapunov exponents computed from altimetry in the Algerian basin, *Deep-Sea Res. Pt. I*, 56, 15–31, <https://doi.org/10.1016/j.dsr.2008.07.014>, 2009.
- Fabro, E., Almandoz, G. O., Krock, B., and Tillmann, U.: Field observations of the dinoflagellate genus *Azadinium* and azaspiracid toxins in the south-west Atlantic Ocean, *Mar. Freshw. Res.*, 71, 832–843, <https://doi.org/10.1071/MF19124>, 2019.
- Ferronato, C., Berden, G., Rivarossa, M., and Guinder, V. A.: Wind-driven currents and water masses shape spring phytoplankton distribution and composition in hydrologically complex, productive shelf waters, *Limnol. Oceanogr.*, 68, 2195–2210, <https://doi.org/10.1002/lno.12413>, 2023.
- Ferronato, C., Guinder, V. A., Gilabert, A., Lopez-Abbate, C., Nocera, A., Loizaga, R., and D’Agostino, V.: Hydrographic data during Ana Maria Gayoso cruise in the South west-

- ern Atlantic during late spring 2021, PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.971564>, 2024.
- Ferronato, C., Guinder, V. A., Rivarossa, M., Saraceno, M., Ibarbalz, F., Tillmann, U., Almandoz, G., Bourdin, G., D'Agostino, V., Gilabert, A., Loizaga, R., López Abbate, C., Nocera, A. C., Silva, R., and Flombaum, P.: Insights into protistan plankton blooms in the highly dynamic Patagonian shelf and adjacent ocean basin in the Southwestern Atlantic, *J. Geophys. Res.-Ocean.*, 130, e2024JC021412, <https://doi.org/10.1029/2024JC021412>, 2025.
- Frey, D. I., Piola, A. R., and Morozov, E. G.: Convergence of the Malvinas current branches near 44° S, *Deep-Sea Res. Pt. I*, 196, 104023, <https://doi.org/10.1016/j.dsr.2023.104023>, 2023.
- García, V. M. T., García, C. A. E., Mata, M. M., Pollery, R. C., Piola, A. R., Signorini, S. R., McClain, C. R., and Iglesias-Rodríguez, M. D.: Environmental factors controlling the phytoplankton blooms at the Patagonia shelf break in spring, *Deep-Sea Res. Pt. I*, 55, 1150–1166, <https://doi.org/10.1016/j.dsr.2008.04.011>, 2008.
- GEBCO Compilation Group: General Bathymetric Chart of the Ocean, GEBCO 2021 grid, [https://www.gebco.net/data\\_and\\_products/historical\\_data\\_sets/#gebco\\_2021](https://www.gebco.net/data_and_products/historical_data_sets/#gebco_2021) (last access: 19 September 2023), 2021.
- Genitsaris, S., Stefanidou, N., Sommer, U., and Moustaka-Gouni, M.: Phytoplankton blooms, red tides and mucilaginous aggregates in the urban Thessaloniki Bay, Eastern Mediterranean, *Diversity*, 11, 136, <https://doi.org/10.3390/d11080136>, 2019.
- Giddings, S. N., MacCready, P., Hickey, B. M., Banas, N. S., Davis, K. A., Siedlecki, S. A., Trainer, V. J., Kudela, R. M., Pelland, N. A., and Connolly, T. P.: Hindcasts of potential harmful algal bloom transport pathways on the Pacific Northwest coast, *J. Geophys. Res.-Ocean.*, 119, 2439–2461, <https://doi.org/10.1002/2013JC009622>, 2014.
- Gilabert, A. S., López-Abbate, C., Flombaum, P., Unrein, F., Arbilla, L. A., Garzón-Cardona, J. E., Martínez, A. M., Ibarbalz, F. M., Vincent, F., Saraceno, M., Ruiz-Etcheverry, L. A., Ferronato, C., Guinder, V. A., Silva, R., Ubrig, R. A., D'Agostino, V., Loizaga, R., and Lara, R. J.: Planktonic drivers of carbon transformation during different stages of the spring bloom at the Patagonian Shelf-break front, Southwestern Atlantic Ocean, *Biogeochemistry*, 168, 1–25, <https://doi.org/10.1007/s10533-024-01192-6>, 2025.
- Glibert, P. M.: Margalef revisited: a new phytoplankton mandala incorporating twelve dimensions, including nutritional physiology, *Harmful Algae*, 55, 25–30, <https://doi.org/10.1016/j.hal.2016.01.008>, 2016.
- Guinder, V. A.: Finite Size Lyapunov Exponents on the Patagonian shelf and their role in retaining phytoplankton blooms, SEANO Sea Scientific Open Data Publication [video], <https://doi.org/10.17882/107582>, 2025.
- Guinder, V. A., Ferronato, C., Segura, V., Dogliotti, A., and Lutz, V.: The phytoplankton of the Patagonian shelf break Front, in: *The Patagonian shelf-break Front. Ecology, Fisheries, Wildlife Conservation*, edited by: Acha, M., Iribarne, O., and Piola, A., Aquatic Ecology Series, 13, Springer, Cham, [https://doi.org/10.1007/978-3-031-71190-9\\_3](https://doi.org/10.1007/978-3-031-71190-9_3), 2024.
- Guinder, V. A., Tillmann, U., Krock, B., Delgado, A., Garzón-Cardona, J. E., Katja, M., López Abbate, M. C., Silva, R., and Lara, R.: Plankton multiproxy analyses in the Northern Patagonian Shelf, Argentina: community structure, phycotoxins and characterization of toxic *Alexandrium* strains, *Front. Mar. Sci.*, 5, 394, <https://doi.org/10.3389/fmars.2018.00394>, 2018.
- Guinder, V. A., Malits, A., Ferronato, C., Krock, B., Garzón-Cardona, J. E., and Martínez, A.: Microbial plankton configuration in the epipelagic realm from the Beagle Channel to the Burdwood Bank, a Marine Protected Area in Sub-Antarctic waters, *Plos One*, 15, e0233156, <https://doi.org/10.1371/journal.pone.0233156>, 2020.
- Hallegraeff, G. M., Anderson, D. M., Belin, C., Bottein, M. Y. D., Bresnan, E., Chinain, M., Enevoldsen, H., Iwataki, M., Karlson, B., McKenzie, C. H., Sunesen, I., Pitcher, G. C., Provoost, P., Richardson, A., Schweibold, L., Tester, P. A., Trainer, V. L., Yniguez, A. T., and Zingone, A.: Perceived global increase in algal blooms is attributable to intensified monitoring and emerging bloom impacts, *Commun. Earth Environ.*, 2, 1–10, <https://doi.org/10.1038/s43247-021-00178-8>, 2021.
- Haller, G. and Beron-Vera, F. J.: Geodesic theory of transport barriers in two-dimensional flows, *Phys. D*, 241, 1680–1702, <https://doi.org/10.1016/j.physd.2012.06.012>, 2012.
- Hansen, H. P. and Koroleff, F.: Determination of Nutrients, in: *Methods of Seawater Analysis*, edited by: Grasshoff, K., Kremling, K., and Ehrhardt, M., Wiley, 159–228, <https://doi.org/10.1002/9783527613984.ch10>, 1999.
- Hasle, G. R.: The Inverted-Microscope Methods, in: *Phytoplankton Manual*, edited by: Sournia, A., United Nations Educational, Scientific, and Cultural Organization, Paris, 88–96, 1978.
- Hernández-Carrasco I., Alou-Font E., Dumont P. A., Cabornero A., Allen J., and Orfila A.: Lagrangian flow effects on phytoplankton abundance and composition along filament-like structures, *Prog. Oceanogr.*, 189, 102469, <https://doi.org/10.1016/j.pocean.2020.102469>, 2020.
- Iriarte, J. L., Pizarro, G., and Frangopulos, M.: Harmful algal blooms in Patagonian fjords and channels systems: Recent advances, gaps, and priorities in a changing ocean, *Prog. Oceanogr.*, 217, 103087, <https://doi.org/10.1016/j.pocean.2023.103087>, 2023.
- Jeffrey S. T. and Humphrey G. F.: New spectrophotometric equations for determining chlorophylls *a*, *b*, *c*<sub>1</sub> and *c*<sub>2</sub> in higher plants, algae and natural phytoplankton, *Biochem. Physiol. Pflanz.*, 167, 191–194, [https://doi.org/10.1016/S0015-3796\(17\)30778-3](https://doi.org/10.1016/S0015-3796(17)30778-3), 1975.
- Kelley, D. E., Richards, C., and Layton, C.: oce: An R package for oceanographic analysis, *J. Open Source Softw.*, 7, 3594, <https://doi.org/10.21105/joss.03594>, 2022.
- Krock, B., Tillmann, U., Tebben, J., Trefault, N., and Gu, H.: Two novel azaspiracids from *Azadinium poporum*, and a comprehensive compilation of azaspiracids produced by Amphidomataceae, (Dinophyceae), *Harmful Algae*, 82, 1–8, <https://doi.org/10.1016/j.hal.2018.12.005>, 2019.
- Kuwata, K., Lum, W. M., Takahashi, K., Benico, G., Takahashi, K., Lim, P. T., Leaw, C. P., Uchida, H., Ozawa, M., Matsushima, R., Watanabe, R., Suzuki, T., and Iwataki, M.: Phylogeny and ultrastructure of a non-toxic dinoflagellate *Amphidoma fulgens* sp. nov. (Amphidomataceae, Dinophyceae), with a wide distribution across Asian Pacific, *Harmful Algae*, 138, 102701, <https://doi.org/10.1016/j.hal.2024.102701>, 2024.
- Lago, L. S., Saraceno, M., Piola, A. R., and Ruiz-Etcheverry, L. A.: Volume transport variability on the northern argentine continental shelf from in situ and satellite altimetry



- try data, *J. Geophys. Res.-Ocean.*, 126, e2020JC016813, <https://doi.org/10.1029/2020JC016813>, 2021.
- Lehahn, Y., d'Ovidio, F., Lévy, M., and Heifetz, E.: Stirring of the northeast Atlantic spring bloom: A Lagrangian analysis based on multisatellite data, *J. Geophys. Res.-Ocean.*, 112, C08005, <https://doi.org/10.1029/2006JC003927>, 2007.
- Lehahn, Y., Koren, I., Sharoni, S., d'Ovidio, F., Vardi, A., and Boss, E.: Dispersion/dilution enhances phytoplankton blooms in low-nutrient waters, *Nat. Commun.*, 8, 14868, <https://doi.org/10.1038/ncomms14868>, 2017.
- Lévy, M., Franks, P. J., and Smith, K. S.: The role of submesoscale currents in structuring marine ecosystems, *Nat. Commun.*, 9, 4758, <https://doi.org/10.1038/s41467-018-07059-3>, 2018.
- Lin, S., Litaker, R. W., and Sunda, W. G.: Phosphorus physiological ecology and molecular mechanisms in marine phytoplankton, *J. Phycol.*, 52, 10–36, <https://doi.org/10.1111/jpy.12365>, 2016.
- Liu, M., Tillmann, U., Ding, G., Wang, A., and Gu, H.: Metabarcoding revealed a high diversity of Amphidomataceae (Dinophyceae) and the seasonal distribution of their toxigenic species in the Taiwan Strait, *Harmful Algae*, 124, 102404, <https://doi.org/10.1016/j.hal.2023.102404>, 2023.
- López-Rivera, A., O'callaghan, K., Moriarty, M., O'Driscoll, D., Hamilton, B., Lehane, M., James, K. J., and Furey, A.: First evidence of azaspiracids (AZAs): a family of lipophilic polyether marine toxins in scallops (*Argopecten purpuratus*) and mussels (*Mytilus chilensis*) collected in two regions of Chile, *Toxicon*, 55, 692–701, <https://doi.org/10.1016/j.toxicon.2009.10.020>, 2010.
- Mahadevan, A.: The impact of submesoscale physics on primary productivity of plankton, *Annu. Rev. Mar. Sci.*, 8, 161–184, <https://doi.org/10.1146/annurev-marine-010814-015912>, 2016.
- Mangolte, I., Lévy, M., Haëck, C., and Ohman, M. D.: Sub-frontal niches of plankton communities driven by transport and trophic interactions at ocean fronts, *Biogeosciences*, 20, 3273–3299, <https://doi.org/10.5194/bg-20-3273-2023>, 2023.
- Maritorena, S., d'Andon, O. H. F., Mangin, A., and Siegel, D. A.: Merged satellite ocean color data products using a bio-optical model: Characteristics, benefits and issues, *Remote Sens. Environ.*, 114, 1791–1804, <https://doi.org/10.1016/j.rse.2010.04.002>, 2010.
- Martinetto, P., Alemany, D., Botto, F., Mastrángelo, M., Falabella, Acha, E. M., Antón, G., Bianchi, A., Campagna, C., Cañete, G., Filippo, P., Iribarne, O., Laterra, P., Martínez, P., Negri, R., Piola, A. R., Romero, S. I., Santos, D., and Saraceno, M.: Linking the scientific knowledge on marine frontal systems with ecosystem services, *Ambio*, 49, 541–556, <https://doi.org/10.1007/s13280-019-01222-w>, 2019.
- Matano, R. P., Palma, E. D., and Piola, A. R.: The influence of the Brazil and Malvinas Currents on the Southwestern Atlantic Shelf circulation, *Ocean Sci.*, 6, 983–995, <https://doi.org/10.5194/os-6-983-2010>, 2010.
- McGirr, S., Clarke, D., Kilcoyne, Silke, J., and Touzet, N.: Co-localisation of Azaspiracid Analogs with the Dinoflagellate Species *Azadinium spinosum* and *Amphidoma languida* in the Southwest of Ireland, *Microb. Ecol.*, 83, 635–646, <https://doi.org/10.1007/s00248-021-01777-w>, 2022.
- Palma, E. D., Matano, R. P., and Piola, A. R.: A numerical study of the southwestern Atlantic shelf circulation: Stratified Ocean response to local and offshore forcing, *J. Geophys. Res.*, 113, C11010, <https://doi.org/10.1029/2007jc004720>, 2008.
- Piola, A. R., Bodnariuk, N., Combes, V., Franco, B. C., Matano, R. P., Palma, E. D., Romero, S. I., Saraceno, M., and Urricariet, M. M.: Anatomy and dynamics of the Patagonia shelf break front, in: *The Patagonian shelf-break Front. Ecology, Fisheries, Wildlife Conservation*, edited by: Acha, M., Iribarne, O., and Piola, A., Springer Nature, ISBN: 978-3-031-71189-3, 2024.
- Pitcher, G. C., Figueiras, F. G., Hickey, B. M., and Moita, M. T.: The physical oceanography of upwelling systems and the development of harmful algal blooms, *Prog. Oceanogr.*, 85, 5–32, <https://doi.org/10.1016/j.pocean.2010.02.002>, 2010.
- Ramírez, F. J., Guinder, V. A., Ferronato, C., and Krock, B.: Increase in records of toxic phytoplankton and associated toxins in water samples in the Patagonian Shelf (Argentina) over 40 years of field surveys, *Harmful Algae*, 118, 102317, <https://doi.org/10.1016/j.hal.2022.102317>, 2022.
- Raine, R.: A review of the biophysical interactions relevant to the promotion of HABs in stratified systems: the case study of Ireland, *Deep-Sea Res. Pt. II*, 101, 21–31, <https://doi.org/10.1016/j.dsr2.2013.06.021>, 2014.
- Rossi, R., Dell'Aversano, C., Krock, B., Ciminiello, P., Percopo, I., Tillmann, U., Soprano, V., and Zingone, A.: Mediterranean *Azadinium dexteroporum* (Dinophyceae) produces AZA-35 and six novel azaspiracids: a structural study by a multi-platform mass spectrometry approach, *Anal. Bioanal. Chem.*, 409, 1121–1134, <https://doi.org/10.1007/s00216-016-0037-4>, 2017.
- Salas, R., Tillmann, U., Gu, H., Wietkamp, S., Krock, B., and Clarke, D.: Morphological and molecular characterization of multiple new *Azadinium* strains revealed a high diversity of non-toxicogenic species of Amphidomataceae (Dinophyceae) including two new *Azadinium* species in Irish waters, North East Atlantic, *Phycol. Res.*, 69, 88–115, <https://doi.org/10.1111/pre.12448>, 2021.
- Salyuk, P. A., Mosharov, S. A., Frey, D. I., Kasyan, V. V., Ponomarev, V. I., Kalinina, O. Y., Morozov, E. G., Latushkin, A. A., Sapozhnikov, P. V., Ostroumova, S. A., Lipinskaya, N. A., Budyansky, M. V., Chukmasov, P. V., Krechik, V. A., Uleysky, M. Y., Fayman, P. A., Mayor, A. Y., Mosharova, I. V., Chernetsky, A. D., Shkorba, S. P., and Shved, N. A.: Physical and biological features of the waters in the outer Patagonian shelf and the Malvinas Current, *Water*, 14, 3879, <https://doi.org/10.3390/w14233879>, 2022.
- Saraceno, M., Bodnariuk, N., Ruiz-Etcheverry, L. A., Berta, M., Simionato, C. G., Beron-Vera, F. J., and Olascoaga, M. J.: Lagrangian characterization of the southwestern Atlantic from a dense surface drifter deployment., *Deep-Sea Res. Pt. I*, 208, 104319, <https://doi.org/10.1016/j.dsr.2024.104319>, 2024.
- Ser-Giacomi, E., Martinez-Garcia, R., Dutkiewicz, S., and Follows, M. J. A.: Lagrangian model for drifting ecosystems reveals heterogeneity-driven enhancement of marine plankton blooms, *Nat. Commun.*, 14, 6092, <https://doi.org/10.1038/s41467-023-41469-2>, 2023.
- Smayda, T. J.: Turbulence, watermass stratification and harmful algal blooms: an alternative view and frontal zones as “pelagic seed banks”, *Harmful Algae*, 1, 95–112, [https://doi.org/10.1016/S1568-9883\(02\)00010-0](https://doi.org/10.1016/S1568-9883(02)00010-0), 2002.
- Sunesen, I., Méndez, S. M., Mancera-Pineda, J. E., Dechraoui-Bottein, M. Y., and Enevoldsen, H.: The Latin America and Caribbean HAB status report based on OBIS and HAE-

- DAT maps and databases. *Harmful Algae* 102, 101920, <https://doi.org/10.1016/j.hal.2020.101920>, 2021.
- Tillmann, U.: Amphidomataceae, in: *Harmful Algal Blooms: A compendium Desk Reference*, edited by: Shumway, S. E., Burkholder, J. M., and Morton, S. L., Wiley, Singapore, 575–582, ISBN: 978-1-118-99467-2, 2018a.
- Tillmann, U.: Electron microscopy of a 1991 spring plankton sample from the Argentinean shelf reveals the presence of four new species of Amphidomataceae (Dinophyceae), *Phycol. Res.*, 66, 269–290, <https://doi.org/10.1111/pre.12225>, 2018b.
- Tillmann, U., Salas, R., Gottschling, M., Krock, B., O'Driscoll, D., and Elbrächter, M.: *Amphidoma languida* sp. nov. (Dinophyceae) reveals a close relationship between *Amphidoma* and *Azadinium*, *Protist*, 163, 701–719, <https://doi.org/10.1016/j.protis.2011.10.005>, 2012.
- Tillmann, U. and Akselman, R.: Revisiting the 1991 algal bloom in shelf waters off Argentina: *Azadinium luciferelloides* sp. nov. (Amphidomataceae, Dinophyceae) as the causative species in a diverse community of other amphidomataceans, *Phycol. Res.*, 64, 160–175, <https://doi.org/10.1111/pre.12133>, 2016.
- Tillmann, U., Trefault, N., Krock, B., Parada-Pozo, G., De la Iglesia, R., and Vásquez, M.: Identification of *Azadinium poporum* (Dinophyceae) in the Southeast Pacific: morphology, molecular phylogeny, and azaspiracid profile characterization, *J. Plank. Res.*, 39, 350–367, <https://doi.org/10.1093/plankt/fbw099>, 2017a.
- Tillmann, U., Sánchez-Ramírez, S., Krock, B., and Bernal-Jiménez, A.: A bloom of *Azadinium polongum* in coastal waters off Peru, *Rev. Biol. Mar. Oceanogr.*, 52, 591–610, <https://doi.org/10.4067/S0718-19572017000300015>, 2017b.
- Tillmann, U., Gottschling, M., Guinder, V., and Krock, B.: *Amphidoma parvula* (Amphidomataceae), a new planktonic dinophyte from the Argentine Sea, *Eur. J. Phycol.*, 53, 14–28, <https://doi.org/10.1080/09670262.2017.1346205>, 2018.
- Tillmann, U., Gottschling, M., Krock, B., Smith, K. F., and Guinder, V. A.: High abundance of Amphidomataceae (Dinophyceae) during the 2015 spring bloom of the Argentinean Shelf and a new, non-toxicogenic ribotype of *Azadinium spinosum*, *Harmful Algae*, 84, 244–260, <https://doi.org/10.1016/j.hal.2019.01.008>, 2019.
- Tillmann, U., Wietkamp, S., Krock, B., Tillmann, A., Voss, D., and Gu, H.: Amphidomataceae (Dinophyceae) in the western Greenland area, including the description of *Azadinium perforatum* sp. nov., *Phycologia*, 59, 63–88, <https://doi.org/10.1080/00318884.2019.1670013>, 2020.
- Tillmann, U., Wietkamp, S., Gu, H., Krock, B., Salas, R., and Clarke, D.: Multiple new strains of Amphidomataceae (Dinophyceae) from the North Atlantic revealed a high toxin profile variability of *Azadinium spinosum* and a new non-toxicogenic Az. cf. *spinosum*, *Microorganisms*, 9, 134, <https://doi.org/10.3390/microorganisms9010134>, 2021.
- Turner, A. D. and Goya, A. B.: Occurrence and profiles of lipophilic toxins in shellfish harvested from Argentina, *Toxicon*, 102, 32–42, <https://doi.org/10.1016/j.toxicon.2015.05.010>, 2015.
- Wells, M. L., Karlson, B., Wulff, A., Kudela, R., Trick, C., Asnaghi, V., Berdalet, E., Cochlan, W., Davidson, K., De Rijcke, M., Dutkiewicz, S., Hallegraeff, G., Flynn, K. J., Legrand, C., Paerl, H., Silke, J., Suikkanen, S., Thompson, P., and Trainer, V. L.: Future HAB science: Directions and challenges in a changing climate, *Harmful algae*, 91, 101632, <https://doi.org/10.1016/j.hal.2019.101632>, 2020.
- Wietkamp, S., Krock, B., Gu, H., Voß, D., Klemm, K., and Tillmann, U.: Occurrence and distribution of Amphidomataceae (Dinophyceae) in Danish coastal waters of the North Sea, the Limfjord, and the Kattegat/Belt area, *Harmful Algae*, 88, 101637, <https://doi.org/10.1016/j.hal.2019.101637>, 2019.
- Wietkamp, S., Krock, B., Clarke, D., Voß, D., Salas, R., Kilcoyne, J., and Tillmann, U.: Distribution and abundance of azaspiracid-producing dinophyte species and their toxins in North Atlantic and North Sea waters in summer 2018, *PLoS One*, 15, e0235015, <https://doi.org/10.1371/journal.pone.0235015>, 2020.
- Wilson, C., Sastre, A. V., Hoffmeyer, M., Rowntree, V. J., Fire, S. E., Santinelli, N. H., Ovejero, S. D., D'Agostino, V., Marón, C. F., Doucette, G., J., Broadwater, M. H., Wang, Z., Montoya, N., Seger, J., Adler, F. R., Sironi, M., and Uhart, M. M.: Southern right whale (*Eubalaena australis*) calf mortality at Península Valdés, Argentina: Are harmful algal blooms to blame?, *Mar. Mammal. Sci.*, 32, 423–451, <https://doi.org/10.1111/mms.12263>, 2015.