



Impacts of eutrophication and deoxygenation on the sediment biogeochemistry in the Sea of Marmara

İsmail Akçay^{1,2}, Süleyman Tuğrul¹, Hasan Örek¹, Devrim Tezcan¹, Koray Özhan¹, Barış Salihoğlu¹, and Mustafa Yücel¹

¹Institute of Marine Sciences, Middle East Technical University, P.O. Box 28, 33731 Erdemli-Mersin, Türkiye

²Department of Basic Sciences of Fisheries, Mersin University-Faculty of Fisheries, Yenişehir-Mersin, Türkiye

Correspondence: İsmail Akçay (iakcay@mersin.edu.tr)

Received: 17 March 2025 – Discussion started: 25 March 2025

Revised: 29 July 2025 – Accepted: 29 July 2025 – Published: 25 September 2025

Abstract. The biogeochemistry of seafloor sediments can be significantly altered in response to deoxygenation and eutrophication-driven organic carbon production, resulting in increased benthic fluxes of dissolved nutrients (such as ammonia and phosphate) and metals. The Sea of Marmara, which has also faced large-scale mucilage outbreaks in recent years, is undergoing severe eutrophication and deoxygenation, but the consequences on sediment biogeochemistry and benthic feedback have not been studied so far. This study aims to understand the impacts of deoxygenation and coastal eutrophication on sedimentary biogeochemical processes in the Sea of Marmara, which experiences varying degrees of anthropogenic pressure along with natural inputs from the adjacent Black Sea via Bosphorus surface inflows. Multicore-obtained undisturbed sediment core samples indicate that oxic respiration no longer plays a significant role in Marmara sediments, but denitrification, metal reduction, and sulfate respiration are prevalent as respiratory pathways. The deep-water sediments become more reducing in the Eastern Marmara compared to the Western part of the sea. Cores from İzmit Bay, the Easternmost region in Marmara, exhibit permanently sulfidic conditions with anaerobic oxidation of methane (AOM) controlling downcore sulfide profiles. Calculated diffusive benthic nutrient fluxes show markedly high phosphate and ammonium fluxes into the near-bottom waters of highly eutrophic areas of the Eastern Marmara, which are expected to enhance primary production in the upper layer during the dry season. On the other hand, these sediments are a net sink for nitrate due to the denitrification. Benthic nutrient dynamics contribute to the accumulation of organic matter as well as shifting N/P ratios and the development

of a steep hypoxic zone at halocline boundary depths. Additionally, we show that sediments are already influenced by widespread hypoxia in the Sea of Marmara and benthic-pelagic coupling has enhanced the existing eutrophication problem, analogous to the benthic “vicious cycle” observed in the shallower Baltic Sea. We conclude that the Sea of Marmara is now on a clear path towards being included within the list of famous “dead zones” of the Earth’s oceans, such as the Baltic Sea, the Gulf of Mexico, or Chesapeake Bay. For the marine management efforts and ecosystem modeling studies, hypoxia-induced benthic biogeochemical processes and benthic-pelagic coupled cycling of nutrients in the Sea of Marmara should be considered.

1 Introduction

The Sea of Marmara is a connecting basin between the Black Sea and the Mediterranean, but it is in a state of ecosystem decline. Increasing eutrophication and decreasing deep-water oxygen levels, exacerbated by recent large-scale mucilage outbreaks (Yücel et al., 2021; Savun-Hekimoglu and Gazioglu, 2021), have a common cause of excessive nutrient levels in the system. Despite this trend, only a few studies were performed to understand seawater nutrient cycling and organic matter dynamics (Tuğrul and Morkoç, 1989; Tuğrul and Polat, 1995; Polat and Tuğrul, 1995; Polat et al., 1998; Tuğrul et al., 2002; Ediger et al., 2016; Yalçın et al., 2017). The connection between benthic fluxes and eutrophication has received little attention, with most studies on sedimentary biogeochemical processes focused on the effect of the

tectonics (Le Pichon et al., 2001; Géli et al., 2008), and gas-induced porewater geochemistry (Evans et al., 1989; Halbach et al., 2002; Çağatay et al., 2004; Sarı and Çağatay, 2010; Tryon et al., 2010; Ruffine et al., 2018; Yang et al., 2018). Here, we present the first process-based study performed in the Sea of Marmara to understand impacts of eutrophication and rapid deoxygenation on the sediment porewater nutrient dynamics and solid-state geochemistry in the context of coupled biogeochemical cycling of key nutrients (N, P) coupled to oxygen and metal (Fe, Mn) cycles (Williams, 1987; Jørgensen, 1996).

The Sea of Marmara, a two-layer semi-enclosed sea, connects the Black Sea to the Mediterranean via the two shallow and narrow straits, called Istanbul (Bosphorus) and Çanakkale (Dardanelles) Straits (Ünlüata et al., 1990; Beşiktepe et al., 1994; Ediger et al., 2016) (Fig. 1). Marmara's two-layer ecosystem has distinct biogeochemical properties due to water exchanges between the Black Sea and the Mediterranean Sea (Polat and Tuğrul, 1995; Tuğrul and Polat, 1995; Tuğrul et al., 2002). The Sea of Marmara has been highly influenced by the Black Sea inflow and anthropogenic activities, including urban effluents, agricultural run-offs, industrial inputs, fishing, and shipping in recent decades (Tuğrul and Morkoç, 1989; Tuğrul and Polat, 1995; Albayrak et al., 2006; Ediger et al., 2016; Yalçın et al., 2017; Tan and Aslan, 2020). During the 1970s, intensive eutrophication in the Northwestern Black Sea collapsed the Black Sea ecosystem and fisheries (Mee, 1992; Tuğrul et al., 2014), leading to similar changes in the Sea of Marmara during the 1980s (Polat and Tuğrul, 1995; Ediger et al., 2016; Yalçın et al., 2017). Marmara's two-layer density gradient, established permanently between 10–30 m depths, limits physical mixing (Beşiktepe et al., 1994), leading to limited exchange of dissolved nutrients and oxygen through the steep pycnocline (Tuğrul et al., 2002; Ediger et al., 2016; Yalçın et al., 2017). As a result, dissolved oxygen (DO) concentrations, at near-saturation in the productive upper layer of the Black Sea origin, decrease rapidly in the permanent steep pycnocline (15–20 m) coinciding with the nutricline, allowing limited influx of oxygen to upper depths of sub-halocline waters though ship-induced turbulence may occur in the northeast part of the Sea of Marmara due to regional high traffic densities (Nylund et al., 2021). Thus, a major oxygen source for the Sea of Marmara lower layer is the oxygen-rich Mediterranean inflow via Dardanelles undercurrent. This leads to a west-to-east decrease of the dissolved oxygen of the deep Marmara waters (Tuğrul and Polat, 1995; Tuğrul et al., 2002; Ediger et al., 2016; Yalçın et al., 2017). This oxygen deficiency has reached hypoxic ($< 2 \text{ mg L}^{-1}$, or $< 65 \text{ } \mu\text{M}$) to near anoxic ($< 0.5 \text{ mg L}^{-1}$, $< 10 \text{ } \mu\text{M}$) levels in the deep basin (1200 m) of the Eastern Marmara (Yücel et al., 2021; ÇŞİDB, ODTÜ-DBE, 2022; Yücel et al., 2023b) and central part of the İzmit Bay (depth $< 150 \text{ m}$) due to increasing inputs of nutrients of different origin from Marmara basin, leading to development of eutrophication in the bays, coastal and open waters

of Marmara Sea (Fig. 1). Enhanced particulate organic matter (POM) export from these regions to the deeper waters has steadily consumed limited oxygen there, with eastern locations such as İzmit Bay and in the Çınarcık basin progressively becoming anoxic since the late 1980s (Yücel et al., 2023b). These long-term changes are very likely to induce previously insignificant redox-dependent processes in the oxygen-deficient deep waters and muddy sediments (Ediger et al., 2016; Yalçın et al., 2017; Tan and Aslan, 2020).

As a result, Marmara is now a strong candidate to be included on the list of Earth's well-known hypoxic to anoxic zones such as Gulf of Mexico, Chesapeake Bay or the Baltic Sea; but the consequences of this phenomenon on ecosystem functioning, including within the sedimentary system, has been poorly tackled so far. Here, we aim to fill a significant gap by investigating the consequences of recent eutrophication and deoxygenation on the sediment biogeochemistry of the Sea of Marmara. The specific aims of this study are (i) to determine the impacts of eutrophication and development of hypoxic and anoxic conditions (deoxygenation) in the near bottom water on the porewater nutrient dynamics, (ii) to understand solid state organic matter (C, N) geochemistry and carbon burial, (iii) to describe early diagenetic processes in the porewaters and sediments and (iv) to determine the porewater diffusive nutrient (Si, N, P) fluxes in the three different regions of the Marmara Sea: İzmit Bay, Çınarcık Basin and the southern Marmara Sea.

2 Methodology

2.1 Materials and Methods

Main physical and biogeochemical parameters were measured at 97 stations in the Sea of Marmara (Fig. 1) in the winter and summer of 2019 by R/V *Bilim-2* of METU-IMS. Sediment core samples were obtained with a multicorer (Octopus, Kiel) from 13 selected stations in the İzmit Bay, Çınarcık Basin, and southern Marmara Sea. The upper parts of the sediment columns were generally brown in color with a fine-grained sediment texture. However, at below 6–8 cm depth, the sediments, specifically in the Çınarcık Basin and İzmit Bay, became darker suggesting the reducing conditions. Core slices were used to measure porewater nutrients (PO_4 , Si, $\text{NO}_3 + \text{NO}_2$, NH_4), major ions (Cl, SO_4 , Br, Li, Na, K, Mg, Ca), hydrogen sulfide (H_2S), dissolved iron (dFe), solid-state total carbon / organic carbon (TC / TOC) and nitrogen (TN) concentrations.

The temperature and salinity of the water column were measured *in situ* by a SEABIRD 9plus CTD probe that was coupled to a 12-Niskin bottle Rosette System. The Secchi Disk Depth (SDD), an estimation of water transparency, was collected from each water column station (Fig. 1). In addition to the CTD sensor-derived measurements, dissolved oxygen measurements were also carried out by the Win-




Figure 1. The Sea of Marmara with (a) the inset, showing the tectonic framework leading to the formation of three deep basins (a; Tryon et al., 2010) and (b) main panel, outlining the water column sampling locations in winter and summer of 2019. Red circles represent sediment core stations. Source: Reiner Schlitzer, Ocean Data View, <https://odv.awi.de/>, last access: 18 December 2024

kler titration (UNEP/MAP, 2005). Total phosphorus concentrations were determined by the wet digestion methods and using the conventional colorimetric detection at 880 nm wavelength (Strickland and Parsons, 1972; Grasshoff et al., 1983) after persulfate oxidation of seawater samples (Menzel and Corwin, 1965). Water column and porewater dissolved inorganic nutrients (nitrate + nitrite, ammonium, phosphate, and silicate) were measured using a Bran + Luebbe Model four-channel Autoanalyzer (Grasshoff et al., 1983). The detection limits of dissolved nutrients were 0.05, 0.04, 0.01, 0.04, and 0.03 μM for nitrate + nitrite, ammonium, phosphate, silicate, and total phosphorus, respectively. Chlorophyll *a* (Chl *a*) measurements were carried out by the conventional spectrofluorometric method after digestion of filtered samples by 90 % (*v/v*) acetone solution using a HITACHI model F-2500 fluorescence spectrophotometer (Strickland and Parsons, 1972; UNEP/MAP, 2005). Hydrogen sulfide ($\text{H}_2\text{S}/\text{HS}^-$) concentrations in seawater and porewater samples were measured on board by the spectrophotometric methylene blue method at 670 nm wavelength (Cline, 1969). Porewater dissolved iron (sum of Fe^{2+} and Fe^{3+}) concentrations were measured on board by spectrophotometry, after reducing all $\text{Fe}(\text{III})$ to $\text{Fe}(\text{II})$ by dithionite, followed by ferrozine-based assay and detection of the color-producing Fe complex at 562 nm wavelength (Stookey, 1970; Jeitner, 2014).

Sediment core samples were obtained by a multiple-corer (Oktopus, Kiel). Each sediment core was sliced on board under anoxic conditions using N_2 or Ar gas. After obtaining sliced sediments, each sample was put in a 50 mL falcon tube and centrifuged at 6000 rpm for 20 min. Then, the porewa-

ter samples were extracted using syringe-coupled GF/F filters through 0.45 μm . Dissolved iron and hydrogen sulfide concentrations of porewater sub-samples were immediately determined on board by the methods mentioned above, and the remained sub-samples were placed in 15 mL falcon tubes and stored at -20°C for the analysis of dissolved nutrients and major ions. Analyses of porewater nutrients were carried out by the same method as water column samples. Analysis of major ions (Cl^- , SO_4^{2-} , Li^+ , Na^+ , K^+ , Mg^{2+} , Ca^{2+}) in porewater samples was conducted by ion chromatography.

Solid phase sediment samples were also stored for the determination of TC, TOC, and TN concentrations and surface sediment porosity measurements. Sediment samples for the TC, TOC, and TN measurements were initially freeze-dried. Dry sediments were then powdered and sieved on 63 μm pore size for the homogenization of the samples. For TOC analysis, about 30 mg of dry and homogeneous sediment samples were put into the silver cups, which had been pre-combusted at 400°C for 6 h. Then, 10 μL of distilled water was added into each silver cup to wet the samples. After the addition of distilled water, 10 μL of 20 % HCl (*v/v*) was added to remove inorganic carbon from the sediment samples. The HCl additions were continued until all weak-acid soluble carbonates were removed. Then, the samples were dried at $60\text{--}70^\circ\text{C}$ for 24 h. After drying the sediment samples, silver cups were compacted and put into the autosampler of the CHN analyzer. TC and TN analyses in sediments were performed by the same method as for the TOC analysis, but without acid addition. TN concentrations were also determined on HCl-added sediment samples, and there was no significant difference between HCl-treated and untreated samples, indicat-

ing that a significant fraction of the TN pool was in the organic nitrogen form. Therefore, TN concentrations of HCl-untreated sediment samples were reported in this study. Total carbon (TC), organic carbon (TOC), and nitrogen (TN) concentrations were determined by the dry oxidation method using the Vario El Cube Elementar Model CHN Analyzer (UNEP/MAP, 2006). Porosity of wet sediments was determined by the displacement method as described by Mu et al. (2017).

2.2 Calculation of Diffusive Nutrients (Si, N, P) Fluxes

We have calculated the diffusive nutrient fluxes based on Fick's first law of diffusion (Cheng et al., 2014; Mu et al., 2017):

$$F = \phi D_s (dC/dz) \quad (1)$$

Where F corresponds to the diffusive flux across the sediment-water interface, dC/dz to the concentration gradient of nutrients across the sediment-water interface, ϕ to the porosity of sediment, and D_s to the actual molecular diffusion coefficient corrected for the sediment tortuosity.

Ullman and Aller (1982) proposed an empirical formula related to the actual diffusion coefficient, D_s , and porosity, ϕ :

$$\begin{aligned} D_s &= \phi D_0 & (\phi < 0.7) \\ D_s &= \phi^2 D_0 & (\phi > 0.7) \end{aligned} \quad (2)$$

where D_0 is the self-diffusion coefficient of ions at infinite dilution corrected by *in situ* temperature. For phosphate, nitrate, nitrite and ammonium, D_0 values were taken from Li and Gregory (1974) and for silicate, D_0 values were taken from Rebreanu et al. (2008) as $7.34 \times 10^{-6} \text{ cm}^2 \text{ s}^{-1}$ for PO_4 , $19.0 \times 10^{-6} \text{ cm}^2 \text{ s}^{-1}$ for NO_3 , $19.1 \times 10^{-6} \text{ cm}^2 \text{ s}^{-1}$ for NO_2 , $19.8 \times 10^{-6} \text{ cm}^2 \text{ s}^{-1}$ for NH_4 and $11.7 \times 10^{-6} \text{ cm}^2 \text{ s}^{-1}$ for Si at 25 °C, respectively.

3 Results and Discussion

3.1 General Oceanographic State in the Sea of Marmara in 2019

We describe the general oceanography of the system based on the two expeditions performed in the Sea of Marmara in the summer and winter periods of 2019 by R/V *Bilim-2*. Physical and chemical oceanographic properties of the two-layer Marmara Sea ecosystem are principally dominated by the brackish (S : 17.0–17.5) Black Sea and the salty Mediterranean inflow (S : 35–38) through the Dardanelles (Ünlüata et al., 1990; Beşiktepe et al., 1994; Polat and Tuğrul, 1995; Tuğrul and Polat, 1995; Polat et al., 1998; Tuğrul et al., 2002). Driven by the physical properties of the exchange flows, temperatures of the upper layer waters ranged from 6.93–10.98 °C in winter to 19.06–26.17 °C in summer for

the study period. The surface water salinity values increased from 17.29 in the Bosphorus exit to 29.40 in the entry to Dardanelles (Table 1, Fig. 2). Water transparency increased from the Eastern Marmara (SDD 4–5 m) towards Dardanelles (SDD 17–18 m) in the summer months. The winter SDDs displayed less pronounced spatial variations, ranging between 4 and 11 m (Table 1). Lowest values were recorded in the Eastern Marmara, notably in the Çınarcık Basin and the İzmit Bay (Fig. 2).

Altogether, our observations on the state of the Sea of Marmara as of 2019 are in line with the previous reports, pointing out to the combined effect of brackish Black Sea inflow and land-based sources (Tuğrul and Morkoç, 1989; Polat and Tuğrul, 1995; Tuğrul and Polat, 1995; Polat et al., 1998; Ediger et al., 2016; Yalçın et al., 2017; Tan and Aslan, 2020). We have found that the brackish surface waters of Black Sea origin were relatively rich in NO_x (NO_x : $\text{NO}_2 + \text{NO}_3$) and PO_4 concentrations in winter, and this enrichment is due to enhanced inputs of nutrients from the lower layer by vertical mixing processes (Tuğrul et al., 2002; Ediger et al., 2016; Yalçın et al., 2017). Similar seasonal and regional variations were also recorded in the concentrations of NH_4 and reactive Si, with the maximum values recorded at the Bosphorus southern exit and Eastern Marmara Sea; İzmit Bay and Çınarcık Basin (Fig. 2). The surface water chlorophyll a (Chl a) concentrations increased towards eastern basin (Fig. 2) and from summer ($0.01\text{--}1.81 \mu\text{g L}^{-1}$) to winter ($0.26\text{--}3.77 \mu\text{g L}^{-1}$) due to enhanced inputs of inorganic nutrients from lower layer and Black Sea in winter (Tuğrul and Polat, 1995). According to eutrophication assessment based on the Chl a values (Simboura et al., 2005), trophic status of the Eastern Marmara upper layer and bays were classified as “mesotrophic/moderate” to “dystrophic/bad” levels of ecological status. Enhanced primary productivity (in terms of Chl a) in the upper layer has led to development of eutrophication and hypoxia in the lower layer over the basin and development of anoxic conditions in deep waters of the Eastern Marmara Sea and İzmit Bay (Fig. 2). In the Çınarcık Basin of Eastern Marmara Sea (max depth 1270 m), dissolved oxygen was entirely depleted and anoxia developed in the deep waters particularly below 750 m. Development of similar anoxic conditions were consistently observed in the central deep waters of the İzmit Bay in the last two decades as well. Here, limited dissolved oxygen inputs from the oxygen-depleted Eastern Marmara deep waters remained insufficient for aerobic oxidation of POM, leading to denitrification and sulfate reduction processes and thus hydrogen sulfide accumulation in the bottom waters of the bay during summer months (Yücel et al., 2023a).

3.2 Nutrient Trends in the Deep Water Column and Sediment Porewaters

A comparison of averages of data from > 1000 m water depth in the Çınarcık Basin indicated an accumulation of

Table 1. Summary of surface water physical and biogeochemical parameters in the Sea of Marmara in the winter and summer of 2019.

Winter	Temperature (°C)	Salinity	TP (μM)	PO ₄ (μM)	NO _x (μM)	NH ₄ (μM)	Si (μM)	DO (μM)	Chl <i>a</i> ($\mu\text{g L}^{-1}$)	SDD (m)
Mean	8.80	24.15	0.48	0.19	2.19	1.11	5.56	296.3	1.24	8.2
SD	1.28	4.39	0.23	0.18	2.55	1.46	4.10	31.6	0.79	1.8
Min.	6.93	17.96	0.16	0.02	0.09	0.15	0.85	195.3	0.26	4.0
Max.	10.98	28.83	1.11	0.72	9.97	8.84	16.19	344.4	3.77	11.0
N	39	39	39	39	39	39	39	39	39	12
Summer	Temperature (°C)	Salinity	TP (μM)	PO ₄ (μM)	NO _x (μM)	NH ₄ (μM)	Si (μM)	DO (μM)	Chl <i>a</i> ($\mu\text{g L}^{-1}$)	SDD (m)
Mean	24.27	23.00	0.53	0.14	0.92	0.30	1.54	235.0	0.23	11.9
SD	1.37	2.73	0.46	0.40	2.16	0.10	1.26	9.7	0.24	2.9
Min.	19.06	17.29	0.14	0.02	0.05	0.02	0.20	188.1	0.01	4.5
Max.	26.17	29.40	3.45	3.27	9.28	0.64	7.08	254.1	1.81	17.5
N	84	84	84	84	84	84	84	84	84	45

reactive Si and PO₄ but consumption of NO_x (via denitrification) in the oxygen-depleted deep waters of Marmara (Fig. 3). These findings indicate enhanced inputs of POM from the highly productive layer to the lower layer during at least the last three decades, and subsequent burial in the sediment phase.

In order to address the apparent long-term changes in the nutrient and organic matter cycles in the oxygen-depleted deep-waters (Fig. 3), sediment core samples were obtained from three key regions in the Marmara: the highly eutrophic İzmit Bay, eutrophic Çınarcık Basin displaying permanently anoxic properties in the deep waters, and from the mesotrophic/eutrophic southern Marmara Sea shelf with still-oxygen-rich bottom waters. The bottom water physical and biogeochemical variables of the core stations presented in Table 2 and also surface water spatial variability of the parameters (Fig. 2) indicated that the selected regions displayed significantly different physico-chemical and biogeochemical properties, allowing the discerning of redox- and organic matter-driven effects on sediment biogeochemistry.

Porewater nutrient concentrations throughout the obtained sediment core samples ranged between 0.47 and 58.2 μM for PO₄, 0.34 and 91.5 μM for NO₃, 0.17 to 666 μM for NH₄ and 9.02 to 335 μM for Si, respectively (Fig. 4). Biogeochemical cycling of key nutrients (N, P) is highly linked to dissolved oxygen concentrations and metal (Mn, Fe) cycles due to rapid degradation of labile organic matter by the redox-processes as denitrification, manganese and iron reduction as well as sulfate reduction and methanogenesis. These respiratory pathways in the uppermost centimeters of sediment column result in net nutrient efflux from the sediments (Williams, 1987; Christensen et al., 1988; Jørgensen, 1996; Ignatieva, 1999; Rasheed, 2004; Al-Rousan et al., 2004; Hille et al., 2005; Rasheed et al., 2006; Rydin et al., 2011; Cheng et al., 2014; Mu et al., 2017). Expectedly, maximum porewater PO₄, NH₄, and Si concentrations were mea-

sured in the İzmit Bay (Fig. 4), with the vertical profiles of nutrients displaying an increasing trend with depth throughout the POM-enriched muddy sediment. Porewater NO₃ concentrations were highest in the porewaters of core samples (0–5 cm) from the southern Marmara Sea having oxic conditions in the deep waters and lower surface water primary productivity (in terms of Chl *a*) (Fig. 2). Vertical profiles of porewater NO₃ concentrations in the studied sites suggested that denitrification process dominated the uppermost millimeters of sediments of the reducing İzmit Bay and Çınarcık Basin while oxic respiration was the major process for the organic matter remineralization in the upper 15–20 cm of sediment cores obtained in southern Marmara Sea.

Diffusive nutrient fluxes across the sediment-water interface were calculated based on the Fick's first law of diffusion in the Sea of Marmara (Table 3). Calculated porewater diffusive fluxes showed spatial variability in the Sea of Marmara. In all sites, sediments acted as a source for PO₄, NH₄ and Si, whilst sediments in the highly eutrophic and anoxic/sulfidic sites of the İzmit Bay acted as a sink for NO_x (referred to NO₃ + NO₂) due to rapid utilization of NO₃ ion as a terminal electron acceptor by the denitrification process. Porewater diffusive PO₄, NH₄ and Si fluxes from the sediment to bottom waters were highest in the İzmit Bay with the anoxic bottom waters whereas the negative sign of the diffusive NO_x fluxes here indicated NO_x loss by denitrification. The diffusive fluxes of PO₄, NH₄ and Si to the bottom water decreased in the Çınarcık basin, and further minimized in the southern Marmara Sea sediments in line with increasing bottom water dissolved oxygen levels (Table 3). These flux estimations indicate that the increase in diffusive fluxes of dissolved inorganic nutrients (PO₄, NH₄, Si), reactive iron and manganese in the organic-rich muddy shelf sediments of the anoxic regions are likely to lead to nutrient accumulation in bottom waters and contribute to the upper layer chemical fluxes through vertical mixing across the studied basins, in-




Figure 2. Distributions of surface water salinity, dissolved nutrients, Secchi Disk Depth (SDD) and deep (near-bottom) water dissolved oxygen content (DWDO) in the Sea of Marmara in the summer of 2019. Source: Reiner Schlitzer, Ocean Data View, <https://odv.awi.de/>, last access: 18 December 2024


tensifying during winter (Konovalov et al., 2007; Noffke et al., 2012; Mu et al., 2017).

The relationship between calculated diffusive nutrient fluxes and the deep water dissolved oxygen concentrations clearly indicated the redox-dependent benthic nutrient mobilization in the Sea of Marmara (Fig. 5); the lower deep water dissolved oxygen concentrations resulted in the higher porewater diffusive PO_4 , NH_4 , and Si fluxes in the POM-rich muddy sediments. The apparent denitrification in sediment pore waters must have contributed to the apparent deep-water nitrate loss signal in the Çınarcık Basin during the last decades (Fig. 3). Furthermore, the diffusive nutrient fluxes also in the anoxic benthic interface of the highly produc-

tive NE shelf region of the Marmara Sea and eutrophic inner İzmit bay might further enhance eutrophication in the NE Marmara shelf and bays by vertical mixing processes (Table 3). The Si/N and N/P molar ratios of the measured nutrients in the shelf and deep basin waters of Marmara Sea ranged between 1.8 and 5.5 for Si/N and 5.6–12 for N/P, respectively. The Si/N molar ratios of the diffusive nutrient fluxes (Si/N: 1.0–5.5) were similar to the ratios in deep waters of the studied sites whereas the N/P molar ratios of diffusive nutrient fluxes (8.4–31) were greater than deep water nutrient (N/P) molar composition; but comparable with the N/P molar ratios calculated in the diffusive nutrient fluxes in

Table 2. Bottom water physico-chemical properties of the sediment core stations.

Region	Station	Date	Depth (m)	Temperature (°C)	Salinity	TP (μM)	PO ₄ (μM)	NO _x (μM)	Si (μM)	NH ₄ (μM)	DO (μM)
İzmit Bay	IZ-30	5 Jan 2019	29	15.88	38.63	0.99	0.96	8.44	20.35	0.78	54.1
	1	24 Jul 2019	63	15.41	38.72	1.15	1.06	9.45	21.39	0.54	45.6
	2	24 Jul 2019	111	15.27	38.77	1.76	1.03	10.32	23.08	0.43	35.9
	IZ-2	5 Jan 2019	157	15.63	38.76	1.34	1.13	8.34	26.75	1.18	30.3
	3	24 Jul 2019	238	14.56	38.71	1.52	1.04	7.66	33.34	0.40	7.8
	4	24 Jul 2019	425	14.46	38.70	1.28	0.85	5.42	26.12	0.28	10.3
Çınarcık Basin	5	25 Jul 2019	98	15.32	38.80	0.95	0.85	9.06	20.79	0.22	57.5
	6	25 Jul 2019	190	14.73	38.73		0.97	9.62	31.02	0.36	17.2
	7	25 Jul 2019	288	14.51	38.70		1.06	8.32	35.69	0.17	7.2
	8	25 Jul 2019	407	14.47	38.70		1.29	7.00	40.02	0.24	6.3
Southern Marmara	EK-12	6 Jan 2019	45	16.22	38.68	0.53	0.40	4.31	8.65	0.47	164.1
	EK-10	6 Jan 2019	56	15.80	38.84	0.78	0.66	6.49	15.69	0.56	118.4
	MD-18	6 Jan 2019	145	15.24	38.80	1.09	0.69	6.49	17.21	0.19	51.6

**Figure 3.** Summary of the deep water (> 1000 m, averages) biogeochemical shift in the Çınarcık Basin, Eastern Marmara Sea (Yücel et al., 2021; ÇŞİDB, ODTÜ-DBE, 2022; Yücel et al., 2023b).

the Baltic Sea (N/P: 20–30) (Ignatieva, 1999; van Helmond et al., 2020).

3.3 Sediment Organic Matter Geochemistry

Close examination of vertical profiles for the geochemical variables clearly showed that solid-state total carbon (TC), total organic carbon (TOC) and total nitrogen (TN) concentrations in the upper 40–50 cm of sediment from different regions of the Sea of Marmara displayed spa-

tial variability (Fig. 6). TC concentrations varied regionally from 1.97 to 4.12 mmol g⁻¹ dw (dry weight) in the upper 3 cm of the obtained core samples and decreased to 0.92–3.25 mmol g⁻¹ dw levels below 30 cmbs (Fig. 6). Maximum concentrations of TOC and TN were determined in the surface sediments from the İzmit Bay where anoxic and sulfidic conditions have developed in the deep waters (Fig. 6). TOC values were in the range of 0.47–2.70 mmol g⁻¹ dw while TN values varied between 0.04 and 0.23 mmol g⁻¹ dw. The apparent decrease in the TOC and TN concentrations with the increasing depth throughout the sediment column indicated organic matter deposition exceeding the decomposition in the top 8–20 cm of the sediment column. Below this depth, TOC and TN concentrations were relatively constant and did not change noticeably, suggesting the dominance of refractory organic matter below 20 cmbs buried in the anoxic sediments. The Çınarcık and İzmit bay bottom sediments had higher TOC levels compared to other regions in the Sea of Marmara, in line with their more reducing deep-water conditions, also in agreement with the previous studies (Evans et al., 1989; Çağatay et al., 2004; Sarı and Çağatay, 2010; Ruffine et al., 2018; Yang et al., 2018; Tan and Aslan, 2020). Maximum TOC and TN concentrations were detected in the surface sediments of the İzmit Bay due to excess anthropogenic inputs from point and diffuse land sources (Tuğrul and Morkoç, 1989; Yalçın et al., 2017; Tan and Aslan, 2020). The study findings further show the development of anoxic conditions in the bottom waters and the organic matter degradation mainly by denitrification and sulfate reduction in the bottom water and underlying sediments in the Çınarcık Basin and İzmit Bay. Our results also report for the first time the accumulation of H₂S (starting from 0.52 μM at 29 m) in the bottom waters of the İzmit inner bay (depth > 150 m) in summer months, showing that the bottom waters have shifted to




Figure 4. Porewater nutrient concentrations in the sediment cores from the Sea of Marmara.

Table 3. Diffusive nutrient fluxes estimated from the porewater profiles of different regions having distinct trophic and redox states.

Region	Trophic Status	Deep Water Redox State	PO ₄ (mmol m ⁻² yr ⁻¹)	NO _x (mmol m ⁻² yr ⁻¹)	NH ₄ (mmol m ⁻² yr ⁻¹)	Si (mmol m ⁻² yr ⁻¹)
İzmit Bay ^a	Eutrophic to Dystrophic	Suboxic/Anoxic/Sulfidic	1.81–9.76	–34.2–7.65	18.7–93.5	34.8–128.4
Çınarcık Basin ^a	Eutrophic	Suboxic/Anoxic	0.51–3.03	–9.78–8.98	1.45–25.6	44.1–114
Southern Marmara ^a	Mesotrophic to Eutrophic	Oxic	0.11–3.22	0.65–40.2	10.93–25.4	31.7–44.0
Eastern Mediterranean ^b	Oligotrophic	Oxic	0.60	4.14	6.79	34.51
Red Sea ^c	Oligotrophic	Oxic	1.28	6.32	2.08	20.04
Peruvian OMZ ^d	Eutrophic to Dystrophic	Suboxic/Anoxic	2.3–227.6			
Baltic Sea ^e	Eutrophic to Dystrophic	Suboxic/Anoxic/Sulfidic	0.47–67.82		15.64–1441.23	
Çınarcık Basin ^f	Mesotrophic to Eutrophic	Suboxic	0.60		2.04	

^a This study; ^b Christensen et al. (1988); ^c Rasheed et al. (2006); ^d Noffke et al. (2012); ^e Ignatieva (1999); ^f Çağatay et al. (2004)

reducing conditions as a result of the general environmental degradation in the Sea of Marmara.

The porewater and solid-state results altogether show that maximum sediment TOC and TN concentrations (Fig. 6) in the İzmit Bay might further enhance eutrophication in the

Sea of Marmara, analogous to benthic “vicious cycle” in the much shallower Baltic Sea due to release of increased amount of dissolved nutrients, biologically available for the phytoplankton (Ignatieva, 1999; van Helmond et al., 2020). In the Çınarcık Basin, relatively lower concentrations of TC,




Figure 5. Relationship between calculated diffusive nutrient fluxes and deep-water concentrations of dissolved oxygen and nutrients in the Sea of Marmara. A flux with a positive sign indicated that sediment acts as a source of that particular constituent, whereas a negative sign indicated that sediment acts as a sink.

TOC, and TN were measured compared to eastern İzmit Bay, but comparable to those measured in the western İzmit Bay, having similar deep-water physical and biogeochemical properties (Table 2), indicating redox-dependency on the benthic organic matter dynamics. In the southern Marmara Sea, levels of TOC and TN were also lower compared to eastern İzmit Bay. However, the TC concentrations were comparable with the İzmit Bay and Çınarcık Basin. Maximum porewater nutrients (Si, PO_4 , NH_4) and sediment TOC and TN concentrations were measured in the anoxic and sulfidic İzmit Bay and Çınarcık Basin of the Sea of Marmara, having sedimentation rates ranging between 70 and 190 cm per 1000 years (Ergin and Yörük, 1990; Ergin et al., 1994; Çağatay et al., 2004). In the southern Marmara Sea, with the lower Chl *a* concentrations in the surface waters and more oxygenated deep waters, relatively low concentrations of porewater nutrients and sediment organic matter were measured with a moderately low sedimentation rate of 45 cm per 1000 years (Çağatay et al., 2004). The lower primary productivity (in terms of Chl *a*) and also lower sedimentation rate (Çağatay et al., 2004) in the southern Marmara Sea resulted in low organic matter accumulation at the sediment and the higher deep water dissolved oxygen concentrations (Fig. 2, Table 2) prevented the release of reactive phosphate, silicate, and ammonium fluxes from the surface sediments. There-

fore, development of eutrophication and deep-water hypoxia/anoxia in the Çınarcık Basin and İzmit Bay caused accumulation of excess amount of organic matter in the upper sediment column highly enhancing the release of dissolved nutrients whereas in the oxic southern Marmara Sea, lower sediment organic matter (C, N) concentrations resulted in lower porewater nutrients concentrations and their diffusive fluxes due to high concentrations of deep water dissolved oxygen.

Benthic nutrient fluxes play a critical role in nutrient cycling and productivity in marine ecosystems. Climate change can intensify deoxygenation and enhance ocean stratification, which limits vertical mixing. These changes may disrupt nutrient dynamics and primary productivity more profoundly than diffusive benthic nutrient fluxes. However, hypoxia and anoxia potentially lead to the long-term release and transport of nutrients to the surface layer. Additionally, although natural eutrophication may occur in certain regions, such as Çınarcık Basin due to Black Sea inflow, human-induced nutrient fluxes may further enhance eutrophication. In the context of marine management efforts, it is essential to consider both benthic nutrient fluxes and larger-scale climate impacts to effectively address nutrient dynamics and sustain ecosystem health.




Figure 6. Vertical profiles of TC, TOC and TN in the sediment cores from the Sea of Marmara.

3.4 Sedimentary Early Diagenetic Processes in the Sea of Marmara

In order to complement the above-described solid state and porewater parameters, we further describe and discuss in this section additional parameters that enables us to present a fuller picture of the early diagenetic processes in the Sea of Marmara. Concentrations of Cl, SO_4 , H_2S , Li, Na, K, Mg, Ca and dFe in porewaters of the collected core samples were determined. Although the measured concentrations of the selected ions exhibited minimal spatial variability within the study region, they demonstrated significant variations with depth (Fig. 7). The ranges of the measured concentrations throughout the sediment column were 326–747 mM for Cl, 0.84–32.6 mM for SO_4 , 1.6–3204 μM for H_2S , 8.64–31.7 μM for Li, 313–614 mM for Na, 9.2–15.2 mM for K,

43.9–77 mM for Mg, 3.3–15.8 mM for Ca and 0.2–31.8 μM for dFe, respectively (Fig. 7), similar to previous studies (Çağatay et al., 2004; Tryon et al., 2010; Ruffine et al., 2018). The variations in the porewater concentrations of dissolved nutrients (Fig. 4) and major ions (Fig. 7) throughout the sediment column were affected by a series of microbially mediated diagenetic reactions modifying the geochemical composition of the porewaters in the studied sites.

An important process that needs to be considered while interpreting major ion data in the porewaters of Marmara's surface sediments is the presence of older, lake period-derived porewaters. It was shown that there is a downward freshening gradient of major ions (specifically Cl and Na) due to mixing with the buried brackish water from the Marmara Lake which was observed at all sites in the Sea of Marmara with the exception of Western High and this freshening was more




Figure 7. Vertical profiles of Cl, Na, Ca, Mg, K, Li, dFe, SO_4 and H_2S in the sediment porewaters from the obtained core samples in the Sea of Marmara.

pronounced below 250–300 cmbs in the Marmara Sea (Tryon et al., 2010). As shown in Fig. 7, the lowermost part of our Na and Cl profile (after 35–40 cmbs) started displaying a decreasing trend, supporting the phenomenon of a downcore freshening in the Marmara Sea (Fig. 7). Similar trends were obtained for Mg and K as well, while the decrease in Li and Ca was less pronounced. Still, these data show that most of the porewaters studied in the upper 50 cm of Marmara's current seafloor are well under the influence of the current deep Marmara waters, which are Mediterranean in origin. Hence, we can interpret these data as well as other porewater data in the framework of marine sediment diagenesis.

In that context of major ions discussed above, we can conclude that porewater geochemistry is under the main influence of the series of microbially mediated organic matter degradation reactions such as respiration by NO_3 (denitrification), Fe/Mn oxides, and SO_4 , as well as the processes of the SO_4 depletion by anaerobic oxidation of methane (AOM). Carbonate precipitation and the silicate diagenesis also shaped porewater profiles. Vertical profiles of porewater nutrients (Fig. 4) and solid-state TOC / TN (Fig. 6) concentrations indicated that organic matter degradation processes take place in the upper 10–20 cmbs, introducing high concentrations of porewater reactive Si, PO_4 , and NH_4 and re-

moving oxidized nitrogen (NO_3) through denitrification. The decrease in SO_4 concentrations with depth in the sediments was due to sulfate reduction as well as the AOM, releasing H_2S , Mg, and Ca in the eastern İzmit Bay. The release of Mg and Ca, then, resulted in carbonate precipitation reactions reducing the porewater concentrations of these ions with depth throughout the sediment column in the eastern İzmit Bay (Fig. 7). Similar chemical composition and diagenetic reactions in the porewaters in different regions of the Sea of Marmara have been observed by the studies performed by Halbach et al. (2002), Çağatay et al. (2004), Tryon et al. (2010), Ruffine et al. (2018) and the strong correlation between gas emissions and fast depletion of sulfate in the upper sedimentary column of the Sea of Marmara suggested the important role of AOM for preventing complete releases of methane originating from the North Anatolian Fault (N.A.F.) and associated faults (Géli et al., 2008, Ruffine et al., 2018). The porewater geochemistry of the obtained core samples from this study and previous studies in different regions of the Sea of Marmara (mainly Çınarcık, Central and Tekirdağ Basins, southern Marmara Sea, İzmit Bay) revealed that the sulfate-depletion zone, corresponding to the sulfate-methane transition zone (SMTZ), ranged from a few meters to less than 15 cm below seafloor (Halbach et al., 2002; Çağatay et al., 2004; Tryon et al., 2010; Ruffine et al., 2018). The findings of this study also showed a shallower SMTZ, less than 50 cm, in the eastern İzmit Bay (Fig. 7) due to an increase in AOM reaction rates by intense emissions of methane resulting in fast depletion of sulfate in the upper sedimentary column (Ruffine et al., 2018). It should be also noted that the SMTZ depth in the Sea of Marmara is controlled by multiple factors in addition to the dissolved oxygen of the bottom waters. These controls include the upward flux of methane (from deeper, lake-derived sediments) and sedimentation rates. The upward methane flux is in turn related to tectonic activity and gas hydrate dissociation, which might be controlled by warming of the bottom waters and hydrostatic pressure changes. As reported previously, spatial variability in methane flux is evident across different sub-basins; in particular, the Çınarcık Basin and İzmit Bay exhibit elevated fluxes in proximity to active fault zones, where the SMTZ occurs at or near the seafloor (Bourry et al., 2009; Crémère et al., 2012; Çağatay et al., 2018).

Organic matter degradation by the Fe/Mn oxides released the reduced forms of these metals into the sediment porewaters (Jørgensen, 1996). Though the dissolved manganese was not measured in this study, previous studies performed in the oxic, suboxic, and anoxic regions showed that anoxic conditions support the accumulation of dissolved manganese, whereas oxic sediments become depleted in manganese (Jørgensen, 1996). In the oxic region of the Black Sea, for example, redox transformations and physical processes caused redistribution of manganese from its initial sources across the bottom water, leading to accumulation of manganese in oxidized form in the upper sedimentary column (Konovalov et

al., 2007). A similar process was observed with the relative depletion of reduced iron in sediments in the Southern Marmara Sea having oxic deep waters and its accumulation in the sediments under hypoxic conditions in the Çınarcık and İzmit Bay as also experienced in the oxic, suboxic and anoxic marine environments (Jørgensen, 1996; Konovalov et al., 2007; Yücel et al., 2010) (Fig. 7). According to a study performed in the sediment cores obtained from oxic, suboxic, anoxic/sulfidic parts of the Black Sea (Konovalov et al., 2007), FeS production is faster than pyrite reacting with zerovalent sulfur or H_2S to form FeS_2 . If the Fe(III) is available in the sulfidic sediments, Fe(III) reacts with the H_2S to form Fe(II), which in turn reacts with H_2S to form FeS phases. In the Sea of Marmara, the obtained geochemical profiles indicated that the H_2S produced by AOM formed iron-sulfide precipitates in the presence of reduced iron (Halbach et al., 2002). In this study, dFe (referred to sum of Fe(II) + Fe(III)) concentrations ranged from 0.2 to 31.8 μM in the porewaters of the obtained sediment core samples, displaying a maximum in the upper sedimentary column, decreasing to undetectable values below 10–15 cmbs, indicating iron-sulfur precipitation reactions, as also experienced in the anoxic/sulfidic sediments of the Sea of Marmara (Halbach et al., 2002) and Black Sea (Konovalov et al., 2007). The decrease in SO_4 concentrations with depth in the sediments was due to sulfate reduction as well as the AOM, releasing H_2S , Mg, and Ca, specifically in the eastern İzmit Bay as discussed above. The release of Mg and Ca, then, results in carbonate precipitation reactions reducing the porewater concentrations of these ions with depth throughout the sediment column (Fig. 7). However, the release of Ca might also be the result of low-temperature silicate diagenetic processes as also shown in a recent study performed in the Sea of Marmara (Tryon et al., 2010) since low-temperature silicate diagenetic processes depend not only on the availability of the silicate minerals but also available cations such as K, B, Li, Mg and Na. The end products of these diagenetic reactions, Ca and Sr, are released to the porewaters in the Sea of Marmara (Tryon et al., 2010, and references therein). The downcore decreases in the concentrations of Mg, K, Na and Li obtained from this study were also associated with the low-temperature silicate diagenetic processes in the Sea of Marmara.

4 Conclusions

Our results showed that the dynamics of porewater nutrients, redox-sensitive elements, major ions involved in the diagenetic processes and also solid-state geochemistry in the Sea of Marmara have been influenced by eutrophication and bottom water hypoxia/anoxia. High concentrations of nutrients enhanced eutrophication and hence increased the organic matter (C, N) concentrations accumulated at the sediment. It should also be noted that the sediment organic matter degradation processes differed in the three distinct regions

that we focused in this work. The denitrification and sulfate reduction processes were observed in the Çınarcık Basin and İzmit Bay while oxic respiration was the major process for the organic matter remineralization in the uppermost sedimentary column of the southern Marmara Sea. Moreover, the distribution of porewater sulfate, hydrogen sulfide, and major elements throughout the sediment column obtained specifically from the İzmit Bay, suggested principal biogeochemical and early diagenetic processes such as anaerobic oxidation of methane (AOM), carbonate precipitation, Fe-reduction, Fe-S precipitation, and low-temperature silicate diagenesis.

The dynamics of porewater nutrients, redox-sensitive elements, major ions involved in the diagenetic processes, and also solid-state geochemistry in the Sea of Marmara were found to be clearly influenced by eutrophication and the subsequent bottom water hypoxia/anoxia. Development of eutrophication and deep water suboxia/anoxia has been observed in the Çınarcık Basin and İzmit Bay due to enhanced terrestrial pressures during the last several decades. Porewater nutrients (Si, N, P) results indicated that maximum concentrations of porewater reactive silicate, phosphate and ammonium were measured in the eutrophic and suboxic/anoxic İzmit Bay, having higher primary production (in terms of chlorophyll *a*) resulted in higher diffusive fluxes of nutrients from sediments to the deep waters. Lower porewater nutrient concentrations were recorded in the southern Marmara Sea, having lower primary production and more oxygenated deep waters. Porewater nitrate, sulfate, and hydrogen sulfide concentrations showed that organic matter degradation processes in the upper 20–30 cmbs have occurred by denitrification and sulfate reduction, whilst organic matter decomposition was limited by oxic respiration in the upper centimeters of the sedimentary column in the southern Marmara Sea. The depth distribution of TOC concentrations in the sediment core samples and the calculated redox-dependent benthic nutrient fluxes in the Sea of Marmara indicate high rates of organic matter decomposition and limited trapping of nutrients in the benthic interface. The increased diffusive nutrient inputs, especially in the shelf zone of the eastern basin in recent decades, is expected to have further enhanced eutrophication in the Sea of Marmara, analogous to the benthic “vicious cycle” in the much shallower Baltic Sea. The increased nitrate loss in the lower layer of deep basins and the higher rates of diffusive fluxes of phosphate, ammonium, and reactive silicate in the shelf zones may affect the water column nutrient concentrations and thus their molar ratios. These changes naturally altered the ratios of nutrients reaching the upper layer by physical mixing in the Bosphorus junction and over the basin, leading to further ecological problems such as enhanced eutrophication, deep water anoxia, changes in phytoplankton abundance and composition, and mucilage formation recently observed in the Sea of Marmara. We conclude that if sufficient measures are not implemented, the Sea of Marmara is now on a clear path towards being included within the list of famous “dead zones”

of the Earth, such as the Baltic Sea, Gulf of Mexico or Chesapeake Bay.

Data availability. The data are available upon reasonable request.

Author contributions. İA, ST, HÖ, DT, KÖ, BS, and MY conceptualized and conducted the field measurements. İA and MY conducted the data curation, formal analysis, and visualization. MY supervised this study. The paper was prepared by İA with contributions from all co-authors.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher’s note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors. Also, please note that this paper has not received English language copy-editing.

Acknowledgements. This study has been supported by DEKOSIM (Centre for Marine Ecosystem and Climate Research, Project Code BAP-08-11-DPT.2012K120880) Project, the Scientific and Technological Research Council of Turkey (TUBİTAK-2247, 119C027) Project, The Marmara Integrated Modelling System Project (MARMOD), and TUBA-GEBIP Program of the Turkish Academy of Sciences. We would like to thank the scientific cruise participants, crew of R/V *Bilim-2*, and METU-IMS technical personnel for helping biogeochemical sampling and analysis.

Review statement. This paper was edited by Mark Lever and reviewed by Sergey Konovalov, M. Namık Cagatay, and one anonymous referee.

References

- Albayrak, S., Balkis, H., Zenetos, A., Kurun, A., and Kubanç, C.: Ecological quality status of coastal benthic ecosystems in the Sea of Marmara, *Marine Pollution Bulletin*, 52, 790–799, <https://doi.org/10.1016/j.marpolbul.2005.11.022>, 2006.
- Al-Rousan, S., Rasheed, M., and Badran, M.: Nutrient diffusive fluxes from sediments in the northern Gulf of Aqaba, Red Sea, *Scientia Marina*, 68, 483–490, <https://doi.org/10.3989/scimar.2004.68n4483>, 2004.
- Beşiktepe, S., Sur, H. İ., Özsoy, E., Latif, M. A., Oğuz, T., and Ünlüata, Ü.: Circulation and hydrography of the Marmara Sea, *Prog. Oceanogr.*, 34, 285–334, [https://doi.org/10.1016/0079-6611\(94\)90018-3](https://doi.org/10.1016/0079-6611(94)90018-3), 1994.

- Bourry, C., Chazallon, B., Charlou, J. L., Pierre Donval, J., Ruffine, L., Henry, P., Geli, L., Çağatay, M. N., Inan, S., and Moreau, M.: Free gas and gas hydrates from the Sea of Marmara, Turkey: Chemical and structural characterization, *Chem. Geol.*, 264, 197–206, <https://doi.org/10.1016/j.chemgeo.2009.03.007>, 2009.
- Çağatay, M. N., Yıldız, G., Bayon, G., Ruffine, L., and Henry, P.: Seafloor authigenic carbonate crusts along the submerged part of the North Anatolian Fault in the Sea of Marmara: Mineralogy, geochemistry, textures and genesis, *Deep-Sea Res. Pt. II*, 153, 92–109, <https://doi.org/10.1016/j.dsr2.2017.09.003>, 2018.
- Çağatay, M. N., Özcan, M., and Güngör, E.: Pore-water and sediment geochemistry in the Marmara Sea (Turkey): early diagenesis and diffusive fluxes, *Geochem. Explor. Environ. Anal.*, 4, 213–225, <https://doi.org/10.1144/1467-7873/04-202>, 2004.
- Cheng, X., Zeng, Y., Guo, Z., and Zhu, L.: Diffusion of nitrogen and phosphorus across the sediment-water interface and in seawater at aquaculture areas of Daya Bay, China, *Int. J. Environ. Res. Public Health*, 11, 1557–1572, <https://doi.org/10.3390/ijerph110201557>, 2014.
- Christensen, J. P., Goldsmith, V., Walline, P., Schneller, A., and El Sayed, S. Z.: Sedimentary nutrient regeneration on the oligotrophic eastern Mediterranean continental shelf, *Oceanol. Acta*, Special Issue, 219–234, 1988.
- Cline, J. D.: Spectrophotometric determination of hydrogen sulfide in natural waters, *Limnol. Oceanogr.*, 14, 454–458, 1969.
- Crémière, A., Pierre, C., Blanc-Valleron, M. M., Zitter, T., Çağatay, M. N., and Henry, P.: Methane-derived authigenic carbonates along the North Anatolian fault system in the Sea of Marmara (Turkey), *Deep-Sea Res. Pt. I*, 66, 114–130, <https://doi.org/10.1016/j.dsr.2012.03.014>, 2012.
- ÇŞİDB, ODTÜ-DBE: Marmara Denizi Bütünleşik Modelleme Sistemi FAZ II Projesi (MARMOD – FAZ II) 2022 Yılı Değerlendirme Raporu, Ankara, 2022.
- Ediger, D., Beken, Ç., Yüksek, H., and Tugrul, S.: Eutrophication in the Sea of Marmara, in: *The Sea of Marmara; Marine Biodiversity, Fisheries, Conservation and Governance*, edited by: Özsoy, E., Çağatay, M. N., Balkis, N., Balkis, N., and Ozturk, B., Turkish Marine Research Foundation (TUDAV), Publication No: 42, Istanbul, Turkey, 723–736, 2016.
- Ergin, M. and Yörük, R.: Distribution and texture of the bottom sediments in a semi-enclosed coastal inlet, the Izmit Bay from the Eastern Sea of Marmara (Turkey), *Estuar. Coast. Shelf Sci.*, 30, 647–654, [https://doi.org/10.1016/0272-7714\(90\)90100-6](https://doi.org/10.1016/0272-7714(90)90100-6), 1990.
- Ergin, M., Bodur, M. N., Yıldız, M., Ediger, D., Ediger, V., Yemenicioğlu, S., and Yücesoy, F.: Sedimentation rates in the Sea of Marmara: a comparison of results based on organic carbon-primary productivity and ²¹⁰Pb dating, *Cont. Shelf Res.*, 14, 1371–1387, [https://doi.org/10.1016/0278-4343\(94\)90054-X](https://doi.org/10.1016/0278-4343(94)90054-X), 1994.
- Evans, G., Erten, H., Alavi, S. N., Von Gunten, H. R., and Ergin, M.: Superficial deep-water sediments of the Eastern Marmara basin, *Geo-Mar. Lett.*, 9, 27–36, <https://doi.org/10.1007/BF02262815>, 1989.
- Géli, L., Henry, P., Zitter, T., Dupré, S., Tryon, M., Çağatay, M. N., de Lépinay, B. M., Le Pichon, X., Şengör, A. M. C., Görür, N., Natalin, B., Uçarkuş, G., Özeren, S., Volker, D., Gasperini, L., Burnard, P., Bourlange, S., and the Marnaut Scientific Party: Gas emissions and active tectonics within the submerged section of the North Anatolian Fault zone in the Sea of Marmara, *Earth Planet. Sc. Lett.*, 274, 34–39, <https://doi.org/10.1016/j.epsl.2008.06.047>, 2008.
- Grasshoff, K., Ehrhardt, M., and Kremling, K.: Determination of nutrients, in: *Methods of Seawater Analysis*, 2nd edn., Verlag Chemie GmbH, Weinheim, Germany, 125–188, 1983.
- Halbach, P., Kuşçu, İ., Inthorn, M., Kuhn, T., Pekdeğer, A., and Seifert, R.: Methane in sediments of the deep Marmara Sea and its relation to local tectonic structures, In: *Integration of Earth Science Research on the Turkish and Greek 1999 Earthquakes*, 71–85, Springer, Dordrecht, 2002.
- Hille, S., Nausch, G., and Leipe, T.: Sedimentary deposition and reflux of phosphorus (P) in the Eastern Gotland Basin and their coupling with P concentrations in the water column, *Oceanologia*, 47, 663–679, 2005.
- Ignatieva, N. V.: Nutrient exchange across the sediment-water interface in the eastern Gulf of Finland, *Boreal Environ. Res.*, 4, 295–306, 1999.
- Jeitner, T. M.: Optimized ferrozine-based assay for dissolved iron, *Anal., Biochem.*, 454, 36–37, <https://doi.org/10.1016/j.ab.2014.02.026>, 2014.
- Jørgensen, B. B.: Material flux in the sediment, in: *Eutrophication in Coastal Marine Ecosystems*, edited by: Jørgensen, B. B., and Richardson, K., Washington, D.C.: American Geophysical Union, 52, 115–135, <https://doi.org/10.1029/CE052p0115>, 1996.
- Konovalov, S. K., Luther III, G. W., and Yücel, M.: Porewater redox species and processes in the Black Sea sediments, *Chem. Geol.*, 245, 254–274, <https://doi.org/10.1016/j.chemgeo.2007.08.010>, 2007.
- Le Pichon, X., Sengör, A. M. C., Demirbag, E., Rangin, C., Imren, C., Armijo, R., Görür, N., Çağatay, N., Mercier de Lépinay, B., Meyer, B., Saatçılar, R., and Tok, B.: The active Main Marmara fault, *Earth Planet. Sc. Lett.*, 192, 595–616, [https://doi.org/10.1016/S0012-821X\(01\)00449-6](https://doi.org/10.1016/S0012-821X(01)00449-6), 2001.
- Li, Y.-H. and Gregory, S.: Diffusion of ions in sea water and in deep-sea sediments, *Geochim. Cosmochim. Ac.*, 38, 703–714, [https://doi.org/10.1016/0016-7037\(74\)90145-8](https://doi.org/10.1016/0016-7037(74)90145-8), 1974.
- Mee, L.: The Black Sea in crisis: A need for concerted international action, *Ambio*, 21, 278–286, 1992.
- Menzel, D. W. and Corwin, N.: The measurement of total phosphorus in seawater based on the liberation of organically bound fractions by persulfate oxidation, *Limnol. Oceanogr.*, 10, 280–282, <https://doi.org/10.4319/lo.1965.10.2.0280>, 1965.
- Mu, D., Yuan, D., Feng, H., Xing, F., Teo, F. Y., and Li, S.: Nutrient fluxes across sediment-water interface in Bohai Bay Coastal Zone, China, *Mar. Pollut. Bull.*, 114, 705–714, <https://doi.org/10.1016/j.marpolbul.2016.10.056>, 2017.
- Noffke, A., Hensen, C., Sommer, S., Scholz, F., Bohlen, L., Mosch, T., Graco, M., and Wallmann, K.: Benthic iron and phosphorus fluxes across the Peruvian oxygen minimum zone, *Limnol. Oceanogr.*, 57, 851–867, <https://doi.org/10.4319/lo.2012.57.3.0851>, 2012.
- Nylund, A. T., Arneborg, L., Tengberg, A., Mallast, U., and Hassellöv, I.-M.: In situ observations of turbulent ship wakes and their spatiotemporal extent, *Ocean Sci.*, 17, 1285–1302, <https://doi.org/10.5194/os-17-1285-2021>, 2021.
- Polat, Ç. and Tuğrul, S.: Nutrient and Organic Carbon Exchanges between the Black and Marmara Seas through the Bosphorus Strait, *Cont. Shelf Res.*, 15, 1115–1132, [https://doi.org/10.1016/0278-4343\(94\)00064-T](https://doi.org/10.1016/0278-4343(94)00064-T), 1995.

- Polat, C., Tuğrul, S., Çoban, Y., Bastürk, O., and Salihoğlu, İ.: Elemental composition of seston and nutrient dynamics in the Sea of Marmara, *Hydrobiologia*, 363, 157–167, <https://doi.org/10.1023/A:1003117504005>, 1998.
- Rasheed, M., Al-Rousan, S., Manasrah, R., and Al-Horani, F.: Nutrient fluxes from deep sediment support nutrient budget in the oligotrophic waters of the Gulf of Aqaba, *J. Oceanogr.*, 62, 83–89, <https://doi.org/10.1007/s10872-006-0034-x>, 2006.
- Rasheed, M.: Nutrient Fluxes from sediments of the northern Gulf of Aqaba under various anthropogenic activities, *Lebanese Sci. J.*, 5, 3–16, 2004.
- Rebreanu, L., Vanderborght, J. P., and Chou, L.: The diffusion coefficient of dissolved silica revisited, *Mar. Chem.*, 112, 230–233, <https://doi.org/10.1016/j.marchem.2008.08.004>, 2008.
- Ruffine, L., Ondreas, H., Blanc-Valleron, M. M., Teichert, B. M. A., Scalabrin, C., Rinnert, E., Birot, D., Croguennec, C., Ponzevera, E., Pierre, C., Donval, J. P., Alix, A. S., Germain, Y., Bignon, L., Etoubleau, J., Caprais, J. C., Knoery, J., Lesongeur, F., Thomas, B., Roubi, A., Legoix, L., Burnard, P., Chevalier, N., Lu, H., Dupré, S., Fontanier, C., Dissard, D., Olgun, N., Yang, H., Strauss, H., Özaksoy, V., Perchoc, J., Podeur, C., Tarditi, C., Özbeki, E., Guyader, V., Marty, B., Madre, D., Pitel-Roudaut, M., Grall, C., Embriaco, D., Polonia, A., Gasperini, L., Çağatay, M. N., Henry, P., and Géli, L.: Multidisciplinary investigation on cold seeps with vigorous gas emissions in the Sea of Marmara (MarsiteCruise): Strategy for site detection and sampling and first scientific outcome, *Deep-Sea Res. Pt. II*, 153, 36–47, <https://doi.org/10.1016/j.dsr2.2018.03.006>, 2018.
- Rydin, E., Malmäus, J. M., Karlsson, O. M., and Jonsson, P.: Phosphorus release from coastal Baltic Sea sediments as estimated from sediment profiles, *Estuar. Coast. Shelf Sci.*, 92, 111–117, <https://doi.org/10.1016/j.ecss.2010.12.020>, 2011.
- Sarı, E. and Çağatay, N.: Sediment core studies on the North Anatolian Fault Zone in the Eastern Sea of Marmara: Evidence of Sea Level Changes and Fault Activity, *Bull. Miner. Res. Explor.*, 140, 1–18, 2010.
- Savun-Hekimoğlu, B. and Gazioglu, C.: Mucilage Problem in the Semi-Enclosed Seas: Recent Outbreak in the Sea of Marmara, *Int. J. Environ. Geoinformatics* 8, 402–413, 2021.
- Simboura, N., Panayotidis, P., and Papathanassiou, E.: A synthesis of the biological quality elements for the implementation of the European Water Framework Directive in the Mediterranean ecoregion: the case of Saronikos Gulf, *Ecol. Indic.*, 5, 253–266, <https://doi.org/10.1016/j.ecolind.2005.03.006>, 2005.
- Stookey, L. L.: Ferrozine-A new spectrophotometric reagent for iron, *Anal. Chem.*, 42, 779–781, 1970.
- Strickland, J. D. H. and Parsons, T. R.: A Practical Handbook of Seawater Analysis, 2nd edn., *Bull. Fish. Res. Bd. Can.*, 167, 310 pp., 1972.
- Tan, İ. and Aslan, E.: Metal pollution status and ecological risk assessment in marine sediments of the inner İzmit Bay, *Reg. Stud. Mar. Sci.*, 33, 100850, <https://doi.org/10.1016/j.rsma.2019.100850>, 2020.
- Tryon, M. D., Henry, P., Çağatay, M. N., Zitter, T. A. C., Geli, L., Gasperini, L., Burnard, P., Bourlange, S., and Grall, C.: Pore fluid chemistry of the North Anatolian Fault Zone in the Sea of Marmara: a diversity of sources and processes, *Geochem. Geophys. Geosys.*, 11, 1–22, <https://doi.org/10.1029/2010GC003177>, 2010.
- Tuğrul, S. and Morkoç, E.: Oceanographic Characteristics of İzmit Bay, NATO TU-WATERS Project, Technical Report, TÜBİTAK, MRC Publ, Kocaeli, Turkey, 1989.
- Tuğrul, S. and Polat, Ç.: Quantitative comparison of the influxes of nutrients and organic carbon into the Sea of Marmara both from anthropogenic sources and from the Black Sea, *Water Sci. Technol.*, 32, 115–121, [https://doi.org/10.1016/0273-1223\(95\)00576-9](https://doi.org/10.1016/0273-1223(95)00576-9), 1995.
- Tuğrul, S., Beşiktepe, T., and Salihoğlu, İ.: Nutrient exchange fluxes between the Aegean and Black Seas through the Marmara Sea, *Medit. Mar. Sci.*, 3, 33–42, <https://doi.org/10.12681/mms.256>, 2002.
- Tuğrul, S., Murray, J. W., Friederich, G. E., and Salihoğlu, İ.: Spatial and temporal variability in the chemical properties of the oxic and suboxic layers of the Black Sea, *J. Mar. Syst.*, 135, 29–43, <https://doi.org/10.1016/j.jmarsys.2013.09.008>, 2014.
- Ullman, W. J. and Aller, R. C.: Diffusion coefficients in nearshore marine sediments, *Limnol. Oceanogr.*, 27, 552–556, <https://doi.org/10.4319/lo.1982.27.3.0552>, 1982.
- UNEP/MAP: Sampling and analysis techniques for the eutrophication monitoring strategy of MED POL, MAP Technical Rep. Ser., No. 163, Athens, 2005.
- UNEP/MAP: Methods for sediment sampling and analysis, Review Meeting of MED POL – Phase III Monitoring Activities, Athens, Palermo (Sicily), Italy, 12–15 December 2005, <https://sednet.org/download/wg-282-inf-5-rev-1.pdf>, 2006.
- Ünlüata, Ü., Oğuz, T., Latif, M. A., and Özsoy, E.: On the Physical Oceanography of the Turkish Straits, In: *The Physical Oceanography of Sea Straits*, edited by: Pratt, L. J., NATO/ASI Series, Kluwer, Dordrecht, 25–60, 1990.
- van Helmond, N. A. G. M., Robertson, E. K., Conley, D. J., Hermans, M., Humborg, C., Kubeneck, L. J., Lenstra, W. K., and Slomp, C. P.: Removal of phosphorus and nitrogen in sediments of the eutrophic Stockholm archipelago, Baltic Sea, *Biogeosciences*, 17, 2745–2766, <https://doi.org/10.5194/bg-17-2745-2020>, 2020.
- Williams, G. R.: The coupling of biogeochemical cycles of nutrients, *Biogeochemistry*, 4, 61–75, [doi:10.1007/BF02187362](https://doi.org/10.1007/BF02187362), 1987.
- Yalçın, B., Artüz, M. L., Pavlidou, A., Çubuk, S., and Dassenakis, M.: Nutrient dynamics and eutrophication in the Sea of Marmara: Data from recent oceanographic research, *Sci. Total Environ.*, 601, 405–424, <https://doi.org/10.1016/j.scitotenv.2017.05.179>, 2017.
- Yang, H., Lu, H., and Ruffine, L.: Geochemical characteristics of iron in sediments from the Sea of Marmara, *Deep-Sea Res. Pt. II*, 153, 121–130, <https://doi.org/10.1016/j.dsr2.2018.01.010>, 2018.
- Yücel, M., Konovalov, S. K., Moore, T. S., Janzen, C. P., and Luther III, G. W.: Sulfur speciation in the upper Black Sea sediments, *Chem. Geol.*, 269, 364–375, <https://doi.org/10.1016/j.chemgeo.2009.10.010>, 2010.
- Yücel, M., Özkan, K., Örek, H., Mantıkçı, M., Tezcan, D., Akçay, İ., Özhan, K., Arkin, S., Tuğrul, S., and Salihoğlu, B.: The sea snout outbreak in the Sea of Marmara: Biogeochemical transformations of the sea, modern-day pressures and a roadmap for the way forward. *Ecology of the Marmara Sea: Formation and Interactions of Marine Mucilage, and Recommendations for Solutions*, edited by: Öztürk, İ. and Şeker, M., Turkish Academy of Sciences, Ankara, Turkey, 249–267, 2021.

Yücel, M., Örek, H., Alımlı, N., Akçay, İ., Mantıkçı, M., Özhan, K., Fach, B., Tezcan, D., Ak Örek, Y., Kalkan Tezcan, E., Arkin, S., Tuğrul, S., and Salihoğlu, B.: Phosphorus build-up, nitrogen loss and sulfide accumulation in response to recent deep-water deoxygenation in the Sea of Marmara, Goldschmidt 2023 Conference, Lyon, France, <https://doi.org/10.7185/gold2023.15805>, 2023a.

Yücel, M., Tezcan, E. K., Örek, H., Fach, B., Tezcan, D., Arkin, S., Sadıhgrad, E., Acar, A. O., Mantıkçı, M., Özkan, M., Özhan, K., Hüsrevoğlu, S., Ak Örek, Tuğrul, S., and Salihoğlu, B.: The study of deoxygenation and mucilage formation in the Marmara Sea using novel oceanographic approaches in the frame of MAR-MOD project. *Journal of Environment, Urbanization and Climate Change*, 2, 82–96, 2023b.