



Supplement of

Terrestrial and marine plastic pollution outlook in the Mediterranean region: a box-model approach based on OECD policy scenarios

Théo Segur and Jeroen E. Sonke

Correspondence to: Théo Segur (theo.segur@get.omp.eu) and Jeroen E. Sonke (jeroen.sonke@get.omp.eu)

The copyright of individual parts of the supplement might differ from the article licence.

I. Code availability

The model code is available at: <https://doi.org/10.5281/zenodo.17100118>

II. Model parameters

Table S1: Model parameters and their probability distribution (miscellaneous parameters). Whenever uncertainties were unavailable, they are set to zero, and parameters treated as constants.

R1: S. Europe, R2: N. Africa & M. East, R3: Nile basin.

Parameter name	Description	Mean $\pm$ sd	Median (IQR 25%-75%)
LB19	Fragmentation rate (3% per year)	$3.0\text{E-}2 \pm 5.7\text{E-}3$	$3.0\text{E-}2$ ($2.6\text{E-}2$ - $3.4\text{E-}2$)
f_LMP	Fraction of LMP in waste	$1.4\text{E-}1 \pm 2.7\text{E-}2$	$1.4\text{E-}1$ ($1.2\text{E-}1$ - $1.6\text{E-}1$)
f_burn_R1	Fraction of mismanaged plastic waste that is open burned each year (from OECD)	$3.1\text{E-}1 \pm 0.0\text{E+}0$	$3.1\text{E-}1$ ($3.1\text{E-}1$ - $3.1\text{E-}1$)
f_burn_R2		$3.1\text{E-}1 \pm 0.0\text{E+}0$	$3.1\text{E-}1$ ($3.1\text{E-}1$ - $3.1\text{E-}1$)
f_burn_R3		$3.1\text{E-}1 \pm 0.0\text{E+}0$	$3.1\text{E-}1$ ($3.1\text{E-}1$ - $3.1\text{E-}1$)
f_inc_random	Normal randomisation of waste fractions	$1.0\text{E+}0 \pm 4.9\text{E-}2$	$1.0\text{E+}0$ ($9.7\text{E-}1$ - $1.0\text{E+}0$)
f_mism_random		$1.0\text{E+}0 \pm 5.0\text{E-}2$	$1.0\text{E+}0$ ($9.7\text{E-}1$ - $1.0\text{E+}0$)
f_rec_random		$1.0\text{E+}0 \pm 4.9\text{E-}2$	$1.0\text{E+}0$ ($9.7\text{E-}1$ - $1.0\text{E+}0$)
f_runoff_R1	Fraction of total runoff attributed to each region	$1.1\text{E-}1 \pm 0.0\text{E+}0$	$1.1\text{E-}1$ ($1.1\text{E-}1$ - $1.1\text{E-}1$)
f_runoff_R2		$7.5\text{E-}1 \pm 0.0\text{E+}0$	$7.5\text{E-}1$ ($7.5\text{E-}1$ - $7.5\text{E-}1$)
f_runoff_R3		$1.5\text{E-}1 \pm 0.0\text{E+}0$	$1.5\text{E-}1$ ($1.5\text{E-}1$ - $1.5\text{E-}1$)
f_soil_R1	Fraction of each region surface considered as remote soil	$5.7\text{E-}1 \pm 0.0\text{E+}0$	$5.7\text{E-}1$ ($5.7\text{E-}1$ - $5.7\text{E-}1$)
f_soil_R2		$6.6\text{E-}1 \pm 0.0\text{E+}0$	$6.6\text{E-}1$ ($6.6\text{E-}1$ - $6.6\text{E-}1$)
f_soil_R3		$8.0\text{E-}1 \pm 0.0\text{E+}0$	$8.0\text{E-}1$ ($8.0\text{E-}1$ - $8.0\text{E-}1$)
f_ssed	Fraction of seafloor considered as shelf sediment	$1.8\text{E-}1 \pm 0.0\text{E+}0$	$1.8\text{E-}1$ ($1.8\text{E-}1$ - $1.8\text{E-}1$)

10

Table S2: Model parameters and their probability distribution (fluxes between boxes).

R1: S. Europe, R2: N. Africa & M. East, R3: Nile basin. Note that the “Terrestrial” compartment incorporates the terrestrial mismanaged and the remote land compartment.

Parameter name	Compartment from	Compartment to	Plastic size category	Mean $\pm$ sd	Median (IQR 25%-75%)
----------------	------------------	----------------	-----------------------	---------------	----------------------

k_R1_P_terr_to_surf	Terrestrial (R1)	Sea surface	P	2.5E-3 ± 1.6E-3	2.1E-3 (1.2E-3 - 3.4E-3)
k_R1_LMP_terr_to_surf	Terrestrial (R1)	Sea surface	LMP	7.2E-2 ± 1.0E-1	2.6E-2 (4.7E-3 - 9.7E-2)
k_R1_sMP_terr_to_surf	Terrestrial (R1)	Sea surface	SMP	4.4E-3 ± 1.3E-3	4.2E-3 (3.4E-3 - 5.2E-3)
k_R1_sMP_terr_to_atmo	Terrestrial (R1)	Atmosphere	SMP	2.4E-3 ± 1.0E-3	2.3E-3 (1.6E-3 - 3.0E-3)
k_R2_P_terr_to_surf	Terrestrial (R2)	Sea surface	P	3.1E-4 ± 2.0E-4	2.6E-4 (1.5E-4 - 4.2E-4)
k_R2_LMP_terr_to_surf	Terrestrial (R2)	Sea surface	LMP	9.1E-3 ± 1.3E-2	3.3E-3 (6.0E-4 - 1.2E-2)
k_R2_sMP_terr_to_surf	Terrestrial (R2)	Sea surface	SMP	8.9E-4 ± 2.6E-4	8.6E-4 (6.9E-4 - 1.1E-3)
k_R2_sMP_terr_to_atmo	Terrestrial (R2)	Atmosphere	SMP	2.4E-3 ± 1.0E-3	2.3E-3 (1.6E-3 - 3.0E-3)
k_R3_P_terr_to_surf	Terrestrial (R3)	Sea surface	P	4.7E-6 ± 3.0E-6	3.9E-6 (2.3E-6 - 6.2E-6)
k_R3_LMP_terr_to_surf	Terrestrial (R3)	Sea surface	LMP	1.4E-4 ± 2.0E-4	4.9E-5 (8.9E-6 - 1.8E-4)
k_R3_sMP_terr_to_surf	Terrestrial (R3)	Sea surface	SMP	3.0E-2 ± 5.7E-3	3.0E-2 (2.6E-2 - 3.4E-2)
k_R3_sMP_terr_to_atmo	Terrestrial (R3)	Atmosphere	SMP	3.0E-2 ± 5.7E-3	3.0E-2 (2.6E-2 - 3.4E-2)
k_R1_sMP_atmo_to_terr	Atmosphere	Terrestrial (R1)	SMP	5.7E+1 ± 1.8E+1	5.5E+1 (4.3E+1 - 7.1E+1)
k_R2_sMP_atmo_to_terr	Atmosphere	Terrestrial (R2)	SMP	5.7E+1 ± 1.8E+1	5.5E+1 (4.3E+1 - 7.1E+1)
k_R3_sMP_atmo_to_terr	Atmosphere	Terrestrial (R3)	SMP	5.7E+1 ± 1.8E+1	5.5E+1 (4.3E+1 - 7.1E+1)
k_sMP_atmo_to_surf	Atmosphere	Sea surface	SMP	4.0E+1 ± 2.4E+1	3.5E+1 (2.1E+1 - 5.4E+1)
k_sMP_surf_to_atmo	Sea surface	Atmosphere	SMP	1.3E+0 ± 9.2E-1	1.0E+0 (5.6E-1 - 1.7E+0)
k_P_surf_to_sand	Sea surface	Sandy beach	P	2.3E-1 ± 9.0E-3	2.3E-1 (2.2E-1 - 2.4E-1)
k_LMP_surf_to_sand	Sea surface	Sandy beach	LMP	0.0E+0 ± 0.0E+0	0.0E+0 (0.0E+0 - 0.0E+0)
k_sMP_surf_to_sand	Sea surface	Sandy beach	SMP	0.0E+0 ± 0.0E+0	0.0E+0 (0.0E+0 - 0.0E+0)
k_P_surf_to_ssed	Sea surface	Shelf sediment	P	2.0E+2 ± 1.8E+1	2.0E+2 (1.9E+2 - 2.1E+2)
k_LMP_surf_to_ssed	Sea surface	Shelf sediment	LMP	1.5E+4 ± 2.4E+4	5.1E+3 (7.9E+2 - 1.9E+4)
k_sMP_surf_to_ssed	Sea surface	Shelf sediment	SMP	1.2E+2 ± 2.5E+1	1.2E+2 (9.9E+1 - 1.4E+2)
k_P_surf_to_wcol	Sea surface	Water column	P	0.0E+0 ± 0.0E+0	0.0E+0 (0.0E+0 - 0.0E+0)
k_LMP_surf_to_wcol	Sea surface	Water column	LMP	2.5E+2 ± 6.6E+0	2.4E+2 (2.0E+2 - 2.9E+2)
k_sMP_surf_to_wcol	Sea surface	Water column	SMP	3.0E-2 ± 5.7E-3	3.0E-2 (2.6E-2 - 3.4E-2)
k_LMP_wcol_to_dsed	Water column	Deep sediment	LMP	1.0E-3 ± 6.5E-19	1.0E-3 (1.0E-3 - 1.0E-3)
k_sMP_wcol_to_dsed	Water column	Deep sediment	SMP	3.0E-2 ± 5.7E-3	3.0E-2 (2.6E-2 - 3.4E-2)

15

III.

Per capita plastic waste generation




Figure S1: Per capita plastic waste generation by region.

Dots are data points from OECD reports (OECD, 2022). Quadratic regression lines with y=0 at x=1950 were used to extrapolate these points to the full 1950 to 2060 range.

20

Region 1: S. Europe, Region 2: N. Africa & M. East, Region 3: Nile basin.

IV. Demography data per country




Figure S2: Fraction of the country population living inside the Mediterranean (Med.) catchment and 50 km coastline from 2000 to 2020. Obtained using georeferenced population data from WorldPop database (WorldPop, 2020).

This fraction is constant from 2000 to 2020 for most countries, that is why we decided to consider it constraint for the whole study timeframe. Increasing fraction observed for low population countries (such as Albania or Montenegro) should have a negligible impact on the overall population estimation.




Figure S3: Population inside the Mediterranean catchment. Obtained using UN World Population Prospect 2022 (United Nations, 2022), combined with extrapolation from Figure S.4.1.

35 V. Sediment extrapolation

To estimate the quantity of LMP accumulated from 1950 ($t=0$) to 2015 ($t=T[y]$) in the marine sediment, an extrapolation method was used. The available observations provide plastic concentrations q [g.m^{-3}] in the d [m] first meter of sediment, d being the sampling depth of the core.

The d first meters of sediment were deposited on the seafloor during an accumulation time Δt [y]:

$$40 \quad \Delta t = \frac{d}{S_r} \quad \text{with } S_r [\text{m.y}^{-1}] \text{ the sedimentation rate of the study area.}$$

From 1950 ($t=0$) to 2015 ($t=T$), a sediment layer of height D [m] was deposited on the seafloor:

$$D = T \cdot S_r$$

Let's convert the volumetric concentration q [g.m^{-3}] into an observed concentration per surface unit c_{obs} [g.m^{-2}]:

$$45 \quad c_{obs} = q \cdot d$$

This quantity c_{obs} corresponds to the plastic mass accumulated on 1m^2 of seafloor during the Δt years before 2015.

Let's introduce a plastic sedimentation rate S [$\text{g.m}^{-2}.\text{y}^{-1}$], i.e. a plastic flux from the water column to 1m^2 of seafloor. The function S is unknown, but we can suppose that it is approximated by a quadratic equation according to the model of Sonke et al., 2022 (Sonke et al., 2022). Furthermore, $S(t \leq 0) = 0$ since no plastic contaminated the environment before 1950. Therefore:

$$S(t \geq 0) = a \cdot t^2 \quad \text{and} \quad S(t < 0) = 0 \quad \text{with } a \text{ a parameter to determine.}$$

Let's compute the parameter a in order to observe a concentration c_{obs} during the last Δt years. The quantity of plastic precipitated over one unit of surface between $T-\Delta t$ and T is can be expressed as:

$$c_{obs} = \int_{t=T-\Delta t}^{t=T} S(t).dt$$

- If $d \leq D$ (the core only sampled sediment deposited after 1950):

$$c_{obs} = \int_{t=T-\Delta t}^{t=T} S(t).dt = \int_{t=T-\Delta t}^{t=T} a \cdot t^2 \cdot dt = \frac{a}{3} [t^3]_{T-\Delta t}^T = \frac{a}{3} [T^3 - (T - \Delta t)^3]$$

Knowing the apparent surface concentration c_{obs} , we can then find the parameter a that define the precipitation function S :

$$a = c_{obs} \cdot \frac{3}{T^3 - (T - \Delta t)^3}$$

We can then compute C [g.m^{-2}], the plastic mass accumulated on 1m^2 of seafloor from 1950 ($t=0$) to 2015 ($t=T$) :

$$C = \int_{t=0}^{t=T} S(t).dt = a \cdot \int_{t=0}^{t=T} t^2 \cdot dt = \frac{a}{3} [t^3]_0^T = \frac{a}{3} \cdot T^3 = c_{obs} \cdot \frac{T^3}{T^3 - (T - \Delta t)^3}$$

Expressing T and Δt according to D and d , we have:

$$C = c_{obs} \frac{1}{1 - (1 - \frac{d}{D})^3}$$

- if $d > D$ (the core sampled some sediment deposited before 1950 that did not contain any plastics):

$$c_{obs} = \int_{t=T-\Delta t}^{t=T} S(t).dt = \int_{t=T-\Delta t}^{t=0} S(t).dt + \int_{t=0}^{t=T} S(t).dt = 0 + \int_{t=0}^{t=T} S(t).dt$$

since S is null before 1950 (no plastic contamination yet)

$$\Rightarrow c_{obs} = a \cdot \int_{t=0}^{t=T} t^2 \cdot dt = \frac{a}{3} [t^3]_0^T = \frac{a}{3} \cdot T^3 \quad \Rightarrow a = c_{obs} \cdot \frac{3}{T^3}$$

$$\Rightarrow C = \int_{t=0}^{t=T} S(t).dt = a \cdot \int_{t=0}^{t=T} t^2 \cdot dt = \frac{a}{3} [t^3]_0^T = \frac{a}{3} \cdot T^3 = c_{obs}$$

Therefore:

$$C = q \cdot \frac{d}{1 - (1 - \frac{d}{D})^3} \quad \text{if } d \leq D \quad (S1)$$

$$C = q \cdot d \quad \text{if } d > D$$

75 To estimate M [Mt] the quantity of plastic stoked in the Mediterranean seafloor, we multiply C by the surface of the sediment A [km²]:

$$M = A \cdot C \cdot 10^6$$

80

85 VI. SMP extrapolation

Facing the lack of observation of SMP in the Mediterranean region, the Mediterranean SMP stock for sea surface (surf), water column (wcol), shelf sediment (ssed) and sandy beach (sand) were extrapolated from a similar box model by Sonke et al. (2025) to perform model calibration. Table S3 presents the values used to compute the extrapolated SMP stocks for the Mediterranean Sea.

90 **Table S3:** SMP extrapolation from Sonke et al. (2025)

	Surface [km ²]	Mean depth [km]	Volume [km ³]	Stock [Tg]	SD [Tg]
Sea surface					
Global	3.61E+08**	1.00E-01	3.61E+07	0.11**	0.05**
Med. Sea	2.52E+06*	1.00E-01	2.52E+05	7.7E-04	3.5E-04
Water column					
Global	3.61E+08**	3.59E+00**	1.30E+09	41**	17**
Med. Sea	2.52E+06*	1.45E+00*	3.64E+06	1.2E-01	4.8E-02
Shelf sediment					
Global	3.57E+07**			14**	6.7**
Med. Sea	4.51E+05*			0.18	0.08
Sandy beach					
Global	2.63E+05**			0.20**	0.19**
Med. Sea	1.09E+03*			8.3E-04	7.8E-04

*Computed from Poulos, (2020)

**Values from Sonke et al. (2025)

VII. Geographical distribution of plastic runoff

95 **Table S4:** Comparison between different studies providing geographical distribution of plastic runoff to sea.

	Lebreton et al. (2017)	Meijer et al. (2021)	Nyberg et al. (2023)	Weiss et al. (2025)	Cózar et al. (2024)
# of river (global)	40 760	31 819	156 163	9 988	-
# of river (Med.)	2 724 (6.7%)	1 328 (4.2%)	4 004 (2.6%)	549 (5.5%)	-
Global flux (Tg/y)	1.41	1.00	19.19	0.006	-
Med. flux (Tg/y)	0.001 (0.1%)	0.02 (2.3%)	1.24 (6.5%)	8.0 10 ⁻⁵	-
Regional contribution to plastic runoff from land to sea	S.Eu : 16.3% N. Af : 60.1% Nile : 23.7%	S.Eu : 5.6 % N. Af : 82.3 % Nile : 12.1 %	S.Eu : 11.0 % N. Af : 78.5 % Nile : 10.5 %	S.Eu : 79.5 % N. Af : 14.0 % Nile : 6.5 %	S.Eu : 11.0 % N. Af : 78.5 % Nile : 10.5 %
Calibration	13 rivers 30 records No African river	67 rivers 136 records No African river	No calibration	89 rivers 127 sampling location 340 records No African river	Calibration with Mediterranean data over 75 months
Plastic type	P + LMP	P	Not mentioned	LMP	P
Runoff	Monthly runoff used Dams and floods are modelled	Annual wind and precipitation used, Land use. Don't account for dams, nor for flood	Do not account for water discharge change, but for river connectivity	Account for drainage intensity	Account for precipitation and runoff (at least monthly)
Method	Population density + MMPW statistics generates flux that follows gradients and can be blocked by a dam. Monthly runoff is used	Probability model P(MMPW emission to the ocean) = P(mobilization on land) * P(transport from land to river) * P(transport from river to ocean)	MMPW input per km ² is computed using the Lebreton and Andrady method. Connectivity of river is computed and maximum discharge estimated	Population density and precipitation in the catchment are the two predictors used for the regression.	Optimize plastic runoff distribution around the mediterranean coast with hotspots of floating macroplastics identified via satellite imaging.
Positive point	Include dams Include monthly runoff and some data during floods	Include land use, wind and annual runoff	High number of rivers in results	High number of river used in calibration	Calibrated on Mediterranean observation Accounts for precipitation and runoff
Negative point	Low number of calibration points (30) Calibrated globally No african rivers Only 1 Mediterranean river (Po)	Don't account for dams nor floods. Calibrated globally No african rivers Only 2 Mediterranean rivers for calibration (and 2 others for validation)	No calibration Don't account for runoff Calibrated globally	Only 4 mediterranean rivers in calibration Calibrated globally Assumes a constant mass per particle	Still a proof of concept Calibrated on proxy (macro litter hotspots)

100 **VIII. Per capita results**

Table S5. Comparison of per capita plastic leakage with Global estimate and between Global estimations and Mediterranean regions.

		Global (from Sonke et al. (2025))		Mediterranean region		S. Europe		N. Africa & M. East		Nile basin	
	Unit \ Year	2023	2060	2023	2060	2023	2060	2023	2060	2023	2060
Population	M capita	8119	10067	536	775	124	107	140	186	273	482
Plastic waste generation	Mt/y	272	656	26	59	16	26	6	16	4	17
	kg/y/capita	34	65	49	76	129	239	40	86	15	35
Mismanaged plastic waste generation	Mt/y	75	148	5.5	11.5	0.8	0.2	2.8	5.5	2.0	5.8
	kg/y/capita	9	15	10.3	14.8	6.5	1.9	20.0	29.6	7.3	12.0
Plastic runoff to sea	Mt/y	18	43	0.4	0.5	0.30	0.27	0.1	0.2	0.0005	0.0027
	kg/y/capita	2.2	4.3	0.7	0.6	2.4	2.5	0.4	1.1	0.0018	0.0055

IX. Total marine plastic




Figure S4: Total marine plastic in 2060 according to the 4 modelled scenarios. Letters indicate significance according to the Dunn post-hoc test. Dotted lines represent the mean, and solid lines the median.

REFERENCES

- 110 Cózar, A., Arias, M., Suaria, G., Viejo, J., Aliani, S., Koutroulis, A., Delaney, J., Bonnery, G., Macías, D., De Vries, R., Sumerot, R., Morales-Caselles, C., Turiel, A., González-Fernández, D., and Corradi, P.: Proof of concept for a new sensor to monitor marine litter from space, *Nat Commun*, 15, 4637, <https://doi.org/10.1038/s41467-024-48674-7>, 2024.
- 115 Lebreton, L. C. M., van der Zwet, J., Damsteeg, J.-W., Slat, B., Andrady, A., and Reisser, J.: River plastic emissions to the world's oceans, *Nat Commun*, 8, 15611, <https://doi.org/10.1038/ncomms15611>, 2017.
- Meijer, L. J. J., van Emmerik, T., van der Ent, R., Schmidt, C., and Lebreton, L.: More than 1000 rivers account for 80% of global riverine plastic emissions into the ocean, *Sci. Adv.*, 7, eaaz5803, <https://doi.org/10.1126/sciadv.aaz5803>, 2021.
- 120 Nyberg, B., Harris, P. T., Kane, I., and Maes, T.: Leaving a plastic legacy: Current and future scenarios for mismanaged plastic waste in rivers, *Science of The Total Environment*, 869, 161821, <https://doi.org/10.1016/j.scitotenv.2023.161821>, 2023.
- OECD: Global Plastics Outlook: Policy Scenarios to 2060, 2022.
- 125 Poulos, S. E.: The Mediterranean and Black Sea Marine System: An overview of its physico-geographic and oceanographic characteristics, *Earth-Science Reviews*, 200, 103004, <https://doi.org/10.1016/j.earscirev.2019.103004>, 2020.
- Sonke, J. E., Koenig, A., Segur, T., and Yakovenko, N.: Global environmental plastic dispersal under OECD policy scenarios toward 2060, *Science Advances*, 11, eadu2396, <https://doi.org/10.1126/sciadv.adu2396>, 2025.
- United Nations: World Population Prospects 2022, 2022.
- 130 Weiss, L., Estournel, C., Marsaleix, P., Mikolajczak, G., Constant, M., and Ludwig, W.: From source to sink: part 1—characterization and Lagrangian tracking of riverine microplastics in the Mediterranean Basin, *Environ Sci Pollut Res*, 32, 10081–10104, <https://doi.org/10.1007/s11356-024-34635-6>, 2025.
- 135 WorldPop: Unconstrained individual countries 2000-2020 UN adjusted (1km resolution), 2020.