



Supplement of

BrGDGT-based palaeothermometer in drylands: the necessity to constrain aridity and salinity as confounding factors to ensure the robustness of calibrations

Lucas Dugerdil et al.

Correspondence to: Lucas Dugerdil (lucas.dugerdil@ens-lyon.fr) and Guillemette Ménot (guillemette.menot@ens-lyon.fr)

The copyright of individual parts of the supplement might differ from the article licence.

S1. Supplementary Figures




Figure S1. Climographs for four regions of the ACA: Azerbaijan, Uzbekistan, Mongolia and China. The climate parameters were extracted from interpolated climate data from `worldclim2.1` (Fick and Hijmans, 2017) at the sample location of the ACADB sites. The estimated Growth Season (GS) was roughly determined with $T > 5\text{ }^{\circ}\text{C}$ and $P > 2 \times T$. Climate data represent averages over the period ca. 1970–2000 (Fick and Hijmans, 2017).




Figure S2. Graphical determination of the more efficient pH threshold affecting the bimodal distribution of CBT' against pH for soil samples. The pH threshold values were implemented from 4 to 11 for each 0.01 step. The maximum multiple R^2 is found for $\text{pH} = 7.3$. The lower panel shows the number of samples for each cluster. In blue are given the results for the *acidic* soils (i.e., the samples below the threshold), in orange the *alkaline* soils (above the threshold), and in black the overall multiple R^2 .




Figure S3. Effect of the salinity and the aridity on the sum of 5-, 6-, 7-methyl and 6+7-methyl FAs for soil and lacustrine samples.




Figure S4. Boxplot for the main brGDGT-based ratios presented in this study clustered by the four grouping factors (i.e., alkalinity, aridity, salinity, and sample type). This figure is associated with the MANOVA and ANOVA results of Table 3.




Figure S5. Relationships between the Aridity Index (AI) and Methylation indices, i.e. MBT'_{5Me} (A), MBT'_{6Me} (B), and $\Delta(MBT'_{5Me}, MBT'_{6Me})$ (C) linear relationships with AI. The regressions were tested for soil (orange) and lacustrine (blue) samples. Solely, the R^2 for significant regression (p -value < 0.001) were displayed.




Figure S6. Sensitivity analyses to determinate the salinity threshold values affecting the bimodal distribution of IR_{6Me} (A) and IR_{7Me} (B) for lacustrine samples. The salinity threshold values were implemented from 30 to 15,000 $mg.L^{-1}$ for each 100 $mg.L^{-1}$. The lower panels shows the number of samples for each cluster. In blue are given the results for the samples below the threshold), in orange the samples above the threshold), and in black the overall multiple R^2 .




Figure S7. Effect of the alkalinity (i.e., the pH of each surface sample) on the linear relation between temperature (here MAAT) and the MBT'_{5Me}. Based on the alkalinity classes (i.e., acid, neutral, and alkaline), different temperature calibration MBT'_{5Me}-based are proposed following $MAAT = a \times MBT'_{5Me} + b$. Using the z -statistic with its p -value (Clogg et al., 1995), the significance of the difference between the slopes (a) of the linear regression is evaluated. Similarly, the z -statistic was used for the intercept (b) differences. For the p -values we have *** for $p \leq 0.01$, ** for $p \leq 0.05$ and * for $p \leq 0.1$.




Figure S8. Effect of the aridity (i.e. the AI of each surface sample) on the linear relation between temperature (here MAAT) and the MBT'_{5Me}. Based on the aridity classes (i.e., hyper-arid, arid, semi-arid, dry sub-humid, and humid), different temperature calibration MBT'_{5Me}-based are proposed following $MAAT = a \times MBT'_{5Me} + b$. Using the z -statistic with its p -value (Clogg et al., 1995), the significance of the difference between the slopes (a) of the linear regression is evaluated. Similarly, the z -statistic was used for the intercept (b) differences. For the p -values we have *** for $p \leq 0.01$, ** for $p \leq 0.05$, and * for $p \leq 0.1$.




Figure S9. Effect of the sample type on the linear relation between temperature (here MAAT) and the MBT'_{5Me} . Based on the sample types (i.e., soil or lacustrine), different temperature calibration MBT'_{5Me} -based are proposed following $MAAT = a \times MBT'_{5Me} + b$. Using the z -statistic with its p -value (Clogg et al., 1995), the significance of the difference between the slopes (a) of the linear regression is evaluated. Similarly, the z -statistic was used for the intercept (b) differences. For the p -values we have *** for $p \leq 0.01$, ** for $p \leq 0.05$, and * for $p \leq 0.1$.

Table S1: Geographical and biological presentation of the surface sites of the ACADB analysed in this study and previously published in (Dugerdil et al., 2025).

Sites	Longitude	Latitude	Altitude	Date	Country	Sample type	Ecosystem	pH	Salinity
MAZT1C02	46.99	40.99	109	2021	Azerbaijan	Lacustrine	Halophytic desert	8.11	124839.00
MAZT1C06	45.77	41.19	550	2021	Azerbaijan	Lacustrine	Shrub desert	8.53	2016.95
MAZT1M01	46.99	40.99	112	2021	Azerbaijan	Soil	Halophytic desert	8.37	2249.00
MAZT1M03	47.00	41.01	138	2021	Azerbaijan	Soil	Halophytic desert	8.41	1001.00
MAZT1M04	45.77	41.18	597	2021	Azerbaijan	Soil	Shrub desert	5.82	2314.00
MAZT1M07	45.74	41.06	246	2021	Azerbaijan	Soil	Shrub desert	7.18	3185.00
MAZT1M08	45.79	41.01	288	2021	Azerbaijan	Soil	Shrub desert	7.34	2522.00
MAZT1M09	47.78	40.71	222	2022	Azerbaijan	Soil	Thermophilous woodland	7.61	2008.50
MAZT1M10	46.91	41.21	391	2022	Azerbaijan	Soil	Thermophilous woodland	7.09	1111.50
MAZT1M11	46.05	40.70	668	2022	Azerbaijan	Soil	Thermophilous woodland	6.81	2405.00
MAZT1S05	45.77	41.19	552	2021	Azerbaijan	Soil	Shrub desert	7.67	1612.00
MAZT2C02	46.26	40.37	1722	2021	Azerbaijan	Lacustrine	Deciduous forest	6.81	507.00
MAZT2C05	46.59	40.64	273	2021	Azerbaijan	Lacustrine	Halophytic desert	7.57	3302.00
MAZT2M01	46.28	40.40	1412	2021	Azerbaijan	Soil	Deciduous forest	6.79	1469.00
MAZT2M03	46.27	40.37	1820	2021	Azerbaijan	Soil	Deciduous forest	6.49	1313.00
MAZT2M04	46.36	40.46	1232	2021	Azerbaijan	Soil	Deciduous forest	7.50	1274.00
MAZT2M06	46.96	40.55	96	2021	Azerbaijan	Soil	Shrub desert	7.96	1092.00
MAZT2M07	45.75	40.54	1660	2022	Azerbaijan	Soil	Grasslands	7.11	1229.80
MAZT3C01	47.76	40.11	-10	2021	Azerbaijan	Lacustrine	Halophytic desert	7.96	4399.20
MAZT3C03	47.61	39.89	15	2021	Azerbaijan	Lacustrine	Halophytic desert	8.98	90530.05
MAZT3M04	47.61	39.90	15	2021	Azerbaijan	Soil	Halophytic desert	7.45	877.50
MAZT3M06	47.28	39.97	86	2022	Azerbaijan	Soil	Halophytic desert	7.26	2620.15
MAZT3M07	47.99	39.88	-2	2022	Azerbaijan	Soil	Halophytic desert	9.11	4134.00
MAZT3S02	47.59	40.09	-4	2021	Azerbaijan	Soil	Halophytic desert	8.90	16705.00
MAZT3S05	47.60	39.90	12	2021	Azerbaijan	Soil	Halophytic desert	9.05	23029.50
MAZT4M01	48.88	40.70	846	2022	Azerbaijan	Soil	Steppe	7.80	942.50
MAZT4M02	48.57	40.85	1485	2022	Azerbaijan	Soil	Steppe	7.63	926.90
MAZT4M03	47.97	40.87	522	2022	Azerbaijan	Soil	Deciduous forest	7.74	1072.50
MAZT4M04	47.90	41.05	1956	2022	Azerbaijan	Soil	Deciduous forest	8.21	1638.00
MAZT4M05	47.81	41.09	1732	2022	Azerbaijan	Soil	Deciduous forest	7.83	390.00
MAZT4M06	47.43	41.06	517	2022	Azerbaijan	Soil	Thermophilous woodland	7.39	1519.05
MAZT4M07	47.13	41.30	898	2022	Azerbaijan	Soil	Deciduous forest	6.18	1318.20
MAZT4M08	46.56	41.70	626	2022	Azerbaijan	Soil	Deciduous forest	8.58	2496.00
MAZT4M09	46.47	41.63	242	2022	Azerbaijan	Soil	Deciduous forest	6.86	1462.50
MAZT4M10	46.77	41.58	643	2022	Azerbaijan	Soil	Deciduous forest	6.57	1092.65
MAZT4M11	46.79	41.31	204	2022	Azerbaijan	Soil	Deciduous forest	6.48	741.00
MAZT4M12	48.14	41.18	2150	2022	Azerbaijan	Soil	Steppe	6.96	487.50
MAZT4M13	48.44	41.26	900	2022	Azerbaijan	Soil	Deciduous forest	7.39	2262.00
MAZT4M14	48.71	40.81	1382	2022	Azerbaijan	Soil	Steppe	7.93	936.00
MAZT5M03	49.15	39.42	-28	2022	Azerbaijan	Soil	Shrub desert	7.51	1209.00
MAZT5M04	48.91	39.88	-22	2022	Azerbaijan	Soil	Halophytic desert	7.96	955.50
MAZT5M05	48.95	40.08	58	2022	Azerbaijan	Soil	Shrub desert	7.51	682.50
MAZT5S01	48.48	39.27	19	2022	Azerbaijan	Soil	Shrub desert	7.56	799.50
MAZT5S02	49.21	39.28	-28	2022	Azerbaijan	Soil	Halophytic desert	7.75	14872.65
MAZT5S06	49.34	40.90	-24	2022	Azerbaijan	Soil	Halophytic desert	7.28	7936.50
MAZT5S07	49.22	40.57	493	2022	Azerbaijan	Soil	Shrub desert	4.17	4368.00
MAZT6M01	48.66	38.86	85	2022	Azerbaijan	Soil	Hyrcanian forest	6.11	546.00
MAZT6M02	48.74	38.77	84	2022	Azerbaijan	Soil	Hyrcanian forest	6.35	390.00
MCNT1M01	100.54	36.32	3012	2016	China, Qinghai	Soil	<i>Achnatherum splendens</i> steppe	5.98	2262.00
MCNT1M04	100.59	36.41	3545	2016	China, Qinghai	Soil	<i>Kobresia pygmaea</i> alpine meadow	7.17	1917.50
MCNT1M05	100.47	36.54	3736	2016	China, Qinghai	Soil	<i>Stipa breviflora</i> desert steppe	7.01	3861.00
MCNT1M06	100.47	36.55	3519	2016	China, Qinghai	Soil	<i>Stipa breviflora</i> desert steppe	7.21	5660.20
MCNT1M07	100.47	36.55	3519	2016	China, Qinghai	Soil	<i>Stipa breviflora</i> desert steppe	7.76	1622.40
MCNT1M09	100.72	36.54	3205	2016	China, Qinghai	Soil	<i>Achnatherum splendens</i> steppe	7.75	1501.50
MCNT1M11	100.79	36.81	3240	2016	China, Qinghai	Soil	<i>Achnatherum splendens</i> steppe	7.91	745.55
MCNT1M13	100.62	37.09	3329	2016	China, Qinghai	Soil	<i>Kobresia pygmaea</i> alpine meadow	7.89	1625.65
MCNT1M14	100.64	37.25	3393	2016	China, Qinghai	Soil	<i>Stipa purpurea</i> alpine steppe		
MCNT1M15	100.68	37.33	3511	2016	China, Qinghai	Soil	<i>Dasiphora fruticosa</i> scrub	7.52	4395.30
MCNT1M16	100.75	37.32	3613	2016	China, Qinghai	Soil	<i>Dasiphora fruticosa</i> scrub		
MCNT1M17	100.80	37.33	3724	2016	China, Qinghai	Soil	<i>Dasiphora fruticosa</i> scrub	7.81	4894.50
MCNT1M18	100.80	37.36	3811	2016	China, Qinghai	Soil	<i>Dasiphora fruticosa</i> scrub	7.53	5005.00
MCNT1M19	100.82	37.38	4038	2016	China, Qinghai	Soil	<i>Saussurea</i> spp. sparse vegetation	7.45	2496.00
MCNT1M20	100.82	37.39	4038	2016	China, Qinghai	Soil	<i>Saussurea</i> spp. sparse vegetation		
MCNT1M21	100.82	37.39	4038	2016	China, Qinghai	Soil	<i>Saussurea</i> spp. sparse vegetation		
MCNT1M22	101.10	37.48	4012	2016	China, Qinghai	Soil	<i>Dasiphora fruticosa</i> scrub		
MCNT1M23	101.10	37.48	3936	2016	China, Qinghai	Soil	<i>Dasiphora fruticosa</i> scrub		
MCNT1M24	101.11	37.49	3936	2016	China, Qinghai	Soil	<i>Dasiphora fruticosa</i> scrub	4.41	9607.65
MCNT1M25	101.11	37.50	3814	2016	China, Qinghai	Soil	<i>Dasiphora fruticosa</i> scrub	5.34	5239.00
MCNT1M26	101.12	37.51	3670	2016	China, Qinghai	Soil	<i>Kobresia pygmaea</i> alpine meadow	6.40	3744.00
MCNT1M27	101.15	37.51	3541	2016	China, Qinghai	Soil	<i>Kobresia pygmaea</i> alpine meadow	6.00	4430.40
MCNT1M28	101.19	37.50	3326	2016	China, Qinghai	Soil	<i>Kobresia pygmaea</i> alpine meadow	6.15	9701.25
MCNT1M31	101.32	37.66	3322	2016	China, Qinghai	Soil	<i>Dasiphora fruticosa</i> scrub	7.10	8927.10
MCNT1M32	101.20	37.77	3418	2016	China, Qinghai	Soil	<i>Dasiphora fruticosa</i> scrub	6.87	5265.00
MCNT1M33	101.14	37.83	3671	2016	China, Qinghai	Soil	<i>Kobresia pygmaea</i> alpine meadow	7.64	5705.70
MCNT1M34	101.11	37.84	3758	2016	China, Qinghai	Soil	<i>Kobresia pygmaea</i> alpine meadow	6.82	4664.40
MCNT1M35	101.07	37.86	3590	2016	China, Qinghai	Soil	<i>Kobresia pygmaea</i> alpine meadow	6.55	9272.25
MCNT1M36	100.94	38.16	3049	2016	China, Qinghai	Soil	<i>Salix gilashanica</i> scrub	6.96	7616.70
MCNT1M37	100.93	38.19	3005	2016	China, Qinghai	Soil	<i>Salix gilashanica</i> scrub	7.18	6674.20
MCNT1M38	100.93	38.25	2780	2016	China, Qinghai	Soil	<i>Kobresia</i> spp., <i>Carex</i> spp. alpine meadow	8.20	1131.00
MCNT1M39	100.94	38.30	2646	2016	China, Qinghai	Soil	<i>Stipa breviflora</i> , <i>S. bungeana</i> steppe	7.40	4290.00
MCNT1M40	100.91	38.37	2492	2016	China, Qinghai	Soil	<i>Stipa breviflora</i> , <i>S. bungeana</i> steppe	8.50	1657.50
MCNT1M41	101.04	38.39	2420	2016	China, Qinghai	Soil	<i>Stipa breviflora</i> , <i>S. bungeana</i> steppe	7.75	12421.50
MCNT1M42	101.09	38.46	2265	2016	China, Qinghai	Soil	<i>Stipa breviflora</i> , <i>S. bungeana</i> steppe	7.94	1565.20

MCNT1M47	101.11	38.87	2276	2016	China, Qinghai	Soil	Dwarf needlegrass desert steppe	8.79	1039.35
MCNT1S02	100.59	36.37	3148	2016	China, Qinghai	Soil	<i>Kobresia pygmaea</i> alpine meadow	7.68	273.00
MCNT1S03	100.59	36.39	3357	2016	China, Qinghai	Soil	<i>Kobresia pygmaea</i> alpine meadow	7.54	399.75
MCNT1S10	100.79	36.71	3235	2016	China, Qinghai	Soil	<i>Artemisia arenaria</i> desert	8.97	104.00
MCNT1S12	100.58	37.04	3198	2016	China, Qinghai	Soil	<i>Achnatherum splendens</i> steppe	9.13	5038.80
MCNT1S29	101.23	37.52	3127	2016	China, Qinghai	Soil	<i>Dasiphora fruticosa</i> scrub	9.10	1296.75
MCNT1S30	101.25	37.62	3161	2016	China, Qinghai	Soil	<i>Dasiphora fruticosa</i> scrub	8.53	1344.20
MCNT1S45	101.09	38.84	1903	2016	China, Qinghai	Soil	Dwarf needlegrass desert steppe	9.30	390.00
MCNT1S48	101.09	38.88	2235	2016	China, Qinghai	Soil	<i>Stipa breviflora</i> , <i>S. bungeana</i> steppe	9.48	358.80
MCNT1S49	101.11	38.90	2057	2016	China, Qinghai	Soil	<i>Stipa breviflora</i> , <i>S. bungeana</i> steppe	9.04	2639.00
MCNT1S50	101.18	38.96	1885	2016	China, Qinghai	Soil	Dwarf needlegrass desert steppe	9.56	1394.90
MCNT1S53	101.39	39.13	1499	2016	China, Qinghai	Soil	<i>Reaumuria soongorica</i> desert	10.35	304.20
MCNT1S54	101.50	39.19	1435	2016	China, Qinghai	Soil	<i>Reaumuria soongorica</i> desert	9.91	764.40
MTAT1M01	71.52	38.86	2624	2021	Tajikistan	Soil	<i>Yailau</i> cryophilous open woodlands	7.24	666.90
MTAT1M02	71.60	38.81	2799	2021	Tajikistan	Soil	<i>Yailau</i> cryophilous open woodlands	6.76	702.00
MTAT1M03	71.69	38.76	3046	2021	Tajikistan	Soil	<i>Yailau</i> cryophilous open woodlands	6.82	382.20
MTAT1M06	71.88	38.51	3984	2021	Tajikistan	Soil	<i>Yailau</i> mesic grasslands	5.92	2068.30
MTAT1M07	71.88	38.45	3164	2021	Tajikistan	Soil	<i>Yailau</i> cryophilous open woodlands	7.18	2359.50
MTAT1M08	71.83	38.38	2472	2021	Tajikistan	Soil	<i>Yailau</i> cryophilous open woodlands	6.95	2223.00
MTAT1M09	71.79	38.37	2397	2021	Tajikistan	Soil	<i>Yailau</i> cryophilous open woodlands	7.41	838.50
MTAT2M01	68.23	38.65	1220	2022	Tajikistan	Soil	<i>Tau</i> xeric shrublands	6.56	6006.00
MTAT2M02	69.98	38.11	1251	2022	Tajikistan	Soil	<i>Tau</i> xeric shrublands	7.41	915.20
MTAT2M03	69.97	38.86	1150	2022	Tajikistan	Soil	<i>Tau</i> xeric shrublands	7.01	1774.50
MTAT2M04	70.03	38.86	1198	2022	Tajikistan	Soil	<i>Tau</i> xeric shrublands	7.23	1150.50
MTAT2M05	68.52	39.21	2052	2022	Tajikistan	Soil	<i>Tau</i> steppes	6.91	4712.50
MUZT1C04	69.84	41.40	1418	2021	Uzbekistan	Lacustrine	<i>Tau</i> xeric shrublands	7.83	2574.00
MUZT1C05	69.85	41.42	1272	2021	Uzbekistan	Lacustrine	<i>Tau</i> open woodlands	8.19	6006.00
MUZT1M01	69.84	41.40	1420	2021	Uzbekistan	Soil	<i>Tau</i> open woodlands	7.86	546.00
MUZT1M06	70.14	41.15	1646	2022	Uzbekistan	Soil	<i>Tau</i> open woodlands	7.06	1433.25
MUZT1S02	69.85	41.42	1291	2021	Uzbekistan	Soil	<i>Tau</i> steppes	8.00	741.00
MUZT1S03	69.85	41.42	1284	2021	Uzbekistan	Soil	<i>Tau</i> steppes	7.89	2676.70
MUZT2C01	66.88	39.33	1796	2021	Uzbekistan	Lacustrine	<i>Tau</i> steppes	6.66	897.00
MUZT2M02	66.88	39.33	1801	2021	Uzbekistan	Soil	<i>Tau</i> steppes	6.63	312.00
MUZT2M04	67.39	39.83	816	2022	Uzbekistan	Soil	<i>Adyr</i> steppes	7.04	1170.00
MUZT2M06	66.92	39.28	1424	2022	Uzbekistan	Soil	<i>Tau</i> xeric shrublands	7.76	1338.35
MUZT2M07	67.41	39.54	870	2022	Uzbekistan	Soil	<i>Tau</i> xeric shrublands	7.25	526.50
MUZT2M08	66.52	39.50	912	2022	Uzbekistan	Soil	<i>Tau</i> xeric shrublands	7.18	1158.30
MUZT2S03	66.88	39.35	1727	2021	Uzbekistan	Soil	<i>Tau</i> steppes	6.53	251.55
MUZT2S05	66.19	39.86	516	2022	Uzbekistan	Soil	<i>Adyr</i> pseudosteppes	7.80	507.00
MUZT3C02	66.70	38.13	1646	2021	Uzbekistan	Lacustrine	<i>Tau</i> open woodlands	8.18	1014.00
MUZT3C04	66.71	38.12	1847	2021	Uzbekistan	Lacustrine	<i>Tau</i> open woodlands	7.98	9165.00
MUZT3C19	66.73	38.08	1672	2023	Uzbekistan	Lacustrine	<i>Tau</i> open woodlands		
MUZT3M03	66.70	38.12	1740	2021	Uzbekistan	Soil	<i>Tau</i> open woodlands	7.71	624.00
MUZT3M05	67.76	38.62	1582	2021	Uzbekistan	Soil	<i>Tau</i> open woodlands	7.38	760.50
MUZT3M07	67.69	38.66	3038	2021	Uzbekistan	Soil	<i>Yailau</i> mesic grasslands	7.51	1384.50
MUZT3M08	67.66	38.64	3121	2021	Uzbekistan	Soil	<i>Yailau</i> mesic grasslands	7.45	841.75
MUZT3M09	66.91	38.21	1236	2022	Uzbekistan	Soil	<i>Adyr</i> steppes	7.22	1228.50
MUZT3M10	66.73	38.09	1709	2022	Uzbekistan	Soil	<i>Tau</i> open woodlands	7.11	2457.00
MUZT3M11	66.77	38.68	1308	2022	Uzbekistan	Soil	<i>Tau</i> xeric shrublands	6.73	5253.95
MUZT3M12	67.03	38.21	1077	2022	Uzbekistan	Soil	<i>Tau</i> xeric shrublands	7.31	663.00
MUZT3M13	66.73	37.61	786	2022	Uzbekistan	Soil	<i>Adyr</i> pseudosteppes	7.30	1228.50
MUZT3M14	66.76	37.77	958	2022	Uzbekistan	Soil	<i>Adyr</i> steppes	6.86	3178.50
MUZT3M15	66.74	37.63	664	2022	Uzbekistan	Soil	<i>Adyr</i> steppes	7.28	838.50
MUZT3M16	66.79	37.79	936	2022	Uzbekistan	Soil	<i>Adyr</i> steppes	7.74	2905.50
MUZT3M17	66.69	37.63	713	2022	Uzbekistan	Soil	<i>Adyr</i> steppes	7.47	897.00
MUZT3M18	66.70	37.78	1130	2022	Uzbekistan	Soil	<i>Tau</i> xeric shrublands	7.49	1774.50
MUZT3S06	67.71	38.65	2442	2021	Uzbekistan	Soil	<i>Yailau</i> cryophilous open woodlands	6.87	1625.00
MUZT4C02	64.22	39.64	200	2021	Uzbekistan	Lacustrine	<i>Chul</i> deserts	9.02	121649.45
MUZT4C04	64.24	39.63	208	2021	Uzbekistan	Lacustrine	<i>Chul</i> deserts	8.89	122995.60
MUZT4M05	64.91	40.09	292	2022	Uzbekistan	Soil	<i>Adyr</i> pseudosteppes	7.43	526.50
MUZT4M09	64.31	39.11	272	2022	Uzbekistan	Soil	<i>Chul</i> deserts	7.54	604.50
MUZT4S01	65.10	39.33	280	2021	Uzbekistan	Soil	<i>Chul</i> deserts	7.28	11778.00
MUZT4S03	64.22	39.64	204	2021	Uzbekistan	Soil	<i>Chul</i> deserts	8.48	141255.40
MUZT4S06	64.86	38.96	272	2022	Uzbekistan	Soil	<i>Chul</i> deserts	7.91	273.00
MUZT4S07	65.05	39.06	288	2022	Uzbekistan	Soil	<i>Chul</i> deserts	8.24	1111.50
MUZT5C02	63.21	41.77	53	2021	Uzbekistan	Soil	<i>Chul</i> deserts	8.33	57943.60
MUZT5M03	63.85	42.52	366	2022	Uzbekistan	Soil	<i>Chul</i> deserts	7.55	916.50
MUZT5M04	64.15	42.93	257	2022	Uzbekistan	Soil	<i>Chul</i> deserts	7.75	320.45
MUZT5M05	62.82	42.31	97	2022	Uzbekistan	Soil	<i>Chul</i> deserts	7.80	487.50
MUZT5M06	63.43	42.20	194	2022	Uzbekistan	Soil	<i>Chul</i> deserts	7.54	604.50
MUZT5S08	63.27	40.27	180	2022	Uzbekistan	Soil	<i>Chul</i> deserts	8.34	390.00
MUZT6C02	66.59	40.55	1644	2021	Uzbekistan	Lacustrine	<i>Tau</i> steppes	7.75	3336.45
MUZT6M03	66.63	40.57	1166	2021	Uzbekistan	Soil	<i>Tau</i> xeric shrublands	7.66	955.50
MUZT6M04	66.66	40.58	836	2021	Uzbekistan	Soil	<i>Tau</i> xeric shrublands	7.92	2145.00
MUZT6S01	66.62	40.57	1460	2021	Uzbekistan	Soil	<i>Tau</i> steppes	7.62	487.50
MUZT7C01	58.99	44.86	38	2023	Uzbekistan	Lacustrine	<i>Chul</i> deserts	9.11	237994.90
MUZT7C02	58.49	44.80	29	2023	Uzbekistan	Lacustrine	<i>Chul</i> deserts	8.25	77231.70
MUZT7C03	58.65	44.73	31	2023	Uzbekistan	Lacustrine	<i>Chul</i> deserts	8.82	36718.50
MUZT7C04	58.73	44.73	29	2023	Uzbekistan	Lacustrine	<i>Chul</i> deserts	8.72	241833.80

Table S2: List of the different references compiled mainly from Raberg et al. (2022) plus other studies; to build the ACADB and the WDB used in this study.

DB	Reference	N _{site}	N _{lac.}	N _{soil}	Geographical area	Citation
ACA	Chen et al. (2021)	53	0	53	Tajikistan	Chen, C., Bai, Y., Fang, X., Zhuang, G., Khodzhev, A., Bai, X., and Murodov, A.: Evaluating the Potential of Soil Bacterial Tetraether Proxies in Westerlies Dominating Western Pamirs, Tajikistan and Implications for Paleoenvironmental Reconstructions, <i>Chemical Geology</i> , 559, 119 908, https://doi.org/10.1016/j.chemgeo.2020.119908 , 2021
ACA	Dang et al. (2018)	10	10	0	China	Dang, X., Ding, W., Yang, H., Pancost, R. D., Naafs, B. D. A., Xue, J., Lin, X., Lu, J., and Xie, S.: Different Temperature Dependence of the Bacterial brGDGT Isomers in 35 Chinese Lake Sediments Compared to That in Soils, <i>Organic Geochemistry</i> , 119, 72–79, https://doi.org/10/gdfs3x , 2018
ACA	Cromartie et al. (2025)	22	0	22	Armenia	Cromartie, A., De Jonge, C., Ménot, G., Robles, M., Dugerdil, L., Peyron, O., Rodrigo-Gámiz, M., Camuera, J., Ramos-Roman, M. J., Jiménez-Moreno, G., Colombié, C., Sahakyan, L., and Joannin, S.: Utilizing Probability Estimates from Machine Learning and Pollen to Understand the Depositional Influences on Branched GDGT in Wetlands, Peatlands, and Lakes, https://doi.org/10.5194/egusphere-2025-526 , 2025
ACA	Dearing Crampton-Flood et al. (2020)	15	0	15	Global BayMBT Soils	Dearing Crampton-Flood, E., Tierney, J. E., Peterse, F., Kirkels, F. M. S. A., and Sinninghe Damsté, J. S.: BayMBT: A Bayesian Calibration Model for Branched Glycerol Dialkyl Glycerol Tetraethers in Soils and Peats, <i>Geochimica et Cosmochimica Acta</i> , 268, 142–159, https://doi.org/10/gg9758 , 2020
ACA	Duan et al. (2020)	18	0	18	China, Tian Shan	Duan, Y., Sun, Q., Werne, J. P., Yang, H., Jia, J., Wang, L., Xie, H., and Chen, F.: Soil pH Dominates the Distributions of Both 5- and 6-Methyl Branched Tetraethers in Arid Regions, <i>Journal of Geophysical Research: Biogeosciences</i> , 125, e2019JG005 356, https://doi.org/10.1029/2019jg005356 , 2020
ACA	Duan et al. (2021)	48	0	48	Northern Iran	Duan, Y.: Northern Iran and Global Soil brGDGT Dataset, https://doi.org/10.11888/Paleoenv.tpd.271742 , 2021
ACA	Dugerdil et al. (2021a)	44	2	42	Mongolia, Baikal	Dugerdil, L., Joannin, S., Peyron, O., Jouffroy-Bapicot, I., Vanni�re, B., Boldgiv, B., Unkelbach, J., Behling, H., and M�not, G.: Climate Reconstructions Based on GDGT and Pollen Surface Datasets from Mongolia and Baikal Area: Calibrations and Applicability to Extremely Cold–Dry Environments over the Late Holocene, <i>Climate of the Past</i> , 17, 1199–1226, https://doi.org/10.5194/cp-17-1199-2021 , 2021
ACA	Guo et al. (2021)	32	0	32	China, Inner Mong.	Guo, J., Ma, T., Liu, N., Zhang, X., Hu, H., Ma, W., Wang, Z., Feng, X., and Peterse, F.: Branched Tetraether Lipids and Bacterial Communities along an Aridity Soil Transect in Inner Mongolia, Northern China, https://doi.org/10.1594/PANGAEA.938067 , 2021
ACA	Kou et al. (2022)	129	107	22	China, Tibet	Kou, Q., Zhu, L., Ju, J., Wang, J., Xu, T., Li, C., and Ma, Q.: Influence of Salinity on Glycerol Dialkyl Glycerol Tetraether-Based Indicators in Tibetan Plateau Lakes: Implications for Paleotemperature and Paleosalinity Reconstructions, <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 601, 111 127, https://doi.org/10.1016/j.palaeo.2022.111127 , 2022
ACA	Li et al. (2017)	11	0	11	China, Inner Mong.	Li, J., Naafs, B. D. A., Pancost, R. D., Yang, H., Liu, D., and Xie, S.: Distribution of Branched Tetraether Lipids in Ponds from Inner Mongolia, NE China: Insight into the Source of brGDGTs, <i>Organic Geochemistry</i> , 112, 127–136, https://doi.org/10.1016/j.orggeochem.2017.07.005 , 2017
ACA	Naafs et al. (2017a)	48	0	48	Global soils	Naafs, B., Gallego-Sala, A., Inglis, G., and Pancost, R.: Refining the Global Branched Glycerol Dialkyl Glycerol Tetraether (brGDGT) Soil Temperature Calibration, <i>Organic Geochemistry</i> , 106, 48–56, https://doi.org/10/gbjssd , 2017a
ACA	Wang et Liu (2021)	16	0	16	China	Wang, H. and Liu, W.: Soil Temperature and brGDGTs along an Elevation Gradient on the Northeastern Tibetan Plateau: A Test of Soil brGDGTs as a Proxy for Paleoelevation, <i>Chemical Geology</i> , 566, 120 079, https://doi.org/10.1016/j.chemgeo.2021.120079 , 2021
ACA	Wang et al. (2020)	75	0	75	China	Wang, H., An, Z., Lu, H., Zhao, Z., and Liu, W.: Calibrating Bacterial Tetraether Distributions towards in Situ Soil Temperature and Application to a Loess–Paleosol Sequence, <i>Quaternary Science Reviews</i> , 231, 106 172, https://doi.org/10.1016/j.quascirev.2020.106172 , 2020
ACA	Wang et al. (2021)	52	52	0	ACA lakes	Wang, H., Liu, W., He, Y., Zhou, A., Zhao, H., Liu, H., Cao, Y., Hu, J., Meng, B., Jiang, J., Kolpakova, M., Krivonogov, S., and Liu, Z.: Salinity-Controlled Isomerization of Lacustrine brGDGTs Impacts the Associated MBT5ME' Terrestrial Temperature Index, <i>Geochimica et Cosmochimica Acta</i> , 305, 33–48, https://doi.org/10.1016/j.gca.2021.05.004 , 2021
ACA	Zang et al. (2018)	19	0	19	China	Zang, J., Lei, Y., and Yang, H.: Distribution of Glycerol Ethers in Turpan Soils: Implications for Use of GDGT-based Proxies in Hot and Dry Regions, <i>Frontiers of Earth Science</i> , 12, 862–876, https://doi.org/10/gfb6xq , 2018
World DB	Cao et al. (2020)	6	1	5	China	Cao, J., Rao, Z., Shi, F., and Jia, G.: Ice Formation on Lake Surfaces in Winter Causes Warm-Season Bias of Lacustrine brGDGT Temperature Estimates, <i>Biogeosciences</i> , 17, 2521–2536, https://doi.org/10.5194/bg-17-2521-2020 , 2020
World DB	Dang et al. (2018)	32	32	0	China	Dang, X., Ding, W., Yang, H., Pancost, R. D., Naafs, B. D. A., Xue, J., Lin, X., Lu, J., and Xie, S.: Different Temperature Dependence of the Bacterial brGDGT Isomers in 35 Chinese Lake Sediments Compared to That in Soils, <i>Organic Geochemistry</i> , 119, 72–79, https://doi.org/10/gdfs3x , 2018

World DB	Dang et al. (2018)	32	32	0	Russia, Baikal	Dang, X., Ding, W., Yang, H., Pancost, R. D., Naafs, B. D. A., Xue, J., Lin, X., Lu, J., and Xie, S.: Different Temperature Dependence of the Bacterial brGDGT Isomers in 35 Chinese Lake Sediments Compared to That in Soils, <i>Organic Geochemistry</i> , 119, 72–79, https://doi.org/10/gdfs3x , 2018
World DB	De Jonge et al. (2019)	30	0	30	Iceland ForHot	De Jonge, C., Radujković, D., Sigurdsson, B. D., Weedon, J. T., Janssens, I., and Peterse, F.: Lipid Biomarker Temperature Proxy Responds to Abrupt Shift in the Bacterial Community Composition in Geothermally Heated Soils, <i>Organic Geochemistry</i> , 137, 103 897, https://doi.org/10.1016/j.orggeochem.2019.07.006 , 2019
World DB	De Jonge et al. (2021)	25	0	25	Netherlands	De Jonge, C., Kuramae, E. E., Radujković, D., Weedon, J. T., Janssens, I. A., and Peterse, F.: The Influence of Soil Chemistry on Branched Tetraether Lipids in Mid-and High Latitude Soils: Implications for brGDGT-based Paleothermometry, <i>Geochimica et Cosmochimica Acta</i> , 310, 95–112, https://doi.org/10.1016/j.gca.2021.06.037 , 2021
World DB	De Jonge et al. (2021)	25	0	25	Scotland	De Jonge, C., Kuramae, E. E., Radujković, D., Weedon, J. T., Janssens, I. A., and Peterse, F.: The Influence of Soil Chemistry on Branched Tetraether Lipids in Mid-and High Latitude Soils: Implications for brGDGT-based Paleothermometry, <i>Geochimica et Cosmochimica Acta</i> , 310, 95–112, https://doi.org/10.1016/j.gca.2021.06.037 , 2021
World DB	Dearing Crampton-Flood et al. (2020)	61	0	61	Global BayMBT Soils	Dearing Crampton-Flood, E., Tierney, J. E., Peterse, F., Kirkels, F. M. S. A., and Sinnighe Damsté, J. S.: BayMBT: A Bayesian Calibration Model for Branched Glycerol Dialkyl Glycerol Tetraethers in Soils and Peats, <i>Geochimica et Cosmochimica Acta</i> , 268, 142–159, https://doi.org/10/gg9758 , 2020
World DB	Ding et al. (2018)	22	0	22	Germany	Ding, S., Kohlhepp, B., Trumbore, S., Küsel, K., Totsche, K.-U., Pohnert, G., Gleixner, G., and Schwab, V. F.: In Situ Production of Core and Intact Bacterial and Archaeal Tetraether Lipids in Groundwater, <i>Organic Geochemistry</i> , 126, 1–12, https://doi.org/10.1016/j.orggeochem.2018.10.005 , 2018
World DB	Guo et al. (2020)	74	0	74	England	Guo, J., Glendell, M., Meersmans, J., Kirkels, F., Middelburg, J. J., and Peterse, F.: Assessing Branched Tetraether Lipids as Tracers of Soil Organic Carbon Transport through the Carminow Creek Catchment (Southwest England), <i>Biogeosciences</i> , 17, 3183–3201, https://doi.org/10.5194/bg-17-3183-2020 , 2020
World DB	Kirkels et al. (2019)	28	0	28	Amazon	Kirkels, F. M. S. A., Ponton, C., Galy, V., West, A. J., Feakins, S. J., and Peterse, F.: From Andes to Amazon: Assessing Branched Tetraether Lipids as Tracers for Soil Organic Carbon in the Madre de Dios River System, <i>Journal of Geophysical Research: Biogeosciences</i> , 125, e2019JG005 270, https://doi.org/10.1029/2019JG005270 , 2020
World DB	Kusch et al. (2019)	17	0	17	Siberia	Kusch, S., Winterfeld, M., Mollenhauer, G., Höfle, S. T., Schirrmeister, L., Schwamborn, G., and Rethemeyer, J.: Glycerol Dialkyl Glycerol Tetraethers (GDGTs) in High Latitude Siberian Permafrost: Diversity, Environmental Controls, and Implications for Proxy Applications, <i>Organic Geochemistry</i> , 136, 103 888, https://doi.org/10.1016/j.orggeochem.2019.06.009 , 2019
World DB	Li et al. (2018)	29	0	29	China	Li, Y., Zhao, S., Pei, H., Qian, S., Zang, J., Dang, X., and Yang, H.: Distribution of Glycerol Dialkyl Glycerol Tetraethers in Surface Soils along an Altitudinal Transect at Cold and Humid Mountain Changbai: Implications for the Reconstruction of Paleoalimetry and Paleoclimate, <i>Science China Earth Sciences</i> , 61, 925–939, https://doi.org/10.1007/s11430-017-9168-9 , 2018
World DB	Martínez-Sosa et al. (2021)	158	158	0	Global	Martínez-Sosa, P., Tierney, J. E., Stefanescu, I. C., Crampton-Flood, E. D., Shuman, B. N., and Routson, C.: A Global Bayesian Temperature Calibration for Lacustrine brGDGTs, <i>Geochimica et Cosmochimica Acta</i> , 305, 87–105, https://doi.org/10.1016/j.gca.2021.04.038 , 2021
World DB	Miller et al. (2018)	11	1	10	USA	Miller, D. R., Habicht, M. H., Keisling, B. A., Castañeda, I. S., and Bradley, R. S.: A 900-Year New England Temperature Reconstruction from in Situ Seasonally Produced Branched Glycerol Dialkyl Glycerol Tetraethers (brGDGTs), <i>Climate of the Past</i> , 14, 1653–1667, https://doi.org/10.5194/cp-14-1653-2018 , 2018
World DB	Naafs et al. (2017a)	302	0	302	Global soils	Naafs, B., Gallego-Sala, A., Inglis, G., and Pancost, R.: Refining the Global Branched Glycerol Dialkyl Glycerol Tetraether (brGDGT) Soil Temperature Calibration, <i>Organic Geochemistry</i> , 106, 48–56, https://doi.org/10/gbjssd , 2017a
World DB	Ning et al. (2019)	2	1	1	China	Ning, D., Zhang, E., Shulmeister, J., Chang, J., Sun, W., and Ni, Z.: Holocene Mean Annual Air Temperature (MAAT) Reconstruction Based on Branched Glycerol Dialkyl Glycerol Tetraethers from Lake Ximenglongtan, Southwestern China, <i>Organic Geochemistry</i> , https://doi.org/10.1016/j.orggeochem.2019.05.003 , 2019
World DB	Pei et al. (2019)	10	0	10	China	Pei, H., Wang, C., Wang, Y., Yang, H., and Xie, S.: Distribution of Microbial Lipids at an Acid Mine Drainage Site in China: Insights into Microbial Adaptation to Extremely Low pH Conditions, <i>Organic Geochemistry</i> , 134, 77–91, https://doi.org/10.1016/j.orggeochem.2019.05.008 , 2019
World DB	Pérez-Angel et al. (2020)	36	0	36	Colombia	Pérez-Angel, L. C., Sepúlveda, J., Molnar, P., Montes, C., Rajagopalan, B., Snell, K., Gonzalez-Arango, C., and Dildar, N.: Soil and Air Temperature Calibrations Using Branched GDGTs for the Tropical Andes of Colombia: Toward a Pan-Tropical Calibration, <i>Geochimica et Cosmochimica Acta</i> , 21, e2020GC008 941, https://doi.org/10.1029/2020GC008941 , 2020
World DB	Qian et al. (2019)	6	1	5	China	Qian, S., Yang, H., Dong, C., Wang, Y., Wu, J., Pei, H., Dang, X., Lu, J., Zhao, S., and Xie, S.: Rapid Response of Fossil Tetraether Lipids in Lake Sediments to Seasonal Environmental Variables in a Shallow Lake in Central China: Implications for the Use of Tetraether-Based Proxies, <i>Organic Geochemistry</i> , 128, 108–121, https://doi.org/10.1016/j.orggeochem.2018.12.007 , 2019

World DB	Raberg et al. (2021)	43	43	0	Baffin	Raberg, J. H., Harning, D. J., Crump, S. E., de Wet, G., Blumm, A., Kopf, S., Geirsdóttir, Á., Miller, G. H., and Sepúlveda, J.: Revised Fractional Abundances and Warm-Season Temperatures Substantially Improve brGDGT Calibrations in Lake Sediments, <i>Biogeosciences</i> , 18, 3579–3603, https://doi.org/10.5194/bg-18-3579-2021 , 2021
World DB	Raberg et al. (2021)	43	43	0	Iceland	Raberg, J. H., Harning, D. J., Crump, S. E., de Wet, G., Blumm, A., Kopf, S., Geirsdóttir, Á., Miller, G. H., and Sepúlveda, J.: Revised Fractional Abundances and Warm-Season Temperatures Substantially Improve brGDGT Calibrations in Lake Sediments, <i>Biogeosciences</i> , 18, 3579–3603, https://doi.org/10.5194/bg-18-3579-2021 , 2021
World DB	Russell et al. (2018)	65	65	0	Africa	Russell, J. M., Hopmans, E. C., Loomis, S. E., Liang, J., and Damsté, J. S. S.: Distributions of 5- and 6-Methyl Branched Glycerol Dialkyl Glycerol Tetraethers (brGDGTs) in East African Lake Sediment: Effects of Temperature, pH, and New Lacustrine Paleotemperature Calibrations, <i>Organic Geochemistry</i> , 117, 56–69, https://doi.org/10.1016/j.orggeochem.2017.12.003 , 2018
World DB	Véquaud et al. (2021)	49	0	49	France	Véquaud, P., Derenne, S., Anquetil, C., Collin, S., Poulenard, J., Sabatier, P., and Huguet, A.: Influence of Environmental Parameters on the Distribution of Bacterial Lipids in Soils from the French Alps: Implications for Paleo-Reconstructions, <i>Organic Geochemistry</i> , p. 104194, https://doi.org/10.1016/j.orggeochem.2021.104194 , 2021
World DB	Wang et al. (2018)	28	0	28	China	Wang, M., Zong, Y., Zheng, Z., Man, M., Hu, J., and Tian, L.: Utility of brGDGTs as Temperature and Precipitation Proxies in Subtropical China, <i>Scientific Reports</i> , 8, 1–9, https://doi.org/10.1038/s41598-017-17964-0 , 2018
World DB	Wang et al. (2019)	47	0	47	China	Wang, M., Zheng, Z., Zong, Y., Man, M., and Tian, L.: Distributions of Soil Branched Glycerol Dialkyl Glycerol Tetraethers from Different Climate Regions of China, <i>Scientific reports</i> , 9, 1–8, https://doi.org/10.1038/s41598-019-39147-9 , 2019
World DB	Wang et al. (2020)	74	0	74	China	Wang, H., An, Z., Lu, H., Zhao, Z., and Liu, W.: Calibrating Bacterial Tetraether Distributions towards in Situ Soil Temperature and Application to a Loess-Paleosol Sequence, <i>Quaternary Science Reviews</i> , 231, 106 172, https://doi.org/10.1016/j.quascirev.2020.106172 , 2020
World DB	Wang et al. (2021)	11	11	0	Russia, Baikal	Wang, H. and Liu, W.: Soil Temperature and brGDGTs along an Elevation Gradient on the Northeastern Tibetan Plateau: A Test of Soil brGDGTs as a Proxy for Paleoelevation, <i>Chemical Geology</i> , 566, 120 079, https://doi.org/10.1016/j.chemgeo.2021.120079 , 2021
World DB	Weber et al. (2018)	56	36	20	Switzerland	Weber, Y., Damsté, J. S. S., Zopfi, J., De Jonge, C., Gilli, A., Schubert, C. J., Lepori, F., Lehmann, M. F., and Niemann, H.: Redox-Dependent Niche Differentiation Provides Evidence for Multiple Bacterial Sources of Glycerol Tetraether Lipids in Lakes, <i>Proceedings of the National Academy of Sciences</i> , 115, 10 926–10 931, https://doi.org/10.1073/pnas.1805186115 , 2018

Table S3. Presentation of the different cuff-offs associated with each of the grouping factor used in this study.

	ID	Classes	Cut-off	Reference
Alkalinity	1	Acid	pH < 7	common classification
	2	Neutral	pH ∈ [7; 8[	
	3	Alkaline	pH ≥ 8	
Aridity	1	Hyper-arid	AI < 1500	Nash (1999)
	2	Arid	AI ∈ [1500; 3000[	
	3	Semi-arid	AI ∈ [3000; 3700[	
	4	Dry sub-humid	AI ∈ [3700; 5500[	
	5	Humid	AI ≥ 5500	
Salinity	1	Fresh	Salinity < 700	modified from Rusydi (2018) with a sensitivity analysis
	2	Hyposaline	Salinity ∈ [700; 26000[	
	3	Saline	Salinity ∈ [26; 140000[	
	4	Hypersaline	Salinity ≥ 140000	
Sample type	1	Soil		Martínez-Sosa et al. (2023)
	2	Lacustrine		

Table S4. Results of the Variance Factors analyses (VIF) carried out on the brGDGT FAs vs. climate parameter for model 1 (i.e., all the available climate parameters) and model 2 (after removing the covariant climate parameters owing to only having VIFs < 10).

Climate parameters	VIF (model 1)	VIF (model 2)
AI	21.60	3.00
Altitude	22.20	
MAAT	620.60	5.60
MAF	28.00	4.40
MAP	37.80	
MPCOQ	17.80	
MPWAQ	16.70	3.20
MTCOQ	159.10	
MTWAQ	271.60	
pH	1.70	1.70
Salinity	1.30	1.30

Table S5. Levene’s test results for the two MANOVA models used to evaluate the grouping factors effect across brGDGT FA and indices. The violation of the assumption of homogeneity of variance-covariance is marked by ** (p-values < 0.001) or * (p-values < 0.01)

	Alkalinity	Aridity	Salinity	Sample type
f(IIIa)			**	
f(IIIa')			**	*
f(IIIb)			*	
f(IIIb')				
f(IIIc)				**
f(IIIc')				*
f(IIa)		**	**	
f(IIa')				
f(IIb)				
f(IIb')			**	
f(IIc)				
f(IIc')				
f(Ia)	*	**		
f(Ib)				
f(Ic)				*
MBT' _{5Me}		**	*	
MBT' _{6Me}	**	*		
IR _{6Me}				
IR _{6+7Me}	*			
IR' _{6+7Me}	*		**	
CBT'		*		
CBT' _{5Me}		**		*

Table S6. Main GDGT indices discussed in this study with their formula, proxy interpretation and references.

Index	Formula	RMSE	Proxy interpretation	References
MAAT _{Tjk-Chen}	$= 1.1 + 19.2 \cdot \text{MBT}'_{5\text{Me}}$	2.3	Soil, local (Tajikistan)	Chen et al. (2021)
MAAT _{L-Sun}	$= 3.949 - 5.593 \cdot \text{CBT} + 38.213 \cdot \text{MBT}'_{5\text{Me}}$	4.27	Lacustrine, local (China)	Sun et al. (2011)
MAAT _{mr-DJ}	$= 7.17 + 17.1 \cdot [\text{Ia}] + 25.9 \cdot [\text{Ib}] + 34.4 \cdot [\text{Ic}] - 28.6 \cdot [\text{IIa}]$	4.6	Soil, global	De Jonge et al. (2014)
MAAT _{mrs-DJ}	$= 5.58 + 17.91 \cdot [\text{Ia}] - 18.77 \cdot [\text{IIa}]$	5.0	Soil, global	De Jonge et al. (2014)
MAAT _{MBT-DJ}	$= -8.57 + 31.45 \cdot \text{MBT}'_{5\text{Me}}$	4.8	Soil, global	De Jonge et al. (2014)
MAAT _{MBT-Peterse}	$= 0.81 - 5.67 \cdot \text{CBT} + 31 \cdot \text{MBT}'$	5.0	Soil, global	Peterse et al. (2012)
MAAT _{MBT-Wang}	$= 27.63 \cdot \text{Index}_1 - 5.72$	2.5	Soil, local (north China)	Wang et al. (2016)
MAAT _{S-MBT-Naafs}	$= -15.25 + 40.01 \cdot \text{MBT}'_{5\text{Me}}$	5.3	Soil, global	Naafs et al. (2017a)
MAAT _{S-mr-Naafs}	$= 4.32 + 19.8 \cdot [\text{Ia}] + 31.1 \cdot [\text{Ib}] - 23.4 \cdot [\text{IIa}]$	4.7	Soil, global	Naafs et al. (2017a)
MAAT _{P-MBT-Naafs}	$= -23.05 + 52.18 \cdot \text{MBT}'_{5\text{Me}}$	4.7	Peatland, global	Naafs et al. (2017b)
MAAT _{mr-Yang}	$= 20.9 - 13.4 \cdot [\text{IIa}] - 17.2 \cdot [\text{IIIa}] - 17.5 \cdot [\text{IIb}] + 11.2 \cdot [\text{Ib}]$	1.7	Soil, local (arid China)	Yang et al. (2014)
MAAT _{mr-Thomas}	$= 20.87 - 18.04 \cdot [\text{IIIa}] + 38.88 \cdot [\text{IIIb}] - 12.92 \cdot [\text{IIa}]$ $- 20.41 \cdot [\text{IIb}] - 30.27 \cdot [\text{IIc}] + 12.94 \cdot [\text{Ib}]$	1.6	Soil, local (Tibet)	Thomas et al. (2017)
MAF _{Meth}	$= 92.9 + 63.84 \cdot [\text{Ib}]^2_{\text{m}} - 130.51 \cdot [\text{Ib}]_{\text{m}} - 28.77 \cdot [\text{IIa}]^2_{\text{m}} - 72.28 \cdot [\text{IIb}]^2_{\text{m}}$ $- 5.88 \cdot [\text{IIc}]^2_{\text{m}} + 20.89 \cdot [\text{IIIa}]^2_{\text{m}} - 40.54 \cdot [\text{IIIa}]_{\text{m}} - 80.47 \cdot [\text{IIIb}]_{\text{m}}$	2.14	Soil, global	Raberg et al. (2021)
MAF _{Full}	$= -8.06 + 37.52 \cdot [\text{Ia}] - 266.83 \cdot [\text{Ib}]^2 + 133.42 \cdot [\text{Ib}]$ $+ 100.85 \cdot [\text{IIa}]'^2 + 58.15 \cdot [\text{IIIa}]'^2 + 12.79 \cdot [\text{IIIa}]$	1.97	Soil, global	Raberg et al. (2021)
MAF _{L-Bay}	$= \frac{\text{MBT}'_{5\text{Me}} - 0.075}{0.03}$	0.089	Lacustrine, global	Martínez-Sosa et al. (2021)

References

- Cao, J., Rao, Z., Shi, F., and Jia, G.: Ice Formation on Lake Surfaces in Winter Causes Warm-Season Bias of Lacustrine brGDGT Temperature Estimates, *Biogeosciences*, 17, 2521–2536, <https://doi.org/10.5194/bg-17-2521-2020>, 2020.
- 5 Chen, C., Bai, Y., Fang, X., Zhuang, G., Khodzhiyev, A., Bai, X., and Murodov, A.: Evaluating the Potential of Soil Bacterial Tetraether Proxies in Westerlies Dominating Western Pamirs, Tajikistan and Implications for Paleoenvironmental Reconstructions, *Chemical Geology*, 559, 119908, <https://doi.org/10.1016/j.chemgeo.2020.119908>, 2021.
- Clogg, C. C., Petkova, E., and Haritou, A.: Statistical Methods for Comparing Regression Coefficients between Models, *American Journal Of Sociology*, 100, 1261–1293, <https://doi.org/10.1086/230638>, 1995.
- 10 Cromartie, A., De Jonge, C., Ménot, G., Robles, M., Dugerdil, L., Peyron, O., Rodrigo-Gámiz, M., Camuera, J., Ramos-Roman, M. J., Jiménez-Moreno, G., Colombié, C., Sahakyan, L., and Joannin, S.: Utilizing Probability Estimates from Machine Learning and Pollen to Understand the Depositional Influences on Branched GDGT in Wetlands, Peatlands, and Lakes, <https://doi.org/10.5194/egusphere-2025-526>, 2025.
- Dang, X., Ding, W., Yang, H., Pancost, R. D., Naafs, B. D. A., Xue, J., Lin, X., Lu, J., and Xie, S.: Different Temperature Dependence of the Bacterial brGDGT Isomers in 35 Chinese Lake Sediments Compared to That in Soils, *Organic Geochemistry*, 119, 72–79, <https://doi.org/10/gdfs3x>, 2018.
- 15 De Jonge, C., Hopmans, E. C., Zell, C. I., Kim, J.-H., Schouten, S., and Sinninghe Damsté, J. S.: Occurrence and Abundance of 6-Methyl Branched Glycerol Dialkyl Glycerol Tetraethers in Soils: Implications for Palaeoclimate Reconstruction, *Geochimica et Cosmochimica Acta*, 141, 97–112, <https://doi.org/10/f6h2xw>, 2014.

- 20 De Jonge, C., Radujković, D., Sigurdsson, B. D., Weedon, J. T., Janssens, I., and Peterse, F.: Lipid Biomarker Temperature Proxy Responds to Abrupt Shift in the Bacterial Community Composition in Geothermally Heated Soils, *Organic Geochemistry*, 137, 103 897, <https://doi.org/10.1016/j.orggeochem.2019.07.006>, 2019.
- De Jonge, C., Kuramae, E. E., Radujković, D., Weedon, J. T., Janssens, I. A., and Peterse, F.: The Influence of Soil Chemistry on Branched Tetraether Lipids in Mid-and High Latitude Soils: Implications for brGDGT-based Paleothermometry, *Geochimica et Cosmochimica Acta*, 310, 95–112, <https://doi.org/10.1016/j.gca.2021.06.037>, 2021.
- 25 Dearing Crampton-Flood, E., Tierney, J. E., Peterse, F., Kirkels, F. M. S. A., and Sinninghe Damsté, J. S.: BayMBT: A Bayesian Calibration Model for Branched Glycerol Dialkyl Glycerol Tetraethers in Soils and Peats, *Geochimica et Cosmochimica Acta*, 268, 142–159, <https://doi.org/10/gg9758>, 2020.
- Ding, S., Kohlhepp, B., Trumbore, S., Küsel, K., Totsche, K.-U., Pohnert, G., Gleixner, G., and Schwab, V. F.: In Situ Production of Core and Intact Bacterial and Archaeal Tetraether Lipids in Groundwater, *Organic Geochemistry*, 126, 1–12, <https://doi.org/10.1016/j.orggeochem.2018.10.005>, 2018.
- Duan, Y.: Northern Iran and Global Soil brGDGT Dataset, <https://doi.org/10.11888/Paleoenv.tpd.271742>, 2021.
- Duan, Y., Sun, Q., Werne, J. P., Yang, H., Jia, J., Wang, L., Xie, H., and Chen, F.: Soil pH Dominates the Distributions of Both 5- and 6-Methyl Branched Tetraethers in Arid Regions, *Journal of Geophysical Research: Biogeosciences*, 125, e2019JG005 356, <https://doi.org/10.1029/2019jg005356>, 2020.
- 35 Dugerdil, L., Joannin, S., Peyron, O., Jouffroy-Bapicot, I., Vanni re, B., Boldgiv, B., Unkelbach, J., Behling, H., and M not, G.: Climate Reconstructions Based on GDGT and Pollen Surface Datasets from Mongolia and Baikal Area: Calibrations and Applicability to Extremely Cold–Dry Environments over the Late Holocene, *Climate of the Past*, 17, 1199–1226, <https://doi.org/10.5194/cp-17-1199-2021>, 2021.
- Dugerdil, L., Joannin, S., Peyron, O., Cromartie, A., Robles, M., Duan, Y., Huang, X., Chen, F., and M not, G.: Boosted Regression Trees Machine Learning Method Improves the brGDGT-based Climate Reconstruction in Drylands, *Paleoceanography and Paleoclimatology*, 40, e2025PA005 214, <https://doi.org/10.1029/2025PA005214>, 2025.
- 40 Fick, S. E. and Hijmans, R. J.: WorldClim 2: New 1-Km Spatial Resolution Climate Surfaces for Global Land Areas: NEW CLIMATE SURFACES FOR GLOBAL LAND AREAS, *International Journal of Climatology*, 37, 4302–4315, <https://doi.org/10/gb2jnj>, 2017.
- Guo, J., Glendell, M., Meersmans, J., Kirkels, F., Middelburg, J. J., and Peterse, F.: Assessing Branched Tetraether Lipids as Tracers of Soil Organic Carbon Transport through the Carminowe Creek Catchment (Southwest England), *Biogeosciences*, 17, 3183–3201, <https://doi.org/10.5194/bg-17-3183-2020>, 2020.
- 45 Guo, J., Ma, T., Liu, N., Zhang, X., Hu, H., Ma, W., Wang, Z., Feng, X., and Peterse, F.: Branched Tetraether Lipids and Bacterial Communities along an Aridity Soil Transect in Inner Mongolia, Northern China, <https://doi.org/10.1594/PANGAEA.938067>, 2021.
- Kirkels, F. M. S. A., Ponton, C., Galy, V., West, A. J., Feakins, S. J., and Peterse, F.: From Andes to Amazon: Assessing Branched Tetraether Lipids as Tracers for Soil Organic Carbon in the Madre de Dios River System, *Journal of Geophysical Research: Biogeosciences*, 125, e2019JG005 270, <https://doi.org/10.1029/2019JG005270>, 2020.
- 50 Kou, Q., Zhu, L., Ju, J., Wang, J., Xu, T., Li, C., and Ma, Q.: Influence of Salinity on Glycerol Dialkyl Glycerol Tetraether-Based Indicators in Tibetan Plateau Lakes: Implications for Paleotemperature and Paleosalinity Reconstructions, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 601, 111 127, <https://doi.org/10.1016/j.palaeo.2022.111127>, 2022.
- 55 Kusch, S., Winterfeld, M., Mollenhauer, G., H fle, S. T., Schirrmeister, L., Schwamborn, G., and Rethemeyer, J.: Glycerol Dialkyl Glycerol Tetraethers (GDGTs) in High Latitude Siberian Permafrost: Diversity, Environmental Controls, and Implications for Proxy Applications, *Organic Geochemistry*, 136, 103 888, <https://doi.org/10.1016/j.orggeochem.2019.06.009>, 2019.
- Li, J., Naafs, B. D. A., Pancost, R. D., Yang, H., Liu, D., and Xie, S.: Distribution of Branched Tetraether Lipids in Ponds from Inner Mongolia, NE China: Insight into the Source of brGDGTs, *Organic Geochemistry*, 112, 127–136, <https://doi.org/10.1016/j.orggeochem.2017.07.005>, 2017.
- 60 Li, Y., Zhao, S., Pei, H., Qian, S., Zang, J., Dang, X., and Yang, H.: Distribution of Glycerol Dialkyl Glycerol Tetraethers in Surface Soils along an Altitudinal Transect at Cold and Humid Mountain Changbai: Implications for the Reconstruction of Paleoaltimetry and Paleoclimate, *Science China Earth Sciences*, 61, 925–939, <https://doi.org/10.1007/s11430-017-9168-9>, 2018.
- Mart nez-Sosa, P., Tierney, J. E., Stefanescu, I. C., Crampton-Flood, E. D., Shuman, B. N., and Routson, C.: A Global Bayesian Temperature Calibration for Lacustrine brGDGTs, *Geochimica et Cosmochimica Acta*, 305, 87–105, <https://doi.org/10.1016/j.gca.2021.04.038>, 2021.
- 65 Mart nez-Sosa, P., Tierney, J. E., P rez-Angel, L. C., Stefanescu, I. C., Guo, J., Kirkels, F., Sep lveda, J., Peterse, F., Shuman, B. N., and Reyes, A. V.: Development and Application of the Branched and Isoprenoid GDGT Machine Learning Classification Algorithm (BIGMaC) for Paleoenvironmental Reconstruction, *Paleoceanography and Paleoclimatology*, 38, e2023PA004 611, <https://doi.org/10.1029/2023PA004611>, 2023.
- 70 Miller, D. R., Habicht, M. H., Keisling, B. A., Casta eda, I. S., and Bradley, R. S.: A 900-Year New England Temperature Reconstruction from in Situ Seasonally Produced Branched Glycerol Dialkyl Glycerol Tetraethers (brGDGTs), *Climate of the Past*, 14, 1653–1667, <https://doi.org/10.5194/cp-14-1653-2018>, 2018.

- Naafs, B., Gallego-Sala, A., Inglis, G., and Pancost, R.: Refining the Global Branched Glycerol Dialkyl Glycerol Tetraether (brGDGT) Soil Temperature Calibration, *Organic Geochemistry*, 106, 48–56, <https://doi.org/10/gbjssd>, 2017a.
- 75 Naafs, B. D. A., Inglis, G. N., Zheng, Y., Amesbury, M. J., Biester, H., Bindler, R., Blewett, J., Burrows, M. A., Del Castillo Torres, D., and Chambers, F. M.: Introducing Global Peat-Specific Temperature and pH Calibrations Based on brGDGT Bacterial Lipids, *Geochimica et Cosmochimica Acta*, 208, 285–301, <https://doi.org/10/f99n9b>, 2017b.
- Nash, D. J.: *World atlas of desertification*, vol. 165, Blackwell Publishing Ltd., 1999.
- Ning, D., Zhang, E., Shulmeister, J., Chang, J., Sun, W., and Ni, Z.: Holocene Mean Annual Air Temperature (MAAT) Reconstruction
80 Based on Branched Glycerol Dialkyl Glycerol Tetraethers from Lake Ximenglongtan, Southwestern China, *Organic Geochemistry*, <https://doi.org/10.1016/j.orggeochem.2019.05.003>, 2019.
- Pei, H., Wang, C., Wang, Y., Yang, H., and Xie, S.: Distribution of Microbial Lipids at an Acid Mine Drainage Site in China: Insights into Microbial Adaptation to Extremely Low pH Conditions, *Organic Geochemistry*, 134, 77–91, <https://doi.org/10.1016/j.orggeochem.2019.05.008>, 2019.
- 85 Pérez-Angel, L. C., Sepúlveda, J., Molnar, P., Montes, C., Rajagopalan, B., Snell, K., Gonzalez-Arango, C., and Dildar, N.: Soil and Air Temperature Calibrations Using Branched GDGTs for the Tropical Andes of Colombia: Toward a Pan-Tropical Calibration, *Geochemistry, Geophysics, Geosystems*, 21, e2020GC008941, <https://doi.org/10.1029/2020GC008941>, 2020.
- Peterse, F., van der Meer, J., Schouten, S., Weijers, J. W., Fierer, N., Jackson, R. B., Kim, J.-H., and Damsté, J. S. S.: Revised Calibration of the MBT–CBT Paleotemperature Proxy Based on Branched Tetraether Membrane Lipids in Surface Soils, *Geochimica et Cosmochimica*
90 *Acta*, 96, 215–229, <https://doi.org/10/f4db3v>, 2012.
- Qian, S., Yang, H., Dong, C., Wang, Y., Wu, J., Pei, H., Dang, X., Lu, J., Zhao, S., and Xie, S.: Rapid Response of Fossil Tetraether Lipids in Lake Sediments to Seasonal Environmental Variables in a Shallow Lake in Central China: Implications for the Use of Tetraether-Based Proxies, *Organic Geochemistry*, 128, 108–121, <https://doi.org/10.1016/j.orggeochem.2018.12.007>, 2019.
- Raberg, J. H., Harning, D. J., Crump, S. E., de Wet, G., Blumm, A., Kopf, S., Geirsdóttir, Á., Miller, G. H., and Sepúlveda, J.: Revised
95 Fractional Abundances and Warm-Season Temperatures Substantially Improve brGDGT Calibrations in Lake Sediments, *Biogeosciences*, 18, 3579–3603, <https://doi.org/10.5194/bg-18-3579-2021>, 2021.
- Raberg, J. H., Flores, E., Crump, S. E., de Wet, G., Dildar, N., Miller, G. H., Geirsdóttir, Á., and Sepúlveda, J.: Intact Polar brGDGTs in Arctic Lake Catchments: Implications for Lipid Sources and Paleoclimate Applications, *Journal of Geophysical Research: Biogeosciences*, 127, e2022JG006969, <https://doi.org/10.1029/2022JG006969>, 2022.
- 100 Russell, J. M., Hopmans, E. C., Loomis, S. E., Liang, J., and Damsté, J. S. S.: Distributions of 5- and 6-Methyl Branched Glycerol Dialkyl Glycerol Tetraethers (brGDGTs) in East African Lake Sediment: Effects of Temperature, pH, and New Lacustrine Paleotemperature Calibrations, *Organic Geochemistry*, 117, 56–69, <https://doi.org/10.1016/j.orggeochem.2017.12.003>, 2018.
- Rusydi, A. F.: Correlation between Conductivity and Total Dissolved Solid in Various Type of Water: A Review, *IOP Conference Series: Earth and Environmental Science*, 118, 012019, <https://doi.org/10.1088/1755-1315/118/1/012019>, 2018.
- 105 Sun, Q., Chu, G., Liu, M., Xie, M., Li, S., Ling, Y., Wang, X., Shi, L., Jia, G., and Lü, H.: Distributions and Temperature Dependence of Branched Glycerol Dialkyl Glycerol Tetraethers in Recent Lacustrine Sediments from China and Nepal, *Journal of Geophysical Research: Biogeosciences*, 116, 2011.
- Thomas, E. K., Clemens, S. C., Sun, Y., Huang, Y., Prell, W., Chen, G., Liu, Z., and Loomis, S.: Midlatitude Land Surface Temperature Impacts the Timing and Structure of Glacial Maxima, *Geophysical Research Letters*, 44, 984–992, <https://doi.org/10.1002/2016gl071882>,
110 2017.
- Véquaud, P., Derenne, S., Anquetil, C., Collin, S., Poulenard, J., Sabatier, P., and Huguet, A.: Influence of Environmental Parameters on the Distribution of Bacterial Lipids in Soils from the French Alps: Implications for Paleo-Reconstructions, *Organic Geochemistry*, p. 104194, <https://doi.org/10.1016/j.orggeochem.2021.104194>, 2021.
- Wang, H. and Liu, W.: Soil Temperature and brGDGTs along an Elevation Gradient on the Northeastern Tibetan Plateau: A Test of Soil
115 brGDGTs as a Proxy for Paleoelevation, *Chemical Geology*, 566, 120079, <https://doi.org/10.1016/j.chemgeo.2021.120079>, 2021.
- Wang, H., Liu, W., and Lu, H.: Appraisal of Branched Glycerol Dialkyl Glycerol Tetraether-Based Indices for North China, *Organic Geochemistry*, 98, 118–130, <https://doi.org/10.1016/j.orggeochem.2016.05.013>, 2016.
- Wang, H., An, Z., Lu, H., Zhao, Z., and Liu, W.: Calibrating Bacterial Tetraether Distributions towards in Situ Soil Temperature and Application to a Loess-Paleosol Sequence, *Quaternary Science Reviews*, 231, 106172, <https://doi.org/10.1016/j.quascirev.2020.106172>, 2020.
- 120 Wang, H., Liu, W., He, Y., Zhou, A., Zhao, H., Liu, H., Cao, Y., Hu, J., Meng, B., Jiang, J., Kolpakova, M., Krivonogov, S., and Liu, Z.: Salinity-Controlled Isomerization of Lacustrine brGDGTs Impacts the Associated MBT5ME' Terrestrial Temperature Index, *Geochimica et Cosmochimica Acta*, 305, 33–48, <https://doi.org/10.1016/j.gca.2021.05.004>, 2021.
- Wang, M., Zong, Y., Zheng, Z., Man, M., Hu, J., and Tian, L.: Utility of brGDGTs as Temperature and Precipitation Proxies in Subtropical China, *Scientific Reports*, 8, 1–9, <https://doi.org/10.1038/s41598-017-17964-0>, 2018.

- 125 Wang, M., Zheng, Z., Zong, Y., Man, M., and Tian, L.: Distributions of Soil Branched Glycerol Dialkyl Glycerol Tetraethers from Different Climate Regions of China, *Scientific reports*, 9, 1–8, <https://doi.org/10.1038/s41598-019-39147-9>, 2019.
- Weber, Y., Damsté, J. S. S., Zopfi, J., De Jonge, C., Gilli, A., Schubert, C. J., Lepori, F., Lehmann, M. F., and Niemann, H.: Redox-Dependent Niche Differentiation Provides Evidence for Multiple Bacterial Sources of Glycerol Tetraether Lipids in Lakes, *Proceedings of the National Academy of Sciences*, 115, 10 926–10 931, <https://doi.org/10.1073/pnas.1805186115>, 2018.
- 130 Yang, H., Pancost, R. D., Dang, X., Zhou, X., Evershed, R. P., Xiao, G., Tang, C., Gao, L., Guo, Z., and Xie, S.: Correlations between Microbial Tetraether Lipids and Environmental Variables in Chinese Soils: Optimizing the Paleo-Reconstructions in Semi-Arid and Arid Regions, *Geochimica et Cosmochimica Acta*, 126, 49–69, <https://doi.org/10/f5n3gh>, 2014.
- Zang, J., Lei, Y., and Yang, H.: Distribution of Glycerol Ethers in Turpan Soils: Implications for Use of GDGT-based Proxies in Hot and Dry Regions, *Frontiers of Earth Science*, 12, 862–876, <https://doi.org/10/gfb6xq>, 2018.