



A high-resolution nested model to study the effects of alkalinity additions in Halifax Harbour, a mid-latitude coastal fjord

Arnaud Laurent¹, Bin Wang¹, Dariia Atamanchuk¹, Subhadeep Rakshit^{1,a}, Kumiko Azetsu-Scott², Chris Algar¹, and Katja Fennel¹

¹Department of Oceanography, Dalhousie University, Halifax, Nova Scotia, Canada

²Bedford Institute of Oceanography, Dartmouth, Nova Scotia, Canada

^apresent address: Program in Atmospheric and Oceanic Sciences, Princeton University, Princeton, NJ, USA

Correspondence: Arnaud Laurent (arnaud.laurent@dal.ca)

Received: 15 July 2025 – Discussion started: 22 July 2025

Revised: 26 November 2025 – Accepted: 3 December 2025 – Published: 7 January 2026

Abstract. Surface ocean alkalinity enhancement (OAE), through the release of alkaline materials, is an emerging marine carbon dioxide removal technology that could increase the storage of anthropogenic carbon in the ocean. Observations collected during recent and on-going field trials will provide important information on the feasibility and effects of alkalinity additions on carbon cycling and study ecological responses. However, given the scales involved (24/7 continuous addition, meters to hundreds/thousands of kilometers and minutes to months for alkalinity dispersion) observations, even with the use of autonomous platforms, will remain inherently sparse and limited. Alone, they cannot provide a comprehensive quantification of the effects of OAE on the carbonate system, and ultimately of the net air-sea CO₂ fluxes. Numerical models, informed and validated by field observations, are therefore essential to OAE deployments and the measurement, reporting, and verification (MRV) of any resulting carbon uptake. They can help guide fieldwork design, including optimal design of measurement monitoring networks, provide forecasts of the ocean state, simulate the effects of alkalinity additions on the seawater carbonate system, and allow one to quantify net CO₂ uptake. Here we describe a coupled physical-biogeochemical model that is specifically designed for coastal OAE. The model is an implementation of the Regional Ocean Modelling System (ROMS) in a nested grid configuration with increasing spatial resolution from the Scotian Shelf to Halifax Harbour (coastal fjord, eastern Canada), a current test site for operational alkalinity addition. The biogeochemical model simulates oxygen dynamics, carbonate system processes (includ-

ing air-sea gas exchange), and feedstock properties (dissolution, sinking). We present a multi-year hindcast validated against the long-term weekly time series available for a long-term monitoring station at the deepest part of Halifax Harbour, as well as alkalinity addition simulations at various locations inside and outside the harbour to show the model's capabilities for assessing the effects of OAE at this coastal site. We found that release locations had a strong effect on the outcome of the addition, whereas feedstock types influenced the distribution of the signal. Up to 69 % of the net CO₂ uptake was realized within the model domain, most of it inside the harbour. These results stress the importance of operational design, as well as the use of high-resolution regional models when quantifying additionality.

1 Introduction

Global temperature has increased by more than 1.1 °C from pre-industrial average and is likely to exceed, at least temporarily, the +1.5 °C goal of the Paris agreement in the coming decade (WMO, 2024a, b). At the current rate of greenhouse gas emissions, warming is expected to reach +2.7 °C by the end of the century (UNEP, 2023). Without rapid and significant mitigation actions, changes in the climate system (increasingly irreversible) will severely affect ecosystems, biodiversity, as well as the livelihood of current and future generations (IPCC, 2023). As the urgency increases, carbon dioxide removal (CDR) techniques have been proposed as a supplement to emission reductions. CDR can accelerate

progress toward net zero CO₂ emission by compensating for hard-to-abate residual emissions (e.g. from agriculture, aviation, shipping or industrial sectors) and remove legacy anthropogenic CO₂ from the atmosphere (IPCC, 2023). It is estimated that 10 Gt of CO₂ residual emissions need to be compensated annually by 2050 and 20 Gt by 2100 to achieve net zero CO₂ emissions by the end of the century (NASEM, 2019).

Current CDR measures involve land-based techniques such as afforestation, reforestation or carbon capture and storage (Nabuurs et al., 2023; Riahi et al., 2023). The risks, such as competition for land, non-permanence, energy cost, depend on the method and the scale involved (Fuhrman et al., 2020). Given the large reservoir of carbon in the ocean and its potential for additional storage, marine CDR methods (mCDR) have gained interest in the last decade (Gattuso et al., 2021; Oschlies et al., 2025). Seaweed cultivation, nutrient fertilization (e.g., iron) or artificial upwelling have been proposed to increase the ocean's carbon storage (NASEM, 2022). However, the risks associated to these biotic mCDR technologies may be important because they modify the biological carbon pump and nutrient cycles, with potential important side effects on marine ecosystems (Cullen and Boyd, 2008; Fennel, 2008; Oschlies et al., 2010; Wu et al., 2023). The impact on the deep ocean is of particular concern (Levin et al., 2023). The effectiveness of biotic mCDR technologies is therefore uncertain (Oschlies et al., 2025). Ocean alkalinity enhancement (OAE) is an abiotic method that aims to store additional inorganic carbon in the ocean via the addition of alkalinity at the surface. In theory, this technique has fewer risks to marine ecosystems because it does not modify biological cycles, although the side effects depend on the source material and technology used and possibly on the addition level (Delacroix et al., 2024; Ferderer et al., 2024; Guo et al., 2024). If scaled up to the global ocean, OAE could remove > 1 Gt CO₂ yr⁻¹ from the atmosphere (Renforth and Henderson, 2017).

The challenge for all mCDR methods is to detect and monitor their effects on the carbon pool, and to quantify the carbon removed from the atmosphere (Fennel, 2026). For OAE the effects of the added alkalinity on the solubility pump and the environment need to be quantified from tens of meters near the release site to possibly thousands of kilometers until addition waters are equilibrated with the atmosphere. Moreover, this signal needs to be detected on top of the natural variability in carbonate system parameters, which is challenging especially in coastal settings (Carter et al., 2019). Because the perturbations due to OAE are overlayed on top of natural variability of carbonate system parameters, which is controlled by a combination of biotic and abiotic factors, observations alone are insufficient in characterizing the effects of OAE (Bach et al., 2023).

To quantify how much carbon is removed, an essential aspect of measurement, reporting, and verification (MRV), the addition needs to be monitored and compared with a counterfactual case without addition. By definition, it is not possible to measure the counterfactual (Fennel, 2026). Therefore, fit-for-purpose validated models are essential for quantifying any net carbon uptake resulting from alkalinity addition (Fennel et al., 2023; Ho et al., 2023).

Global ocean models represent the large-scale circulation and therefore have been used to assess the feasibility of OAE and track its effects at the global scale (Burt et al., 2021; Feng et al., 2017; Ilyina et al., 2013; Köhler et al., 2013; Lenton et al., 2018; Palmiéri and Yool, 2024; Tyka, 2025). The limitation of these models is their coarse resolution which does not allow them to represent mesoscale processes and poorly resolves dynamics on coastal and shelf scales (Laurent et al., 2021). For practical reasons, e.g., pre-existing infrastructure such as sewage and cooling outfalls, and to reduce the carbon footprint of alkalinity delivery, nearshore areas will likely be major sites of alkalinity addition in OAE projects (Eisaman et al., 2023). High resolution regional models are better suited to simulate alkalinity addition at these scales (10 m–100 km) than global models. The addition signal can then be passed on to a global model for basin scale tracking of unequilibrated alkalinity or to an earth system model to assess the long-term, global re-equilibration of the mobile carbon pool (Fennel, 2026).

Wang et al. (2025) recently developed a regional model to study OAE in the Halifax Harbour on the eastern coast of Canada. Halifax Harbour has been in operation as an OAE test site since 2023 (Cornwall, 2023). Wang et al.'s (2025) study highlighted the importance of local processes, such as local circulation and residence time, and feedstock characteristics such as its dissolution and sinking rates on the fate of the added material. The likelihood of ecosystem exposure to alkaline waters was also simulated and quantified by Wang et al. (2025). It is clear from Wang et al.'s (2025) results that added alkalinity doesn't simply disperse onto the shelf after addition, as assumed in global models, but is retained for some time inshore, depending on the dissolution and sinking rates of the alkaline material used and seasonal circulation features (i.e., residence time). The model version in Wang et al. (2025) did not include an explicit representation of the carbonate system and therefore was not used to assess addition effects on carbonate system parameters nor to quantify the resulting local net CO₂ uptake. Other regional modelling studies including explicit carbonate system parameters (e.g., Khangaonkar et al., 2024; Li et al., 2025; Wang et al., 2023) have been carried out but not for this location. OAE studies usually use generic, previously validated biogeochemical models that include carbon chemistry but that were not explicitly tailored to OAE research. Most models assume the addition of fully dissolved rather than particulate material (e.g., Kwiatkowski et al., 2023; Nagwekar et al., 2024; Zhou et al., 2025), whereas mineral feedstock particles will dissolve over time (Schulz et al., 2023). Palmiéri and Yool (2024) assumed dissolution but of an inert particulate feedstock at the sediment-water interface. Feedstock characteristics have

a significant effect on the location and time scale of surface expression of the addition signal (Wang et al., 2025). Recent attempts have been made, at the global scale, to adapt biogeochemical models to study OAE. Zhou et al. (2025) used two carbonate systems to simulate the addition and counterfactual cases, but without dissolution kinetics. Fakhraee et al. (2023) developed a biogeochemical-mineral dissolution model but within a coarse physical framework.

To fill the gaps described above, we present a high-resolution, coupled physical-biogeochemical-addition-dissolution model that is designed to support OAE research in coastal environments. The model builds on Wang et al. (2025) to include oxygen and carbonate system chemistry in a realistic alkalinity addition setting. The model is validated and tested for Halifax Harbour. The key question that is investigated with this novel model is how feedstock type and dosing location influence alkalinity dispersion and net CO_2 air-sea flux in the Halifax Harbour and surrounding nearshore waters.

The observations used for the study are described in Sect. 2. The circulation model is presented in Sect. 3 and the results of an eight-year simulation (2016–2023) validated against observations on the shelf and in the harbour are presented. The biogeochemical model is described in Sect. 4 and validated for 2016–2023. The addition model is then presented and a series of OAE simulations carried out in Sect. 5. The overall results are discussed in Sect. 6.

2 Supporting observations

2.1 Station 2 on the Scotian Shelf

Station 2, located on the Scotian Shelf in 150 m water depth ($44^{\circ}16'01''\text{N}$, $63^{\circ}19'01''\text{W}$), is sampled biweekly or monthly by the Atlantic Zone Monitoring Program (AZMP). The observations include CTD casts (temperature, salinity, oxygen) (MEDS, 2024) and Niskin bottle samples at various depths for dissolved inorganic carbon (DIC) and alkalinity (Cyr et al., 2022).

2.2 Compass Buoy station in Halifax Harbour

The Bedford Basin Monitoring Program (BBMP) provides weekly observations, since 1992, at the Compass Buoy station in Halifax Harbour's Bedford Basin (the deep basin at the head of the harbour). This station is at the deepest location of the harbour, in the center of Bedford Basin (70 m depth, $44^{\circ}41'37''\text{N}$, $63^{\circ}38'25''\text{W}$) (Fisheries and Oceans Canada, 2025). The observations include CTD casts (temperature, salinity, oxygen) and Niskin bottle samples (for various biological and chemical variables) at 4 depths (0, 5, 10 and 60 m). Carbonate system measurements (DIC, alkalinity) have been collected since 2016.

pCO_2 was calculated from TA and DIC with CO2SYS (pyCO2SYS, Humphreys et al., 2022) using the “Mehrbach

refit” dissociation constants for carbonic acid (Dickson and Millero, 1987; Mehrbach et al., 1973) and the “Dickson” dissociation constants for bisulfate ions (Dickson, 1990). Bottle samples reported in units of $\mu\text{mol kg}^{-1}$ were converted to mmol m^{-3} with a conversion factor of 1.025 kg m^{-3} (assuming a density of 1025 kg m^{-3} and converting μmol to mmol).

2.3 Benthic fluxes and respiration

Sediment-water fluxes, benthic and water column respiration were measured at several locations within the harbour using a benthic chamber and sediment core profiles (Rakshit et al., 2023). These data were used to parameterize the biogeochemical model (see Sect. 4.1).

3 Circulation model and shelf biogeochemistry

The model is a regional implementation of the Regional Ocean Modelling System (ROMS, version 3.9) for the Scotian Shelf and the Halifax Harbour (Fig. 1). ROMS is a free-surface, primitive equation ocean model with terrain-following vertical coordinates (Haidvogel et al., 2008). The model has 40 vertical layers with increasing resolution near the surface and bottom; the horizontal resolution is different for each nested grid (see below). Advection is prescribed with a third-order upwind (horizontal) and a fourth order centered (vertical) scheme for momentum equations and with the High-order Spatial Interpolation at the Middle Temporal level scheme (HSIMT; Wu and Zhu, 2010) for biological tracers. Vertical mixing is parameterized using the nonlocal K-profile parameterization (Durski et al., 2004; Large and Gent, 1999).

3.1 Nested model grids

The model is setup with 3 nested grids of increasing resolution focussing in on Halifax Harbour (Fig. 1c, d). The coarsest grid, ROMS-H1 (Fig. 1a, c), covers the central Scotian Shelf with an average horizontal resolution of 760 m and a bathymetry interpolated from GEBCO (<https://www.gebco.net>, last access: 17 June 2022). The grid is run with a biogeochemical model that includes the nitrogen cycle (Fennel et al., 2006; Laurent et al., 2021), phosphorus cycling (Laurent et al., 2012), oxygen dynamics (Fennel et al., 2013), and carbonate chemistry (Fennel, 2010; Laurent et al., 2017). This model is referred subsequently as the explicit biogeochemical model. It is forced with tides and reanalysis data at the open boundaries (see Sect. 3.2 below). A similar implementation of the model was used for larger-scale implementations of the northwest North Atlantic shelf and adjacent open ocean and shown to represent physical and biogeochemical dynamics well (Ohashi et al., 2024; Rutherford et al., 2021). The ROMS-H2 domain is nested within ROMS-H1 with a refinement factor of 5 that covers the inner Scotian Shelf and the Halifax Harbour (Fig. 1c). It has a horizontal resolution of




Figure 1. Study area (**A, B**) and nested model grids (**C, D**). (**A**) Bathymetry on the Northwest North Atlantic (NWA; <https://www.gebco.net>, last access: 17 June 2022, color scale as in **C**) including the southern Labrador Sea (LS), the Gulf of Saint Lawrence (GSL) and the Gulf of Maine (GoM). ROMS-H1 is delimited by the black rectangle. (**B**) Satellite view of the Halifax Harbour (<http://www.earth.google.com>, last access: 17 September 2024). (**C**) Bathymetry and outer limits of the nested grids. (**D**) Bathymetry and outer limits of ROMS-H3 plotted on top of ROMS-H2. The red star indicates the location of the compass buoy, the red dots the release locations at Mill Cove (MC), Tufts Cove (TC) and Herring Cove (HC) and the blue dots other WWTPs. The thin grey lines indicate the limits of the 5 areas used for spatial integration, namely Bedford Basin (BB), Inner Harbour (IH), Outer Harbour (OH), Northwest Arm (NA) and Shelf. Satellite image in panel (**B**) was generated with Google Earth. Image credits: Landsat/Copernicus, © 2025 Maxar Technologies, © 2025 CNES/Airbus.

150 m and is forced at the open boundaries by interpolated results from ROMS-H1. The bathymetry is generated from the NONNA-10 and NONNA-100 products of the Canadian Hydrographic Services (<https://open.canada.ca/data/en/dataset/d3881c4c-650d-4070-bf9b-1e00aabf0a1d>, last access: 16 September 2022), supplemented by GEBCO where high-resolution data from the NONNA products are unavailable. ROMS-H3 is the highest-resolution domain, nested within the ROMS-H2 grid with a refinement factor of 3 and covers Halifax Harbour (Fig. 1b–d). The grid has a horizontal resolution of 50 m, and its bathymetry is generated from NONNA-10. ROMS-H2 and ROMS-H3 are coupled with the reduced biogeochemical model (see Sect. 4). ROMS-H3 can be run with two-way nesting or with offline one-way nesting. For the two-way nesting case (called ROMS-H23), ROMS-H2 and ROMS-H3 are run simultaneously and exchange information at every time step. For the offline one-way nesting case, ROMS-H3 open boundary conditions are calculated offline using results from ROMS-H2. Hereafter, most of the model validation and analysis use ROMS-H2. Simulations with ROMS-H3 are presented in Sect. 5.2.4. Results of

ROMS-H23 are not presented here, but are available in Wang et al. (2025).

3.2 Forcing

The 3 ROMS domains share the same atmospheric forcing, prescribed hourly from the ERA5 reanalysis (Hersbach et al., 2023). Initial and open boundary conditions in ROMS-H1 are set by the GLORYS12V1 reanalysis product (<https://doi.org/10.48670/moi-00021>; Lellouche et al., 2021) for the physical variables and with a daily climatology calculated from the multiyear simulation of Rutherford et al. (2021) for the biogeochemical variables. ROMS-H1 is also forced by tides using 9 tidal constituents extracted from the ADCIRC Tidal Database version EC2001_v2e (<https://adcirc.org/products/adcirc-tidal-databases/>, last access: 2 May 2024; Mukai et al., 2002) and has a surface heat flux correction towards the Multi-scale Ultra-high Resolution (MUR) sea surface temperature (SST) daily product (<https://doi.org/10.5067/GHGMR-4FJ04>; Chin et al., 2017) with a relaxation time scale of 10 d.

River forcing is added using daily discharge observations from the National Water Data Archive (HYDAT) database

(<https://wateroffice.ec.gc.ca>, last access: 23 October 2024) for the LaHave River (01EF001; ROMS-H1) and Sackville River (01EJ001; all grids). The Halifax Harbour includes additional freshwater sources from five wastewater treatment plants (WWTP, Fig. 1d) and 11 other minor inflows. Freshwater discharge at Mill Cove WWTP uses a daily climatology calculated with a multiannual daily discharge time series from Halifax Water. Other WWTP are scaled from Mill Cove using annual discharge information available from Halifax Water, following Wang et al. (2025). Minor inflows are scaled to the Sackville River using average discharge information from Buckley and Winters (1992), summarized in Table 1 of Shan (2010). River nutrients and carbonate data are set constant based on observations collected in the Sackville River (Sackville River + minor inflows) and at Mill Cove (McGinn et al., 2012; WWTPs).

Atmospheric pCO_2 is parameterized using a fit to the monitoring observations from Sable Island collected by the Environment Canada Greenhouse Gas Measurement Program and available in the GLOBALVIEW database (Schuldt et al., 2023):

$$\text{pCO}_{2,\text{air}} = a_0 + a_1 t + a_2 t^2 + 1.25 a_3 \times \log(\sin(2\pi t + a_4) + a_5), \quad (1)$$

where t is the fractional year and the coefficients are $a_0 = 72134.3$, $a_1 = -73.6115$, $a_2 = 0.01886$, $a_3 = 5.47721$, $a_4 = 0.64527$ and $a_5 = 1.12893$ ($r^2 = 0.95$, $p < 0.001$). Due to the lack of local observations, this estimate does not include an urban effect (local increase in atmospheric pCO_2) around the Halifax Harbour.

3.3 Validation of model physics

A multiyear simulation was initialized from BBMP data for Halifax Harbour and GLORYS and climatological data for the shelf (1 January 2015) and spun up for 1 year. The period 2016–2023 was used for comparison with surface and in-situ observations on the shelf for ROMS-H1 and at the compass buoy for ROMS-H2 and ROMS-H3 (Sect. 5.2.4). Simulated surface temperature in ROMS-H1 agrees well with observed seasonal patterns over the shelf ($r^2 > 0.78$) and at the 3 open boundaries of ROMS-H2 ($r^2 = 0.99$) (Fig. S3, Table 2). The only discrepancy not captured by the model is the large high-frequency variability in fall/winter surface temperature. Surface and in-situ temperature (Fig. S4) and salinity (Fig. S5) are also in good agreement with the CTD profiles from Station 2 on the shelf ($r^2 = 0.77$ and 0.74, respectively, Table 2). At the Compass Buoy station in Bedford Basin, surface temperature (Fig. S11) and salinity (Fig. S12) simulated with ROMS-H2 are close to observations, with some overestimation in spring and summer for temperature and a slight underestimation for salinity. Vertical profiles agree well with observed temperature, although the agreement is not as good for salinity ($r^2 = 0.78$ and 0.46, respectively, Table 2). At

the surface, temperature is somewhat overestimated in spring and summer, while salinity is slightly underestimated. The profile comparisons indicate that the biases at the surface may be driven by the too shallow pycnocline. ROMS-H2 also tends to overmix in the fall (Fig. S11).

Temperature and salinity profiles at Compass Buoy site indicate that the model is able to reproduce the timing of most of the discontinuities in water column properties, driven by convection in winter and intrusions of shelf waters into the Bedford Basin in summer and fall (Garcia Larez, 2023). Sharp and concomitant increases in deep water temperature and salinity at this site are a typical signal of intrusions events (Figs. S11 and S12). Despite the agreement in timing, the magnitude of some events is overestimated by the model. The sharp surface pycnocline (Fig. S11), which likely enhances the estuarine circulation, may play a role in this mismatch.

Overall, the circulation model simulates the physical dynamics of the Halifax Harbour – Scotian Shelf system well (Table 2). Thus, we deem the model as an appropriate framework for simulating the dynamics of oxygen and carbonate system parameters in this area.

3.4 Shelf biogeochemistry

ROMS-H1 used the explicit biogeochemical model described in Sect. 3.1 but only oxygen and carbon chemistry were activated in the higher-resolution ROMS-H2 so we focus on these variables hereafter. Simulated oxygen compares well with the seasonal patterns observed at Station 2 ($r^2 = 0.74$, Table 2, Fig. S6). The main difference is that the model overestimates deep oxygen concentrations (below ~ 100 m) and underestimates, at times, surface oxygen concentrations during the spring bloom. Despite these mismatches, oxygen variability is well represented in the model.

In Halifax Harbour, observations of alkalinity are strongly correlated with salinity (Atamanchuk et al., 2024; Fig. S10). Since the initial and boundary conditions for alkalinity and salinity in ROMS-H1 have different sources (climatology and GLORYS, respectively), the simulated alkalinity (TA) versus salinity (S) relationship does not match the relationship observed in Halifax Harbour. Therefore, to calculate initial and boundary conditions for alkalinity in ROMS-H2 the following observed relationship from the Halifax Harbour was used:

$$\text{TA}_{\text{fit}} = 63.417 \times S + 235.135 \text{ mmol m}^{-3} \quad (2)$$

To avoid an artificial disequilibrium in pCO_2 (due to modified TA but not DIC), initial and boundary conditions for DIC were also recalculated with CO2SYS using simulated salinity, simulated pCO_2 and TA_{fit} . These calculated TA and DIC values are in good agreement with observations at Station 2 (Figs. S7 and S8, $r^2 = 0.84$ and 0.85, respectively, Table 2). The seasonal variability of surface DIC is also well captured (Fig. S8). Similar to oxygen, deep TA (below ~ 100 m), and somewhat DIC, were generally overestimated.

Table 1. Summary description of simulations. H1 indicates a simulation carried out with ROMS-H1, H2 with ROMS-H2 and H3 with ROMS-H3.

Dosing location	Feedstock type		Time range	BGC		Purpose
	–	Particulate		Dissolve	Full	
No dosing	H1	–	–	Start: 2016-01-01	Full	BC for H2, validation
	H2	–	–	End: 2023-12-31	Reduced	Validation
Mill Cove (MC)	–	H2, H3	H2, H3	Start: 2017-07-01	Reduced + Addition	1. Test addition model
Tufts Cove (TC)	–	H2, H3	H2, H3	End: 2018-12-31		2. Effects of dosing type + location
Herring Cove (HC)	–	H2, H3	H2, H3			3. BC for H3 (H2 only)

Table 2. Model-observations comparison statistics for Figs. S3–15.

Grid	Location	Variable	Type	R^2	bias	RMSE
H1	Shelf (maps)	SST (seasonal)	DJF	0.981	-0.15	0.23
			JJA	0.781	-0.09	0.23
			MAM	0.940	-0.12	0.21
			SON	0.944	-0.10	0.14
	Shelf (series)	SST (H2 boundaries)	South	0.993	0.01	0.53
			West	0.989	-0.01	0.66
			East	0.991	0.06	0.62
	Profiles Station 2	Temperature	CTD	0.773	0.75	1.89
		Salinity		0.741	0.28	0.76
		Oxygen		0.738	0.19	30.3
	H2 BBMP	DIC*	Water samples	0.853	11.0	40.3
		TA*		0.842	37.9	56.2
		Temperature	CTD	0.777	0.45	1.66
		Salinity	Profiles	0.457	-0.23	0.64
		Oxygen		0.554	-19.2	65.2
		DIC	Water samples	0.585	24.3	109
		TA		0.520	15.4	63.3
		pCO ₂		0.408	38.6	562

Seasonal averages: DJF: December–February, JJA: June–August, MAM: March–May, SON: September–November. Units: Temperature: °C, Oxygen, DIC, TA: mmol m⁻³, pCO₂: µatm.

* Corrected values based on the TA versus salinity relationship (see Sect. 3.4).

4 Reduced complexity biogeochemical model

The model is based on the explicit biogeochemical model used in ROMS-H1 (Fennel, 2010; Fennel et al., 2006, 2013; Laurent et al., 2012, 2017, 2021) but simplified to improve computational efficiency for studying the carbonate system in the context of alkalinity enhancement. This fit-for-purpose modelling approach delivers what we need in terms of describing the seasonal cycle of background DIC without the computational cost of additional biogeochemical tracers. Since oxygen is measured routinely, in contrast to carbonate variables, and has sources and sinks similar to DIC (with opposite sign) the model includes oxygen as a diagnostic variable. Reduced complexity biogeochemical models have pre-

viously proven useful in studies of hypoxia in Chesapeake Bay (Scully, 2013) and in the northern Gulf of Mexico (Yu et al., 2015).

4.1 Model description

The model state variables are TA, DIC and oxygen (O₂; Fig. 2). The general model equations are available in the supplement information. The main differences to the original biogeochemical model used in ROMS-H1 are the removal of state variables for nutrients (NO₃, PO₄, NH₄) and particulate organic matter (phytoplankton, zooplankton, small and large detritus) and their replacement with parameterizations




Figure 2. Schematic of the biogeochemical model. Dark grey boxes represent the main variables of the biogeochemical model (control). Pink boxes and arrows indicate the addition model variables and processes, respectively. Dashed boxes indicate calculated variables (rather than state variables).

of primary production and respiration in water column and sediment.

The ROMS-H2 domain includes the inner Scotian Shelf, where biological conditions are typical of those on the northwest North Atlantic shelves with biannual blooms, and the inner Halifax Harbour (including the Bedford Basin) where conditions are highly eutrophic due to the input of anthropogenic nutrients from wastewater treatment plants. Primary production and respiration are therefore parameterized differently in the harbour and on the shelf. The harbour parameterizations are also used in ROMS-H3.

4.1.1 Primary production

A multiyear simulation (2017–2022) was run for ROMS-H2 using the same explicit biogeochemical model (ROMS-H2-BGC) as in ROMS-H1. The results were used to calculate time varying, but spatially averaged climatologies of surface primary production for the Scotian Shelf and Halifax Harbour (Fig. S1). We then parameterized primary production on the Scotian Shelf (PP_S) and in Halifax Harbour (PP_H) with a fit to climatology such that it is a function of time:

$$PP_x(t) = a_1 + b_1 \times (\cos(b_2 t - 1) + 1)^{b_3} + (\sin(c_1 t - 1) + 1)^{c_2} + (\sin(d_1 t - 1) + 1)^{d_2} + (\sin(e_1 t - 1) + 1)^{e_2} \quad (3)$$

where x represents the Halifax Harbour (PP_H) or the shelf (PP_S), t is the fractional year, and PP has units of mmol

$\text{O}_2 \text{ m}^{-3} \text{ d}^{-1}$. These parameterizations are applied over the upper 30 m on the shelf and 15 m in Halifax Harbour to account for the higher light attenuation in the harbour (Craig et al., 2012). The coefficients for each location are listed in Supplement Table S2. The parameterizations are compared to climatology in Supplement Fig. S1.

4.1.2 Water column respiration

The spatially and temporally averaged water column respiration in ROMS-H2-BGC for the shelf (WR_S ; $\text{mmol m}^{-3} \text{ d}^{-1}$) and the inner harbour (WR_H ; $\text{mmol m}^{-3} \text{ d}^{-1}$) areas were approximated with a temperature dependency to parameterize water column respiration, such that:

$$WR_x = a_x \times \theta_T \quad (4)$$

where x represents the Halifax Harbour (WR_H ; $a_H = 1.39403$) or the shelf (WR_S ; $a_S = 1.31349$), and the temperature (T ; $^{\circ}\text{C}$) dependency $\theta_T = 0.59 \times 1.066^T$, as in the explicit biogeochemical model (Laurent et al., 2021). We assume a respiratory quotient of 1 such that water column respiration has the same magnitude (with opposite sign) for O_2 and DIC .

4.1.3 Sediment respiration

On the shelf, sediment respiration (SR_S ; $\text{mmol m}^{-2} \text{ d}^{-1}$) is parameterized as a temporally and spatially varying approximation of ROMS-H2-BGC results, such that:

$$SR_S = b_0 + c_0 \text{lon} + d_0 \text{lat} + \sum_{i=1}^5 (e_0^i x_i + e_1^i x_i^2 + e_2^i x_i^3) \quad (5)$$

where lon is longitude, lat is latitude, and x_i represents bathymetry ($i = 1$), surface and bottom temperature ($i = 2, 3$), and surface and bottom salinity ($i = 4, 5$). The coefficients for each variable are listed in Supplement Table S2. As for water column respiration, we assume a respiratory quotient of 1. For the inner harbour, sediment flux measurements at 6 stations (Bedford Basin and Narrows) in fall 2022 and spring 2023 show a relationship between dissolved inorganic carbon flux (F_H^{DIC} ; $\text{mmol C m}^{-2} \text{ d}^{-1}$) and depth (z ; m) such that sediment respiration (SR_H ; $\text{mmol m}^{-2} \text{ d}^{-1}$) is:

$$SR_H = F_H^{\text{DIC}} = 3.37 \times e^{0.0033 \cdot z} \quad (6)$$

Measurements in the Bedford Basin showed a decrease in oxygen flux at depth ($F_H^{O_2}$; $\text{mmol O}_2 \text{ m}^{-2} \text{ d}^{-1}$) under hypoxic conditions and therefore an oxygen (O_2 ; mmol m^{-3}) dependency was added to the sediment flux such that:

$$F_H^{O_2} = SR_H \times (1 - e^{-O_2/30}) \quad (7)$$

4.2 Biogeochemical model validation

ROMS-H2 is able to simulate the seasonal variations of oxygen at the Compass Buoy station, including the seasonal sub-

surface depletion driven by respiration and occasional renewal of deep water oxygen during summer and fall intrusions ($r^2 = 0.55$, Table 2, Fig. S13). Surface oxygen (upper 1 m) is generally underestimated, particularly during the annual spring bloom and in summer. As mentioned in Sect. 3.3, a few of the simulated intrusions were not visible in the observations (e.g. in fall 2019), generating some mismatches between simulated and observed subsurface oxygen. Another source of discrepancy was the overestimation of vertical mixing in fall that increased subsurface and bottom oxygen. This overestimation is reduced in ROMS-H3 (see Sect. 5.2.4).

At the Compass Buoy station, carbonate system observations are available at the surface (~ 1 – 3 m) and 60 m depth for comparison with the model (Table 2, Figs. S14 and S15). Surface alkalinity and DIC are in good agreement with observations. Surface variability, largely associated with river discharge, is particularly well simulated. At 60 m, the dynamics differ for simulated DIC and alkalinity. DIC accumulates in summer due to respiration, whereas alkalinity remains relatively constant. As for oxygen, the DIC accumulation is interrupted during deep water renewal by intrusions or deep mixing events (Fig. S15). The rate and magnitude of the DIC increase in bottom waters is in good agreement with observations (e.g., 2017–2018), indicating that respiration is parameterized appropriately. The occasional mismatches with observations can have at least two causes: an overestimation of vertical mixing that counteracts the accumulation of DIC in late fall, which is improved in ROMS-H3 (see Sect. 5.2.4), and a mismatch in the occurrence or effect of intrusion events. The latter is more pronounced in the second part of the simulation.

The magnitude of simulated alkalinity agrees with observations at 60 m (Fig. S14) with some mismatches in late fall (i.e. 2019–2022). The observed increase in alkalinity likely originates from benthic alkalinity fluxes associated with alkalinity production in the sediment (e.g., sulphate reduction), which are not yet parameterized in the model.

$p\text{CO}_2$ at the Compass Buoy station (calculated from TA and DIC with CO2SYS) has similar patterns as DIC (Fig. S16). At 60 m, the rate of change of $p\text{CO}_2$ is close to the observations, similar to the variability in deep DIC (accumulation, mixing/intrusions). At the surface, $p\text{CO}_2$ is within the observed range but the variability is larger in the model, probably driven by rapid changes in the location of the Sackville River plume. Simulated surface $p\text{CO}_2$ also tends to be elevated in the fall compared to observations, which is likely associated with overmixing at this time in ROMS-H2.

ROMS-H3 better represents vertical mixing in the fall (more stratification, Figs. S21–22). With this higher-resolution model deep oxygen and DIC do not level off as in ROMS-H2 and therefore are better represented in late summer and fall (i.e. Figs. S13, S23, S15, S25). Alkalinity is still underestimated at depth (bias: $-36.7 \text{ mmol m}^{-3}$), likely due to the missing source from the sediment in the model (see above).

The model-observation comparison presented above indicates that both ROMS-H2 and ROMS-H3 represent well the background carbonate system dynamics in the Halifax Harbour.

5 Alkalinity addition module

To simulate alkaline feedstock addition (often in a liquid form, e.g. slurry) and the effect of added alkalinity on the carbonate system, several tracers were added to the biogeochemical model. An overview of this model, including the addition module, is presented in Fig. 2. The alkalinity addition module is described in detail below. It was designed so that a single simulation can be used to calculate the difference between the realistic addition case and counterfactual.

5.1 Description

The carbonate system model simulates TA and DIC as state variables and computes the CO_2 air-sea flux online at every timestep. New tracers were added for ΔTA (mmol m^{-3}) and ΔDIC (mmol C m^{-3}), which represent the added alkalinity and the resulting increase in DIC, respectively. The model includes a point source for feedstock addition that can be set constant or to vary with time using a forcing file. Since the feedstock is often partly in particulate form (delivered as a slurry), a tracer TAP (mmol m^{-3}) is added to represent the particulate phase of the added feedstock, which dissolves into ΔTA and sinks at specified rates. The addition rate of feedstock is TA_{in} (mmol s^{-1}). The allocation of TA_{in} into the dissolved (ΔTA) and particulate (TAP) pools is set by the parameter $\theta_{\text{P:D}}$. This representation of dissolution of the particulate stock builds on the formulation in Wang et al. (2025). The value of $\theta_{\text{P:D}}$ varies for each feedstock and needs to be assessed prior to dosing. Simulation results (see below) from experiments with fully dissolved ($\theta_{\text{P:D}} = 0$) or fully particulate ($\theta_{\text{P:D}} = 1$) indicate the maximum effect of $\theta_{\text{P:D}}$. All tracers are also subject to advection and mixing. To calculate air-sea gas exchange with ($F_{\text{CO}_2}^{\text{a}}$; $\text{mmol C m}^{-2} \text{ d}^{-1}$) and without ($F_{\text{CO}_2}^{\text{c}}$; $\text{mmol C m}^{-2} \text{ d}^{-1}$) added alkalinity, surface $p\text{CO}_2$ is calculated from TA and DIC (counterfactual case, no added alkalinity, $p\text{CO}_2^{\text{c}}$) and from TA+ ΔTA and DIC+ ΔDIC (case with added alkalinity, $p\text{CO}_2^{\text{a}}$).

The formulations of the biogeochemical source and sink terms are as follows:

$$\frac{\partial \text{TAP}}{\partial t} = \theta_{\text{P:D}} \frac{\text{TA}_{\text{in}}}{V} - k_{\text{diss}} \text{TAP} - w_{\text{P}} \frac{\partial \text{TAP}}{\partial z} \quad (8)$$

$$\frac{\partial \Delta\text{TA}}{\partial t} = (1 - \theta_{\text{P:D}}) \cdot \frac{\text{TA}_{\text{in}}}{V} + k_{\text{diss}} \text{TAP} \quad (9)$$

where k_{diss} (d^{-1}) is the dissolution rate, w_{P} the sinking rate of TAP and V is the volume of the addition grid cell. $\text{TA}_{\text{in}} > 0$ at the addition location only. Although not size dependent, the exponential decay formulation of particulate feedstock dissolution is a first-order representation of dissolution that can

be parameterized to best match observations (see Sect. 5.2 below).

At the surface, ΔDIC varies due to the additional air-sea flux of CO_2 (ΔF_{CO_2} ; $\text{mmol C m}^{-2} \text{d}^{-1}$) into the top layer as follows:

$$\frac{\partial \Delta\text{DIC}}{\partial t} \Big|_{z=0} = F_{\text{CO}_2}^{\text{a}} - F_{\text{CO}_2}^{\text{c}} \quad (10)$$

$$F_{\text{CO}_2}^{\text{c}} = \nu k_{\text{CO}_2} K_{\text{CO}_2} (\text{pCO}_{2,\text{air}} - \text{pCO}_2^{\text{c}}) \quad (11)$$

$$F_{\text{CO}_2}^{\text{a}} = \nu k_{\text{CO}_2} K_{\text{CO}_2} (\text{pCO}_{2,\text{air}} - \text{pCO}_2^{\text{a}}) \quad (12)$$

where K_{CO_2} ($\text{mmol C m}^{-2} \text{atm}^{-1}$) is the solubility of CO_2 and νk_{CO_2} is the gas transfer velocity of CO_2 . The gas transfer velocity (cm h^{-1}) is parameterized using the gas transfer relationship in Wanninkhof (2014):

$$\nu k_{\text{CO}_2} = 0.251 \cdot u_{10}^2 \sqrt{\frac{660}{S_{\text{CO}_2}}} \quad (13)$$

where S_{CO_2} is the Schmidt number of CO_2 and u_{10} (m s^{-1}) is the wind velocity at 10 m height. The Schmidt number varies with temperature according to Wanninkhof (2014). The sensitivity of uptake to air-sea gas exchange was tested in Laurent et al. (2024b).

In the bottom layer N , sinking material is immediately re-suspended and therefore accumulates. Instead of a sinking loss term there, we assume that a constant fraction of TA_p present in the bottom layer (θ_{loss}) and thus in contact with the sediment is incorporated into the sediment through bioturbation and thus lost from the system so that:

$$\frac{\partial \Delta\text{TA}}{\partial t} \Big|_{z=N} = (1 - \theta_{\text{loss}}) \cdot k_{\text{diss}} \text{TA}_p \quad (14)$$

Given the lack of available information, we set $\theta_{\text{loss}} = 0$ in the simulations. Sediment-feedstock interaction is currently studied in the Halifax Harbour, which will help constrain this parameter in the future. At the release location an additional source term represents the feedstock addition so that $\theta_{\text{P:D}} \frac{\text{TA}_{\text{in}}}{V}$ is added to the TA_p pool and $(1 - \theta_{\text{P:D}}) \cdot \frac{\text{TA}_{\text{in}}}{V}$ to ΔTA .

5.2 Alkalinity addition simulations

ROMS-H2 was run for 1.5 years (July 2017 to December 2018) with constant addition in the first two months (60 d) at a rate of 1.29 mol s^{-1} for a total alkalinity addition of $6665 \times 10^3 \text{ mol}$ (Fig. S2). This corresponds to 194 t of magnesium hydroxide (Mg(OH)_2) and approximately 250 t of brucite (including impurities), the mineral added in Halifax Harbour in 2023 (Cornwall, 2023). The model ran in six different configurations: three outfall locations in the inner (Mill Cove, location of a waste water treatment plant), mid (Tufts Cove, location of a cooling pipe from power plant) and outer (Herring Cove, location of a waste water treatment plant) harbour (Fig. 1d) each with two different feedstocks, either fully dissolved ($\theta_{\text{P:D}} = 0$) or fully particulate

($\theta_{\text{P:D}} = 1$). We assumed a particulate feedstock with a particle size of $12 \mu\text{m}$, dissolution rate of 0.015 d^{-1} and sinking rate of 5.5 m^{-1} . The dissolution and sinking characteristics of the particulate feedstock were chosen to be similar to the brucite used in Halifax Harbour in 2023. Dissolution and sinking of brucite's full particle size distribution were first calculated using a 1D model and dissolution experiments in the lab. Bulk parameters for the one particle case of the reduced model were then estimated to match the time and depth profiles of the full particle size distribution. The addition occurred via a point source and distributed uniformly over the upper 5 m. The cases with addition inside the harbour (Tufts Cove and Mill Cove) were repeated with the higher-resolution ROMS-H3 (Sect. 5.2.4). For analysis, we used 5 areas for spatial integration (Fig. 1d), namely Bedford Basin (BB; 17.3 km^2), Inner Harbour (IH; 8.2 km^2), Outer Harbour (OH; 8.1 km^2), Northwest Arm (NA; 2.6 km^2) and Shelf (1785.9 km^2).

5.2.1 Tracking the dispersion and effects of alkalinity addition

We first describe the simulation with addition of dissolved feedstock at Tufts Cove (mid harbour). The sensitivity to feedstock and release location will be discussed in the following sections.

The location, magnitude and temporal evolution of ΔTA (Fig. 3–5) indicate the fate of added alkalinity. Tufts Cove is located on the northern side of The Narrows where circulation is controlled by tides, wind, and freshwater discharge from the Sackville River (Shan et al., 2011). Depending on the conditions (tide phase, wind direction, river discharge) ΔTA is advected either into Bedford Basin toward the head of the estuary or the outer harbour and the open shelf (Fig. 5). The average surface concentration of ΔTA in the harbour increased to $\sim 10 \text{ mmol m}^{-3}$ by the end of the release (late August 2017) and dropped to $< 5 \text{ mmol m}^{-3}$ a month later (Fig. 5). Once the release had stopped, ΔTA was mainly located in the Bedford Basin where residence time is longer (Wang et al., 2025). The Bedford Basin is also where the largest fraction of ΔTA was found during the simulation (Fig. 4). After the release, it took about 4 months to flush most of ΔTA out of the harbour (Fig. 4). Once advected to the inner shelf, ΔTA left the model domain rapidly. ΔDIC gradually increased at the surface during the release because of net ΔCO_2 uptake from the atmosphere (Fig. 6). After the release, ΔDIC decreased rapidly and, as for ΔTA , remained longer in the Bedford Basin before being flushed out.

The spatial patterns of ΔF_{CO_2} illustrate the location of the uptake. For the Tufts Cove addition, the magnitude of the uptake was largest in the Narrows, where the feedstock was released, followed by Bedford Basin, where ΔTA had a longer residence time (Fig. 7). CO_2 uptake happened quickly for the dissolved feedstock with 44 % of the total uptake within the model domain (59.2 t) occurring during the 2 months of




Figure 3. Time-averaged (July 2017–December 2018), vertically integrated ΔTA in the harbour and surrounding areas during each simulation (excluding offshore areas where $\Delta\text{TA} \sim 0$). Dosing locations are indicated with a red dot.




Figure 5. Snapshots of surface ΔTA in the simulation with the release of fully dissolved feedstock from Tufts Cove (offshore areas were excluded from the map). Dosing locations are indicated with a red dot. Satellite image was generated with MapBox[©] (<https://www.mapbox.com/about/maps>, last access: 19 June 2024).




Figure 4. Area-integrated time series of ΔTA in the simulations with dissolved and particulate feedstock for the four regions of the model domain. The Northwest Arm region is not shown because area-integrated ΔTA is small there. The spatial limits of the regions are indicated on Fig. 1d (BB: Bedford Basin, IH: Inner Harbour, OH: Outer Harbour, NA: Northwest Arm).

release and 95 % (127.5 t) within the first 6 months of the simulation (Figs. 7 and 8, orange line).

Maximum achievable net CO_2 uptake from the alkalinity addition was estimated to be 261 t using a theoretical uptake efficiency of 1.34 g CO_2 per g $\text{Mg}(\text{OH})_2$. The calculation assumed a potential CO_2 uptake of 0.89 mol CO_2 per mol TA , derived with CO2SYS using volume averaged local properties in ROMS-H2 (i.e., temperature, salinity, alkalinity and TIC). This value varies regionally (Zhou et al., 2025), with a global potential CO_2 uptake of ~ 0.85 mol CO_2 per mol

TA (Tyka, 2025). Total uptake over the course of the simulation was 134.5 t so the realized uptake within the ROMS-H2 model domain was 51.5 % of the achievable uptake (Tufts Cove/dissolved case, Fig. 9, Table 3). As discussed above, uptake varied spatially. Since the Inner Harbour (including The Narrows) is relatively small compared to the adjacent areas, total uptake in this area amounted to only 21.3 %. The largest fractions of the total uptake were in the Outer Harbour (30.7 %) and Bedford Basin (29.1 %). Despite its large area, the shelf accounted for only 17.6 % of the total uptake. The Northwest Arm had a negligible contribution (1.4 %).

5.2.2 Effect of release locations

Next, we describe the effect of release location for the simulations with addition of dissolved feedstock. The sensitivity to feedstock type is discussed in the next section. The outcome of the alkalinity addition is sensitive to the release location in the harbour and has potential implications when considering future deployments. For addition at the inshore site (Mill Cove), ΔTA concentration is much larger in the Bedford Basin even a month after the end of the release (Fig. 4, see also Fig. S17). The subsequent flushing is similar to the Tufts Cove case and most of the ΔTA was gone 4 months after the end of the release. For addition at the coastal site (Herring Cove), the amount of ΔTA is much lower ($< 5 \text{ mmol m}^{-3}$), most of it found near the release location in the Outer Harbour and on the shelf (Fig. 4, see also Fig. S18). Since residence times in the outer harbour and on

Table 3. Net ΔCO_2 uptake and realized uptake for each simulation with ROMS-H2, as shown in Figs. 8 and 9. Areas of the model grid are Bedford Basin (BB), Northwest Arm (NA), Inner Harbour (IH), Outer Harbour (OH) and Shelf. The total represents the sum of the five areas (all grid cells). Maximum achievable net CO_2 uptake from the alkalinity addition was estimated to be 261 t using a theoretical uptake efficiency of 1.34 g CO_2 per g $\text{Mg}(\text{OH})_2$ (see Sect. 5.2.1). The realized uptake is the relative difference between maximum achievable net uptake and simulated net uptake.

Dosing location	Feedstock type	Net ΔCO_2 uptake (t)					Realized uptake (%)					
		BB	NA	IH	OH	Shelf	Total	BB	NA	IH	OH	Shelf
Mill Cove (MC)	Particulate	90.8	1.3	26.6	32.7	14.7	166.1	34.8	0.5	10.2	12.5	5.6
	Dissolved	90.9	1.5	28.4	33.7	15.4	169.9	34.8	0.6	10.9	12.9	5.9
Tufts Cove (TC)	Particulate	49.9	3.5	20.8	32.6	18.5	125.3	19.1	1.3	8.0	12.5	7.1
	Dissolved	39.1	1.9	28.6	41.3	23.6	134.5	15.0	0.7	11.0	15.8	9.1
Herring Cove (HC)	Particulate	14.5	4.4	6.8	15.2	12.2	53.1	5.5	1.7	2.6	5.8	4.7
	Dissolved	3.4	1.4	3.1	40.9	30.9	79.7	1.3	0.5	1.2	15.7	11.8




Figure 6. Snapshots of surface ΔDIC in the simulation with the release of fully dissolved feedstock from Tufts Cove (offshore areas were excluded from the map). Dosing locations are indicated with a red dot. Satellite image was generated with MapBox© (<https://www.mapbox.com/about/maps>).

the shelf are much lower, ΔTA was transported rapidly out of the ROMS-H2 domain. Nearly all ΔTA was gone 1 month after the end of the release (Fig. 4, see also Fig. S18).

Figure S19 (left panels) indicates the location of ΔTA transport out of the eastern and western boundaries.

The magnitude of the cumulative net CO_2 uptake is similar after 2 months of addition in the three simulations (Tufts Cove: 59.2 t, Mill Cove: 64.8 t, Herring Cove: 58.2 t). The start of CO_2 uptake was somewhat delayed in July 2017 (by ~ 2 weeks) in the Mill Cove release case but then followed the same trajectory as the other cases. Local conditions re-

lated to the influence of the Sackville River (shallow stratification, discharge pulses) may have limited the onset of CO_2 uptake at the beginning of the simulation. Subsequently, the difference between the three release cases was in the location of the uptake (Fig. 7). The net CO_2 uptake was largest either in the Bedford Basin (Mill Cove release), the Narrows (Tufts Cove release) or the outer Harbour (Herring Cove release). After the initial release period (September 2017 onward), the trajectories of net CO_2 uptake changed drastically between the three cases, which highlight the importance of residence time for the realized net CO_2 uptake. When alkalinity was added at the coastal site (Herring Cove), most of the uptake (73.0 %) happened during the two months of feedstock release (92.6 % was completed after the initial 3 months). ΔTA was dispersed rapidly out of the domain when the release stopped and therefore net uptake was limited then. With an inshore release (Mill Cove), 56.0 % of the uptake occurred after the initial two months (the release period).

Accordingly, spatially integrated net uptake (Fig. 7, Table 3) and the realized net uptake (Fig. 9, Table 3) diverged significantly for the three release sites. The realized net uptake was maximum for the inshore release (Mill Cove), reaching 65.1 % by the end of the simulation, whereas only 30.5 % of the achievable uptake was realized for coastal release site (Herring Cove). In all cases > 95 % of the uptake occurred within the first 6 months of the simulation. These numbers are in line with the spatial redistribution of the uptake. For a release in Mill Cove, 70.2 % of the uptake occurred inshore (Bedford Basin: 53.5 %, Inner Harbour: 16.7 %), whereas it was only 8.2 % for a release in Herring Cove (Bedford Basin: 4.3 %, Inner Harbour: 3.9 %). In this latter case, 51.3 % of the uptake was in the outer harbour (where alkalinity is released) and 38.8 % on the shelf (Fig. 9).

5.2.3 Effect of feedstock type

The addition of alkalinity as particulate material at the 3 locations resulted in a delayed availability of ΔTA , controlled




Figure 7. Maps of time-averaged (July 2017–December 2018) net ΔCO_2 uptake during the simulations (excluding offshore areas where ΔCO_2 uptake ~ 0). Dosing locations are indicated with a red dot.




Figure 8. Time series of spatially integrated net CO_2 uptake in the simulations with dissolved (top) and particulate (bottom) feedstock.

by the feedstock's dissolution kinetics, in comparison to the dissolved feedstock addition (Figs. S20 and 4). The spatial distribution of ΔTA was also modified because a large fraction of the particulate feedstock sank to the bottom layer. There, it was entrained in the return flow of the estuarine circulation toward the Bedford Basin where it accumulated and dissolved (Fig. 10). ΔTA was relocated toward the Bedford Basin in this case (Fig. 3).

With particulate feedstock release at the 3 locations, only a small fraction of the uptake occurred during the release period (7.3 %–10.9 %, Figs. 7 and 8). After 6 months, only 71.5 %–78.3 % of the achievable uptake was realized and, for the releases at Mill Cove or Tufts Cove, it took the whole simulation (1.5 years) to reach uptake values close to those

in the case with fully dissolved feedstock (93.1 %–97.7 %, Figs. 7 and 9). These delays are the result of feedstock dissolution as well as the time it takes for the ΔTA signal to re-emerge at the surface and produce CO_2 uptake. The spatial distribution of net CO_2 uptake did not change from the dissolved feedstock case for the Mill Cove release but a relocation towards the Bedford Basin was observed for the Tufts Cove release case (Fig. 9).

For a release at Herring Cove, total uptake by the end of the simulation was much lower for particulate than dissolved feedstock (53.1 t, 66.6 % of dissolved case, Table 3). When particulate feedstock is released at Herring Cove, material that is not entrained in a return flow towards the head of the harbour is moved downcoast by the coastal current on




Figure 9. Realized net CO_2 uptake for each simulation. The colors in the legend indicate the total uptake in each area of the model grid. Pink and olive-green lines under the Tufts Cove – dissolved bar indicate the realized uptake in ROMS-H3 (inside harbour) and in the outer area of ROMS-H2 (outside harbour), respectively (see Fig. 1d). Maximum achievable net CO_2 uptake from the alkalinity addition was estimated to be 261 t using a theoretical uptake efficiency of 1.34 g CO_2 per g $\text{Mg}(\text{OH})_2$ (see Sect. 5.2.1). The realized uptake is the relative difference between maximum achievable net uptake and simulated net uptake.




Figure 10. Average distribution of particulate feedstock during the simulations with addition at the three locations (red dots).

the shelf and ultimately leaves the domain relatively quickly (Fig. S19). In this case, the potential uptake is not realized within the model domain.

5.2.4 High-resolution Halifax Harbour model

The capability of the addition model was tested above within ROMS-H2 which has 150 m horizontal resolution in the harbour and on the shelf. The third level nest ROMS-H3, with 51 m resolution, is better able to represent small scale features in the harbour. The high resolution also improves simulation of vertical mixing (Wang et al., 2025). ROMS-H3 was run for the same period as the addition experiments in ROMS-H2 (July 2017 to December 2018).

Alkalinity addition in ROMS-H3 was tested for the four cases where release occurred inside ROMS-H3, namely Tufts Cove and Mill Cove each with either dissolved or particulate feedstock. Boundary conditions at the entrance of the harbour were provided by the corresponding ROMS-H2 simulations (see Sect. 5.2.2 and 5.2.3). ROMS-H3 simulates small scale features in the harbour, driven by local wind and tides, especially around the release location (Fig. 11). High resolution was particularly important to represent spatial features in The Narrows for the Tufts Cove release (Fig. 11b) and in Bedford Basin for the Mill Cove release (Fig. 11c). The dispersion of feedstock outside the harbour was tracked, as previously discussed, with ROMS-H2 (Fig. 11a, d).

Net CO_2 fluxes in the three-level nest were consistent with ROMS-H2 (Fig. S26), especially for a release from Tufts Cove (Fig. S27a, b). The main temporal difference occurred with particulate feedstock during a period of intrusion/mixing from late December 2017 to January 2018. In Bedford Basin, the intrusion brought to the surface deep ΔTA (dissolved from particulate feedstock in the bottom layer) and was followed by a period of enhanced net CO_2 fluxes in ROMS-H3. The differences were more pronounced with a release at Mill Cove (Fig. S27c, d). For dissolved feedstock, net CO_2 flux was larger in late summer and fall, consistent with the better representation of vertical mixing in ROMS-H3 (Fig. S27c). For particulate feedstock, the net CO_2 flux was also higher following the intrusion (Fig. S27d). Integrated in time, total net CO_2 flux did not change significantly with ROMS-H3 for a release at Tufts Cove. For Mill Cove, net CO_2 flux increased only slightly with particulate feedstock, whereas the realized uptake increased to 68.8 % with dissolved feedstock (+3.1 %). This increase is associated with enhanced fluxes in the Bedford Basin (Fig. S26).

6 Discussion

6.1 Model performance

The circulation model is modified from the nested model of Wang et al. (2025) using rotated and refined grids. Not surprisingly, the physical model had similar performance to Wang et al.'s version when compared with observations (Table 2). The main difference to Wang et al. (2025) and earlier circulation models of the Halifax Harbour (Shan, 2010; Sui et al., 2024) was the coupling with biogeochemistry. The simplified biology, based on observed respiration rates in the Bedford Basin and simulated rates on the shelf, was sufficient to reproduce the characteristic benthic respiration-driven deep-water depletion/accumulation of oxygen and DIC in the Bedford Basin (Burt et al., 2013; Rakshit et al., 2023, 2025). The model also simulated the gradient in oxygen and inorganic carbon variables between the harbour and the shelf as seen when comparing the time series at the Compass Buoy station and Station 2. Using parameterizations




Figure 11. Snapshots of surface ΔTA in the nested simulations with ROMS-H2 and ROMS-H3 for a release of fully dissolved feedstock at Tufts Cove (a, b) and at Mill Cove (c, d). Panels (b) and (c) show a zoom in on the inner Harbour (b) and the Bedford Basin (c). Dosing locations are indicated with a red dot. Satellite images were generated with MapBox© (<https://www.mapbox.com/about/maps>, last access: 19 June 2024).

rather than full descriptions of biological sources and sinks allowed for efficient carbonate system dynamics in Halifax. In this context the simplified model, which is meant to represent the background state and counterfactual case during TA addition experiments, seems fit for purpose.

Winter mixing and water intrusions from the shelf (Garcia Larez, 2023; Sui et al., 2024) are the two physical processes controlling subsurface water renewal in Bedford Basin (Burt et al., 2013; Haas et al., 2021; Rakshit et al., 2023). The two processes are important in the context of alkalinity enhancement because they lower residence time (Wang et al., 2025) and bring subsurface waters into contact with the atmosphere. Deep water renewal from convective mixing and intrusions are simulated well by the model, with some discrepancies to observations. More specifically, the frequency and magnitude of intrusions are sometimes overestimated in the baseline simulation (see Garcia Larez, 2023 for a detailed statistical comparison with the Wang et al., 2025, model version). OAE simulations were carried out over a period when convective mixing and intrusions were well represented in both ROMS-H2 and ROMS-H3 so their background dynamics can be assumed correct. The impact of intrusion events is illustrated, for example, by the enhanced net CO_2 uptake from December 2017 to January 2018 in the case with particulate feedstock where an intrusion redistributed ΔTA from the bottom layer in late December (Fig. 8).

Validating the simulated circulation and background carbonate chemistry is an important step prior to simulating alkalinity additions in the harbour as it gives confidence in the

model results. Local circulation controls the distribution of added alkalinity, whereas background carbonates can influence the timing of net CO_2 uptake (Fennel, 2026). However, other coastal systems where the model may be relocated won't necessarily have long-term time series that can be used for validation. In such cases, available information about general circulation features, satellite SST and any prior data should be used. Comparison with climatological carbonate data or targeted sampling are needed to increase confidence in the model outputs.

6.2 Alkalinity addition module

The model with alkalinity addition module was developed with the goal of supporting realistic OAE field trials and associated research and operations in Halifax Harbour (Laurent et al., 2024a) as well as to provide more general insights into the potential of coastal OAE, the optimal design of field operations, and as a flexible and freely available capability for MRV in this and other coastal systems. The key characteristics of the addition module are: (1) its capability to simulate realistic time series of point source addition, (2) tuneable feedstock properties (i.e. dissolved and particulate fractions, dissolution and sinking rates), (3) spatially and temporally resolved tracking of feedstock effects on the carbonate system, (4) quantification of the additional CO_2 air-sea flux using a unified code for addition and counterfactual cases, and (5) relocatability with, arguably, minor effort.

The capability for point source addition was included in the model to enable time-varying rates of feedstock dosing rates, which is relevant for real-world operations. This level of detail is especially important in dynamic tidal environments, such as the Halifax Harbour, where the timing of the release with respect to the tidal phase influences the spatial dispersion pattern of the feedstock (Wang et al., 2025). Fine-scale dispersion features, as seen in ROMS-H3 (Fig. 11) depend on the accurate timing of additions. They also depend on the characteristics of the feedstock. Many types of alkaline material are available for OAE (Caserini et al., 2022; Eisaman et al., 2023; Renforth, 2019) with a range of specific properties (e.g., dissolution kinetics, particle size). Wang et al. (2025) showed that feedstock properties have a strong effect on exposure and detectability of OAE signals in Halifax Harbour. Building on this prior work, the addition module presented here shows examples of the effects of feedstock properties on the carbonate system and net CO₂ uptake in the Halifax Harbour (Figs. 8 and 9; see Sect. 6.3 below).

Coupled, high-resolution, spatially resolved regional models are essential for MRV of OAE (Fennel et al., 2023). They provide the counterfactual case that simply cannot be obtained with in-situ measurements (Fennel, 2026). The simulations presented here show that for a release in Halifax Harbour, depending on the feedstock and release location, 20 %–69 % of the potential CO₂ uptake was realized inside or in direct surroundings of the harbour (Fig. 9), mostly within six months of release (Figs. 7 and 8). One can expect comparable results for realized uptake in other coastal systems with similar characteristics. Residence time is a particularly relevant factor (Wang et al., 2025). Despite their importance, inshore/coastal areas are poorly resolved, if at all, in global ocean models (Laurent et al., 2021) and their residence time can be underestimated significantly in coarse models (Rutherford and Fennel, 2018). Global models therefore may be more appropriate for assessing first-order, large-scale patterns of coastal OAE potential and for simulating far-field and global responses of the Earth system to coastal additions (Feng et al., 2017; He and Tyka, 2023). Regional models, such as the one presented here, should be deemed essential for a comprehensive verification of net CO₂ uptake resulting from coastal additions. Our results indicated that the multi-nest ROMS model with TA addition module has the appropriate level of skill and thus capability for that task. The two or three nested levels provided sufficient resolution for representing the background state of the carbonate system and for assessing additionality at local to regional scale.

One of the features of the addition module is its unified source code for the paired, simultaneous simulation of the addition and counterfactual cases providing computational efficiency and simplicity. Zhou et al. (2025) took a similar approach in their global model. Adding ΔTA , ΔDIC and particulate feedstock as state variables in the addition module provides a simple and efficient way to track OAE-related perturbations of the background carbonate system. An optional,

diagnostic state variable was added to keep track of the alkalinity originating from in-situ dissolution of feedstock in contrast to feedstock that was added directly in dissolved form. It should be relatively easy to relocate the framework presented here to other coastal systems. At a minimum, relocation requires knowledge of the local alkalinity to salinity relationship and local sources and sinks of alkalinity and DIC from rivers, sewage, and sediments. Ideally, information would also be available to tune the parameterizations of primary production and water-column and sediment respiration. Feedstock should also be characterized well.

6.3 Simulating additionality in the Halifax Harbour

The detectability of ΔTA during addition operations and the quantification of its effects on air-sea fluxes is a challenge for OAE (Ho et al., 2023). This is particularly true in systems with large seasonal variations of the background carbonate system variables such as Halifax Harbour (Fig. S14–16). In this system, Wang et al. (2025) demonstrated that detectability varies in time and space and provided insights for maximizing ΔTA detection while minimizing exposure risk. Our results on the distribution of ΔTA are in line with these findings. The mean estuarine circulation of Halifax Harbour results in retention of the ΔTA signal in the Bedford Basin. This area may be an ideal spot for maximizing the detectability of ΔTA and its effects on CO₂ uptake although, as noted by Wang et al. (2025), increased detectability goes hand in hand with an increased risk of ecosystem exposure. The latter would also be most acute if the release occurred at the head of the basin (Figs. 10 and 11), especially with long term, continuous dosing as alkalinity would accumulate there, raising the background pH. In the case of a particulate feedstock, it would also increase the interaction with the sediment. In a follow up study, we will explore the maximum dosing allowed in the Halifax Harbour to meet the local environmental regulations, i.e. pH < 9 (Bach et al., 2025).

The timescale of CO₂ equilibration is an important parameter for OAE because it influences the location of net CO₂ uptake as well as the maximum achievable uptake if the residence time of ΔTA in the surface mixed layer is shorter than the equilibration timescale. In the open ocean, the timescale of CO₂ equilibration is 3 to 6 months on average (Jones et al., 2014). Following an alkalinity addition, the time scale will vary with the thickness of the surface layer and the initial disequilibrium with the atmosphere (Fennel, 2026). In the simulations presented here, the time evolution of net CO₂ uptake is consistent with these previously reported timescales. Note that residence time is relatively similar to the equilibration time scale in the entire Bedford Basin (*e*-folding flushing time ~ 90 d, relevant for the particulate feedstock), smaller in the upper Bedford Basin only (0–20 m, *e*-folding flushing time ~ 39 d, relevant for the dissolved feedstock) but much shorter elsewhere (< 5 d) (Shan and Sheng, 2012; Wang et al., 2025).

Release location is a main factor controlling ΔTA retention in the Bedford Basin (Fig. 4) and the magnitude of net CO_2 uptake occurring in the harbour (Figs. 7 and 9, Table 3). A Mill Cove release resulted in the most uptake within the model domain, independently of the feedstock (166–170 t CO_2 uptake). 53 % of uptake occurred in the Bedford Basin in this case. Another reason to carry out sensitivity simulations is to understand the effect of local circulation features on the dispersion of feedstock. Wang et al. (2025) found that feedstock characteristics affect the residence time of the addition signal. We also found a redistribution of the ΔTA signal, controlled by the estuarine return flow, toward the Bedford Basin (Fig. 3) and a delay in the uptake due to the feedstock dissolution kinetics (Fig. 9). However, despite the spatial and temporal redistribution of net CO_2 uptake, feedstock characteristics had a limited effect on total uptake by the end of the simulation when release occurred in the Bedford Basin (Mill Cove) and the inner harbour (Tufts Cove).

Tufts Cove is the location for alkalinity addition for on-going OAE operations in the Halifax Harbour. This is a practical choice and as discussed above total uptake is significant in this case. Our simulations also provide insights into potential alternative addition sites at WWTPs, which have been suggested for cost effective operations (Cai and Jiao, 2022). Additional factors such as the differences between the outfall types, their discharge depth and operational constraints need to be considered in each case of the future proposed sites. Aside from Mill Cove, two WWTPs are in the inner harbour (similar to Tufts Cove) and two in the outer harbour, including Herring Cove (Fig. 1d). Since feedstock disperses rapidly at a coastal release site, MRV would be more challenging for an outer harbour WWTP addition. In this case, a larger regional or global model would be essential to track net uptake.

The difference in horizontal model resolution between ROMS-H2 and ROMS-H3 (i.e. 150 m versus 50 m) did not have a systematic effect on net fluxes and therefore on the outcome of the additions. Nonetheless, higher resolution resulted in fine scale features of ΔTA and ΔDIC (Fig. 11), knowledge of which are informative to guide field sampling, validate simulated feedstock dissolution and dispersion, and therefore provide confidence in the estimates of net CO_2 air-sea fluxes. ROMS-H3 better resolved vertical mixing in the Bedford Basin, which ultimately influenced the residence time of ΔTA in the Bedford Basin, as in Wang et al. (2025). The resulting relocation of net CO_2 uptake toward the Bedford Basin was small but not negligible for a release at Mill Cove (Fig. S26).

6.4 Current limitations and future development

Several processes are either poorly constrained or not represented in the addition module as presented here. For instance, a constant sediment loss term that mimics the incorporation and loss of feedstock into the sediment is included in the ad-

dition module, but given the lack of observation-based constraints it was set to zero in our simulations. Earlier sensitivity tests indicated that results scale with the magnitude of this loss term (Laurent et al., 2024a, c). This parameterization provides a first-order representation of sediment loss but should be refined further to include possible feedback between feedstock deposition and benthic alkalinity flux (e.g., Bach, 2024). A better understanding of feedstock-sediment interaction will be necessary to simulate this feedback.

Particulate feedstock is represented with one tracer that dissolves with an exponential decay formulation. Given the high spatial model resolution, simulating a particle size spectrum was not feasible as it would be computationally too expensive. Although this bulk representation seemed appropriate for brucite, it may not be the case for feedstocks with a wider or non-gaussian particle size spectrum. Splitting TA_{in} into several tracers to better represent particulate feedstock should be considered in future model development. Size dependent dissolution should also be considered and may be implemented from multiple tracers.

BBMP observations indicate a source of TA in the deep layer of Bedford Basin (Fig. S14). Burt et al. (2013) and Rakshit et al. (2025) suggested the occurrence of sediment denitrification but they did not observe or report a matching TA flux. The source of TA accumulation at depth should be investigated and then parameterized accordingly in the model. This is the subject of an on-going study. Other sources of alkalinity and DIC were also poorly constrained in the model due to a lack of observations. For example, additional carbonate data should be collected in the Sackville River and WWTPs effluents, to better constrain natural sources of DIC and alkalinity in the model.

7 Conclusions

We developed a reduced-complexity biogeochemical module within a high-resolution nested model of the Halifax Harbour. The model was able to simulate the variability of the carbonate system parameters observed in the harbour. To simulate OAE, we included an option for alkalinity addition that tracks particulate feedstock, added alkalinity, and the resulting net CO_2 uptake through an additional DIC pool. The model provides the capability for testing the outcome of OAE scenarios, to quantify additionality and realized uptake within the model domain, as well as to operationally support OAE field trials. It is fully relocatable, as a module within the ROMS framework. We tested the addition module for the Halifax Harbour to assess its capability and support OAE activities in the harbour. Release locations had a strong effect on the outcome of the addition, whereas feedstock types influenced the distribution of the signal. The simulations showed that up to 69 % of the net CO_2 uptake was realized within the model domain, most of it inside the harbour. These results stress the importance of operational de-

sign, such as dosing location, type (dissolved, particulate), slurry properties (dissolution rate, sinking speed), as well as the use of high-resolution regional models when quantifying additionality. In the future, the nested model will be applied to on-going OAE operations in the Halifax Harbour, to further validate the addition module, understand the fate of the alkalinity added to the harbour and to quantify net CO₂ uptake.

Code and data availability. The ROMS code with reduced BGC and alkalinity addition is available at <https://doi.org/10.5281/zenodo.17871564> (Laurent et al., 2025a). Model outputs used in the analysis are available from the Federated Research Data Repository at <https://doi.org/10.20383/103.01525> (Laurent et al., 2025b).

Supplement. The supplement related to this article is available online at <https://doi.org/10.5194/bg-23-115-2026-supplement>.

Author contributions. AL, BW, and KF developed the model. AL performed the simulations. AL and KF designed the study, performed the analysis and drafted the manuscript. Co-authors provided data, edited and commented on the manuscript.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Acknowledgements. We thank the four anonymous reviewers and the editor for their valuable time and their constructive comments.

Financial support. This research has been supported by the Carbon to Sea grant Ocean Alk-Align, the NSERC Alliance Missions grant ALLRP 570525-2021, a grant from the ICONIQ Impact Ocean Co-Lab, and the NSERC Discovery grant RGPIN-2022-688-02975.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors. Views expressed in the text are those of the authors and do not necessarily reflect the views of the publisher.

Review statement. This paper was edited by Jack Middelburg and reviewed by four anonymous referees.

References

Atamanchuk, D., Normandeau, C., Oberlander, J., Rondón, A. N., Wallace, D. W., and Fradette, C.: Carbonate system parameters (pH_{tot}, Total Alkalinity, Dissolved Inorganic Carbon), salinity, nutrients, in situ sensor-based measurements of temperature and depth collected from Bedford Basin/Halifax Harbour for the period of January to August 2023, PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.973118>, 2024.

Bach, L., Tyka, M., Wang, B., and Fennel, K.: Lethal by design? Guiding environmental assessments of ocean alkalinity enhancement toward realistic contextualization of the alkalinity perturbation, Preprint – CDRXIV, <https://doi.org/10.70212/cdrxiv.2025457.v1>, 2025.

Bach, L. T.: The additionality problem of ocean alkalinity enhancement, *Biogeosciences*, 21, 261–277, <https://doi.org/10.5194/bg-21-261-2024>, 2024.

Bach, L. T., Ho, D. T., Boyd, P. W., and Tyka, M. D.: Toward a consensus framework to evaluate air-sea CO₂ equilibration for marine CO₂ removal, *Limnol. Oceanogr. Lett.*, 8, 685–691, <https://doi.org/10.1002/lol2.10330>, 2023.

Buckley, D. E. and Winters, G. V.: Geochemical characteristics of contaminated surficial sediments in Halifax Harbour: impact of waste discharge, *Can. J. Earth Sci.*, 29, 2617–2639, <https://doi.org/10.1139/e92-208>, 1992.

Burt, D. J., Fröb, F., and Ilyina, T.: The sensitivity of the marine carbonate system to regional ocean alkalinity enhancement, *Frontiers in Climate*, 3, <https://doi.org/10.3389/fclim.2021.624075>, 2021.

Burt, W. J., Thomas, H., Fennel, K., and Horne, E.: Sediment-water column fluxes of carbon, oxygen and nutrients in Bedford Basin, Nova Scotia, inferred from ²²⁴Ra measurements, *Biogeosciences*, 10, 53–66, <https://doi.org/10.5194/bg-10-53-2013>, 2013.

Cai, W.-J. and Jiao, N.: Wastewater alkalinity addition as a novel approach for ocean negative carbon emissions, *The Innovation*, 3, 100272, <https://doi.org/10.1016/j.innn.2022.100272>, 2022.

Carter, B. R., Williams, N. L., Evans, W., Fassbender, A. J., Barbero, L., Hauri, C., Feely, R. A., and Sutton, A. J.: Time of Detection as a Metric for Prioritizing Between Climate Observation Quality, Frequency, and Duration, *Geophys. Res. Lett.*, 46, 3853–3861, <https://doi.org/10.1029/2018GL080773>, 2019.

Caserini, S., Storni, N., and Grossi, M.: The Availability of Limestone and Other Raw Materials for Ocean Alkalinity Enhancement, *Global Biogeochem. Cy.*, 36, <https://doi.org/10.1029/2021GB007246>, 2022.

Chin, T. M., Vazquez-Cuervo, J., and Armstrong, E. M.: A multi-scale high-resolution analysis of global sea surface temperature, *Remote Sens. Environ.*, 200, 154–169, <https://doi.org/10.1016/j.rse.2017.07.029>, 2017.

Cornwall, W.: An alkaline solution, *Science*, 382, 988–992, <https://doi.org/10.1126/science.adn1880>, 2023.

Craig, S. E., Jones, C. T., Li, W. K. W., Lazin, G., Horne, E., Caverhill, C., and Cullen, J. J.: Deriving optical metrics of coastal phytoplankton biomass from ocean colour, *Remote Sens. Environ.*, 119, 72–83, <https://doi.org/10.1016/j.rse.2011.12.007>, 2012.

Cullen, J. J. and Boyd, P. W.: Predicting and verifying the intended and unintended consequences of large-scale ocean iron fertilization, *Mar. Ecol. Prog. Ser.*, 364, 295–301, <https://doi.org/10.3354/meps07551>, 2008.

Cyr, F., Gibb, O., Azetsu-Scott, K., Chassé, J., Galbraith, P., Maillet, G., Pepin, P., Punshon, S., and Starr, M.: Ocean carbonate parameters on the Canadian Atlantic Continental Shelf, Federated Research Data Repository [data set], <https://doi.org/10.20383/102.0673>, 2022.

Delacroix, S., Nystuen, T. J., Tobiesen, A. E. D., King, A. L., and Höglund, E.: Ocean alkalinity enhancement impacts: regrowth of marine microalgae in alkaline mineral concentrations simulating the initial concentrations after ship-based dispersions, *Biogeosciences*, 21, 3677–3690, <https://doi.org/10.5194/bg-21-3677-2024>, 2024.

Dickson, A. G.: Standard potential of the reaction: $\text{AgCl}(\text{s}) + 0.5 \text{H}_2(\text{g}) = \text{Ag}(\text{s}) + \text{HCl}(\text{aq})$, and the standard acidity constant of the ion HSO_4^- in synthetic sea water from 273.15 to 318.15 K, *Journal of Chemical Thermodynamics*, 22, 113–127, [https://doi.org/10.1016/0021-9614\(90\)90074-Z](https://doi.org/10.1016/0021-9614(90)90074-Z), 1990.

Dickson, A. G. and Millero, F. J.: A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media, *Deep-Sea Res. Pt. A*, 34, 1733–1743, [https://doi.org/10.1016/0198-0149\(87\)90021-5](https://doi.org/10.1016/0198-0149(87)90021-5), 1987.

Durski, S. M., Glenn, S. M., and Haidvogel, D. B.: Vertical mixing schemes in the coastal ocean: Comparison of the level 2.5 Mellor-Yamada scheme with an enhanced version of the K profile parameterization, *J. Geophys. Res.-Ocean.*, 109, <https://doi.org/10.1029/2002JC001702>, 2004.

Eisaman, M. D., Geilert, S., Renforth, P., Bastianini, L., Campbell, J., Dale, A. W., Foteinis, S., Grasse, P., Hawrot, O., Löscher, C. R., Rau, G. H., and Rønning, J.: Assessing the technical aspects of ocean-alkalinity-enhancement approaches, in: Guide to Best Practices in Ocean Alkalinity Enhancement Research, edited by: Oschlies, A., Stevenson, A., Bach, L. T., Fennel, K., Rickaby, R. E. M., Satterfield, T., Webb, R., and Gattuso, J.-P., Copernicus Publications, State Planet, 2-oea2023, 3, <https://doi.org/10.5194/sp-2-oea2023-3-2023>, 2023.

Fakhraee, M., Li, Z., Planavsky, N. J., and Reinhard, C. T.: A biogeochemical model of mineral-based ocean alkalinity enhancement: impacts on the biological pump and ocean carbon uptake, *Environmental Research Letters*, 18, 044047, <https://doi.org/10.1088/1748-9326/acc9d4>, 2023.

Feng, E. Y., Koeve, W., Keller, D. P., and Oschlies, A.: Model-based assessment of the CO_2 sequestration potential of coastal ocean alkalization, *Earth's Future*, 5, 1252–1266, <https://doi.org/10.1002/2017EF000659>, 2017.

Fennel, K.: Widespread implementation of controlled upwelling in the North Pacific Subtropical Gyre would counteract diazotrophic N_2 fixation, *Mar. Ecol. Prog. Ser.*, 371, 301–303, <https://doi.org/10.3354/meps07772>, 2008.

Fennel, K.: The role of continental shelves in nitrogen and carbon cycling: Northwestern North Atlantic case study, *Ocean Sci.*, 6, 539–548, <https://doi.org/10.5194/os-6-539-2010>, 2010.

Fennel, K.: The Verification Challenge of Marine Carbon Dioxide Removal, *Ann. Rev. Mar. Sci.*, 18, 6.1–24, <https://doi.org/10.1146/annurev-marine-032123-025717>, 2026.

Fennel, K., Wilkin, J., Levin, J., Moisan, J., O'Reilly, J., and Haidvogel, D.: Nitrogen cycling in the Middle Atlantic Bight: Results from a three-dimensional model and implications for the North Atlantic nitrogen budget, *Global Biogeochem. Cy.*, 20, GB3007, <https://doi.org/10.1029/2005GB002456>, 2006.

Fennel, K., Hu, J., Laurent, A., Marta-Almeida, M., and Hetland, R.: Sensitivity of hypoxia predictions for the northern Gulf of Mexico to sediment oxygen consumption and model nesting, *J. Geophys. Res.-Ocean.*, 118, 990–1002, <https://doi.org/10.1002/jgrc.20077>, 2013.

Fennel, K., Long, M. C., Algar, C., Carter, B., Keller, D., Laurent, A., Mattern, J. P., Musgrave, R., Oschlies, A., Ostiguy, J., Palter, J. B., and Whitt, D. B.: Modelling considerations for research on ocean alkalinity enhancement (OAE), in: Guide to Best Practices in Ocean Alkalinity Enhancement Research, edited by: Oschlies, A., Stevenson, A., Bach, L. T., Fennel, K., Rickaby, R. E. M., Satterfield, T., Webb, R., and Gattuso, J.-P., Copernicus Publications, State Planet, 2-oea2023, 9, <https://doi.org/10.5194/sp-2-oea2023-9-2023>, 2023.

Ferderer, A., Schulz, K. G., Riebesell, U., Baker, K. G., Chase, Z., and Bach, L. T.: Investigating the effect of silicate- and calcium-based ocean alkalinity enhancement on diatom silicification, *Biogeosciences*, 21, 2777–2794, <https://doi.org/10.5194/bg-21-2777-2024>, 2024.

Fisheries and Oceans Canada: Bedford Basin Monitoring Program, <https://www.bio.gc.ca/science/monitoring-monitorage/bbmp-pobb/bbmp-pobb-en.php> (last access: 7 January 2025), 2025.

Fuhrman, J., McJeon, H., Patel, P., Doney, S. C., Shobe, W. M., and Clarens, A. F.: Food-energy–water implications of negative emissions technologies in a +1.5 °C future, *Nat. Clim. Change*, 10, 920–927, <https://doi.org/10.1038/s41558-020-0876-z>, 2020.

Garcia Larez, E.: Using observations and model simulations to characterize the effects of lateral intrusion events in Bedford Basin, MS thesis, Dalhousie University, Halifax, <http://hdl.handle.net/10222/83180> (last access: 13 January 2025), 2023.

Gattuso, J.-P., Williamson, P., Duarte, C. M., and Magnan, A. K.: The Potential for Ocean-Based Climate Action: Negative Emissions Technologies and Beyond, *Frontiers in Climate*, 2, <https://doi.org/10.3389/fclim.2020.575716>, 2021.

Guo, J. A., Strzepek, R. F., Swadling, K. M., Townsend, A. T., and Bach, L. T.: Influence of ocean alkalinity enhancement with olivine or steel slag on a coastal plankton community in Tasmania, *Biogeosciences*, 21, 2335–2354, <https://doi.org/10.5194/bg-21-2335-2024>, 2024.

Haas, S., Robicheau, B. M., Rakshit, S., Tolman, J., Algar, C. K., LaRoche, J., and Wallace, D. W. R.: Physical mixing in coastal waters controls and decouples nitrification via biomass dilution, *Proceedings of the National Academy of Sciences*, 118, <https://doi.org/10.1073/pnas.2004877118>, 2021.

Haidvogel, D. B., Arango, H., Budgell, W. P., Cornuelle, B. D., Curchitser, E., Lorenzo, E. Di, Fennel, K., Geyer, W. R., Hermann, A. J., Lanerolle, L., Levin, J., McWilliams, J. C., Miller, A. J., Moore, A. M., Powell, T. M., Shchepetkin, A. F., Sherwood, C. R., Signell, R. P., Warner, J. C., and Wilkin, J.: Ocean forecasting in terrain-following coordinates: Formulation and skill assessment of the Regional Ocean Modeling System, *J. Comput. Phys.*, 227, 3595–3624, <https://doi.org/10.1016/j.jcp.2007.06.016>, 2008.

He, J. and Tyka, M. D.: Limits and CO_2 equilibration of near-coast alkalinity enhancement, *Biogeosciences*, 20, 27–43, <https://doi.org/10.5194/bg-20-27-2023>, 2023.

Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I.,

Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 hourly data on single levels from 1940 to present, Climate Data Store, <https://doi.org/10.24381/cds.adbb2d47>, 2023.

Ho, D. T., Bopp, L., Palter, J. B., Long, M. C., Boyd, P. W., Neukermans, G., and Bach, L. T.: Monitoring, reporting, and verification for ocean alkalinity enhancement, in: Guide to Best Practices in Ocean Alkalinity Enhancement Research, edited by: Oschlies, A., Stevenson, A., Bach, L. T., Fennel, K., Rickaby, R. E. M., Satterfield, T., Webb, R., and Gattuso, J.-P., Copernicus Publications, State Planet, 2-oea2023, 12, <https://doi.org/10.5194/sp-2-oea2023-12-2023>, 2023.

Humphreys, M. P., Lewis, E. R., Sharp, J. D., and Pierrot, D.: PyCO2SYS v1.8: marine carbonate system calculations in Python, *Geosci. Model Dev.*, 15, 15–43, <https://doi.org/10.5194/gmd-15-15-2022>, 2022.

IPCC: Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Core Writing Team, Lee, H., and Romero, J., IPCC, Geneva, Switzerland, 35–115, <https://doi.org/10.59327/IPCC/AR6-9789291691647>, 2023.

Ilyina, T., Wolf-Gladrow, D., Munhoven, G., and Heinze, C.: Assessing the potential of calcium-based artificial ocean alkalization to mitigate rising atmospheric CO₂ and ocean acidification, *Geophys. Res. Lett.*, 40, 5909–5914, <https://doi.org/10.1002/2013GL057981>, 2013.

Jones, D. C., Ito, T., Takano, Y., and Hsu, W.: Spatial and seasonal variability of the air-sea equilibration timescale of carbon dioxide, *Global Biogeochem. Cy.*, 28, 1163–1178, <https://doi.org/10.1002/2014GB004813>, 2014.

Khangaonkar, T., Carter, B. R., Premathilake, L., Yun, S. K., Ni, W., Stoll, M. M., Ward, N. D., Hemery, L. G., Torres Sanchez, C., Subban, C. V., Ringham, M. C., Eisaman, M. D., Pelman, T., Tallam, K., and Feely, R. A.: Mixing and dilution controls on marine CO₂ removal using alkalinity enhancement, *Environmental Research Letters*, 19, 104039, <https://doi.org/10.1088/1748-9326/ad7521>, 2024.

Köhler, P., Abrams, J. F., Völker, C., Hauck, J., and Wolf-Gladrow, D. A.: Geoengineering impact of open ocean dissolution of olivine on atmospheric CO₂, surface ocean pH and marine biology, *Environmental Research Letters*, 8, 014009, <https://doi.org/10.1088/1748-9326/8/1/014009>, 2013.

Kwiatkowski, L., Berger, M., Bopp, L., Doléac, S., and Ho, D. T.: Contrasting carbon dioxide removal potential and nutrient feedbacks of simulated ocean alkalinity enhancement and macroalgae afforestation, *Environmental Research Letters*, 18, 124036, <https://doi.org/10.1088/1748-9326/ad08f9>, 2023.

Large, W. G. and Gent, P. R.: Validation of Vertical Mixing in an Equatorial Ocean Model Using Large Eddy Simulations and Observations, *J. Phys. Oceanogr.*, 29, 449–464, [https://doi.org/10.1175/1520-0485\(1999\)029<0449:VOVMIA>2.0.CO;2](https://doi.org/10.1175/1520-0485(1999)029<0449:VOVMIA>2.0.CO;2), 1999.

Laurent, A., Fennel, K., Hu, J., and Hetland, R.: Simulating the effects of phosphorus limitation in the Mississippi and Atchafalaya River plumes, *Biogeosciences*, 9, 4707–4723, <https://doi.org/10.5194/bg-9-4707-2012>, 2012.

Laurent, A., Fennel, K., Cai, W.-J., Huang, W.-J., Barbero, L., and Wanninkhof, R.: Eutrophication-induced acidification of coastal waters in the northern Gulf of Mexico: insights into origin and processes from a coupled physical-biogeochemical model, *Geophys. Res. Lett.*, 44, 946–956, <https://doi.org/10.1002/2016GL071881>, 2017.

Laurent, A., Fennel, K., and Kuhn, A.: An observation-based evaluation and ranking of historical Earth system model simulations in the northwest North Atlantic Ocean, *Biogeosciences*, 18, 1803–1822, <https://doi.org/10.5194/bg-18-1803-2021>, 2021.

Laurent, A., Wang, B., and Fennel, K.: Hindcast of the alkalinity addition field trial in Halifax Harbour, Canada (Fall 2023), *Federated Research Data Repository* [data set], <https://doi.org/10.20383/103.0987>, 2024a.

Laurent, A., Wang, B., and Fennel, K.: Hindcast of the alkalinity addition field trial in Halifax Harbour, Canada (Fall 2023) – Sensitivity tests: Air-sea flux, *Federated Research Data Repository* [data set], <https://doi.org/10.20383/103.0992>, 2024b.

Laurent, A., Wang, B., and Fennel, K.: Hindcast of the alkalinity addition field trial in Halifax Harbour, Canada (Fall 2023) – Sensitivity tests: Seabed loss, *Federated Research Data Repository* [data set], <https://doi.org/10.20383/103.0991>, 2024c.

Laurent, A., Wang, B., and Fennel, K.: alaurent101/ROMS_OAE: ROMS biogeochemical model with alkalinity addition (multiple feedstocks) (v2.1.0), *Zenodo* [code], <https://doi.org/10.5281/zenodo.17871564>, 2025a.

Laurent, A., Wang, B., and Fennel, K.: A high-resolution nested model to study the effects of alkalinity additions in Halifax Harbour: dataset of simulations, *Federated Research Data Repository* [data set], <https://doi.org/10.20383/103.01525>, 2025b.

Lellouche, J.-M., Greiner, E., Bourdallé-Badie, R., Garric, G., Melet, A., Drévillon, M., Bricaud, C., Hamon, M., Le Galloudec, O., Regnier, C., Candela, T., Testut, C.-E., Gasparin, F., Ruggiero, G., Benkiran, M., Drillet, Y., and Le Traon, P.-Y.: The Copernicus Global 1/12° Oceanic and Sea Ice GLORYS12 Reanalysis, *Front. Earth Sci.*, 9, <https://doi.org/10.3389/feart.2021.698876>, 2021.

Lenton, A., Matear, R. J., Keller, D. P., Scott, V., and Vaughan, N. E.: Assessing carbon dioxide removal through global and regional ocean alkalization under high and low emission pathways, *Earth System Dynamics*, 9, 339–357, <https://doi.org/10.5194/esd-9-339-2018>, 2018.

Levin, L. A., Alfaro-Lucas, J. M., Colaço, A., Cordes, E. E., Craik, N., Danovaro, R., Hoving, H.-J., Ingels, J., Mestre, N. C., Seabrook, S., Thurber, A. R., Vivian, C., and Yasuhara, M.: Deep-sea impacts of climate interventions, *Science*, 379, 978–981, <https://doi.org/10.1126/science.adc7521>, 2023.

Li, M., Chen, Y., Doyle, R., Testa, J. M., Gagnon, A., Bott, C., and Cai, W.-J.: Wastewater alkalinity enhancement for carbon emission reduction and marine CO₂ removal, *Environmental Research Letters*, 20, 044041, <https://doi.org/10.1088/1748-9326/adc1e3>, 2025.

McGinn, P. J., Dickinson, K. E., Park, K. C., Whitney, C. G., MacQuarrie, S. P., Black, F. J., Frigon, J.-C., Guiot, S. R., and O'Leary, S. J. B.: Assessment of the bioenergy and bioremediation potentials of the microalga *Scenedesmus* sp. AMDD cultivated in municipal wastewater effluent in batch and continuous mode, *Algal. Res.*, 1, 155–165, <https://doi.org/10.1016/j.algal.2012.05.001>, 2012.

MEDS: Maritimes Region Atlantic Zone Monitoring Program (AZMP) CTD Profiles [Data set], <https://catalogue.cioosatlantic>.

ca/dataset/ca-cioos_9a4bd73f-12a2-40ff-a7c7-b961a1d11311 (last access: 20 August 2024), 2024.

Mehrback, C., Culberson, C. H., Hawley, J. E., and Pytkowicz, R. M.: Measurement of the apparent dissociation constants of carbonic acid in seawater at atmospheric pressure, *Limnol. Oceanogr.*, 18, 897–907, <https://doi.org/10.4319/lo.1973.18.6.0897>, 1973.

Mukai, A. Y., Westerink, J. J., Luettich, R. A., and Mark, D. J.: Eastcoast 2001, A Tidal Constituent Database for Western North Atlantic, Gulf of Mexico, and Caribbean Sea, Technical report ERDC/CHL TR-02-24, U.S. Army Corps of Engineers, https://coast.nd.edu/reports_papers/2001-Mukai-TR-02-24.pdf (last access: 27 May 2024), 2002.

Nabuurs, G.-J., Mrabet, R., Abu Hatab, A., Bustamante, M., Clark, H., Havlík, P., House, J., Mbow, C., Ninan, K. N., Popp, A., Roe, S., Sohngen, B., and Towprayoon, S.: Agriculture, Forestry and Other Land Uses (AFOLU), in: IPCC, 2022: Climate Change 2022 – Mitigation of Climate Change. Contribution of Working Group III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Shukla, P. R., Skea, J., Slade, R., Al Khourdajie, A., van Diemen, R., McCollum, D., Pathak, M., Some, S., Vyas, P., Fradera, R., Belkacemi, M., Hasija, A., Lisboa, G., Luz, S., and Malley, J., Cambridge University Press, 747–860, <https://doi.org/10.1017/9781009157926.009>, 2023.

Nagwekar, T., Nissen, C., and Hauck, J.: Ocean Alkalinity Enhancement in Deep Water Formation Regions Under Low and High Emission Pathways, *Earth's Future*, 12, <https://doi.org/10.1029/2023EF004213>, 2024.

NASEM: Negative Emissions Technologies and Reliable Sequestration: A Research Agenda, National Academies Press, Washington, DC, <https://doi.org/10.17226/25259>, 2019.

NASEM: A Research Strategy for Ocean-based Carbon Dioxide Removal and Sequestration, National Academies Press, Washington, DC, <https://doi.org/10.17226/26278>, 2022.

Ohashi, K., Laurent, A., Renkl, C., Sheng, J., Fennel, K., and Oliver, E.: DaROMS-NWA12 v1.0, a coupled circulation–ice–biogeochemistry modelling system for the northwest Atlantic Ocean: development and validation, *Geosci. Model Dev.*, 17, 8697–8733, <https://doi.org/10.5194/gmd-17-8697-2024>, 2024.

Oschlies, A., Pahlow, M., Yool, A., and Matear, R. J.: Climate engineering by artificial ocean upwelling: Channelling the sorcerer's apprentice, *Geophys. Res. Lett.*, 37, <https://doi.org/10.1029/2009GL041961>, 2010.

Oschlies, A., Bach, L. T., Fennel, K., Gattuso, J.-P., and Mengis, N.: Perspectives and challenges of marine carbon dioxide removal, *Frontiers in Climate*, 6, <https://doi.org/10.3389/fclim.2024.1506181>, 2025.

Palmiéri, J. and Yool, A.: Global-scale evaluation of coastal ocean alkalinity enhancement in a fully coupled earth system model, *Earth's Future*, 12, <https://doi.org/10.1029/2023EF004018>, 2024.

Rakshit, S., Dale, A. W., Wallace, D. W., and Algar, C. K.: Sources and sinks of bottom water oxygen in a seasonally hypoxic fjord, *Front. Mar. Sci.*, 10, <https://doi.org/10.3389/fmars.2023.1148091>, 2023.

Rakshit, S., Glock, N., Dale, A. W., Armstrong, M. M. L., Scholz, F., Mutzberg, A., and Algar, C. K.: Foraminiferal denitrification and deep bioirrigation influence benthic biogeochemical cycling in a seasonally hypoxic fjord, *Geochim. Cosmochim. Acta*, 388, 268–282, <https://doi.org/10.1016/j.gca.2024.10.010>, 2025.

Renforth, P.: The negative emission potential of alkaline materials, *Nat. Commun.*, 10, 1401, <https://doi.org/10.1038/s41467-019-09475-5>, 2019.

Renforth, P. and Henderson, G.: Assessing ocean alkalinity for carbon sequestration, *Reviews of Geophysics*, 55, 636–674, <https://doi.org/10.1002/2016RG000533>, 2017.

Riahi, K., Schaeffer, R., Arango, J., Calvin, K., Guivarch, C., Hasegawa, T., Jiang, K., Kriegler, E., Matthews, R., Peters, G. P., Rao, A., Robertson, S., Sebbit, A. M., Steinberger, J., Tavoni, M., and van Vuuren, D. P.: Mitigation Pathways Compatible with Long-term Goals, in: IPCC, 2022: Climate Change 2022 - Mitigation of Climate Change. Contribution of Working Group III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Shukla, P. R., Skea, J., Slade, R., Al Khourdajie, A., van Diemen, R., McCollum, D., Pathak, M., Some, S., Vyas, P., Fradera, R., Belkacemi, M., Hasija, A., Lisboa, G., Luz, S., and Malley, J., Cambridge University Press, 295–408, <https://doi.org/10.1017/9781009157926.005>, 2023.

Rutherford, K. and Fennel, K.: Diagnosing transit times on the northwestern North Atlantic continental shelf, *Ocean Sci.*, 14, 1207–1221, <https://doi.org/10.5194/os-14-1207-2018>, 2018.

Rutherford, K., Fennel, K., Atamanchuk, D., Wallace, D., and Thomas, H.: A modelling study of temporal and spatial CO₂ variability on the biologically active and temperature-dominated Scotian Shelf, *Biogeosciences*, 18, 6271–6286, <https://doi.org/10.5194/bg-18-6271-2021>, 2021.

Schuldt, K. N., Mund, J., Aalto, T., Abshire, J. B., Aikin, K., Allen, G., Arlyn, A., Apadula, F., Arnold, S., Baier, B., Bakwin, P., Bäni, L., Bartyzel, J., Bentz, G., Bergamaschi, P., Beyersdorf, A., Biermann, T., Biraud, S. C., and Blanc, P.-E.: Multi-laboratory compilation of atmospheric carbon dioxide data for the period 1957–2022, *obspack_co2_1_GLOBALVIEWplus_v9.1_2023-12-08*, Global Monitoring Laboratory [Data set], <https://doi.org/10.25925/20231201>, 2023.

Schulz, K. G., Bach, L. T., and Dickson, A. G.: Seawater carbonate chemistry considerations for ocean alkalinity enhancement research: theory, measurements, and calculations, in: Guide to Best Practices in Ocean Alkalinity Enhancement Research, edited by: Oschlies, A., Stevenson, A., Bach, L. T., Fennel, K., Rickaby, R. E. M., Satterfield, T., Webb, R., and Gattuso, J.-P., Copernicus Publications, State Planet, 2-oea2023, 2, <https://doi.org/10.5194/sp-2-oea2023-2-2023>, 2023.

Scully, M. E.: Physical controls on hypoxia in Chesapeake Bay: A numerical modeling study, *J. Geophys. Res.-Ocean.*, 118, 1–18, <https://doi.org/10.1002/jgrc.20138>, 2013.

Shan, S.: Numerical Study of Three-dimensional Circulation and Hydrography in Halifax Harbour Using a Nested-grid Ocean Circulation Model, PhD thesis, Dalhousie University, Halifax, <http://hdl.handle.net/10222/13173> (23 September 2024), 2010

Shan, S. and Sheng, J.: Examination of circulation, flushing time and dispersion in Halifax Harbour of Nova Scotia, *Water Quality Research Journal*, 47, 353–374, <https://doi.org/10.2166/wqrjc.2012.041>, 2012.

Shan, S., Sheng, J., Thompson, K. R., and Greenberg, D. A.: Simulating the three-dimensional circulation and hydrography of Halifax Harbour using a multi-nested coastal

ocean circulation model, *Ocean. Dyn.*, 61, 951–976, <https://doi.org/10.1007/s10236-011-0398-3>, 2011.

Sui, Y., Sheng, J., Lu, Y., and Chen, S.: Intense lateral intrusion of offshore sub-surface waters in Halifax Harbour, *Cont. Shelf Res.*, 277, 105245, <https://doi.org/10.1016/j.csr.2024.105245>, 2024.

Tyka, M. D.: Efficiency metrics for ocean alkalinity enhancements under responsive and prescribed atmospheric CO₂ conditions, *Biogeosciences*, 22, 341–353, <https://doi.org/10.5194/bg-22-341-2025>, 2025.

UNEP: Emissions Gap Report 2023: Broken Record – Temperatures hit new highs, yet world fails to cut emissions (again), United Nations Environment Programme, Nairobi, <https://doi.org/10.59117/20.500.11822/43922>, 2023.

Wang, B., Laurent, A., Pei, Q., Sheng, J., Atamanchuk, D., and Fennel, K.: Maximizing the Detectability of Ocean Alkalinity Enhancement (OAE) While Minimizing Its Exposure Risks: Insights From a Numerical Study, *Earths Future*, 13, <https://doi.org/10.1029/2024EF005463>, 2025.

Wang, H., Pilcher, D. J., Kearney, K. A., Cross, J. N., Shugart, O. M., Eisaman, M. D., and Carter, B. R.: Simulated Impact of Ocean Alkalinity Enhancement on Atmospheric CO₂ Removal in the Bering Sea, *Earths Future*, 11, <https://doi.org/10.1029/2022EF002816>, 2023.

Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean revisited, *Limnol. Oceanogr. Methods*, 12, 351–362, <https://doi.org/10.4319/lom.2014.12.351>, 2014.

WMO: State of the Global Climate 2023, WMO-No. 1347, WMO, ISBN 978-92-63-11347-4, <https://library.wmo.int/idurl/4/68835> (last access: 28 October 2024), 2024a.

WMO: WMO Global Annual to Decadal Climate Update 2024–2028, WMO, <https://library.wmo.int/idurl/4/68910> (last access: 28 October 2024), 2024b.

Wu, H. and Zhu, J.: Advection scheme with 3rd high-order spatial interpolation at the middle temporal level and its application to saltwater intrusion in the Changjiang Estuary, *Ocean Modelling*, 33, 33–51, <https://doi.org/10.1016/j.ocemod.2009.12.001>, 2010.

Wu, J., Keller, D. P., and Oschlies, A.: Carbon dioxide removal via macroalgae open-ocean mariculture and sinking: an Earth system modeling study, *Earth Syst. Dynam.*, 14, 185–221, <https://doi.org/10.5194/esd-14-185-2023>, 2023.

Yu, L., Fennel, K., and Laurent, A.: A modeling study of physical controls on hypoxia generation in the northern Gulf of Mexico, *J. Geophys. Res. Pt. C*, 120, 5019–5039, <https://doi.org/10.1002/2014JC010634>, 2015.

Zhou, M., Tyka, M. D., Ho, D. T., Yankovsky, E., Bachman, S., Nicholas, T., Karspeck, A. R., and Long, M. C.: Mapping the global variation in the efficiency of ocean alkalinity enhancement for carbon dioxide removal, *Nat. Clim. Change*, 15, 59–65, <https://doi.org/10.1038/s41558-024-02179-9>, 2025.