



Subsoils, but not toeslopes, store millennia-old PyC in a gently sloping catchment under temperate climate after centuries of cultivation

Johanne Lebrun Thauront¹, Philippa Ascough², Sebastian Doetterl³, Negar Haghipour^{4,5}, Pierre Barré¹, Christian Walter⁶, and Samuel Abiven^{1,7}

¹Laboratoire de Géologie de l'ENS, ENS-PSL, CNRS, Paris, France

²NEIF Radiocarbon Laboratory, Scottish Universities Environmental Research Centre, East Kilbride, UK

³Soil Resources Group, ETHZ, Zurich, Switzerland

⁴Geological Institute, ETHZ, Zurich, Switzerland

⁵Laboratory of Ion Beam Physics, ETHZ, Zurich, Switzerland

⁶SAS, INRAE, Institut Agro Rennes Angers, Rennes, France

⁷CEREEP-Ecotron Île-de-France, ENS-PSL, CNRS, Saint-Pierre-lès-Nemours, France

Correspondence: Johanne Lebrun Thauront (johanne.lebrun.thauront@ens.psl.eu)

Received: 7 June 2025 – Discussion started: 26 June 2025

Revised: 4 November 2025 – Accepted: 4 December 2025 – Published: 8 January 2026

Abstract. Pyrogenic carbon (PyC) is the carbonaceous solid residue of incomplete combustion of biomass. It is a continuum of mostly condensed and aromatic molecules. PyC persists for longer in soils relative to non-PyC organic carbon. However, estimates of PyC residence time vary greatly. The time and spatial scales investigated are not always adapted to the long-residence time and vertical and lateral mobility of PyC in the soil profile and the landscape. In addition, agricultural land-use and shallow slopes are under-represented in the PyC literature.

We measured the concentrations and stocks of PyC down to 60 cm along three toposequences in a small agricultural catchment with shallow slopes and homogeneous soil parent material in the west of France. We used two methods (chemothermal oxidation – CTO and hydropyrolysis – HyPy) of PyC quantification that cover the intermediate to highly condensed part of the PyC continuum, and also measured the radiocarbon values in both total soil organic carbon (SOC) and the PyC fraction. There was likely little persistent PyC inputs to the catchment in the last 150 years which gave us access to the resultant, long term PyC distribution in the landscape. In particular, we aimed to investigate whether the vertical and horizontal distribution of PyC were similar or differed from SOC and whether they were affected by the soil types along the slope.

Topographic position was not the main driver of PyC stocks in this landscape. The stock of PyC_{CTO} to 60 cm depth averaged $2.5 \pm 0.22 \text{ t ha}^{-1}$ across topographic positions. PyC stocks were the highest in a Solimovic Cambisol at the toeslope ($3.3 \pm 0.26 \text{ t ha}^{-1}$), likely formed following changes in erosion dynamics with land-use. Contrary to previous reports, erosion redistributed already aged PyC without enrichment or depletion. PyC_{HyPy} concentrations in the topsoil decreased from upslope (median = 1.6, IQR = 0.22 g C kg^{-1} soil) to downslope positions (median = 1.10, IQR = 0.40 g C kg^{-1} soil), which we tentatively attribute to PyC_{HyPy} leaching following the destabilisation of mineral associations with iron oxides in the water-table affected portion of the transects. The subsoil (30–60 cm) represented between 37 % and 51 % of the PyC_{CTO} stock. PyC_{HyPy} proportion in SOC increased with depth and reached an average of $11 \pm 3.3 \%$ at 50–60 cm depth. PyC_{HyPy} had an uncalibrated radiocarbon age of 2520 to 9600 years BP at this depth, significantly older than bulk SOC at the same depth and than PyC_{HyPy} at 0–10 cm (1530 to 2630 years BP). These results confirm the long persistence of PyC in soils and point to a slow advection of PyC towards the soil depth under the pedoclimatic conditions of our study area.

Future studies should assess whether erosion modalities and age and quality of PyC affect its fate during erosion events. Identifying the proportion of PyC produced which is quickly transported away from the watershed and that which remains and is stabilised in soils for millennia after a fire is an important knowledge gap that still needs to be investigated to close the terrestrial PyC budget.

1 Introduction

Pyrogenic carbon (PyC) is the carbonaceous solid residue of incomplete combustion or pyrolysis of biomass and fossil fuels. It is present in all compartments of the earth system (Santín et al., 2016; Jones et al., 2019; Wagner et al., 2018). PyC is a continuum of molecules in which the degree of aromaticity and condensation is positively correlated with production temperature (Keiluweit et al., 2010; Wiedemeier et al., 2015). Different PyC detection and quantification methods cover a different part of this continuum (Hammes et al., 2007; Hammes and Abiven, 2013). Across methods, PyC represents on average 15 % of soil organic carbon (SOC) (Reisser et al., 2016).

The stable chemical structure of PyC allows most of it to persist for longer in soils relative to non-PyC, however the range of PyC residence time estimates vary widely (Bird et al., 2015). Large scale modelling of PyC fluxes (Bowring et al., 2022) indicates a persistence in the order of 5000 years. Methods based on radiocarbon dating of charcoal fragments (Carcaillet, 2001; Hajdas et al., 2007; Gavin et al., 2023; Liang et al., 2008) or of the PyC fraction of soil organic carbon (Butnor et al., 2017; Schiedung et al., 2024) have shown that PyC can remain in soils for millennia and up to more than 10 000 years. However, much younger ^{14}C ages have also been reported (Kane et al., 2010; Ohlson et al., 2009; Krull et al., 2006; Czimczik et al., 2005; Bellè, 2023). These could be due to the rejuvenation of the PyC pool by recent fires or high fire frequency (Kane et al., 2010; Bellè, 2023), or to methods of PyC isolation and detection that may exclude older PyC (Ohlson et al., 2009; Krull et al., 2006), but a centennial PyC turnover rate cannot be excluded. Extrapolations based on laboratory or field incubations (see Azzi et al., 2024, for an extensive list) give estimates in the range of decades to millennia, with significant methodological drawbacks related to model choice and non-representative conditions and time scales (Leng et al., 2019; Azzi et al., 2024). Long term field studies with re-sampling (Hammes et al., 2008b; Lutfalla et al., 2017) or based on chronosequences (Alexis et al., 2012; Lehdorff et al., 2014; Selvalakshmi et al., 2018; Nguyen et al., 2009; Cheng et al., 2008) tend to yield shorter estimates, from decades to centuries. However, PyC residence times estimated from the measurement of PyC losses at the plot scale do not provide true estimates of residence time at the landscape scale. In addition to min-

eralisation, which represents a net loss of PyC at all scales, PyC may also disappear from the monitored plot via fragmentation to smaller particles and decomposition to less condensed, more oxidised molecules that can evade PyC quantification (Krull et al., 2006), downward vertical transport out of the soil profile, and lateral transport (erosion by surface run-off and leaching by subsurface flow) to nearby or further away locations (Abney and Berhe, 2018). PyC that has been displaced or transformed to another form of soil carbon still resides in the landscape, although its fate may be altered (i.e. it may become more or less subject to mineralisation depending on its chemistry and the environmental conditions at its new location).

Downward vertical transport of PyC has been observed in controlled experiments (Schiedung et al., 2020; Santos et al., 2022; Hilscher and Knicker, 2011) and in the field under different pedo-climatic contexts (Maestrini et al., 2014; Singh et al., 2014, 2015; Schiedung et al., 2023; Dai et al., 2005; Alexis et al., 2012; Bellè, 2023; Bonhage et al., 2022; Vasilyeva et al., 2011). In addition, many studies found that the proportion of PyC in SOC is greater in the subsoil relative to the topsoil, which is attributed in part to transport of PyC to depth (Velasco-Molina et al., 2016; Rodionov et al., 2006; Soucémarianadin et al., 2019). The processes by which PyC can be redistributed vertically in the soil profile have been reviewed by Hobley (2019). Leaching in dissolved form is usually limited (Abiven et al., 2011; Maestrini et al., 2014; Major et al., 2010; Hilscher and Knicker, 2011; Schiedung et al., 2020; Abney et al., 2024) but transport as colloids and larger particles may be significant, in particular when soil texture is coarse, soil structure is loose, sufficient water is present and/or mixing processes (bioturbation, peloturbation, cryoturbation) are intense (Bellè, 2023; Lehdorff et al., 2016; Schiedung et al., 2020; Rodionov et al., 2006; Leifeld et al., 2007). Whether PyC will be retained in the subsoil or transported further down to the altered bedrock and groundwaters depends on its interaction with the minerals and its stabilisation in the subsoil (Brodowski et al., 2007; Guggenberger et al., 2008; Vasilyeva et al., 2011; Santos et al., 2017).

Recent reviews have highlighted the role of erosion in redistributing carbon (C) at the slope, watershed, and global scale (Berhe et al., 2018; Van Oost and Six, 2023; Zheng et al., 2025). These processes are likely exacerbated for PyC (Abney and Berhe, 2018). Erosion rates often increase after fires due to removal of the vegetation and litter layer, and in some cases, increased soil hydrophobicity (Belcher, 2013; Girona-García et al., 2024). PyC has a low density which favours its detachment and transport, even in apparently flat landscapes (Pyle et al., 2017). Manipulative rainfall experiments showed that more than 50 % of deposited PyC can be eroded after a single rain event (Bellè et al., 2021; Rumpel et al., 2009). PyC was enriched in eroded sediments relative to the mineral soil at the plot scale (about 1 m^2 , Rumpel et al., 2006b) and at the micro catchment scale (Chaplot et al., 2005; Cotrufo et al., 2016b; Abney et al., 2019) indepen-

dent of the slope, but depleted at the larger catchment scale (Chaplot et al., 2005) showing the existence of deposition and storage areas in the landscape. These areas may include alluvial deposits at riverbanks (Cotrufo et al., 2016b), flat areas along the slope or at the toeslope (Abney et al., 2017), patches presenting high surface roughness favouring infiltration of runoff (Boot et al., 2015), where unburned vegetation may retain the charcoal particles (Galanter et al., 2018), or where moderate to low burn severity did not result in increased erosion rates (Abney et al., 2019). Indeed, slope and burn severity seem to have an interactive effect in controlling PyC redistribution by erosion post-fire (McGuire et al., 2021; Galanter et al., 2018). Most of the above-cited studies took place within 4 years post fire. On the decadal time scale, Güereña et al. (2015) showed erosion-related redistribution of mineral soil PyC 10 years after deforestation associated with on site charcoal production in equatorial humid region, but the absence of a flat toeslope (convex toposequence) may have prevented the accumulation of PyC near the stream 62 years after deforestation. Abney et al. (2017) showed that PyC eroded from the slope and deposited on the surface of a riparian area 1 year post-fire had been either decomposed, buried, or remobilised (laterally or vertically, by erosion and/or leaching) 10 years post-fire. Only a few chronosequences of mineral soil PyC have investigated geomorphic factors and landscape positions on longer time scales (Czimeczik et al., 2005; Sass and Kloss, 2015). Centennial to millennial time scales are relevant as they are commensurate with the estimates of PyC residence time in soils. Studies on large time- and spatial-scales are needed to capture the balance of erosion processes on the C cycle (Van Oost and Six, 2023). Since PyC dynamics is often related to its interactions with the hydrological cycle (Masiello and Berhe, 2020), the watershed seems to be a relevant scale to trace the fate of PyC. PyC dynamics in the landscape have been studied in temperate to subtropical forests, often with monsoonal precipitation regime and/or in mountainous regions (Galanter et al., 2018; Boot et al., 2015; Matosziuk et al., 2020; McGuire et al., 2021; Abney et al., 2017), in grasslands (Dai et al., 2005; Rodionov et al., 2006), in boreal forests (Kane et al., 2007; Guggenberger et al., 2008) or in ecosystems actively managed by fire (Selvalakshmi et al., 2018; Rumpel et al., 2006a; Alexis et al., 2012; Nicolay et al., 2024, see Fig. S1 in the Supplement for a map of existing studies). Data on other climate zones, geomorphic settings, land-use and vegetation, and pyromes are needed to better understand the drivers of PyC redistribution.

In this work, we investigated whether the vertical and horizontal distributions of soil PyC differed from that of SOC in a shallow sloping, agricultural watershed under oceanic temperate climate. Cropland has dominated the catchment for at least 150 years, implying no or little recent PyC inputs and allowing us to study the resultant, long-term PyC distribution in the landscape. We hypothesised that, even on shallow slopes, PyC produced during past fires was trans-

ported downslope by erosion due to its light nature and enhanced post-fire erosion, and deposited at the toeslope where it was protected from further decomposition by burial and/or unfavourable conditions for microbial activity due to frequent water-logging. Owing to the long term stability of PyC, these processes would result, to this day, in higher stocks and higher radiocarbon ages at this position compared to upslope (1). We also made the hypothesis that vertical downward transport of PyC in the soil profile, combined with its high stability, would result in increasing proportions of PyC in SOC with increasing soil depth (2a), and that PyC would be older than SOC at all depths (2b). We measured the concentrations and stocks of PyC down to 60 cm along three toposequences through the convex–concave of the slope. We used two methods of PyC quantification (chemo-thermal oxidation – CTO and hydropyrolysis – HyPy) that cover the intermediate to highly condensed part of the PyC continuum, and also measured the radiocarbon (^{14}C) values in both total SOC and the PyC fraction.

2 Material and methods

2.1 Study site

Our study site is located in a catchment of 120 ha in the region of Brittany in the west of France (Fig. 1). Over the period 1994–2020, the mean annual temperature was 12 °C (January 6.1 °C–July 18.5 °C) and the mean annual precipitation was 1043.7 mm, well distributed over the year with only slightly wetter autumn–winter and drier spring–summer. The Kervidy-Naizin catchment is part of the ORE (Observatoire de Recherche en Environnement) AgrHys and the critical zone observatories network OZCAR (Observatoires de la Zone Critique: Application et Recherche). It has been studied since the 1970's and was chosen to represent shallow sloping catchments on sedimentary basement typical in the Armorican Massif, that have experienced agricultural intensification over the last 70 years. It feeds the Coët-Dan, an intermittent stream that often dries out in summer. Slopes are shallow (median 3.0 %, first quartile (Q1) 1.7 % – third quartile (Q3) 4.1 %) and present a convex–concave profile. Soils are developed on loess derived locally during the Last Glacial Period (~115 000 to 11 700 BP, maximal loess deposition in the Northern European Loess Belt around 21 800 BP, Bosq et al., 2023, see also Appendix A1), overlaying weathered siltstone and sandstone, except in the immediate vicinity of the stream where alluvium is the parent material (Walter and Curmi, 1998). The soil texture is silt loam (FAO soil triangle) and soils are acidic ($\text{pH}_{\text{H}_2\text{O}}$ 3.7–6.9). Soils are organised along the convex–concave slope in a gradient of increasing soil evolution towards the stream, typically following the sequence (from summit to toeslope): Cambisol – Haplic Luvisol – Stagnic (solimovic) Luvisol – Stagnosol – Stagnic Fluvisols (transposed to the World Reference Base IUSS Work-

ing Group WRB (2022) from Walter and Curmi, 1998). This spatial distribution of soils (sequence and extent of each soil type) is representative of catchments with similar topography and geology in Brittany (Chaplot et al., 2003). The up-hill domain is well drained with mainly vertical water flow through the soil volume. Going down slope, the winter water table comes closer to the soil surface, the water flow develops a small horizontal component, and in the soils with stagnic properties, a perched water table above a dense albic horizon can sometimes be observed (Walter and Curmi, 1998). Surface flow and erosion are present in the current agricultural land-use but of limited extent and intensity. There is little export of material out of the catchment (Cros-Cayot, 1996).

Because PyC has a long residence time in soil, we also gathered information on land-use change over several millennia (see also Appendix A1). Little is known about the vegetation in Brittany in the periglacial landscape at the end of the last glacial period (11 700 BP) and how it transitioned to the Atlantic oak forest that was likely predominant in Brittany around 7500 BP, when it started to be opened by early human populations for pasture. It is estimated that most of the primary Atlantic forest had been cleared by the end of the Iron Age (Gaudin et al., 2014). Heath and cropland increased globally from the Bronze Age to the end of the Middle Ages (Gaudin et al., 2014). Fire regimes in these successive landscapes are poorly known (Appendix A1). There are no nowadays equivalent to the Atlantic forest, and although a little more is known about managed fire in heathland (Hobbs and Gimingham, 1987) there are only two values of PyC production for heath in the extensive database from Jones et al. (2019). Fire could also have been used to re-open land in a context where arable plots shifted often (Tonnerre, 1992). The transition from a landscape dominated by heath and secondary growth woodland to one where cropland are predominant occurred towards the end of the 19th century (Astill and Davies, 1997). Unintentional fires are virtually inexistent in European croplands under temperate, oceanic climate (Forrest et al., 2024). Agricultural practices such as spreading of plant ashes (Antoine, 2000) or on-site stubble burning (banned from most of Western Europe in the 1980's) could have contributed relatively labile (low temperature) PyC to croplands (see also Appendix A2). The study site is not located downwind of any major road nor, to the best of our knowledge, any current or historical industry that could have been a source of a significant amount of fossil fuel derived PyC (soot). We assume that the current PyC distribution in the landscape integrates the effects of the changes in land use and fire regime since the deposition of the loess that forms the soil parent material (i.e. in the last $\sim 20\,000$ years, see Appendix A1) and interpret our data accordingly.

Most of the catchment area has been under cultivation or pasture since before 1833, except for a small wood at the south-west border. Plot limits did not vary between 1833 and 1952 and many were delimited by hedgerows and/or paths (see Appendix A2 for details on how we derived this infor-

mation). In the 1970's, plots were pooled together to form larger fields, some paths were abandoned and hedgerows uprooted. This affected especially the area where transect T1 and T3 are located whereas the vicinity of T2 seemed already devoid of hedgerows by 1952 (see Fig. 1 for transects location).

2.2 Study design and sampling

We sampled along three topographic transects (toposequences) on north-east (T1), south to south-east (T2) and south-west (T3) facing slopes, at upslope (6 sites – 2 per transect), mid-slope (4 sites – 2 at T1, 1 at T2 and T3) and downslope (6 sites – 2 per transect) positions (Fig. 1). Upslope sites were located at the summit (T1) or at the end of a long, shallow ($< 2\%$) slope running from the summit, just before or at the shoulder (convex portion, T2 and T3). Mid-slope sites were located along the steepest sloping part of each transect (backslope), whereas downslope sites were located at the foot- and toeslope where the slope becomes shallow again (concave portion, see the elevation profiles along the three transects in Appendix Fig. A1). All sites were located in croplands except for the uppermost and lowermost sites of transect 1 which were in a woodland and a cultivated grassland, respectively. Existing sampling sites from previous field campaigns in the study area were relocated based on their GPS coordinates. The GPS coordinates of new sampling sites were recorded. At each sampling site, one lined core was taken using a hydraulic corer and three hand-held auger cores were made about 1.5 m away from each other around the hydraulic core. The hand auger cores were sampled every 10 cm and composited together. Samples and cores were refrigerated within 48 h and kept at $4\text{ }^{\circ}\text{C}$ until analysis. A field soil description was established based on the hand auger core. Selected soil characteristics are summarised in Table 1.

2.3 Soil characteristics

2.3.1 Sample preparation

The soils were dried at $40\text{ }^{\circ}\text{C}$ for a minimum of 48 h and sieved at 2 mm. A 20 g sub-sample of the fine soil was finely ground ($\leq 200\text{ }\mu\text{m}$) using a ball mill ($1\text{ min at }25\text{ s}^{-1}$). The finely ground samples were used for all carbon analyses.

2.3.2 Physical characteristics

Particle size distribution was determined by the Laboratoire d'analyse des sols (LAS, INRAE) according to NF X31 107. In short, the sample was exposed to heated H_2O_2 to break down organic matter prior to particle size evaluation. Coarse sand ($> 200\text{ }\mu\text{m}$) was separated by dry sieving. The remaining material was dispersed in solution and fine fractions ($< 50\text{ }\mu\text{m}$) were determined using Robinson's pipetting

Table 1. Summary statistics of soil characteristics by slope position and layer. CEC_{Co} – cation exchange capacity determined using a cobalthexamine chloride solution, C_{fine} – concentration of fine soil, Fe_{DCB} – Iron extracted by the dithionate-citrate-bicarbonate solution, Fe_{oxalate} – Iron extracted by the oxalate/oxalic acid buffer solution, Subsoil – > 30 cm depth (50–60 cm for all soils except for the Solimovic Cambisol where the subsoil was sampled at 70–80 cm, and an additional sample at 30–40 cm to account for the strongly differentiated horizons in the Stagnic Luvisols and Stagnosol), min. – minimum, max. – maximum.

Slope position	Layer	n	SOC		pH _{H2O}		CEC _{Co}		C _{fine}					
			(g C kg ⁻¹ soil) median min. max.	median min. max.	(cmol + kg ⁻¹) median min. max.	(g cm ⁻³) median min. max.								
Summit and shoulder	0–10 cm	6	22.3	17.0	114.1	6.2	3.7	6.5	7.3	6.2	9	0.82	0.64	0.91
	Subsoil	6	5.1	3.0	13.1	6.1	4.6	6.6	3.4	2.7	4.1	1.11	0.91	1.17
Midslope	0–10 cm	5	21.6	18.3	43.3	6.4	5.4	6.7	7.4	6.9	8.5	0.74	0.65	0.91
	Subsoil	5	3.2	2.4	6.1	6.4	5.7	6.8	3.6	2.1	4.9	1.11	0.91	1.31
Foot- and toeslope	0–10 cm	6	25.2	21.5	75.6	5.9	5.3	6.9	8.6	6.8	11.8	0.82	0.64	0.91
	Subsoil	10	5.7	3.3	18.1	6.3	5.2	6.9	3	2	8.5	1.05	0.88	1.25
Slope position	Layer	n	Clay		Sand		Fe _{oxalate}		Fe _{DCB}					
			(%) median min. max.	(%) median min. max.	(%) median min. max.	(g C kg ⁻¹ soil) median min. max.	(g C kg ⁻¹ soil) median min. max.							
Summit and shoulder	0–10 cm	6	14	13	23	13	12	14	4.5	3.6	5.4	21.3	16.5	23.1
	Subsoil	6	16	15	18	17	14	21	3.0	2.6	3.7	22.7	19.8	28.9
Midslope	0–10 cm	5	14	13	22	14	12	15	4.7	4.5	5.9	19.2	18.4	23.3
	Subsoil	5	18	14	23	16	13	24	2.1	1.9	2.3	27.6	21.5	56.2
Foot- and toeslope	0–10 cm	6	18	13	31	19	11	23	3.9	1.4	5.6	6.8	5.2	18.9
	Subsoil	10	24	14	26	25	13	31	2.0	0.8	6.1	27.3	17.9	54.9




Figure 1. (a) Location of the three transects (T1 to T3) and individual sampling sites within the study area. Shading represents topography. Sites are colored by soil type and labelled according to their position along the toposequence (Up = summit, flat and shoulder, Mid = midslope, Down = footslope and toeslope). (b) Organisation of the soil horizons along the lower part of a convex–concave toposequence. Modified from Cheverry (1998). Inset: location of the study area in France.

method. Fine sand (between 50 and 200 μm) was separated by wet sieving under water flow.

The concentration of fine soil (< 2 mm) was measured for a representative subset of sites using the hydraulic cores. Compaction during coring was evaluated by comparing the depth of soil in the cores to the recorded coring depth. Compaction was considered homogeneous when < 5 % and increasing with depth otherwise. Depth increments equivalent to 10 cm of uncompacted soil were delimited on the outside of the transparent core liner before cutting it open longitudinally and slices were cut out with a knife following the delimitations. The soil was dried at 105 °C for 24 h and weighed (m_{tot}). Coarse elements > 2 mm were separated from the fine soil by wet sieving, dried again and their weight was recorded (m_{rocks}). The concentration of fine soil (C_{fine} , in g cm^{-3}) was calculated according to Eq. (1). Note that this is not the bulk density of fine soil as we did not correct for the volume of coarse fragments, but it is enough to calculate the correct carbon stocks as per Eqs. (8) and (9) of Poeplau et al. (2017).

$$C_{\text{fine}} = \frac{(m_{\text{tot}} - m_{\text{rocks}})}{V_{\text{eq_slice}}}$$

$$\text{with } V_{\text{eq_slice}} = \pi \cdot (7.54)^2 \cdot 10 \quad (1)$$

C_{fine} was estimated for missing sites using a linear model fitted on the measured values with depth, horizon and transect as predictors ($R^2_{\text{adj}} = 0.47$, residual standard error = 0.15 g cm^{-3} , $p < 0.001$).

2.3.3 pH and cation exchange capacity (CEC)

pH and CEC were determined by the LAS (INRAE) according to NF EN ISO 10390 and ISO 23470, respectively. pH was measured in water and in KCl (1 mol L^{-1}) at a 1 : 5 (v/v)

soil to solution ratio. CEC was determined based on the exchange of cobaltihexamine ions at soil pH. 50 mL of cobaltihexamine solution at 50 mmol L^{-1} was mixed with 2.5 g of soil. The concentration of cobaltihexamine remaining in solution after exchange was measured by spectrophotometry at 475 nm and CEC was calculated by difference with the initial concentration.

2.3.4 Iron and aluminium oxides and total elemental composition

Iron (Fe) and aluminium (Al) were extracted at the LAS (INRAE) by the Tamm method in the dark (also called oxalate extraction, described in Courchesne and Turmel, 2008), the Mehra–Jackson method (also called dithionate–citrate–bicarbonate (DCB) extraction Mehra and Jackson, 1960) and hydrofluoric acid (HF) digestion (NF ISO 14869-1) and measured by ICP-AES. The Tamm method extracts iron in organo-metallic complexes and amorphous or poorly crystalline oxides ($\text{Fe}_{\text{oxalate}}$). It is conducted at pH 3, 20 °C and in the dark to avoid iron reduction by UV radiations. 50 mL of oxalate buffer solution was mixed with 1.25 g of finely ground soil during 4 h. The Mehra–Jackson method extracts iron and aluminium in oxides and oxyhydroxides (amorphous and crystalline, Fe_{DCB}). 0.5 g of finely ground soil was exposed to 25 mL of sodium tri-citrate (0.267 mol L^{-1}) and sodium bicarbonate (0.111 mol L^{-1}) and 1.5 mL of sodium dithionite (200 g L^{-1}) for 30 min at 80 °C. The amount of iron in crystalline oxides was calculated by difference: $\text{Fe}_{\text{crystalline}} = \text{Fe}_{\text{DCB}} - \text{Fe}_{\text{oxalate}}$. The HF digestion breaks down all silicate minerals and gives access to the total iron (Fe_{tot}) and aluminium (Al_{tot}) content of the soil. Other elements measured by ICP-AES on the HF extract include calcium (Ca_{tot}), potassium (K_{tot}), magnesium (Mg_{tot}), man-

ganese (Mn_{tot}), sodium (Na_{tot}) and phosphorus (P). From these data we calculated the weathering index (WI, Eq. 2) and the chemical index of alteration (CIA, Eq. 3). The WI is the ratio of $\text{Fe}_{\text{crystalline}}$ to total iron. Iron forms in soils are strongly related to pedogenetic processes such as alteration of crystalline iron oxides present in the parent rock, clay formation and breakdown and secondary precipitation of more or less crystalline iron oxides. The CIA was proposed by Nesbitt and Young (1982) as an indicator of the alteration of feldspars from the parent rock (low CIA) to clay minerals (high CIA), during which the mobile elements (Na, Ca, K) are lost whereas Al is retained. Note that K fertilisation and liming will modify CIA in the topsoil independent of weathering.

$$\text{WI} = \frac{\text{Fe}_{\text{DCB}} - \text{Fe}_{\text{oxalate}}}{\text{Fe}_{\text{tot}}} \quad (2)$$

$$\begin{aligned} \text{CIA} &= \frac{\text{Al}_2\text{O}_3}{\text{Al}_2\text{O}_3 + \text{Na}_2\text{O} + \text{K}_2\text{O} + \text{CaO}^*} \\ &= \frac{\frac{\text{Al}_{\text{tot}}}{2M(\text{Al})}}{\frac{\text{Al}_{\text{tot}}}{2M(\text{Al})} + \frac{\text{Na}_{\text{tot}}}{2M(\text{Na})} + \frac{\text{K}_{\text{tot}}}{2M(\text{K})} + \left(\frac{\text{Ca}_{\text{tot}}}{M(\text{Ca})} - \frac{\text{CaCO}_3}{M(\text{CaCO}_3)} \cdot 10 \right)} \end{aligned} \quad (3)$$

where X_{tot} are the concentrations of the elements in weight % (or g X/100 g soil), $M(X)$ is the atomic mass of the respective elements in g mol^{-1} , the factors 2 are to convert from elemental to oxide molar content and CaCO_3 is the carbonate content in g kg^{-1} soil.

2.3.5 Soil organic carbon and pyrogenic carbon

SOC content was measured by dry combustion – cavity ring down spectroscopy (Picarro G2101-i Isotopic CO_2 and combustion module Costech) calibrated against an in-house standard. Each sample was measured in triplicate. The soils contained no carbonates so total carbon was considered as organic carbon.

PyC content was determined using the chemo-thermal oxidation (CTO-375) method (Gustafsson et al., 1997; Caria et al., 2011) and the hydrogen pyrolysis (or hydrolysis, HyPy) method (Meredith et al., 2012). The CTO-375 method removes non-pyrogenic organic matter and the least condensed fraction of the PyC continuum and targets soot-like PyC (Hammes et al., 2007; Gustafsson et al., 2001; Elmquist et al., 2004). HyPy isolates PyC of > 7 fused aromatic rings (Meredith et al., 2012; Wurster et al., 2013), which covers a wider range of the PyC continuum than CTO-375 (Hammes and Abiven, 2013).

Each sample was prepared and measured in duplicate by CTO-375. 40 mg of finely ground sample were weighed in two silver capsules (three if $\text{SOC} < 0.2\%$). The samples were placed in a stainless steel plate and heated in a muffle furnace to 350°C at $10^\circ\text{C min}^{-1}$ then to 375°C at 1°C min^{-1} and kept at this temperature for 24 h. This heating program

has been found to prevent in-situ formation of PyC during the procedure (M. Schiedung, personal communication, 2022). The ventilation holes in the oven door were kept open throughout to ensure sufficient oxygen supply for complete combustion of thermally labile organic matter. After cooling to room temperature, the samples were acidified with $20\ \mu\text{L}$ of 1% (v/v) hydrochloric acid (HCl) and exposed to concentrated HCl vapours in a bell jar for 8 h. Samples were then dried successively in a desiccator containing silica beads and under vacuum for 24 h and in a vacuum oven at 40°C for 3 d. The two or three silver capsules per sample were then composited in a tin capsule and analysed for C content (noted PyC_{CTO}) as described above for SOC. An in-house standard was submitted to the same procedure with every batch of samples to check for analytical deviation and calculate the coefficient of variation of the PyC_{CTO} content ($\text{CV} = 11\%$).

Additionally, a subset of samples were analysed by HyPy. We selected the 0–10 cm (top of the A horizon) and 50–60 cm (base of the lowermost B horizon, transition towards the C horizon) depth increments for all profiles except for the Solimovic Cambisol where we selected the base of the solimovic material at 40–50 cm and the base of the structural B horizon at 70–80 cm. To represent the eluvial E horizon an additional intermediate depth increment (20–30 or 30–40 cm) was analysed for mid- and downslope sites presenting this horizon. The finely ground samples were mixed with ammonium dioxodithiomolybdate (5% m/m) and suspended in a 20% (v/v) water-methanol solution to dissolve the catalyst. The solution was then evaporated to dryness at 80°C . This allowed coating of the catalyst on the soil particles for higher contact surface. Samples were then weighed into pre-combusted quartz inserts and were pyrolysed at 550°C under a hydrogen pressure of 150 bar and sweep gas flow of $5\ \text{L min}^{-1}$ (ATP). Full details of the pyrolysis programme, including temperature ramp and hold rates, can be found in Meredith et al. (2012). The carbon content of samples before and after hydrolysis was measured by elemental analysis (EA). The PyC measured via HyPy is expressed as the proportion of SOC that is made up of PyC (noted PyC_{HyPy} (%SOC)). PyC_{HyPy} content was obtained by multiplying the PyC_{HyPy} proportion by SOC content.

2.3.6 PyC and SOC radiocarbon dating

Among the samples selected for HyPy, we further restrained the radiocarbon analysis to upslope and downslope sites of each transect.

For ^{14}C measurement of the PyC fraction, samples were prepared following the HyPy procedure as described above. After hydrolysis, the quartz crucible containing PyC was converted to CO_2 by combustion in a sealed quartz tube, as described in Ascough et al. (2024). The sample CO_2 was cryogenically purified and then reduced to graphite by sequential reaction with Zn and Fe. The sample preparation was carried out at the Environmental Radiocarbon Labo-

ratory (Scottish Universities Environmental Research Center, East Kilbride, UK). The $^{14}\text{C}/^{12}\text{C}$ ratio of the graphitised samples was measured by accelerator mass spectrometry (AMS) at the Keck Carbon Cycle AMS Facility (University of California, Irvine, USA). Stable carbon isotope ratios ($^{12}\text{C}/^{13}\text{C}$) were measured online on the AMS to correct the measured radiocarbon values for kinetic fractionation effects.

For the bulk ^{14}C analysis, the homogenised soils were fumigated with 37 % HCl for 72 h at 60 °C to remove all inorganic carbon. After fumigation, the excess acid was neutralised by placing the samples in a closed bell jar with NaOH pellets at 60 °C for a minimum of 48 h. The samples were subsequently wrapped in tin boats and analysed using a coupled elemental analyser-accelerator mass spectrometer (EA-AMS) system (vario MICRO cube, Elementar; Mini Carbon Dating System MICADAS, Switzerland) described in Synal et al. (2007). The ^{14}C analysis was carried out at the Laboratory of Ion Beam Physics at the Swiss Federal Institute of Technology (ETH), Zürich, Switzerland.

Measured $^{14}\text{C}/^{12}\text{C}$ ratios are reported as $F^{14}\text{C}$, as described in Reimer et al. (2004). We did not calculate calibrated ages for PyC ^{14}C data as PyC_{HyPy} results from the accumulation of fires over time and not a single event that can be dated. We also did not attempt to calculate turnover times for either carbon pool. SOC is not an homogeneous pool with a single turnover time, and even assuming PyC_{HyPy} is a homogeneous pool, the turnover time calculation requires either a steady state assumption (inputs = outputs, unlikely for PyC produced from sparse fire events), known input rates, or several time points (Torn et al., 2009), neither of which is met.

2.4 Data processing and statistical analysis

Slope position and soil type were determined based on cartographic information from SAS laboratory visualised in QGIS v3.34.8 and confirmed by field observations and laboratory analysis. This led us to modify the grouping of sites from the original design. In particular, we singled out the summit site of transect 1 (Hyperdystric Cambisol, Fig. 1), located in a small woodland and strongly acidic (pH 3.7–4.6), from the other upslope sites (Haplic Cambisols, $n = 5$) under cropland since at least the first half of the 19th century and located either at intermediate flat position along a longer slope or at the slope shoulder. We also singled out the lowest downslope site of transect 3, the only soil showing strong colluvial features (Solimovic Cambisol), and we further subdivided the remaining downslope positions into footslope sites ($n = 3$), at the transition from Cambisol to Luvisol and toeslope sites ($n = 2$) in the Stagnic Luvisol and Stagnosol domain. Mid-slope Cambisols were not subdivided ($n = 4$). Elevation above sea-level was determined in QGIS based on the GPS coordinates of the sites and the 1 m resolution digital elevation model (DEM) RGE ALTI[®] (IGN, 2013). For each site, we calculated its altitude relative to the site closest to the stream in the respective transect, as a proxy for the

slope length from that site to the toeslope. Slope was calculated in QGIS from the 25 m resolution DEM BD ALTI[®] (IGN, 2015). We used this lower resolution DEM to calculate slope to smooth out the effect of small, localised topographic features. The overall slope were similar between the 3 transects and local slope at the sampling sites ranged from 0 % to 3 %.

Carbon stocks were calculated for each 10 cm depth increment and summed to 60 cm (total stocks), the deepest common depth to all sites, which corresponds to the apparition of the C horizon (altered parent material) at the shallowest profiles (the deepest profiles extended deeper than 90 cm). When available, analytical error was propagated to the final results. In particular, for total stocks, the error was propagated from each depth increment to the sum then to the mean. The degrees of freedom of the variance obtained from error propagation seemed to be erroneously estimated by the Welch–Satterthwaite equation for our small sample size, as observed by O’Hagan et al. (2021), which prevented us from constructing satisfactory confidence intervals for these quantities. Thus we chose to represent the standard deviation, unless specified otherwise. The effect of slope position (original design) was assessed using the Kruskal–Wallis test, a non parametric alternative to ANOVA, after excluding the only solimovic site. Slope positions were compared by the Wilcoxon–Mann–Whitney test and p -values were adjusted for multiple comparisons with the Bonferroni–Holm correction. Some soil groups from the revised design did not contain enough observations to allow meaningful statistical analysis. Trends were assessed visually, taking into account the propagated error. Linear regression was fitted to the concentration and stock data as a function of altitude relative to the toeslope. Diagnostic plots were examined for strong deviations to the normality and homoscedasticity assumptions and identification of high leverage outliers. Correlations between PyC content, $F^{14}\text{C}$ and other soil properties were assessed by Kendall’s rank correlation coefficient (τ). Significance was assessed at the 95 % confidence level. Analytical and field replicates were averaged as necessary. All data processing, statistical analysis and figure production was done with R version 4.4.0 (R Core Team, 2021) in Rstudio version 2024.04.1+748 (Rstudio, 2021), using the packages tidyverse (Wickham et al., 2019) and corrplot (Wei and Simko, 2021). All data used to produce the figures presented below can be found in Lebrun Thauront et al. (2025).

3 Results

3.1 PyC in the soil profile

The evolution of PyC_{CTO} and PyC_{HyPy} proportion in SOC with depth for the six soil groups is represented in Fig. 2. For all soil groups but the Hyperdystric Cambisol under woodland at the summit, the proportion of PyC_{CTO} in SOC re-

mains the same in the first 20 to 30 cm due to homogenisation of the soil material by ploughing. Below 30 cm, an increasing proportion of SOC is made up by PyC_{CTO} as depth increases – between $3.4 \pm 0.23\%$ and $3.83 \pm 0.26\%$ at 30–40 cm and between $6.7 \pm 0.40\%$ and $15.0 \pm 2.6\%$ at 70–80 cm – except for the Solimovic Cambisol at the hill foot which shows a nearly flat profile down to 60 cm depth. The Stagnic Albic Luvisol and Albic Fluvisol Stagnosol at the toeslope presented a peak at 20–30 cm (albic horizon) due to a sharper drop in SOC than PyC. Although PyC proportion in SOC increases, PyC concentration decreases with depth in all profiles (Fig. S2).

The PyC_{HyPy} profiles show the same trends as PyC_{CTO} profiles, with the proportion of PyC_{HyPy} in SOC higher than the proportion of PyC_{CTO} in SOC by a median of 4.3 %, consistent with the commonly held assumption that the HyPy method measures a wider range of the PyC continuum than the CTO-375 method (Hammes and Abiven, 2013).

3.2 PyC along the toposequence

The amount of PyC in the first 10 cm of the soil, from the toeslope upward, is shown in Fig. 3. PyC_{CTO} (panel a) tended to be higher at upslope and downslope positions (median = 0.5 and 0.6 g C kg^{-1} soil, interquartile range (IQR) = 0.18 and 0.15 g C kg^{-1} soil, respectively) and lower at midslope (median = 0.4 and IQR = 0.04 g C kg^{-1} soil) but these differences were not significant ($\chi^2 = 5.3$, $p = 0.072$). PyC_{CTO} content in the topsoil decreased significantly with increasing slope ($R^2 = 0.29$, $p = 0.02$, Fig. S3). These tendencies remained visible in the subsoil (50–60 cm depth) although the variability was greater (not shown). The PyC_{HyPy} content (panel b) and proportion in SOC (not shown) linearly decreased ($R^2 = 0.46$, $p = 0.01$ and $R^2 = 0.51$, $p = 0.01$, respectively) from the summit to the toeslope in the topsoil, but not in the subsoil (not shown). Slope position was a significant factor for PyC_{HyPy} content ($\chi^2 = 7.2$, $p = 0.027$), with significantly more PyC_{HyPy} in the topsoil at upslope positions (median = 1.6, IQR = 0.22 g C kg^{-1} soil) than at downslope positions (median = 1.10, IQR = 0.40 g C kg^{-1} soil). Contrary to PyC_{CTO} , PyC_{HyPy} content in the topsoil did not vary systematically with slope (Fig. S3).

3.3 PyC stocks

The stocks of PyC_{CTO} for each soil group, in the topsoil (0–30 cm), subsoil (30–60 cm), and over the entire soil profile (0–60 cm), is shown in Fig. 4. Contrary to our hypothesis, slope position was not a significant factor explaining soil PyC stocks in this watershed. On average PyC_{CTO} amounted to $2.5 \pm 0.22 \text{ t ha}^{-1}$. PyC_{CTO} stocks were only slightly lower in a Hyperdystric Cambisol under woodland at the summit ($1.9 \pm 0.22 \text{ t ha}^{-1}$) and higher in a Solimovic Cambisol at the footslope ($3.3 \pm 0.26 \text{ t ha}^{-1}$). The proportion of PyC_{CTO} in total SOC stock varied little, from $1.5 \pm 0.35\%$

to $3.1 \pm 0.48\%$. PyC in the subsoil represented between 37 % and 51 % of the PyC_{CTO} stock in the hyperdystric Cambisol under woodland at the summit and the Solimovic Cambisol at the footslope, respectively, and on average 44 % for the other soil groups. SOC stocks amounted to $100 \pm 10 \text{ t ha}^{-1}$ on average (min = 68, max = 141 t ha^{-1}), with 12 % to 49 % of the stock below 30 cm (Fig. S4).

3.4 Radiocarbon ages of PyC and SOC

There was no significant difference in $F^{14}\text{C}$ with slope position for either SOC or the PyC_{HyPy} fraction in the topsoil (Fig. 5, panel a) and the subsoil (not shown). PyC_{HyPy} in the topsoil was significantly older than SOC independent of slope position (Fig. 5a, $F^{14}\text{C}$ range 0.72–0.83 and 0.99–1.14, respectively, which corresponds to 2630–1530 ^{14}C years BP and 50 BP – modern), and the difference increased with depth (Table S1 in the Supplement). In addition, both the PyC_{HyPy} fraction (Fig. 5b) and the total SOC (Table S1) were systematically older in the subsoil relative to the topsoil ($F^{14}\text{C}$ PyC_{HyPy} subsoil median = 0.60, IQR = 0.039, topsoil median = 0.79, IQR = 0.045) except for one profile at the footslope that showed a reverse “stratigraphy”. We found no trend in PyC_{HyPy} ^{14}C values with altitude relative to the toeslope and/or soil type except for the Solimovic Cambisol which had a young PyC_{HyPy} over the whole profile ($F^{14}\text{C}$ 0.73–0.81). All ^{14}C data can be found in Table S1.

3.5 Correlations between soil characteristics and PyC

Both PyC_{CTO} and PyC_{HyPy} were positively correlated with SOC ($\tau = 0.57$, Fig. 6). This correlation was weak and insignificant in the topsoil (0–10 cm) and of intermediate strength at 50–60 cm depth ($\tau = 0.47$ and 0.45 for PyC_{CTO} and PyC_{HyPy} , respectively). On the contrary, the proportion of PyC_{CTO} and PyC_{HyPy} in SOC was strongly negatively correlated with SOC ($\tau = -0.78$ and -0.62 , respectively) and the C : N ratio ($\tau = -0.66$ and -0.65 , respectively) and positively with $\delta^{13}\text{C}$ ($\tau = 0.49$ and 0.67, respectively). Overall, with the exception of CEC_{Co} and WI, correlations between PyC and variables reflecting soil chemistry (pH, CIA) and soil mineralogy and texture ($\text{Fe}_{\text{oxalate}}$ and $\text{Fe}_{\text{crystalline}}$, clay, silt and sand content) were weaker than those with variables representing SOC quantity and quality, and when significant did not remain so in the 0–10 and 50–60 cm depth intervals separately, indicating that they may result from co-variation with depth rather than actual mechanistic relationship. They were however stronger for PyC_{HyPy} concentration, in particular the correlation with $\text{Fe}_{\text{oxalate}}$ ($\tau = 0.47$). $F^{14}\text{C}$ of PyC_{HyPy} was positively correlated with CEC and $\text{Fe}_{\text{oxalate}}$ ($\tau = 0.66$ and 0.37, respectively) and negatively correlated with clay content ($\tau = -0.40$).




Figure 2. Profiles of PyC_{CTO} (dots) and PyC_{HyPy} (triangles) proportion in SOC with depth by soil group, according to the gradient of soil evolution with slope position. CM – Cambisol, LV – Luvisol, ST – Stagnosol, jd – hyperdystric, ha – haplic, lv – luvic, rw – relectistagnic, st – stagnic, ab – albic, fv – fluvic, sv – solimovic. Error bars represent standard error (typical analytical error when $n = 1$)




Figure 3. (a) PyC_{CTO} and (b) PyC_{HyPy} content in the topsoil (0–10 cm) with altitude relative to the toeslope. Points are colored by soil group, according to the gradient of soil evolution with slope position. CM – Cambisol, LV – Luvisol, ST – Stagnosol, jd – hyperdystric, ha – haplic, lv – luvic, rw – relectistagnic, st – stagnic, ab – albic, fv – fluvic, sv – solimovic. In panel a points joined by a line of the same type belong to the same transect, solid line – transect 1, dashed line – transect 2, dotted line – transect 3. Error bars represent the usual analytical error.

4 Discussion

4.1 Cropland soils store old PyC after centuries of cultivation

Our results attest to the durable presence of PyC in cropland soils even after centuries of cultivation. The transition to continuous cropping may not have completely suppressed fire from the landscape, as stubble burning was practised in Western Europe, but we assumed that ensuing PyC inputs were either small (between 25 % and 50 % of cereal crop biomass is harvested and crop residues have many other uses besides burning, Smil, 1999), or/and of more labile nature due to low ($< 350^\circ\text{C}$) fire temperatures (Rasmussen et al., 1986). PyC_{HyPy} in the first 10 cm of the soil had substantially

lower (i.e. older) F^{14}C than SOC (Fig. 5a), highlighting their longer residence time of PyC in soils, up to millennia. These low F^{14}C values (0.72–0.83, 2630–1530 years BP) further indicate the presence of a non-negligible proportion of old PyC remaining in the topsoil, consistent with the assumption of limited to no PyC inputs in the last two centuries. They are similar to F^{14}C found at 0–15 cm depth in boreal soils with discontinuous permafrost in Canada (0.76–0.86, Schiedung et al., 2024) but significantly higher (i.e. younger) than values for continuous permafrost soils from the same study (0.44–0.56). They are lower (i.e. older) than in the 0–15 cm layer of tropical soils under dry shrubland and forest (0.85–0.95 and 0.89–1.05, respectively, Bellè, 2023). These differences could be explained by a combination of different




Figure 4. PyC_{CTO} stocks in the soil profile down to 60 cm (darker shade), topsoil (0–30 cm, intermediate shade), and subsoil (30–60 cm, lighter shade) by soil group according to the gradient of soil evolution with slope position. CM – Cambisol, LV – Luvisol, ST – Stagnosol, jd – hyperdystric, ha – haplic, lv – luvic, rw – relectistagnic, st – stagnic, ab – albic, fv – fluvic, sv – solimovic. The numbers above the bars are the proportion of the topsoil (resp. subsoil) PyC stock in the total stock, the numbers inside the bars are the proportion of PyC stock in the total SOC stocks. Error bars represent the propagated standard error.




Figure 5. (a) $F^{14}C$ of bulk SOC (white fill) and PyC_{H_{Py}} (grey fill) in the topsoil (0–10 cm) for upslope positions (orange) and downslope positions (purple). Two-sided Wilcoxon–Mann–Whitney test for the difference between C pools (***) – $p < 0.01$ and between slope positions within each pool (ns – not significant, $p > 0.1$). (b) $F^{14}C$ of PyC_{H_{Py}} in topsoil (0–10 cm, circles), intermediate (20 to 40 cm depth, triangles) and subsoil (50 to 70 cm depth, squares) horizons in upslope and downslope positions. brown – Hyperdystric Cambisol at the summit, dark orange – Haplic Cambisols at intermediate flats and shoulders, aquamarine – Haplic Cambisol and relectistagnic Luvisols and the footslope, green – Stagnic Albic Luvisol and Albic Fluvic Stagnosol at the toeslope and pink – Solimovic Cambisol at the footslope. Note the different x-axis scale between the two panels.

PyC mineralisation rates (related to soil temperature and PyC quality) and different fire regimes (more frequent and recent PyC inputs in both climatic zones).

PyC_{CTO} concentrations in the topsoil under cropland (0.4 to 0.7 gC kg⁻¹ at 0–10 cm, Figs. 3 and S2) were very similar to that found in soils of the Région Centre in France (mostly croplands, Q1 = 0.4 to Q3 = 1 gC kg⁻¹ at 0–30 cm,

Paroissien et al., 2012) and in Swiss cropland soils (0.4 to 0.8 gC kg⁻¹ at 0–20 cm, Agarwal and Bucheli, 2011). They were slightly lower than that measured at 0–20 cm in grassland (0.8 to 1.3 gC kg⁻¹) and forest (0.6 to 2.7 gC kg⁻¹) soils across Switzerland (Agarwal and Bucheli, 2011) and at 0–5 cm in remote and rural areas of the UK and Norway (range 0.45 to 3.13 gC kg⁻¹, average 1.27 gC kg⁻¹, Nam et al., 2008). These comparisons point to a potential loss of PyC_{CTO} from the topsoil in cultivated soils.

However, in our study area, PyC_{CTO} stocks over 60 cm were lower in a soil under woodland (1.9 ± 0.22 tha⁻¹, Summit – CM-jd in Fig. 4) than under croplands in similar topographic positions (2.6 ± 0.22 tha⁻¹, Flat and Shoulder – CM-ha in Fig. 4), after more than 150 years of cultivation. To the best of our knowledge, there exist no other study that measured PyC_{CTO} stocks in a similar pedoclimatic context. Silva et al. (2023) found an average PyC_{CTO} stock to 1 m depth of 1.37 ± 0.65 tha⁻¹ in Amazon forest fragments in Brazil. This is similar to our stocks to just 30 cm depth (1.2 to 1.6 tha⁻¹, Fig. 4). Santos et al. (2017) and Soucémariadin et al. (2014) found PyC stocks to the lower limit of the B horizon between 1.75 and 4.9 tha⁻¹ in Podzols of North American forests using a PyC isolation method based on chemical oxidation (KMD method). These higher values could be attributed to different fire regimes, soil properties or a larger coverage of the PyC continuum by the KMD method compared to the CTO-375 method which only detects the most condensed, soot-like PyC (Hammes et al., 2007; Gustafsson et al., 2001; Elmquist et al., 2004). Across pedoclimatic contexts, PyC stocks determined with the BPCA method (Lehndorff et al., 2016; Rodionov et al., 2010) are 2 to 20 times larger than PyC_{CTO} stocks (Schiedung et al., 2024; Bellè, 2023, and this study), which points to method differences as the most likely explanation and prevents meaningful comparisons between the data obtained from these two different methods.

The subsoil (30–60 cm) represented a important share of the total PyC_{CTO} stock (44 %, Fig. 4) in our study area. This proportion was slightly lower (not significant) in the Hyperdystric Cambisol under woodland (37 ± 6.2 %) than in the Cambisols under cropland (44 % to 46 %). Soucémariadin et al. (2014) and Silva et al. (2023) found only 17 % and 30 % of the PyC stock below 30 cm in Podzols and Ferralsols under forest, respectively, whereas under cropland Lehndorff (2016) found more than 50 % of the PyC stock in the subsoil (below the A horizon) for an Andosol, an Alisol, a Cambisol and a Vertisol and less than 50 % for a Fluvisol and another Alisol, when not under paddy rice cultivation. Redistribution of PyC to the subsoil in croplands could be favoured by mixing and fragmentation of PyC particles due to tillage. All soil profiles in recently tilled fields showed a net homogenisation of PyC_{CTO} (Fig. S2), SOC and N content (not shown) in the first 20 to 30 cm of the soil compared to the woodland and grassland sites (summit and toeslope positions in transect 1, respectively). Rodionov et al. (2006) showed that




Figure 6. Kendall's rank correlation between PyC content (CTO and HyPy), PyC proportion in SOC (CTO and HyPy), F¹⁴C (PyC_{HyPy} and bulk SOC) and selected soil characteristics, including SOC and indicators of SOC quality ($\delta^{13}\text{C}$, C : N), soil chemistry (pH_{H2O}, CEC_{Co}, Fe_{oxalate} and Fe_{crystalline}), soil texture (Clay to Sand) and soil evolution (Silt : Clay, weathering index WI and chemical index of alteration CIA), across depths. Significant correlations at the 95 % confidence level are in bold. Correlations that remain significant and of the same sign in at least one depth increment (0–10 or 50–60 cm) are outlined in black.

the fine sand and silt fractions (particle size 2–250 μm) at 0–10 cm depth were enriched in PyC in arable fields compared to adjacent native grassland, which they attributed to the crushing of bigger PyC particles by cultivation practices. Using a soil physics model, Hobley (2019) demonstrated that smaller PyC particles were more likely to be transported to depth, especially when eluviation rate is high relative to decomposition rate, which is likely the case for PyC particles in well drained soils. PyC redistribution to the subsoil, where decomposition rates are usually lower and erosion inexistent, could contribute to its preservation in cropland soils.

Altogether, these results show that in temperate regions and on shallow slopes, PyC can persist in cropland soils for centuries, bearing traces of past fire regimes.

4.2 SOC becomes enriched in PyC with depth

We observed an increasing proportion of PyC in SOC with depth in all profiles and for both PyC determination methods (Fig. 2). This is common for PyC_{CTO} (Qi et al., 2017; Agarwal and Bucheli, 2011; Koele et al., 2017; Silva et al., 2023) and PyC measured using other quantification methods (Andreeva et al., 2011; Brodowski et al., 2005; Butnor et al., 2017; Dai et al., 2005; Hammes et al., 2008a; Matoszuik et al., 2020; Soucémariadin et al., 2019; Güereña et al., 2015; Velasco-Molina et al., 2016), across soil type, vegetation and climate, although constant or decreasing proportion of PyC in SOC with depth have been observed under grasslands (Rodionov et al., 2006), in Andosols and in some cases under paddy rice cultivation (Lehndorff et al., 2014), in Podzols (Soucémariadin et al., 2014), and where there are frequent PyC inputs at the surface from stubble burning (Rumpel et al., 2006a). At the same time, F¹⁴C of PyC_{HyPy} was always lower (older) than F¹⁴C of SOC (Table S1) and decreased sharply with depth in all profiles but two (Fig. 5b). We are the first to report F¹⁴C of PyC_{HyPy} in temperate cropland soils but Butnor et al. (2017) found a regular decrease in the F¹⁴C of the chemically resistant SOC fraction (which

can be equated to PyC) from 0–20 cm (0.88–0.66) down to 100 cm (0.28–0.54) in forest soils in the southern US, in close agreement with our findings.

Several hypotheses have been brought forward by the above-cited authors to explain the increasing proportion of PyC in SOC with depth: (1) PyC is preferentially transported to depth relative to non-PyC, (2) PyC is preferentially preserved at depth (e.g. via retention and stabilisation on mineral surfaces) relative to non-PyC, or (3) PyC has an intrinsically higher biological and/or chemical stability relative to non-PyC, which allows a larger proportion of the inputs to reach and stay in the subsoil before being decomposed by microorganisms or abiotic reactions. Although our study was not designed to answer this question, we examined the three mechanisms of PyC enrichment with depth and argue that higher stability of PyC relative to non-PyC is sufficient to explain the profiles we observe (Fig. 7).

(1) Estimates of downward vertical transport rate of particulate and dissolved PyC range from 2 to 30 mm yr^{-1} (Rumpel et al., 2015). Such fast transport of PyC to depth is not compatible with our data (see also Fig. 5b in Hobley, 2019): at 2 mm yr^{-1} it would take only 250 years for PyC to migrate from the 0–10 to 50–60 cm layer when the age difference we observe is between 1740 and 7410 ¹⁴C years. The sharp increase of PyC proportion in SOC at the E horizon of the Stagnic Albic Luvisol and Albic Fluvisol Stagnosol (Fig. 2), where SOC eluviation is the strongest, is also contrary to the hypothesis of preferential vertical transport of PyC relative to SOC in the pedoclimatic context of our study area.

Fast vertical transport in the soil profile may only apply to a small portion of the PyC pool. Indeed, dissolution and leaching of PyC is usually limited relative to SOC (Abiven et al., 2011; Maestrini et al., 2014; Major et al., 2010; Hilscher and Knicker, 2011; Schiedung et al., 2020; Abney et al., 2024). Alternatively, the elevated rates measured in previous studies may apply only to soils with high porosity (Leifeld et al., 2007), coarse texture (Schiedung




Figure 7. Conceptual, 1D model of PyC and non-PyC persistence in soils based on differences in biological and chemical stability. After a fire some PyC is incorporated in the topsoil (a) and starts to migrate downward into the soil profile by leaching, eluviation and turbation (b). Vegetation regrowth provides non-PyC inputs continuously throughout the post-fire succession (b, c), natural fire – roman type, slash and burn – italic type). Over time, transport and decomposition create a concentration gradient with depth (b, c). The transport rate of PyC may be slower than that of non-PyC, but because it is more stable, a larger proportion of it remains available to be transported and its concentration gradient is less pronounced. On the contrary, a large part of non-PyC is lost via microbial respiration before it can reach the altered bedrock. After several fire cycles and millennia of soil evolution, it results in the observed concentration profiles, under the current agricultural land-use (d).

et al., 2020) and/or where annual rainfall is important (Alexis et al., 2012; Major et al., 2010; Nguyen et al., 2009), conditions that are not met in our study area.

Soil mixing processes (bioturbation, peloturbation, cryoturbation and that induced by windthrow and agricultural practices) can transfer particulate PyC to depth (Lehndorff et al., 2016; Schiedung et al., 2020; Carcaillet, 2001) but are not selective for PyC. As demonstrated by Hobley (2019, see figure 1d), these spatially heterogeneous processes are also inconsistent with the nearly systematic increase of PyC_{HyPy} age with depth that we observe here (Fig. 5) and in other studies (Carcaillet, 2001; Hajdas et al., 2007).

(2) Both PyC and SOC in the subsoil are effectively protected from erosion and usually experience lower microbial activity. The proportion of mineral-associated SOC, a fraction which typically has higher stability, is often found to increase with depth (e.g. Sanderman et al., 2021). PyC was found to form mineral-associations potentially mediated by iron (Solomon et al., 2012; Brodowski et al., 2005; Schiedung et al., 2023). Schiedung et al. (2020) have shown that PyC migrating in a saturated soil column was preferentially found in the mineral associated fraction in a finer textured soil with higher clay and $\text{Fe}_{\text{oxalate}}$ contents, and interaction of PyC with iron (hydr)oxides was suggested as the reason for PyC retention in podzolic Bs horizons (Soucémari-

anadin et al., 2014; Santos et al., 2017), but there is no compelling evidence that PyC is preferentially preserved relative to non-PyC.

In our study, $\text{Fe}_{\text{oxalate}}$ (iron in organo-metallic complexes + poorly crystalline oxides) showed weak, negative correlations with the proportion of PyC_{CTO} and PyC_{HyPy} in SOC and a weak positive correlation with F^{14}C of PyC_{HyPy} (Fig. 6) indicating no preferential preservation of PyC with increasing presence of reactive iron phases. The proportion of PyC_{HyPy} in SOC presented an intermediate, positive correlation with iron in crystalline oxides overall (Fig. 6) but not in the 50–60 cm layer ($\tau = -0.09$, $p = 0.73$), indicating that this form of iron was not a factor of PyC stabilisation in the subsoil. In previous studies, PyC content correlated positively with pyrophosphate and Meilich-3 solution extractable iron (iron in organo-metallic complexes, Eckmeier et al., 2010; Soucémariadin et al., 2014; Butnor et al., 2017) but not with $\text{Fe}_{\text{oxalate}}$, Fe_{DCB} and total iron (Qi et al., 2017; Lehndorff et al., 2016; Velasco-Molina et al., 2016), warranting further investigation of PyC interaction with iron in soils.

We also found no significant correlation between the proportion of PyC_{CTO} and PyC_{HyPy} in SOC and clay content (Fig. 6). There was a negative correlation of intermediate strength between PyC_{HyPy} F^{14}C and clay content overall but it disappeared when looking at each depth interval separately, which points to a co-variation with depth rather than a mechanistic relationship. This is not inconsistent with the sparse evidence of correlations between PyC and clay content in the literature (Qi et al., 2017; Velasco-Molina et al., 2016; Butnor et al., 2017; Paroissien et al., 2012; Soucémariadin et al., 2014; Schiedung et al., 2024; Sanderman et al., 2021). In addition, uncertainties remain about whether or not PyC in the clay fraction is actually mineral-associated (Lutfalla et al., 2017; Hilscher and Knicker, 2011; Brodowski et al., 2007; Skjemstad et al., 1996) and whether or not mineral-associated PyC is more stable than its particulate counterpart (Hilscher and Knicker, 2011; Chassé et al., 2021; Vasilyeva et al., 2011).

(3) Millennial persistence of at least some portion of PyC is attested by radiocarbon dating of charcoal fragments (Carcaillet, 2001; Hajdas et al., 2007; Gavin et al., 2023; Ohlson et al., 2009; Liang et al., 2008) and thermally or chemically isolated PyC (Butnor et al., 2017; Schiedung et al., 2024). Long-term field experiments have shown that PyC tends to be more stable than the rest of SOC (Lutfalla et al., 2017; Vasilyeva et al., 2011; Hammes et al., 2008b), which turns over on decadal to centennial time scales (Schmidt et al., 2011; Barré et al., 2016). Our observations support this difference, with modern to ≈ 1000 ^{14}C years old total SOC and ≈ 1500 to $\approx 10\,000$ ^{14}C years old PyC_{HyPy} . Higher stability of PyC in soil would give it time to be transported down the soil profile from the surface where it is produced, and explain its higher proportion in the subsoil relative to SOC, despite potentially large inputs of relatively stable SOC from

roots directly to the subsoil (e.g. Rasse et al., 2006; Sokol et al., 2019). During ageing, PyC is fragmented and oxidised (Hockaday et al., 2006; Sorrenti et al., 2016; Pignatello et al., 2015), its solubility increases (Abiven et al., 2011) and colloids are formed which could move more easily through the soil porosity. These processes could favour PyC transport to depth over time (Carcaillet, 2001). In addition, small, oxidised PyC particles are more prone to form mineral association (Schiedung et al., 2020), which could favour their retention once in the subsoil and contribute to the overall persistence of PyC.

Thus we argue that intrinsically higher stability of PyC relative to SOC, not preferential transport or preservation, explains the large proportion of very old PyC in subsoil SOC stocks in most soils in the Kervidy-Naizin watershed (Fig. 7). Despite good drainage in the upslope domain and noticeable eluviation in the downslope domain (Walter and Curmi, 1998), vertical transport of PyC likely remains slow, consistent with the large $F^{14}C$ difference between topsoil and subsoil PyC and the lingering presence of old PyC in the topsoil.

4.3 Slope position does not control stocks and age of PyC

4.3.1 Toeslopes do not accumulate PyC in the long term

The hypothesis that in this watershed, toeslopes would represent large stocks of old PyC, is refuted. On the contrary, PyC_{CTO} stocks were relatively homogeneous along the gradient of soil evolution with slope position (Fig. 4), around 2.5 t ha^{-1} . In the topsoil, PyC content and proportion in SOC even decreased with elevation relative to the toeslope for PyC_{HyPy} , the fraction of PyC that is 7–14 aromatic rings or bigger, and showed a minimum at midslope for PyC_{CTO} , at the most condensed end of the PyC continuum (Fig. 3). Our hypothesis was based on the assumption that after a fire, PyC characteristics (low bulk density, hydrophobicity) and favourable conditions for erosion (lesser vegetation and litter cover) would lead to the redistribution of PyC to the toeslope, where the shallow slope and riparian vegetation would stop it from reaching the stream. Its incorporation into the soil profile would then protect it from further transport and decomposition. To explain our results, we suggest that PyC produced by past fires was either not significantly redistributed along the slope, washed out to the stream, or lost from the toeslope since then.

If evidence of (preferential) PyC erosion post fire is numerous (Rumpel et al., 2009; Cotrufo et al., 2016a; McGuire et al., 2021), most PyC erosional studies took place on steep slopes (Cotrufo et al., 2016a; Rumpel et al., 2006b; Güereña et al., 2015), or under high rainfall intensity on bare soil (Rumpel et al., 2009; Bellè et al., 2021). In case of high severity fires, PyC can be eroded independent of slope if sufficient rainfall occurs (McGuire et al., 2021; Boot et al., 2015). On shallow slopes and flat areas or where burn sever-

ity is lower or patchy, PyC may only be redistributed locally (Pyle et al., 2017; Galanter et al., 2018). Little is known about past fire regimes in the region, let alone fire severity (Appendix A1). However, under the current agricultural land-use which leaves the soil bare a large part of the year, erosion proceeds in successive steps, with overland flow infiltration several times along the slope and without formation of rills and gullies able to carry material far away (Cros-Cayot, 1996). This suggests that post-fire transport of PyC to the toeslope could have been limited.

Even where there was transport of PyC to the toeslope, several studies have failed to detect long term accumulation of PyC in low topographic positions (Rumpel et al., 2006a; Galanter et al., 2018; Güereña et al., 2015; Abney et al., 2017). Galanter et al. (2018) suggested that PyC directly flowed over the riparian area and into the stream, whereas Güereña et al. (2015) advocated that PyC was only temporarily stored in the lower part of a convex slope before being picked up by the river in later erosion/flood events. The same process could explain the disappearance of PyC at the bottom of a concave slope 10 years post-fire observed by Abney et al. (2017), although the absence of visible erosion features led them to favour the burial or decomposition hypotheses. Here, the presence of a dense albic eluvial layer under the A horizon in the Stagnic Luvisols and Stagnosols near the stream (Fig. 1) reduces infiltration (Curmi et al., 1998; Cros-Cayot, 1996), favouring surface flow that could take away light, freshly deposited PyC to the stream and prevent its incorporation deeper into the soil profile. Indeed, in another study based in the same watershed, Vongvixay et al. (2018) have shown that if riparian trees and grass buffer strips are efficient at intercepting suspended solids transported by runoff during most of the year, they seem to lose this ability during winter, maybe because the vegetation is submerged. Rumpel et al. (2006b) suggested that because it is preferentially found in the light fraction, PyC is easily eroded and transported out of the watershed. Studies considering different spatial scales showed that PyC was enriched in sediment eroded from the headwater catchment but depleted in the larger order catchment (Chaplot et al., 2005), indicating that PyC was transported away from its production site and redeposited further downstream, for instance at river banks (Cotrufo et al., 2016a).

Finally, PyC at the toeslope may have been lost overtime at a faster rate than in the upslope domain. Studies on SOC and DOC in porewater and in the stream in the Kervidy-Naizin catchment have shown that the fluctuation of the water table in the soil profile affects OC dynamics (Lambert et al., 2013, 2014; Jeanneau et al., 2014). In the area close to the stream where the water table is present to the surface a large part of the year, SOC mineralisation was reduced relative to midslope areas but dissolution was increased which resulted in export of OC from the subsoil. The dissolution of SOC was related to the destabilisation of OC-retaining iron oxides under reducing conditions (Lambert et al., 2013; Jean-

neau et al., 2014). Indeed, the Stagnic Luvisols and Stagnosols at the toeslope were depleted in free and amorphous iron which partly re-precipitated as more crystalline iron oxides in mottles in the illuvial B horizon. PyC_{HyPy} content was significantly positively correlated with intermediate strength to $\text{Fe}_{\text{oxalate}}$ (free and amorphous iron) and strongly to $\text{Fe}_{\text{crystalline}}$ in the topsoil ($\tau = 0.47$ and 0.67 , respectively, $p < 0.01$) but not in the subsoil, indicating that the same dissolution processes may have affected PyC_{HyPy} , preventing its accumulation in toeslope positions.

The difference between PyC_{HyPy} and PyC_{CTO} along the toposequence may result from differences in PyC quality. PyC_{HyPy} represents PyC from 7 aromatic rings in size while PyC_{CTO} only accounts for the most condensed, soot-like PyC. As such, PyC_{HyPy} is likely to be more prone to solubilization and more labile, and thus to depend more on mineral interactions (e.g. with iron oxides, see above) for its preservation, while PyC_{CTO} would behave more like an inert soil particle affected primarily by erosion (see Sect. 4.3.2) and eluviation (see Sect. 4.2).

4.3.2 Recent erosion caused redistribution of PyC with the soil matrix to specific locations

PyC_{CTO} and SOC stocks were higher in the Solimovic Cambisol at the toeslope of transect 3 relative to all other soils along the gradient of soil evolution with slope position (Fig. 4), whereas cross transects, PyC_{CTO} content in the topsoil decreased with increasing slope (Fig. S3). We interpret these observations as signs of recent (relative to PyC residence time) erosion processes that have redistributed topsoil PyC in and out of the watershed.

The soil profile that we classified as a Solimovic Cambisol presented a thick A horizon (55 to 60 cm). The composition of this horizon did not markedly differ from the A horizon of soils higher in the same transect, which supports the local origin of this material. PyC and SOC F^{14}C did not decrease with depth like in the other undisturbed soil profiles but were instead slightly higher at 40–50 cm depth than at 0–10 cm (Table S1), a feature consistent with depositional processes (see also Fig. 2d in Hobley, 2019). Since PyC content in the soil decreased with depth (Fig. S2), a change in the relative soil surface due to erosion at steeper, midslope positions would explain the lower PyC_{CTO} concentrations relative to less eroding, shallower summit and toeslope positions.

Despite shallow slopes, erosion in this landscape is likely favoured by extended periods without vegetation cover, and cultivation in the direction of the slope. Based on the extent, thickness and density of deposits, Cros-Cayot (1996) estimated that within one year of conversion from grassland to maize, 15 t of topsoil were eroded and deposited at the concavity in a 3 ha field located in the same catchment as our transects. Using sediment traps along several toposequences in another catchment in the region (similar geology, climate, topography and soil types), she estimated that 152.5 g m^{-2} of

solids were eroded yearly, of which 17 % (25.9 g m^{-2}) was exported to the river while the rest was redeposited along the slope. These values were considered as an upper limit for the Kervidy-Naizin catchment where structural stability of the soil was higher (Cros-Cayot, 1996). This is in line with the results of Vongvixay et al. (2018), who measured discharge and turbidity in the stream at the outlet of the catchment over 9 years, and calculated annual suspended sediment yields ranging from 3 to 22 g m^{-2} .

Another similar catchment in the region had higher suspended sediment yields despite a greater proportion of woodland and grassland (smaller sediment source area) and a higher density of hedgerows (more obstacles to sediment fluxes), which the authors attributed in part to the retention of sediments by riparian trees and grass buffer strips in the Kervidy-Naizin catchment (Vongvixay et al., 2018). The location of the Solimovic Cambisol we sampled, at the toeslope and about 15 m away from the riparian tree line, is consistent with this interpretation. However, we did not find noticeable amounts of solimovic material at the lowermost sampling site of T1 and T2, which suggest that such accumulation is localised. It may be related to a slightly raised bank providing a barrier between the slope and the stream and/or to the short toeslope at T3. Where the toeslope is longer, material eroded at midslope may be redistributed in a diffuse way, consistent with the stepwise erosion model proposed by Cros-Cayot (1996) for these watersheds. Alternatively, sediments deposited at the toeslope throughout the year may be washed out during winter floods.

In the Solimovic Cambisol at the toeslope, SOC had an F^{14}C of 1.03 at 40–50 cm depth (Table S1), indicative of the presence of a non negligible proportion in the subsoil of organic carbon formed following the bomb peak (circa 1963). Throughout the A horizon, soil chemistry (pH, CEC, C/N ratio, CIA, not shown), SOC and PyC content (Fig. S2) were almost constant, showing that the deposited material did not have time to evolve. On shallow slopes and structurally stable soils, erosion is likely negligible under permanent vegetation cover. Agricultural activity in the area may date back as far as 5000 BP but continuous cultivation is likely much more recent (Astill and Davies, 1997; Tonnerre, 1992). Slope continuity along transect 3 was only established in the 1970's when the hedgerows at the former plot boundaries were removed throughout the Kervidy-Naizin catchment (Appendix Fig. A1). In the following decades, this catchment saw an intensification of agricultural practices (see Appendix A2) with an increasing frequency of maize in the rotation (Bordenave and Merceron, 1999). When maize is planted after a winter crop (e.g. wheat), a common occurrence in the fields where our transects are located, the soil can remain bare for more than 6 months, which favours erosion (Bordenave and Merceron, 1999). This concurs to indicate that the establishment of the Solimovic Cambisol at the toeslope of T3 is recent.

There was no PyC enrichment or depletion during erosion and deposition at T3 (Fig. 3). PyC accumulation at this

site was only related to accumulation of topsoil material, and would have gone undetected if we did not consider the entire soil profile. Aggregate stability measured in previous studies was high (Cros-Cayot, 1996) and soil was likely transported in aggregated form, as evidenced by the absence of textural differentiation in the solimovic material (not shown). Under these conditions, centuries to millennia old PyC was not preferentially eroded, as opposed to what was observed for fresh PyC (Bellè et al., 2021; Cotrufo et al., 2016b; Rumpel et al., 2009; Chaplot et al., 2005). It suggests that during aging, PyC becomes involved in aggregates and mineral interactions which stabilise it with regard to erosion.

5 Conclusions and perspectives

We showed that topographic position was not the main driver of PyC stocks in the landscape, in a shallow sloping watershed dominated by agricultural land-use under temperate oceanic climate, and after hundreds of years of cultivation. If it existed, PyC enrichment at the toeslope would have been only temporary and probably reversed in the decades following fire due to subsequent erosion or conditions favouring dissolution and leaching of PyC. Changes in erosion dynamics related to land-use redistributed already aged PyC to a localised area of soil accumulation without enrichment or depletion of PyC. This could make PyC an interesting indicator to trace erosion in temperate agricultural landscape where fire is rare. More studies are needed to assess whether different erosion modalities (sheet vs. rills and gullies) and aggregate stability (i.e. whether or not the soil disaggregates during erosion) affect the fate of aged PyC. Large stocks of old PyC in the subsoil supports the long persistence of PyC in soils and slow advection towards soil depth under the pedoclimatic conditions of our study area. More than 150 years of cultivation did not seem to deplete PyC stocks relative to an adjacent wooded area but may have favoured PyC redistribution below 30 cm depth, although replication is needed to confirm these findings. We call attention to the importance of subsoil sampling when studying persistent forms of organic carbon and estimating their stocks. These results are of interest in the context of biochar application to agricultural soils, as losses of biochar from the topsoil may be accounted for by lateral and/or vertical transport, and not only by mineralisation, which impacts the carbon budget of the system. Identifying the proportion of PyC produced which is quickly transported away from the watershed and that which remains in soils for millennia after a fire is an important knowledge gap that still needs to be investigated to close the terrestrial PyC budget, but it necessitates a better estimate of PyC production from fires and knowledge of fire history and past fire regimes. Pools of PyC of different quality measured by different PyC quantification methods showed different patterns of concentration in the topsoil along the gradient of soil evolution with slope position. The most condensed, soot-like

form of PyC measured by CTO tended to be depleted at mid-slope, possibly as a result of recent erosion that exposed former subsoils less rich in PyC, whereas the intermediate to highly condensed PyC measured by HyPy was increasingly depleted with proximity to the river, which may be related to changes in iron-mediated mineral interactions favouring its dissolution and leaching. These interactions need to be studied in more details. Multi-method PyC characterisation is an opportunity to study the interplay between PyC quality and soil processes.

Appendix A: Land-use and fire history at the study site

A1 Land-use and fire history in Brittany over the Holocene

The loess that form the parent material of the soils in the Kervidy-Naizin watershed are derived from both local and remote sources and were likely deposited on the periglacial landscape during the last glacial period ($\sim 115\,000$ to $11\,700$ BP, Pellerin and Van Vliet-Lanoë, 1998; Van Vliet-Lanoë et al., 1998). More specifically it is estimated that the loess deposition in the Northern European Loess Belt started at about $32\,000$ BP and that maximum deposition occurred around $21\,800$ BP Bosq et al. (2023), and loess deposits in Northern Brittany have been dated between $15\,900$ and $67\,000$ BP Van Vliet-Lanoë et al. (1997) but no data exists at our study site. To the best of our knowledge, there are little indications as to the vegetation dynamics in the region at the Pleistocene to Holocene transition and into the early Holocene. Paleoenvironmental and archaeological studies have shown that the Atlantic oak forest that was likely predominant in Brittany before 7500 BP started to be opened by early human population during the neolithic period, from the coast inwards. The forest was first cleared for pasture, with only localised cropping. Abandoned land was re-colonised by heath and heliophytes tree species that could be maintained by grazing or fire (Gebhardt and Marguerie, 2006; Briard et al., 1989). It is estimated that most of the primary Atlantic forest had been cleared by the end of the Iron Age (Gaudin et al., 2014). Heath and cropland increased globally from the Bronze Age to the end of the Middle Ages with periods of acceleration or stability (Gaudin et al., 2014). Tonnerre (1992) points out that in the Early Middle Ages (1500 – 1000 BP), small pieces of land were cropped probably for short periods and rotated often, which could have led to frequent slash-and-burn practices. Blank spaces (small woods, some wetlands) were taken into cultivation in the late mediaeval – early modern period but heath and secondary growth woodland dominated the landscape until the end of the 19th century (Astill and Davies, 1997). However, heath was not synonymous of land abandonment as part of the heathland was only temporary (a form of fallow), part was grazed, and part was used as a source of plant material for

animal bedding and fertilization of the fields after “composting” or burning the biomass to ashes (Antoine, 2000, see the definitions of *Étrepage* and *Écobuage*). The division in small plots, delimited by fences, ditches and/or hedgerows, originated in the 16th–17th century and lasted until the 1970’s when plots were pooled together to allow the use of heavier machinery on larger surfaces (Astill and Davies, 1997).

There are no good nowadays equivalent of the Atlantic forest that would allow us to estimate pre-historical fire regimes in the area. Fires during the period of human presence after the first deforestation may have been mostly related to slash and burn practices to maintain the heather or clear the secondary growth forest (Gebhardt and Marguerie, 2006; Briard et al., 1989). Heath and moorland fires can be very intense, although very variable (Hobbs and Gimingham, 1987; Davies et al., 2022), and burn into the moss and litter layer (Davies et al., 2022). However, as it is a multi-millennial landscape shaping practice, users have developed a traditional knowledge on how to minimise soil impacts (in particular erosion) while achieving management goals (Davies et al., 2022). This could have limited the severity of heath fires. In croplands, the addition of ashes resulting from heathland vegetation burning, a practice described by Antoine (2000) for the modern period (late 15th to early 19th century), contributed an unknown amount of PyC to the soil.

A2 Land-use and fire in the study area in the last 150 years

The reconstruction of land-use and plot borders in the catchment area over the last 150 years was based on maps, aerial photographs and farmer’s declarations. You can visualise most of the cited maps on Géoportail ([https://www.geoportail.gouv.fr/carte?c=-2.8423422520194026,48.009738098366256&z=15&l0=GEOGRAPHICALGRIDSYSTEMS.ETATMAJOR40::GEOPORTAIL:OGC:WMTS\(1\)&l1=GEOGRAPHICALGRIDSYSTEMS.MAPS.SCAN50.1950::GEOPORTAIL:OGC:WMTS\(1;h\)&l2=ORTHOIMAGERY.ORTHOPHOTOS.1950-1965::GEOPORTAIL:OGC:WMTS\(1;h\)&l3=GEOGRAPHICALGRIDSYSTEMS.PLANIGNV2::GEOPORTAIL:OGC:WMTS\(1;h\)&l4=ORTHOIMAGERY.ORTHOPHOTOS::GEOPORTAIL:OGC:WMTS\(0.25;h\)&d5=3573961\(1\)&d6=3573952\(1\)&permalink=yes](https://www.geoportail.gouv.fr/carte?c=-2.8423422520194026,48.009738098366256&z=15&l0=GEOGRAPHICALGRIDSYSTEMS.ETATMAJOR40::GEOPORTAIL:OGC:WMTS(1)&l1=GEOGRAPHICALGRIDSYSTEMS.MAPS.SCAN50.1950::GEOPORTAIL:OGC:WMTS(1;h)&l2=ORTHOIMAGERY.ORTHOPHOTOS.1950-1965::GEOPORTAIL:OGC:WMTS(1;h)&l3=GEOGRAPHICALGRIDSYSTEMS.PLANIGNV2::GEOPORTAIL:OGC:WMTS(1;h)&l4=ORTHOIMAGERY.ORTHOPHOTOS::GEOPORTAIL:OGC:WMTS(0.25;h)&d5=3573961(1)&d6=3573952(1)&permalink=yes), last access: November 2025). In the following paragraphs we highlight the main changes and their potential impacts on the watershed hydrology.

The oldest document available for our study area is the land register established in 1833, commonly referred to as “*adastre Napoléonien*” (Ministère des Finances, Direction départementale des contributions directes, Bureau du cadastre, 2007). This register consists of two layers, one that displays the limits of plots of land based on ownership and one that represents land use. The later has unfortunately been

lost for most municipalities in the “*département*” (french administrative division, similar to a county) where our study area is located. However, the military survey map established between 1820 and 1866 (IGN, 2021) indicated that most of the study area was open land (cropland, grassland or heath), apart from a wooded area at the south-west border of the catchment (most upslope sampling location in T1) and localised patches of trees, notably around the upslope area of T3. The stream was surrounded by small wetlands in most of the study area. Comparing the 1833 land register and its updated version of 1952 (Direction départementale des finances publiques, 2020), we observed that plot borders have remained the same over the period. Both registers are publicly available on the website <https://patrimoines-archives.morbihan.fr/> (last access: November 2025) of the *département du Morbihan*.

The aerial photograph taken in 1952 (IGN and GEOPAL, 2011) revealed the presence of hedgerows at most plot borders in the area of T1 and T3 whereas only a sparse hedge separated the top-most sampling site in T2 from the rest of the transect. The wetlands around the stream appeared to have been taken into cultivation whereas the wood at the south-west border was still present (and still is to this day).

For the most part, hedgerows were removed in the 1970’s as part of a government driven process to modernise agriculture via the use of heavier machinery on larger surfaces. Several path cutting through T1 and T3 were abandoned in the process but the roads separating the upslope sites in T2 and T3 from the summit remained in place (Fig. A1). These changes likely increased the possible path length for erosion fluxes along these transects. Drains have been installed in several of the fields cross-cut by T2 in the 1990’s, significantly reducing the waterlogging in the midslope to toeslope section of the transect (C. Walter, personal communication, 2021). Agricultural practices in most of the study area since the 90’s consist of a rotation of maize, cereals and sometimes vegetables. Aerial photographs from 1993, 1999, 2004, 2009, 2010, 2013, 2016 and 2019 show that cultivation was systematically carried in the direction of the slope at the plots where transect 2 and 3 are located, whereas the field in the upper part of transect 1 is cultivated perpendicular to the slope. The three sampling sites in the lower part of transect 1 are located in a field that had been a cultivated grassland since at least 1993 before being turned to maize in 2020. The wetland around the stream has been left fallow or used as grassland, with recent tree plantation in part of the buffer zone.

Hedgerow removal associated with in-situ residue and/or stump burning could have generated localised PyC inputs. Although some of our sampling points are close to former plot boundaries, we have found no charcoal accumulation layer that could corroborate such practice. Fire has been close to inexistent in croplands of north-western Europe (and in particular in Brittany) in the 21st century Forrest et al. (period 2002–2014, 2024, Fig. 4). Stubble burning was likely a common practice in Western Europe until the 1980’s but




Figure A1. Elevation profiles derived from the 1 m resolution digital elevation model RGE ALTI[®] 1M (IGN, 2013) along transects 1 to 3, with location of the sampling points (red crosses), hedges (circles), and roads or paths (diamonds). The evolution of obstacles (roads and hedges) through time is indicated by colours: green – 2021 (observed), pink – 1952 (derived from IGN and GEOPAL, 2011 and Direction départementale des finances publiques, 2020), orange – 1850's (derived from IGN, 2021, and Ministère des Finances, Direction départementale des contributions directes, Bureau du cadastre (2007)).

we found no data that would allow us to characterize it and estimate the PyC inputs it represented. We assumed that, if it took place in our study area, stubble burning produced small amounts of relatively labile PyC due to low fire temperature (< 350 °C, Rasmussen et al., 1986) and the small amounts of biomass involved (part of the biomass was exported before burning). This hypothesis is supported by the old radiocarbon age of the PyC^{HyPy} in our study area (see main text Sect. 3.4 and 4.1).

Data availability. Data used to produce the figures of this article are publicly available online in a Zenodo repository with DOI <https://doi.org/10.5281/zenodo.17936907> (Lebrun Thauront et al., 2025).

Sample availability. Soil samples can be provided on reasonable demand within five years of publication.

Supplement. The supplement related to this article is available online at <https://doi.org/10.5194/bg-23-155-2026-supplement>.

Author contributions. Conceptualisation: SA, CW, PB. Funding acquisition: SA, CW. Resources: CW, PA, SD. Investigation: JLT, NH. Methodology: JLT. Formal analysis, Visualisation, Data curation: JLT. Supervision: SA, CW, PB, PA, SD. Writing – original draft: JLT. Writing – review and editing: All authors have been involved.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. The authors bear the ultimate responsibility for providing appropriate place names. Views expressed in the text are those of the authors and do not necessarily reflect the views of the publisher.

Acknowledgements. Johanne Lebrun Thauront thanks Bertrand Guenet and Lauric Cécillon for their assistance and expertise in the field; Jacques Meriguet for his assistance in preparing soil samples; Dr Xiaomei Xu at the Keck Carbon Cycle AMS Facility, University of California, Irvine, USA for his expertise in analysing small samples; Patrick Meunier for his valuable inputs on the interpretation of data; Dominique Marguerie for his expertise on paleoenvironments in Brittany.

Financial support. This research has been supported by the Centre National de la Recherche Scientifique (MITI CPy paysages).

Review statement. This paper was edited by Edzo Veldkamp and reviewed by Rebecca Abney and one anonymous referee.

References

- Abiven, S., Hengartner, P., Schneider, M. P., Singh, N., and Schmidt, M. W.: Pyrogenic Carbon Soluble Fraction Is Larger and More Aromatic in Aged Charcoal than in Fresh Charcoal, *Soil Biology and Biochemistry*, 43, 1615–1617, <https://doi.org/10.1016/j.soilbio.2011.03.027>, 2011.
- Abney, R., Barnes, M. E., Moss, A., and Santos, F.: Constraints and Drivers of Dissolved Fluxes of Pyrogenic Carbon in Soil and Freshwater Systems: A Global Review and Meta-Analysis, *Global Biogeochemical Cycles*, 38, e2023GB008092, <https://doi.org/10.1029/2023GB008092>, 2024.
- Abney, R. B. and Berhe, A. A.: Pyrogenic Carbon Erosion: Implications for Stock and Persistence of Pyrogenic Carbon in Soil, *Frontiers in Earth Science*, 6, 26, <https://doi.org/10.3389/feart.2018.00026>, 2018.
- Abney, R. B., Sanderman, J., Johnson, D., Fogel, M. L., and Berhe, A. A.: Post-Wildfire Erosion in Mountainous Terrain Leads to Rapid and Major Redistribution of Soil Organic Carbon, *Frontiers in Earth Science*, 5, 99, <https://doi.org/10.3389/feart.2017.00099>, 2017.
- Abney, R. B., Kuhn, T. J., Chow, A., Hockaday, W. C., Fogel, M. L., and Berhe, A. A.: Pyrogenic Carbon Erosion after the Rim Fire, Yosemite National Park: The Role of Burn Severity and Slope, *Journal of Geophysical Research: Biogeosciences*, 124, 432–449, <https://doi.org/10.1029/2018JG004787>, 2019.
- Agarwal, T. and Bucheli, T. D.: Adaptation, Validation and Application of the Chemo-Thermal Oxidation Method to Quantify Black Carbon in Soils, *Environmental Pollution*, 159, 532–538, <https://doi.org/10.1016/j.envpol.2010.10.012>, 2011.
- Alexis, M., Rasse, D., Knicker, H., Anquetil, C., and Rumpel, C.: Evolution of Soil Organic Matter after Prescribed Fire: A 20-Year Chronosequence, *Geoderma*, 189–190, 98–107, <https://doi.org/10.1016/j.geoderma.2012.05.003>, 2012.
- Andreeva, D., Leiber, K., Glaser, B., Hambach, U., Erbajeva, M., Chimitdorgieva, G., Tashak, V., and Zech, W.: Genesis and Properties of Black Soils in Buryatia, Southeastern Siberia, Russia, *Quaternary International*, 243, 313–326, <https://doi.org/10.1016/j.quaint.2010.12.017>, 2011.
- Antoine, A.: *Les bocages de la France de l'Ouest*, Belin, Paris, p. 447, ISBN 978-2-7011-2382-0, 2000.
- Ascough, P., Bompard, N., Garnett, M. H., Gulliver, P., Murray, C., Newton, J.-A., and Taylor, C.: ^{14}C Measurement of Samples for Environmental Science Applications at the National Environmental Isotope Facility (NEIF) Radiocarbon Laboratory, SUERC, UK, *Radiocarbon*, 66, 1020–1031, <https://doi.org/10.1017/RDC.2024.9>, 2024.
- Astill, G. and Davies, W.: *A Breton Landscape*, UCL Press, ISBN 1-85728-452-6, 1997.
- Azzi, E. S., Li, H., Cederlund, H., Karlton, E., and Sundberg, C.: Modelling biochar long-term carbon storage in soil with harmonized analysis of decomposition data, *Geoderma*, 441, 116761, <https://doi.org/10.1016/j.geoderma.2023.116761>, 2024.
- Barré, P., Plante, A. F., Cécillon, L., Lutfalla, S., Baudin, F., Bernard, S., Christensen, B. T., Eglin, T., Fernandez, J. M., Houot, S., Kätterer, T., Le Guillou, C., Macdonald, A., Van Oort, F., and Chenu, C.: The Energetic and Chemical Signatures of Persistent Soil Organic Matter, *Biogeochemistry*, 130, 1–12, <https://doi.org/10.1007/s10533-016-0246-0>, 2016.
- Belcher, C. M. (Ed.): *Fire Phenomena and the Earth System: An Interdisciplinary Guide to Fire Science*, John Wiley & Sons, Ltd, <https://doi.org/10.1002/9781118529539.ch11>, 2013.
- Bellé, S.-L.: *The Fate of Organic and Pyrogenic Carbon in Tropical Plant-Soil Systems across Spatial and Temporal Scales*, PhD thesis, Universität Zürich, Zürich, 2023.
- Bellé, S.-L., Berhe, A. A., Hagedorn, F., Santin, C., Schiedung, M., van Meerveld, I., and Abiven, S.: Key drivers of pyrogenic carbon redistribution during a simulated rainfall event, *Biogeosciences*, 18, 1105–1126, <https://doi.org/10.5194/bg-18-1105-2021>, 2021.
- Berhe, A. A., Barnes, R. T., Six, J., and Marín-Spiotta, E.: Role of Soil Erosion in Biogeochemical Cycling of Essential Elements: Carbon, Nitrogen, and Phosphorus, *Annual Review of Earth and Planetary Sciences*, 46, 521–548, <https://doi.org/10.1146/annurev-earth-082517-010018>, 2018.
- Bird, M. I., Wynn, J. G., Saiz, G., Wurster, C. M., and McBeath, A.: The Pyrogenic Carbon Cycle, *Annual Review of Earth and Planetary Sciences*, 43, 273–298, <https://doi.org/10.1146/annurev-earth-060614-105038>, 2015.
- Bonhage, A., Raab, T., Schneider, A., Fischer, T., Ramezany, S., Ouimet, W., Raab, A., and Hirsch, F.: Vertical SOC Distribution and Aromatic Carbon in Centuries Old Charcoal-rich Technosols, *European Journal of Soil Science*, 73, e13293, <https://doi.org/10.1111/ejss.13293>, 2022.

- Boot, C. M., Haddix, M., Paustian, K., and Cotrufo, M. F.: Distribution of black carbon in ponderosa pine forest floor and soils following the High Park wildfire, *Biogeosciences*, 12, 3029–3039, <https://doi.org/10.5194/bg-12-3029-2015>, 2015.
- Bordenave, P. and Merceron, M.: Présentation des bassins versants ateliers (Coët-Dan et Kerharo) et de la baie de Douarnenez, in: *Pollutions diffuses: du bassin versant au littoral*, Ploufragan, 23–24 Septembre 1999, vol. 24, 107–124, France, <https://archimer.ifremer.fr/doc/00000/2841/>, 1999.
- Bosq, M., Kreutzer, S., Bertran, P., Lanos, P., Dufresne, P., and Schmidt, C.: Last Glacial loess in Europe: luminescence database and chronology of deposition, *Earth Syst. Sci. Data*, 15, 4689–4711, <https://doi.org/10.5194/essd-15-4689-2023>, 2023.
- Bowring, S. P., Jones, M. W., Ciais, P., Guenet, B., and Abiven, S.: Pyrogenic carbon decomposition critical to resolving fire's role in the Earth system, *Nature Geoscience*, 15, 135–142, <https://doi.org/10.1038/s41561-021-00892-0>, 2022.
- Briard, J., Gebhardt, A., Marguerie, D., and Querré, G.: Archéologie et environnement en forêt de Broceliande, un exemple d'études pluridisciplinaires, *Bulletin de la Société préhistorique française*, 86, 397–403, <https://doi.org/10.3406/bspf.1989.9899>, 1989.
- Brodowski, S., Rodionov, A., Haumaier, L., Glaser, B., and Amelung, W.: Revised Black Carbon Assessment Using Benzene Polycarboxylic Acids, *Organic Geochemistry*, 36, 1299–1310, <https://doi.org/10.1016/j.orggeochem.2005.03.011>, 2005.
- Brodowski, S., Amelung, W., Haumaier, L., and Zech, W.: Black Carbon Contribution to Stable Humus in German Arable Soils, *Geoderma*, 139, 220–228, <https://doi.org/10.1016/j.geoderma.2007.02.004>, 2007.
- Butnor, J. R., Samuelson, L. J., Johnsen, K. H., Anderson, P. H., González Benecke, C. A., Boot, C. M., Cotrufo, M. F., Heckman, K. A., Jackson, J. A., Stokes, T. A., and Zarnoch, S. J.: Vertical Distribution and Persistence of Soil Organic Carbon in Fire-Adapted Longleaf Pine Forests, *Forest Ecology and Management*, 390, 15–26, <https://doi.org/10.1016/j.foreco.2017.01.014>, 2017.
- Carcaillet, C.: Soil Particles Reworking Evidences by AMS ^{14}C Dating of Charcoal, *Comptes Rendus de l'Académie des Sciences - Series IIA – Earth and Planetary Science*, 332, 21–28, [https://doi.org/10.1016/S1251-8050\(00\)01485-3](https://doi.org/10.1016/S1251-8050(00)01485-3), 2001.
- Caria, G., Arrouays, D., Dubromel, E., Jolivet, C., Ratié, C., Bernoux, M., Barthès, B. G., Brunet, D., and Grinand, C.: Black Carbon Estimation in French Calcareous Soils Using Chemo-thermal Oxidation Method, *Soil Use and Management*, <https://doi.org/10.1111/j.1475-2743.2011.00349.x>, 333–339, 2011.
- Chaplot, V., Van Vliet-Lanoë, B., Walter, C., Curmi, P., and Cooper, M.: Soil Spatial Distribution in the Armorican Massif, Western France: Effect of Soil-Forming Factors, *Soil Science*, 168, 856–868, <https://doi.org/10.1097/01.ss.0000106405.84926.f9>, 2003.
- Chaplot, V. A. M., Rumpel, C., and Valentin, C.: Water Erosion Impact on Soil and Carbon Redistributions within Uplands of Mekong River, *Global Biogeochemical Cycles*, 19, 2005GB002493, <https://doi.org/10.1029/2005GB002493>, 2005.
- Chassé, M., Lutfalla, S., Cécillon, L., Baudin, F., Abiven, S., Chenu, C., and Barré, P.: Long-term bare-fallow soil fractions reveal thermo-chemical properties controlling soil organic carbon dynamics, *Biogeosciences*, 18, 1703–1718, <https://doi.org/10.5194/bg-18-1703-2021>, 2021.
- Cheng, C.-H., Lehmann, J., Thies, J. E., and Burton, S. D.: Stability of Black Carbon in Soils across a Climatic Gradient, *Journal of Geophysical Research: Biogeosciences*, 113, 2007JG000642, <https://doi.org/10.1029/2007JG000642>, 2008.
- Cotrufo, M. F., Boot, C., Abiven, S., Foster, E. J., Haddix, M., Reisser, M., Wurster, C. M., Bird, M. I., and Schmidt, M. W.: Quantification of Pyrogenic Carbon in the Environment: An Integration of Analytical Approaches, *Organic Geochemistry*, 100, 42–50, <https://doi.org/10.1016/j.orggeochem.2016.07.007>, 2016a.
- Cotrufo, M. F., Boot, C. M., Kampf, S., Nelson, P. A., Brogan, D. J., Covino, T., Haddix, M. L., MacDonald, L. H., Rathburn, S., Ryan-Bukett, S., Schmeer, S., and Hall, E.: Redistribution of Pyrogenic Carbon from Hillslopes to Stream Corridors Following a Large Montane Wildfire, *Global Biogeochemical Cycles*, 30, 1348–1355, <https://doi.org/10.1002/2016GB005467>, 2016b.
- Courchesne, F. and Turmel, M.-C.: Extractable Al, Fe, Mn, and Si, in: *Soil sampling and methods of analysis*, CRC press, Boca Raton (Fla.), ISBN 978-0-8493-3586-0, 2008.
- Cros-Cayot, S.: Distribution spatiale des transferts de surface à l'échelle du versant. Contexte armoricain, Thèse de doctorat, Ecole Nationale Supérieure Agronomique de Rennes, Rennes, 1996.
- Curmi, P., Walter, C., Hallaire, V., Gascuel-Odoux, C., Widiatmaka, Taha, A., and Zida, M.: Les sols du bassin versant du Coët Dan: caractéristiques hydrodynamiques des volumes pédologiques, *Science update*, Institut national de la recherche agronomique, Paris, 1998.
- Czimczik, C. I., Schmidt, M. W. I., and Schulze, E.-D.: Effects of Increasing Fire Frequency on Black Carbon and Organic Matter in Podzols of Siberian Scots Pine Forests, *European Journal of Soil Science*, 56, 417–428, <https://doi.org/10.1111/j.1365-2389.2004.00665.x>, 2005.
- Dai, X., Boutton, T., Glaser, B., Ansley, R., and Zech, W.: Black Carbon in a Temperate Mixed-Grass Savanna, *Soil Biology and Biochemistry*, 37, 1879–1881, <https://doi.org/10.1016/j.soilbio.2005.02.021>, 2005.
- Davies, G. M., Vandvik, V., Marrs, R., and Velle, L. G.: Fire Management in Heather-Dominated Heaths and Moorlands of North-West Europe, in: *Global Application of Prescribed Fire*, edited by: Weir, J. R. and Scasta, J. D., CRC Press, ISBN 978-1-4863-1248-1, 2022.
- Direction départementale des finances publiques: Naizin – 1952 – Section B, in: *Cadastre – Plans rénovés*, edited by: Duflos, M. and Michelli, M., Patrimoine et Archives – Département du Morbihan, 2020.
- Eckmeier, E., Egli, M., Schmidt, M., Schlumpf, N., Nötzli, M., Minikus-Stary, N., and Hagedorn, F.: Preservation of Fire-Derived Carbon Compounds and Sorptive Stabilisation Promote the Accumulation of Organic Matter in Black Soils of the Southern Alps, *Geoderma*, 159, 147–155, <https://doi.org/10.1016/j.geoderma.2010.07.006>, 2010.
- Elmqvist, M., Gustafsson, Ö., and Andersson, P.: Quantification of Sedimentary Black Carbon Using the Chemothermal Oxidation Method: An Evaluation of Ex Situ Pretreatments and Standard Additions Approaches, *Limnology and Oceanography: Methods*, 2, 417–427, <https://doi.org/10.4319/lom.2004.2.417>, 2004.

- Forrest, M., Hetzer, J., Billing, M., Bowring, S. P. K., Koscor, E., Oberhagemann, L., Perkins, O., Warren, D., Arrogante-Funes, F., Thonicke, K., and Hickler, T.: Understanding and simulating cropland and non-cropland burning in Europe using the BASE (Burnt Area Simulator for Europe) model, *Biogeosciences*, 21, 5539–5560, <https://doi.org/10.5194/bg-21-5539-2024>, 2024.
- Galanter, A., Cadol, D., and Lohse, K.: Geomorphic Influences on the Distribution and Accumulation of Pyrogenic Carbon (PyC) Following a Low Severity Wildfire in Northern New Mexico, *Earth Surface Processes and Landforms*, 43, 2207–2218, <https://doi.org/10.1002/esp.4386>, 2018.
- Gaudin, L., Bernard, V., and Marguerie, D.: Forêts, friches, landes, marais... et cultures dans l'ouest de la Gaule: dynamique spatio-temporelle des données archéobotaniques au début du Subatlantique dans le massif Armoricaïn, in: *Silva et saltus en Gaule Romaine: Dynamique et gestion des forêts et des zones rurales marginales*, edited by: Bernard, V., Favory, F., and Fiches, J.-L., Presses universitaires de Franche-Comté, pp. 81–89, <https://doi.org/10.4000/books.pufc.8353>, 2014.
- Gavin, D. G., Brubaker, L. B., and Lertzman, K. P.: Holocene Fire History of a Coastal Temperate Rain Forest Based on Soil Charcoal Radiocarbon Dates, *Ecology*, 84, 186–201, [https://doi.org/10.1890/0012-9658\(2003\)084\[0186:HFHOAC\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2003)084[0186:HFHOAC]2.0.CO;2), 2023.
- Gebhardt, A. and Marguerie, D.: II. Les sols, leur couvert végétal et leur utilisation au Néolithique, *Gallia Préhistoire*, 38, 2006.
- Girona-García, A., Vieira, D., Doerr, S., Panagos, P., and Santín, C.: Into the Unknown: The Role of Post-fire Soil Erosion in the Carbon Cycle, *Global Change Biology*, 30, e17354, <https://doi.org/10.1111/gcb.17354>, 2024.
- Güereña, D. T., Lehmann, J., Walter, T., Enders, A., Neufeldt, H., Odiwour, H., Biwott, H., Recha, J., Shepherd, K., Barrios, E., and Wurster, C.: Terrestrial Pyrogenic Carbon Export to Fluvial Ecosystems: Lessons Learned from the White Nile Watershed of East Africa, *Global Biogeochemical Cycles*, 29, 1911–1928, <https://doi.org/10.1002/2015GB005095>, 2015.
- Guggenberger, G., Rodionov, A., Shibistova, O., Grabe, M., Kanas, O. A., Fuchs, H., Mikheyeva, N., Zrazhevskaya, G., and Flessa, H.: Storage and Mobility of Black Carbon in Permafrost Soils of the Forest Tundra Ecotone in Northern Siberia, *Global Change Biology*, 14, 1367–1381, <https://doi.org/10.1111/j.1365-2486.2008.01568.x>, 2008.
- Gustafsson, Ö., Haghseta, F., Chan, C., MacFarlane, J., and Gschwend, P. M.: Quantification of the Dilute Sedimentary Soot Phase: Implications for PAH Speciation and Bioavailability, *Environmental Science & Technology*, 31, 203–209, <https://doi.org/10.1021/es960317s>, 1997.
- Gustafsson, Ö., Bucheli, T. D., Kukulska, Z., Andersson, M., Largeau, C., Rouzaud, J.-N., Reddy, C. M., and Eglinton, T. I.: Evaluation of a Protocol for the Quantification of Black Carbon in Sediments, *Global Biogeochemical Cycles*, 15, 881–890, <https://doi.org/10.1029/2000GB001380>, 2001.
- Hajdas, I., Schlumpf, N., Minikus-Stary, N., Hagedorn, F., Eckmeier, E., Schoch, W., Burga, C., Bonani, G., Schmidt, M. W., and Cherubini, P.: Radiocarbon Ages of Soil Charcoals from the Southern Alps, Ticino, Switzerland, *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms*, 259, 398–402, <https://doi.org/10.1016/j.nimb.2007.02.075>, 2007.
- Hammes, K. and Abiven, S.: Identification of Black Carbon in the Earth System, in: *Fire Phenomena and the Earth System*, 1 edn., edited by: Belcher, C. M., Wiley, pp. 157–176, <https://doi.org/10.1002/9781118529539.ch9>, 2013.
- Hammes, K., Schmidt, M. W. I., Smernik, R. J., Currie, L. A., Ball, W. P., Nguyen, T. H., Louchouart, P., Houel, S., Gustafsson, Ö., Elmquist, M., Cornelissen, G., Skjemstad, J. O., Masiello, C. A., Song, J., Peng, P., Mitra, S., Dunn, J. C., Hatcher, P. G., Hockaday, W. C., Smith, D. M., Hartkopf-Fröder, C., Böhrer, A., Lier, B., Huebert, B. J., Amelung, W., Brodowski, S., Huang, L., Zhang, W., Gschwend, P. M., Flores-Cervantes, D. X., Largeau, C., Rouzaud, J.-N., Rumpel, C., Guggenberger, G., Kaiser, K., Rodionov, A., Gonzalez-Vila, F. J., Gonzalez-Perez, J. A., De La Rosa, J. M., Manning, D. A. C., López-Capel, E., and Ding, L.: Comparison of Quantification Methods to Measure Fire-derived (Black/Elemental) Carbon in Soils and Sediments Using Reference Materials from Soil, Water, Sediment and the Atmosphere, *Global Biogeochemical Cycles*, 21, 2006GB002914, <https://doi.org/10.1029/2006GB002914>, 2007.
- Hammes, K., Smernik, R. J., Skjemstad, J. O., and Schmidt, M. W.: Characterisation and evaluation of reference materials for black carbon analysis using elemental composition, colour, BET surface area and ¹³C NMR spectroscopy, *Applied Geochemistry*, 23, 2113–2122, <https://doi.org/10.1016/j.apgeochem.2008.04.023>, 2008a.
- Hammes, K., Torn, M. S., Lapenas, A. G., and Schmidt, M. W. I.: Centennial black carbon turnover observed in a Russian steppe soil, *Biogeosciences*, 5, 1339–1350, <https://doi.org/10.5194/bg-5-1339-2008>, 2008b.
- Hilscher, A. and Knicker, H.: Degradation of Grass-Derived Pyrogenic Organic Material, Transport of the Residues within a Soil Column and Distribution in Soil Organic Matter Fractions during a 28month Microcosm Experiment, *Organic Geochemistry*, 42, 42–54, <https://doi.org/10.1016/j.orggeochem.2010.10.005>, 2011.
- Hobbs, R. and Gimingham, C.: Vegetation, Fire and Herbivore Interactions in Heathland, *Advances in Ecological Research*, 16, 87–173, [https://doi.org/10.1016/S0065-2504\(08\)60088-4](https://doi.org/10.1016/S0065-2504(08)60088-4), 1987.
- Hobley, E.: Vertical Distribution of Soil Pyrogenic Matter: A Review, *Pedosphere*, 29, 137–149, [https://doi.org/10.1016/S1002-0160\(19\)60795-2](https://doi.org/10.1016/S1002-0160(19)60795-2), 2019.
- Hockaday, W. C., Grannas, A. M., Kim, S., and Hatcher, P. G.: Direct Molecular Evidence for the Degradation and Mobility of Black Carbon in Soils from Ultrahigh-Resolution Mass Spectral Analysis of Dissolved Organic Matter from a Fire-Impacted Forest Soil, *Organic Geochemistry*, 37, 501–510, <https://doi.org/10.1016/j.orggeochem.2005.11.003>, 2006.
- IGN: RGE ALTI, 2013.
- IGN: BD ALTI, 2015.
- IGN: Carte de l'Etat-Major (1820–1866), in: BD CARTO® État-major, Institut national de l'information géographique et forestière, 2021.
- IGN and GEOPAL: Photographie Aérienne 1952, 2011.
- IUSS Working Group WRB: World Reference Base for Soil Resources. International Soil Classification System for Naming Soils and Creating Legends for Soil Maps, no. 106 in *World Soil Resources Reports*, 4th edition edn., International Union of Soil Sciences (IUSS), Vienna, Austria, ISBN 978-92-5-108370-3, 2022.

- Jeanneau, L., Jaffrezic, A., Pierson-Wickmann, A.-C., Gruau, G., Lambert, T., and Petitjean, P.: Constraints on the Sources and Production Mechanisms of Dissolved Organic Matter in Soils from Molecular Biomarkers, *Vadose Zone Journal*, 13, 1–9, <https://doi.org/10.2136/vzj2014.02.0015>, 2014.
- Jones, M. W., Santín, C., Van Der Werf, G. R., and Doerr, S. H.: Global Fire Emissions Buffered by the Production of Pyrogenic Carbon, *Nature Geoscience*, 12, 742–747, <https://doi.org/10.1038/s41561-019-0403-x>, 2019.
- Kane, E. S., Kasischke, E. S., Valentine, D. W., Turetsky, M. R., and McGuire, A. D.: Topographic Influences on Wildfire Consumption of Soil Organic Carbon in Interior Alaska: Implications for Black Carbon Accumulation, *Journal of Geophysical Research: Biogeosciences*, 112, 2007JG000458, <https://doi.org/10.1029/2007JG000458>, 2007.
- Kane, E. S., Hockaday, W. C., Turetsky, M. R., Masiello, C. A., Valentine, D. W., Finney, B. P., and Baldock, J. A.: Topographic Controls on Black Carbon Accumulation in Alaskan Black Spruce Forest Soils: Implications for Organic Matter Dynamics, *Biogeochemistry*, 100, 39–56, <https://doi.org/10.1007/s10533-009-9403-z>, 2010.
- Keiluweit, M., Nico, P. S., Johnson, M., and Kleber, M.: Dynamic molecular structure of plant biomass-derived black carbon (biochar), *Environmental Science and Technology*, 44, 1247–1253, <https://doi.org/10.1021/es9031419>, 2010.
- Koele, N., Bird, M. I., Haig, J., Marimon-Junior, B. H., Marimon, B. S., Phillips, O. L., De Oliveira, E. A., Quesada, C., and Feldpausch, T. R.: Amazon Basin Forest Pyrogenic Carbon Stocks: First Estimate of Deep Storage, *Geoderma*, 306, 237–243, <https://doi.org/10.1016/j.geoderma.2017.07.029>, 2017.
- Krull, E. S., Swanston, C. W., Skjemstad, J. O., and McGowan, J. A.: Importance of Charcoal in Determining the Age and Chemistry of Organic Carbon in Surface Soils, *Journal of Geophysical Research: Biogeosciences*, 111, 2006JG000194, <https://doi.org/10.1029/2006JG000194>, 2006.
- Lambert, T., Pierson-Wickmann, A.-C., Gruau, G., Jaffrezic, A., Petitjean, P., Thibault, J.-N., and Jeanneau, L.: Hydrologically Driven Seasonal Changes in the Sources and Production Mechanisms of Dissolved Organic Carbon in a Small Lowland Catchment: Seasonal Changes in Doc Dynamics, *Water Resources Research*, 49, 5792–5803, <https://doi.org/10.1002/wrcr.20466>, 2013.
- Lambert, T., Pierson-Wickmann, A.-C., Gruau, G., Jaffrezic, A., Petitjean, P., Thibault, J. N., and Jeanneau, L.: DOC sources and DOC transport pathways in a small headwater catchment as revealed by carbon isotope fluctuation during storm events, *Biogeosciences*, 11, 3043–3056, <https://doi.org/10.5194/bg-11-3043-2014>, 2014.
- Lebrun Thauront, J., Ascough, P., Haghipour, N., and Walter, C.: Pyrogenic carbon and other soil physico-chemical characteristics in a small agricultural watershed in the west of France, *Zenodo* [data set], <https://doi.org/10.5281/zenodo.17936907>, 2025.
- Lehndorff, E., Roth, P. J., Cao, Z. H., and Amelung, W.: Black Carbon Accrual during 2000 Years of Paddy-Rice and Non-Paddy Cropping in the Yangtze River Delta, China, *Global Change Biology*, 20, 1968–1978, <https://doi.org/10.1111/gcb.12468>, 2014.
- Lehndorff, E., Houtermans, M., Winkler, P., Kaiser, K., Kölbl, A., Romani, M., Said-Pullicino, D., Utami, S., Zhang, G., Cao, Z., Mikutta, R., Guggenberger, G., and Amelung, W.: Black Carbon and Black Nitrogen Storage under Long-Term Paddy and Non-Paddy Management in Major Reference Soil Groups, *Geoderma*, 284, 214–225, <https://doi.org/10.1016/j.geoderma.2016.08.026>, 2016.
- Leifeld, J., Fenner, S., and Müller, M.: Mobility of black carbon in drained peatland soils, *Biogeosciences*, 4, 425–432, <https://doi.org/10.5194/bg-4-425-2007>, 2007.
- Leng, L., Xu, X., Wei, L., Fan, L., Huang, H., Li, J., Lu, Q., Li, J., and Zhou, W.: Biochar stability assessment by incubation and modelling: Methods, drawbacks and recommendations, *Science of the Total Environment*, 664, 11–23, <https://doi.org/10.1016/j.scitotenv.2019.01.298>, 2019.
- Liang, B., Lehmann, J., Solomon, D., Sohi, S., Thies, J. E., Skjemstad, J. O., Luizão, F. J., Engelhard, M. H., Neves, E. G., and Wirrick, S.: Stability of Biomass-Derived Black Carbon in Soils, *Geochimica et Cosmochimica Acta*, 72, 6069–6078, <https://doi.org/10.1016/j.gca.2008.09.028>, 2008.
- Lutfalla, S., Abiven, S., Barré, P., Wiedemeier, D. B., Christensen, B. T., Houot, S., Kätterer, T., Macdonald, A. J., Van Oort, F., and Chenu, C.: Pyrogenic Carbon Lacks Long-Term Persistence in Temperate Arable Soils, *Frontiers in Earth Science*, 5, 96, <https://doi.org/10.3389/feart.2017.00096>, 2017.
- Maestrini, B., Abiven, S., Singh, N., Bird, J., Torn, M. S., and Schmidt, M. W. I.: Carbon losses from pyrolysed and original wood in a forest soil under natural and increased N deposition, *Biogeosciences*, 11, 5199–5213, <https://doi.org/10.5194/bg-11-5199-2014>, 2014.
- Major, J., Lehmann, J., Rondon, M., and Goodale, C.: Fate of Soil-applied Black Carbon: Downward Migration, Leaching and Soil Respiration, *Global Change Biology*, 1366–1379, <https://doi.org/10.1111/j.1365-2486.2009.02044.x>, 2010.
- Masiello, C. A. and Berhe, A. A.: First Interactions with the Hydrologic Cycle Determine Pyrogenic Carbon's Fate in the Earth System, *Earth Surface Processes and Landforms*, 45, 2394–2398, <https://doi.org/10.1002/esp.4925>, 2020.
- Matosziuk, L. M., Gallo, A., Hatten, J., Bladon, K. D., Ruud, D., Bowman, M., Egan, J., Heckman, K., SanClements, M., Strahm, B., and Weiglein, T.: Short-Term Effects of Recent Fire on the Production and Translocation of Pyrogenic Carbon in Great Smoky Mountains National Park, *Frontiers in Forests and Global Change*, 3, 6, <https://doi.org/10.3389/ffgc.2020.00006>, 2020.
- McGuire, L. A., Rasmussen, C., Youberg, A. M., Sanderan, J., and Fenerty, B.: Controls on the Spatial Distribution of Near-Surface Pyrogenic Carbon on Hillslopes 1 Year Following Wildfire, *Journal of Geophysical Research: Earth Surface*, 126, e2020JF005996, <https://doi.org/10.1029/2020JF005996>, 2021.
- Mehra, O. P. and Jackson, M. L.: Iron Oxide Removal from Soils and Clays by a Dithionite-Citrate System Buffered with Sodium Bicarbonate, in: *Clays and Clay Minerals*, Elsevier, Washington DC, pp. 317–327, <https://doi.org/10.1016/B978-0-08-009235-5.50026-7>, 1960.
- Meredith, W., Ascough, P. L., Bird, M. I., Large, D., Snape, C., Sun, Y., and Tilston, E.: Assessment of Hydropyrolysis as a Method for the Quantification of Black Carbon Using Standard Reference Materials, *Geochimica et Cosmochimica Acta*, 97, 131–147, <https://doi.org/10.1016/j.gca.2012.08.037>, 2012.
- Ministère des Finances, Direction départementale des contributions directes, Bureau du cadastre: Naizin – 1833 – Section B de Penvern et Section E du Boterf, in: *Plans du cadastre dit*

- “Napoléonien” dans le Morbihan, edited by: Chiron, H. and Pellerin, D., Patrimoine et Archives – Département du Morbihan, 2007.
- Nam, J. J., Gustafsson, O., Kurt-Karakus, P., Breivik, K., Steinnes, E., and Jones, K. C.: Relationships between Organic Matter, Black Carbon and Persistent Organic Pollutants in European Background Soils: Implications for Sources and Environmental Fate, *Environmental Pollution*, 156, 809–817, <https://doi.org/10.1016/j.envpol.2008.05.027>, 2008.
- Nesbitt, H. W. and Young, G. M.: Early Proterozoic Climates and Plate Motions Inferred from Major Element Chemistry of Lutites, *Nature*, 299, 715–717, <https://doi.org/10.1038/299715a0>, 1982.
- Nguyen, B. T., Lehmann, J., Kinyangi, J., Smernik, R., Riha, S. J., and Engelhard, M. H.: Long-Term Black Carbon Dynamics in Cultivated Soil, *Biogeochemistry*, 92, 163–176, 2009.
- Nicolay, R. E., Mkhize, N. R., Tedder, M. J., and Kirkman, K. P.: Fire Suppression Interacts with Soil Acidity to Maintain Stable Recalcitrant Pyrogenic Carbon Fractions in South African Mesic Grasslands Soil, *African Journal of Range & Forage Science*, 41, 204–212, <https://doi.org/10.2989/10220119.2024.2355909>, 2024.
- O’Hagan, A., Cox, M., and Wright, L.: Anomalous Behaviour of the Welch-Satterthwaite Approximation, https://tonyohagan.co.uk/academic/pdf/WS_Anomaly_JAS.pdf (last access: May 2025), 2021.
- Ohlson, M., Dahlberg, B., Økland, T., Brown, K. J., and Halvorsen, R.: The Charcoal Carbon Pool in Boreal Forest Soils, *Nature Geoscience*, 2, 692–695, <https://doi.org/10.1038/ngeo617>, 2009.
- Paroissien, J.-B., Orton, T. G., Saby, N. P. A., Martin, M. P., Jolivert, C. C., Ratie, C., Caria, G., and Arrouays, D.: Mapping Black Carbon Content in Topsoils of Central France, *Soil Use and Management*, 28, 488–496, <https://doi.org/10.1111/j.1475-2743.2012.00452.x>, 2012.
- Pellerin, J. and Van Vliet-Lanoe, B.: Le du bassin versant du Coët-Dan au coeur du massif armoricain. 2. Analyse cartographique de la région de Naizin, in: *Agriculture intensive et qualité des eaux*, edited by: Cheverry, C., Science update, Institut national de la recherche agronomique, Paris, ISBN 978-2-7380-0801-5, 1998.
- Pignatello, J. J., Uchimiya, M., Abiven, S., and Schmidt, M. W. I.: *Evolution of Biochar Properties in Soil*, 2nd edn., Routledge, London, 2015.
- Poeplau, C., Vos, C., and Don, A.: Soil organic carbon stocks are systematically overestimated by misuse of the parameters bulk density and rock fragment content, *SOIL*, 3, 61–66, <https://doi.org/10.5194/soil-3-61-2017>, 2017.
- Pyle, L. A., Magee, K. L., Gallagher, M. E., Hockaday, W. C., and Masiello, C. A.: Short-Term Changes in Physical and Chemical Properties of Soil Charcoal Support Enhanced Landscape Mobility, *Journal of Geophysical Research: Biogeosciences*, 122, 3098–3107, <https://doi.org/10.1002/2017JG003938>, 2017.
- Qi, F., Naidu, R., Bolan, N. S., Dong, Z., Yan, Y., Lamb, D., Bucheli, T. D., Choppala, G., Duan, L., and Semple, K. T.: Pyrogenic Carbon in Australian Soils, *Science of The Total Environment*, 586, 849–857, <https://doi.org/10.1016/j.scitotenv.2017.02.064>, 2017.
- R Core Team: R: A language and environment for statistical computing, <https://www.r-project.org> (last access: 6 January 2026), 2021.
- Rasmussen, P. E., Rickman, R. W., and Douglas, C. L.: Air and Soil Temperatures during Spring Burning of Standing Wheat Stubble, *Agronomy Journal*, 78, 261–263, <https://doi.org/10.2134/agronj1986.00021962007800020009x>, 1986.
- Rasse, D. P., Mulder, J., Moni, C., and Chenu, C.: Carbon Turnover Kinetics with Depth in a French Loamy Soil, *Soil Science Society of America Journal*, 70, 2097–2105, <https://doi.org/10.2136/sssaj2006.0056>, 2006.
- Reimer, P. J., Brown, T. A., and Reimer, R. W.: Discussion: Reporting and Calibration of Post-Bomb ^{14}C Data, *Radiocarbon*, 46, 1299–1304, <https://doi.org/10.1017/S0033822200033154>, 2004.
- Reisser, M., Purves, R. S., Schmidt, M. W. I., and Abiven, S.: Pyrogenic Carbon in Soils: A Literature-Based Inventory and a Global Estimation of Its Content in Soil Organic Carbon and Stocks, *Frontiers in Earth Science*, 4, <https://doi.org/10.3389/feart.2016.00080>, 2016.
- Rodionov, A., Amelung, W., Haumaier, L., Urusevskaja, I., and Zech, W.: Black Carbon in the Zonal Steppe Soils of Russia, *Journal of Plant Nutrition and Soil Science*, 169, 363–369, <https://doi.org/10.1002/jpln.200521813>, 2006.
- Rodionov, A., Amelung, W., Peinemann, N., Haumaier, L., Zhang, X., Kleber, M., Glaser, B., Urusevskaya, I., and Zech, W.: Black Carbon in Grassland Ecosystems of the World, *Global Biogeochemical Cycles*, 24, 2009GB003669, <https://doi.org/10.1029/2009GB003669>, 2010.
- Rstudio: RStudio, <https://posit.co/download/rstudio-desktop/> (last access: 6 January 2026), 2021.
- Rumpel, C., Alexis, M., Chabbi, A., Chaplot, V., Rasse, D., Valentin, C., and Mariotti, A.: Black Carbon Contribution to Soil Organic Matter Composition in Tropical Sloping Land under Slash and Burn Agriculture, *Geoderma*, 130, 35–46, <https://doi.org/10.1016/j.geoderma.2005.01.007>, 2006a.
- Rumpel, C., Chaplot, V., Planchon, O., Bernadou, J., Valentin, C., and Mariotti, A.: Preferential Erosion of Black Carbon on Steep Slopes with Slash and Burn Agriculture, *CATENA*, 65, 30–40, <https://doi.org/10.1016/j.catena.2005.09.005>, 2006b.
- Rumpel, C., Ba, A., Darboux, F., Chaplot, V., and Planchon, O.: Erosion Budget and Process Selectivity of Black Carbon at Meter Scale, *Geoderma*, 154, 131–137, <https://doi.org/10.1016/j.geoderma.2009.10.006>, 2009.
- Rumpel, C., Leifeld, J., Santin, C., and Doerr, S. H.: Movement of Biochar in the Environment, in: *Biochar for environmental management: science, technology and implementation*, edited by: Lehmann, J. and Stephen, J., 2nd edn., Routledge, London, 2015.
- Sanderman, J., Baldock, J. A., Dangal, S. R. S., Ludwig, S., Potter, S., Rivard, C., and Savage, K.: Soil Organic Carbon Fractions in the Great Plains of the United States: An Application of Mid-Infrared Spectroscopy, *Biogeochemistry*, 156, 97–114, <https://doi.org/10.1007/s10533-021-00755-1>, 2021.
- Santín, C., Doerr, S. H., Kane, E. S., Masiello, C. A., Ohlson, M., De La Rosa, J. M., Preston, C. M., and Dittmar, T.: Towards a Global Assessment of Pyrogenic Carbon from Vegetation Fires, *Global Change Biology*, 22, 76–91, <https://doi.org/10.1111/gcb.12985>, 2016.

- Santos, F., Wagner, S., Rothstein, D., Jaffe, R., and Miesel, J. R.: Impact of a Historical Fire Event on Pyrogenic Carbon Stocks and Dissolved Pyrogenic Carbon in Spodosols in Northern Michigan, *Frontiers in Earth Science*, 5, 80, <https://doi.org/10.3389/feart.2017.00080>, 2017.
- Santos, F., Bird, J. A., and Asefaw Berhe, A.: Dissolved Pyrogenic Carbon Leaching in Soil: Effects of Soil Depth and Pyrolysis Temperature, *Geoderma*, 424, 116011, <https://doi.org/10.1016/j.geoderma.2022.116011>, 2022.
- Sass, O. and Kloss, S.: Distribution of Macro Charcoal from Forest Fires in Shallow Soils of the Northern Alps, *Journal of Soils and Sediments*, 15, 748–758, <https://doi.org/10.1007/s11368-014-0954-9>, 2015.
- Schiedung, M., Bellè, S.-L., Sigmund, G., Kalbitz, K., and Abiven, S.: Vertical mobility of pyrogenic organic matter in soils: a column experiment, *Biogeosciences*, 17, 6457–6474, <https://doi.org/10.5194/bg-17-6457-2020>, 2020.
- Schiedung, M., Bellè, S.-L., Hoeschen, C., Schweizer, S. A., and Abiven, S.: Enhanced Loss but Limited Mobility of Pyrogenic and Organic Matter in Continuous Permafrost-Affected Forest Soils, *Soil Biology and Biochemistry*, 178, 108959, <https://doi.org/10.1016/j.soilbio.2023.108959>, 2023.
- Schiedung, M., Ascough, P. L., Bellè, S.-L., Bird, M. I., Bröder, L., Haghipour, N., Hilton, R. G., Lattaud, J., and Abiven, S.: Millennial-Aged Pyrogenic Carbon in High-Latitude Mineral Soils, *Communications Earth & Environment*, 5, 177, <https://doi.org/10.1038/s43247-024-01343-5>, 2024.
- Schmidt, M. W. I., Torn, M. S., Abiven, S., Dittmar, T., Guggenberger, G., Janssens, I. A., Kleber, M., Kögel-Knabner, I., Lehmann, J., Manning, D. A. C., Nannipieri, P., Rasse, D. P., Weiner, S., and Trumbore, S. E.: Persistence of Soil Organic Matter as an Ecosystem Property, *Nature*, 478, 49–56, <https://doi.org/10.1038/nature10386>, 2011.
- Selvalakshmi, S., De La Rosa, J. M., Zhijun, H., Guo, F., and Ma, X.: Effects of Ageing and Successive Slash-and-Burn Practice on the Chemical Composition of Charcoal and Yields of Stable Carbon, *CATENA*, 162, 141–147, <https://doi.org/10.1016/j.catena.2017.11.028>, 2018.
- Silva, L. J. D., Oliveira, D. M. D. S., Nóbrega, G. N., Barbosa, R. I., and Cordeiro, R. C.: Pyrogenic Carbon Stocks and Its Spatial Variability in Soils from Savanna-Forest Ecotone in Amazon, *Journal of Environmental Management*, 340, 117980, <https://doi.org/10.1016/j.jenvman.2023.117980>, 2023.
- Singh, B. P., Fang, Y., Boersma, M., Collins, D., Van Zwieten, L., and Macdonald, L. M.: In Situ Persistence and Migration of Biochar Carbon and Its Impact on Native Carbon Emission in Contrasting Soils under Managed Temperate Pastures, *PLOS ONE*, 10, 1–20, <https://doi.org/10.1371/journal.pone.0141560>, 2015.
- Singh, N., Abiven, S., Maestrini, B., Bird, J. A., Torn, M. S., and Schmidt, M. W. I.: Transformation and Stabilization of Pyrogenic Organic Matter in a Temperate Forest Field Experiment, *Global Change Biology*, 20, 1629–1642, <https://doi.org/10.1111/gcb.12459>, 2014.
- Skjemstad, J., Clarke, P., Taylor, J., Oades, J., and Mcclure, S.: The Chemistry and Nature of Protected Carbon in Soil, *Soil Research*, 34, 251, <https://doi.org/10.1071/SR9960251>, 1996.
- Smil, V.: Crop Residues: Agriculture's Largest Harvest, *BioScience*, 49, 299–308, <https://doi.org/10.2307/1313613>, 1999.
- Sokol, N. W., Kuebbing, S. E., Karlsen-Ayala, E., and Bradford, M. A.: Evidence for the Primacy of Living Root Inputs, Not Root or Shoot Litter, in Forming Soil Organic Carbon, *New Phytologist*, 221, 233–246, <https://doi.org/10.1111/nph.15361>, 2019.
- Solomon, D., Lehmann, J., Wang, J., Kinyangi, J., Heymann, K., Lu, Y., Wirrick, S., and Jacobsen, C.: Micro- and Nano-Environments of C Sequestration in Soil: A Multi-Elemental STXM-NEXAFS Assessment of Black C and Organomineral Associations, *Science of The Total Environment*, 438, 372–388, <https://doi.org/10.1016/j.scitotenv.2012.08.071>, 2012.
- Sorrenti, G., Masiello, C. A., Dugan, B., and Toselli, M.: Biochar Physico-Chemical Properties as Affected by Environmental Exposure, *Science of The Total Environment*, 563–564, 237–246, <https://doi.org/10.1016/j.scitotenv.2016.03.245>, 2016.
- Soucémariadin, L., Reisser, M., Cécillon, L., Barré, P., Nicolas, M., and Abiven, S.: Pyrogenic Carbon Content and Dynamics in Top and Subsoil of French Forests, *Soil Biology and Biochemistry*, 133, 12–15, <https://doi.org/10.1016/j.soilbio.2019.02.013>, 2019.
- Soucémariadin, L. N., Quideau, S. A., and MacKenzie, M. D.: Pyrogenic Carbon Stocks and Storage Mechanisms in Podzolic Soils of Fire-Affected Quebec Black Spruce Forests, *Geoderma*, 217–218, 118–128, <https://doi.org/10.1016/j.geoderma.2013.11.010>, 2014.
- Synal, H.-A., Stocker, M., and Suter, M.: MICADAS: A New Compact Radiocarbon AMS System, *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms*, 259, 7–13, <https://doi.org/10.1016/j.nimb.2007.01.138>, 2007.
- Tonnerre, N.-Y.: Villages et Villageois Dans La Bretagne Du Haut Moyen Âge, in: *Villages et Villageois Au Moyen-Âge*, edited by: Société des historiens médiévistes de l'enseignement supérieur public, Editions de la Sorbonne, ISBN 978-2-85944-220-0, 1992.
- Torn, M. S., Swanston, C. W., Castanha, C., and Trumbore, S.: Storage and Turnover of Organic Matter in Soil, in: *Biophysico-Chemical Processes Involving Natural Nonliving Organic Matter in Environmental Systems*, edited by: Senesi, N., Xing, B., and Huang, P. M., Wiley, <https://doi.org/10.1002/9780470494950>, 2009.
- Van Oost, K. and Six, J.: Reconciling the paradox of soil organic carbon erosion by water, *Biogeosciences*, 20, 635–646, <https://doi.org/10.5194/bg-20-635-2023>, 2023.
- Van Vliet-Lanoe, B., Hallegouet, B., and Monnier, J. L.: The Quaternary of Brittany – Guide Book of the Excursion of the Quaternary Research Association in Brittany, 12–15 September 1997, vol. Special volume of *Travaux Du Laboratoire d'Anthropologie*, Université Rennes 1, 1997.
- Van Vliet-Lanoé, B., Pellerin, J., and Chauvel, J.: Le du bassin versant du Coët-Dan au coeur du massif armoricain. 1. Le cadre géologique et géomorphologique, in: *Agriculture intensive et qualité des eaux*, edited by: Cheverry, C., Science update, Institut national de la recherche agronomique, Paris, 1998.
- Vasilyeva, N. A., Abiven, S., Milanovskiy, E. Y., Hilf, M., Rzhkov, O. V., and Schmidt, M. W.: Pyrogenic Carbon Quantity and Quality Unchanged after 55 Years of Organic Matter Depletion in a Chernozem, *Soil Biology and Biochemistry*, 43, 1985–1988, <https://doi.org/10.1016/j.soilbio.2011.05.015>, 2011.

- Velasco-Molina, M., Berns, A. E., Macías, F., and Knicker, H.: Biochemically Altered Charcoal Residues as an Important Source of Soil Organic Matter in Subsoils of Fire-Affected Subtropical Regions, *Geoderma*, 262, 62–70, <https://doi.org/10.1016/j.geoderma.2015.08.016>, 2016.
- Vongvixay, A., Grimaldi, C., Dupas, R., Fovet, O., Birgand, F., Gilliet, N., and Gascuel-Oudou, C.: Contrasting Suspended Sediment Export in Two Small Agricultural Catchments: Cross-influence of Hydrological Behaviour and Landscape Degradation or Stream Bank Management, *Land Degradation & Development*, 29, 1385–1396, <https://doi.org/10.1002/ldr.2940>, 2018.
- Wagner, S., Jaffé, R., and Stubbins, A.: Dissolved Black Carbon in Aquatic Ecosystems, *Limnology and Oceanography Letters*, 3, 168–185, <https://doi.org/10.1002/lol2.10076>, 2018.
- Walter, C. and Curmi, P.: Les sols du bassin versant du Coët-Dan: organisation, variabilité spatiale et cartographie, in: *Agriculture intensive et qualité des eaux*, edited by: Cheverry, C., Science update, Institut national de la recherche agronomique, Paris, 1998.
- Wei, T. and Simko, V.: R package “corrplot”: Visualization of a Correlation Matrix, <https://github.com/taiyun/corrplot> (last access: 6 January 2026), 2021.
- Wickham, H., Averick, M., Bryan, J., Chang, W., D’Agostino McGowan, L., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T. L., Miller, E., Milton Bache, S., Müller, K., Ooms, J., Robinson, D., Seidel, D. P., Spinu, V., Takahashi, K., Vaughan, D., Wilke, C., Woo, K., and Yutani, H.: Welcome to the tidyverse, *Journal of Open Source Software*, 4, 1686, <https://doi.org/10.21105/joss.01686>, 2019.
- Wiedemeier, D. B., Abiven, S., Hockaday, W. C., Keiluweit, M., Kleber, M., Masiello, C. A., McBeath, A. V., Nico, P. S., Pyle, L. A., Schneider, M. P., Smernik, R. J., Wiesenberger, G. L., and Schmidt, M. W.: Aromaticity and Degree of Aromatic Condensation of Char, *Organic Geochemistry*, 78, 135–143, <https://doi.org/10.1016/j.orggeochem.2014.10.002>, 2015.
- Wurster, C. M., Saiz, G., Schneider, M. P., Schmidt, M. W., and Bird, M. I.: Quantifying pyrogenic carbon from thermosequences of wood and grass using hydrogen pyrolysis, *Organic Geochemistry*, 62, 28–32, <https://doi.org/10.1016/j.orggeochem.2013.06.009>, 2013.
- Zheng, H., Miao, C., Huntingford, C., Tarolli, P., Li, D., Panagos, P., Yue, Y., Borrelli, P., and Van Oost, K.: The Impacts of Erosion on the Carbon Cycle, *Reviews of Geophysics*, 63, e2023RG000829, <https://doi.org/10.1029/2023RG000829>, 2025.