



Supplement of

Tree island area in oil palm agroforests directly and indirectly drives evaporative fraction

Thorge Wintz et al.

Correspondence to: Thorge Wintz (twintz@uni-goettingen.de)

The copyright of individual parts of the supplement might differ from the article licence.

S1. Environmental conditions on selected flight days

Table S1: Environmental conditions on selected flight days (high incoming radiation) vs. other days with available data. Median [IQR] for soil moisture on flight days, aggregated precipitation for 1, 3 and 7 days before selected flight day, air temperature and wind speed on flight days.

Variable	Selected high-radiation flights (n = 6)	Other usable flight days (n = 8)
Soil moisture (%)	32.88 [32.64–33.00]	33.19 [32.71–33.68]
Precipitation preceding 1 d (mm)	0.05 [0.00–1.75]	0.25 [0.00–2.20]
Precipitation preceding 3 d (mm)	3.35 [2.52–4.62]	3.45 [1.42–5.72]
Precipitation preceding 7 d (mm)	8.20 [8.20–43.90]	57.95 [44.30–72.10]
Air temperature, 10–14 UTC+7 (°C)	32.54 [31.70–32.79]	31.28 [30.98–31.96]
Wind speed, 10–14 UTC+7 (m s ⁻¹)	0.60 [0.58–0.61]	0.69 [0.58–0.72]

S2. Structural equation modelling




Figure S1: Conceptual model describing the potential drivers influencing Evaporative fraction.

25

30

Table S2: Mechanistic framework linking studied variables with evaporative fraction.

Path	Relationship	Mechanisms	Prediction
A	Planted tree diversity ↓ EF	I - A higher tree diversity creates complementary effects that lead to a more efficient use of energy and therefore higher EF.	A higher (planted / spontaneous) diversity increases EF (Bigelow, 2001; González-Espinosa et al., 2004; Wang et al., 2021).
O	Observed woody diversity ↓ EF		
B	Island area ↓ EF	I - In a larger tree island, the effect of trees over oil palm is stronger, creating a more forest-like microclimate. II - Greater edge length in larger tree islands increases the area influenced by edge effects, where EF is typically higher.	In larger tree islands, EF is higher (Cregg and Dix, 2001).
C	Planted Tree diversity ↓ Observed woody plant diversity	I - Planted trees attract animals that spread seeds (zoochory) and thereby facilitate natural regeneration. II - Planted trees modify environmental conditions, create niches that support natural regeneration.	Observed woody plant diversity is higher in islands with higher planted tree diversity (Paterno et al., 2016; Tinya et al., 2019).

E	Planted tree diversity ↓ Structural complexity	I - Different species occupy different parts of the growing space.	Structural diversity in oil palm increases through higher planted tree diversity and observed woody plant diversity (Zemp et al., 2019; Kikuchi et al., 2024).
G	Observed woody plant diversity ↓ Structural complexity		
I	Observed woody plant diversity ↓ Tree height variability	I - A higher diversity of woody plants also results in a higher morphological diversity, focusing here on tree height.	Higher observed woody plant diversity leads to higher tree height variability.
J	Island area ↓ Tree height variability	I – Larger tree islands harbor a greater diversity of tree species. II – Higher tree diversity results in an increased variability of tree height.	Higher diversity in larger tree islands (Drakare et al., 2006) leads to higher tree height variability in these tree islands (Marks et al., 2016).
L	Island area ↓ Vegetation health	I - Larger tree islands contribute to more homogeneous microclimatic conditions, thereby fostering enhanced vegetation health. II - Within larger tree islands, there is a greater richness of tree species, thereby amplifying vegetation health.	Larger tree islands increase vegetation health.
D	Island area ↓ Observed woody plant diversity	I - Bigger islands receive more seeds from outside. II - Bigger islands harbor greater tree diversity by providing more niches (Island Biogeography theory).	Larger tree islands have more observed woody plant species than smaller tree islands (Zahawi and Augspurger, 2006; Holl et al., 2020; Zemp et al., 2023; Paterno et al., 2024).
F	Island area ↓	I - Larger tree islands provide greater opportunities for spatial diversification.	Larger tree island have higher structural complexity (Zemp et al., 2019).

	Structural complexity		
H	Planted tree diversity ↓ Tree height variability	I - Higher tree diversity increases tree height variability due to interspecific differences in morphology and growth strategy.	Higher planted tree diversity increases variability in tree height (Marks et al., 2016).
K	Planted tree diversity ↓ Vegetation health	I - A higher diversity of planted trees enhances vegetation health, as a greater number of species increases the potential for beneficial interactions among species.	Vegetation health increases with greater planted tree diversity (Cayuela et al., 2006; Pau et al., 2012; Mapfumo et al., 2016).
M	Structural complexity ↓ EF	I - In a more complex island, there are more layers to capture radiation and transform it into latent heat. II – More complex structures provide more surface area, which increases ET.	Higher structural complexity increases EF (Ehbrecht et al., 2017; Ren et al., 2018; Forzieri et al., 2020; Wang et al., 2021).
N	Tree height variability ↓ EF	I - Higher variability in tree height increases surface roughness. II - Higher surface roughness facilitates heat fluxes.	Higher tree height variability increases EF (Chen et al., 2020; Barbeta et al., 2023).
P	Vegetation health ↓ EF	I - Low health of vegetation reduces the ability to convert radiation into latent heat.	Higher vegetation health increases EF (Yang and Wang, 2011; Er-Raki et al., 2013; Zhou and Wang, 2016).

35

40

S3. Decomposition of SEM effects

45 **Table S3: Estimates are standardized effects from the SEM. Confidence intervals are 95% bias-corrected and accelerated (BCa) intervals based on nonparametric bootstrapping (1000 iterations). Individual indirect effects are pathway-specific products of standardized path coefficients; the combined indirect effect is the sum of all indirect pathways.**

Pathway / effect	Estimate	95% BCa CI
Overall effects		
Combined indirect effect	-0.044	-0.213 to 0.103
Direct effect: island area → EF	0.439	0.281 to 0.639
Total effect of island area on EF	0.395	0.314 to 0.532
Individual indirect pathways		
Island area → structural complexity (SSCI) → EF	-0.066	-0.185 to -0.014
Island area → observed woody plant diversity → EF	-0.051	-0.229 to 0.087
Island area → GNDVI → EF	0.002	-0.019 to 0.053
Island area → height variability → EF	0.009	-0.010 to 0.059
Island area → observed woody plant diversity → structural complexity (SSCI) → EF	0.045	0.010 to 0.130
Island area → observed woody plant diversity → height variability → EF	0.016	-0.015 to 0.077

50 **S4. Scale independence of EF**

Workflow:

1. Select 400 m² (n = 13) and 1600 m² (n = 13) tree islands.
2. Apply negative buffer to simulate smaller tree islands:
 - a. For 1600 m² tree islands:
 - 55 i. 10 m buffer → 400 m²
 - ii. 15 m buffer → 100 m²
 - iii. 17.5 m buffer → 25 m²
 - b. For 400 m² tree islands:
 - i. 5 m buffer → 100 m²
 - 60 ii. 7.5 m buffer → 25 m²

3. For each original plot size group:
 - c. Fit a linear model: $EF \sim \text{size}$
 - d. Fit a mixed-effects model: $EF \sim \text{size} + (1|\text{ID})$ using tree island ID as random effect

65 **Table S4: Scale independence of EF**

Model	Slope	SE	R ² (adj.)	p - value	Random effect variance
1600 m ²	0.0004	0.00062	- 0.012	0.52	
Linear					
1600 m ²	0.0004	0.00029		0.17	0.003
Linear mixed effects					
400 m ²	0.0016	0.00142	0.009	0.26	
Linear					
400 m ²	0.0016	0.001		0.12	0.001
Linear mixed effects					

S5. LST min, LST mean and LST max vs. plot size

- 70 Workflow:
1. Use tree island polygons to clip land surface temperature (LST) orthomosaics.
 2. For each tree island, extract:
 - a. Mean LST of all pixels
 - b. Mean of the 1% highest LST values
 - 75 c. Mean of the 0.5% lowest LST values
 3. Test for differences between tree island area groups using the Kruskal-Wallis test (non-parametric).




Figure S2: Mean land surface temperature of the 0.5% coldest pixels within each tree island, grouped by tree island area. No significant differences were found between island area groups (Kruskal-Wallis test, $p = 0.48$).




Figure S3: Mean land surface temperature of all pixels within each tree island, grouped by tree island area. No significant differences were found between island area groups (Kruskal-Wallis test, $p = 0.5$).




Figure S4: Mean land surface temperature of the 1% hottest pixels within each tree island, grouped by tree island area. No significant differences were found between island area groups (Kruskal-Wallis test, $p = 0.48$).


S6. Edge gradients of EF

Workflow:


1. Generate concentric bands (0.5 m width) for each tree island of 100 m², 400 m² and 1600 m².
2. Calculate the area of each band.
3. Use the bands as spatial masks to extract EF values from QWater Model raster outputs.
4. Fit a linear mixed-effects model with edge distance as fixed effect, tree island ID as a random effect and band area as weights:
(EF ~ edge distance + (1|Plot), weights = 1 / area).

100 **Table S5: Mixed random effects model results.**


Tree island area	Edge gradient	EF ~ Distance + (1 Plot), weights = 1/area	Marginal R ²	Conditional R ²
100 m ² + 400 m ² + 1600 m ²	0 – 5 m	Distance: n.s.		
400 m ² + 1600 m ²	0 – 10 m	Distance: p < 0.05	0.011	0.943
100 m ²	0 – 4.5 m	Distance: n.s.		
400 m ²	0 – 9.5 m	Distance: p < 0.001	0.074	0.934
1600 m ²	0 – 17.5 m	Distance: p < 0.05	0.003	0.993



105 **Figure S5: Predicted evaporative fraction for 400 m² and 1600 m² as function of edge distance. Predictions based on a linear mixed-effects model with distance as fixed effect and tree island as random intercept (EF = 0.748 + -0.0021 * Distance). EF decreased significantly with distance (-0.0021 ± 0.001 SE, p = 0.0386). The model explained 1.1% of the variance with fixed effects (marginal R²), and 94.3% including random effects (conditional R²).**



110 **Figure S6: Predicted evaporative fraction for 400 m² as function of edge distance. Predictions based on a linear mixed-effects model with distance as fixed effect and tree island as random intercept ($EF = 0.746 + -0.0052 * \text{Distance}$). EF decreased significantly with distance (-0.0052 ± 0.00125 SE, $p = 0.00004$). The model explained 7% of the variance with fixed effects (marginal R^2), and 93.4% including random effects (conditional R^2).**



115 **Figure S7: Predicted evaporative fraction for 1600 m² as function of edge distance. Predictions based on a linear mixed-effects model with distance as fixed effect and tree island as random intercept ($EF = 0.769 + -6e-04 * \text{Distance}$). EF decreased significantly with distance (-0.00058 ± 0.00024 SE, $p = 0.016$). The model explained 0.3% of the variance with fixed effects (marginal R^2), and 99.3% including random effects (conditional R^2).**

120

References

- Barbeta, A., Miralles, D. G., Mendiola, L., Gimeno, T. E., Sabaté, S., and Carnicer, J.: Disentangling the Role of Forest Structure and Functional Traits in the Thermal Balance of the Mediterranean–Temperate Ecotone, *JGR Biogeosciences*, 125 128, <https://doi.org/10.1029/2022JG007264>, 2023.
- Bigelow, S.: Evapotranspiration modelled from stands of three broad-leaved tropical trees in Costa Rica, *Hydrological Processes*, 15, 2779–2796, <https://doi.org/10.1002/hyp.268>, 2001.
- Cayuela, L., Benayas, J. M. R., Justel, A., and Salas-Rey, J.: Modelling tree diversity in a highly fragmented tropical montane landscape, *Global Ecology and Biogeography*, 15, 602–613, <https://doi.org/10.1111/j.1466-8238.2006.00255.x>, 130 2006.
- Chen, J., Jin, S., and Du, P.: Roles of horizontal and vertical tree canopy structure in mitigating daytime and nighttime urban heat island effects, *International Journal of Applied Earth Observation and Geoinformation*, 89, 102060, <https://doi.org/10.1016/j.jag.2020.102060>, 2020.
- Cregg, B. and Dix, M.: Tree Moisture Stress and Insect Damage in Urban Areas in Relation to Heat Island Effects, *AUF*, 27, 135 8–17, <https://doi.org/10.48044/jauf.2001.002>, 2001.
- Drakare, S., Lennon, J. J., and Hillebrand, H.: The imprint of the geographical, evolutionary and ecological context on species-area relationships, *Ecology letters*, 9, 215–227, <https://doi.org/10.1111/j.1461-0248.2005.00848.x>, 2006.
- Ehbrecht, M., Schall, P., Ammer, C., and Seidel, D.: Quantifying stand structural complexity and its relationship with forest management, tree species diversity and microclimate, *Agricultural and Forest Meteorology*, 242, 1–9, 140 <https://doi.org/10.1016/j.agrformet.2017.04.012>, 2017.
- Er-Raki, S., Rodriguez, J. C., Garatuza-Payan, J., Watts, C. J., and Chehbouni, A.: Determination of crop evapotranspiration of table grapes in a semi-arid region of Northwest Mexico using multi-spectral vegetation index, *Agricultural Water Management*, 122, 12–19, <https://doi.org/10.1016/j.agwat.2013.02.007>, 2013.
- Forzieri, G., Miralles, D. G., Ciais, P., Alkama, R., Ryu, Y., Duveiller, G., Zhang, K., Robertson, E., Kautz, M., Martens, B., 145 Jiang, C., Arneth, A., Georgievski, G., Li, W., Ceccherini, G., Anthoni, P., Lawrence, P., Wiltshire, A., Pongratz, J., Piao, S., Sitch, S., Goll, D. S., Arora, V. K., Lienert, S., Lombardozzi, D., Kato, E., Nabel, J. E. M. S., Tian, H., Friedlingstein, P., and Cescatti, A.: Increased control of vegetation on global terrestrial energy fluxes, *Nat. Clim. Chang.*, 10, 356–362, <https://doi.org/10.1038/s41558-020-0717-0>, 2020.
- González-Espinosa, M., María Rey-Benayas, J., Ramírez-Marcial, N., Huston, M. A., and Golicher, D.: Tree diversity in the northern Neotropics: regional patterns in highly diverse Chiapas, Mexico, *Ecography*, 27, 741–756, 150 <https://doi.org/10.1111/j.0906-7590.2004.04103.x>, 2004.
- Holl, K. D., Reid, J. L., Cole, R. J., Oviedo-Brenes, F., Rosales, J. A., and Zahawi, R. A.: Applied nucleation facilitates tropical forest recovery: Lessons learned from a 15-year study, *Journal of Applied Ecology*, 57, 2316–2328, <https://doi.org/10.1111/1365-2664.13684>, 2020.

- 155 Kikuchi, T., Seidel, D., Ehbrecht, M., Zemp, D. C., Brambach, F., Irawan, B., Sundawati, L., Hölscher, D., Kreft, H., and Paterno, G. B.: Combining planting trees and natural regeneration promotes long-term structural complexity in oil palm landscapes, *Forest Ecology and Management*, 569, 122182, <https://doi.org/10.1016/j.foreco.2024.122182>, 2024.
- Mapfumo, R. B., Murwira, A., Masocha, M., and Andriani, R.: The relationship between satellite-derived indices and species diversity across African savanna ecosystems, *International Journal of Applied Earth Observation and Geoinformation*, 52, 306–317, <https://doi.org/10.1016/j.jag.2016.06.025>, 2016.
- 160 Marks, C. O., Muller-Landau, H. C., and Tilman, D.: Tree diversity, tree height and environmental harshness in eastern and western North America, *Ecology letters*, 19, 743–751, <https://doi.org/10.1111/ele.12608>, 2016.
- Paterno, G. B., Brambach, F., Guerrero-Ramírez, N., Zemp, D. C., Cantillo, A. F., Camarretta, N., Moura, C. C. M., Gailing, O., Ballauff, J., Polle, A., Schlund, M., Erasmi, S., Iddris, N. A., Khokthong, W., Sundawati, L., Irawan, B., Hölscher, D., and Kreft, H.: Diverse and larger tree islands promote native tree diversity in oil palm landscapes, *Science* (New York, N.Y.), 386, 795–802, <https://doi.org/10.1126/science.ado1629>, 2024.
- 165 Paterno, G. B., Siqueira Filho, J. A., and Ganade, G.: Species-specific facilitation, ontogenetic shifts and consequences for plant community succession, *J Vegetation Science*, 27, 606–615, <https://doi.org/10.1111/jvs.12382>, 2016.
- Pau, S., Gillespie, T. W., and Wolkovich, E. M.: Dissecting NDVI–species richness relationships in Hawaiian dry forests, *Journal of Biogeography*, 39, 1678–1686, <https://doi.org/10.1111/j.1365-2699.2012.02731.x>, 2012.
- 170 Ren, Z., He, X., Pu, R., and Zheng, H.: The impact of urban forest structure and its spatial location on urban cool island intensity, *Urban Ecosyst*, 21, 863–874, <https://doi.org/10.1007/s11252-018-0776-4>, 2018.
- Tinya, F., Márialigeti, S., Bidló, A., and Ódor, P.: Environmental drivers of the forest regeneration in temperate mixed forests, *Forest Ecology and Management*, 433, 720–728, <https://doi.org/10.1016/j.foreco.2018.11.051>, 2019.
- 175 Wang, X., Dallimer, M., Scott, C. E., Shi, W., and Gao, J.: Tree species richness and diversity predicts the magnitude of urban heat island mitigation effects of greenspaces, *The Science of the total environment*, 770, 145211, <https://doi.org/10.1016/j.scitotenv.2021.145211>, 2021.
- Yang, J. and Wang, Y.: Estimating evapotranspiration fraction by modeling two-dimensional space of NDVI/albedo and day–night land surface temperature difference: A comparative study, *Advances in Water Resources*, 34, 512–518, <https://doi.org/10.1016/j.advwatres.2011.01.006>, 2011.
- 180 Zahawi, R. A. and Augspurger, C. K.: Tropical Forest Restoration: Tree Islands As Recruitment Foci In Degraded Lands Of Honduras, *Ecological applications a publication of the Ecological Society of America*, 16, 464–478, [https://doi.org/10.1890/1051-0761\(2006\)016\[0464:TFRTIA\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2006)016[0464:TFRTIA]2.0.CO;2), 2006.
- Zemp, D. C., Guerrero-Ramirez, N., Brambach, F., Darras, K., Grass, I., Potapov, A., Röhl, A., Arimond, I., Ballauff, J., 185 Behling, H., Berkelmann, D., Biagioni, S., Buchori, D., Craven, D., Daniel, R., Gailing, O., Ellsäßer, F., Fardiansah, R., Hennings, N., Irawan, B., Khokthong, W., Krashevskaya, V., Krause, A., Kückes, J., Li, K., Lorenz, H., Maraun, M., Merk, M. S., Moura, C. C. M., Mulyani, Y. A., Paterno, G. B., Pebrianti, H. D., Polle, A., Di Prameswari, A., Sachsenmaier, L., Scheu, S., Schneider, D., Setiajiati, F., Setyaningsih, C. A., Sundawati, L., Tschardtke, T., Wollni, M.,

- Hölscher, D., and Kreft, H.: Tree islands enhance biodiversity and functioning in oil palm landscapes, *Nature*, 618, 316–
190 321, <https://doi.org/10.1038/s41586-023-06086-5>, 2023.
- Zemp, D. C., Ehbrecht, M., Seidel, D., Ammer, C., Craven, D., Erkelenz, J., Irawan, B., Sundawati, L., Hölscher, D., and
Kreft, H.: Mixed-species tree plantings enhance structural complexity in oil palm plantations, *Agriculture, Ecosystems
& Environment*, 283, 106564, <https://doi.org/10.1016/j.agee.2019.06.003>, 2019.
- Zhou, C. and Wang, K.: Biological and Environmental Controls on Evaporative Fractions at AmeriFlux Sites, *Journal of*
195 *Applied Meteorology and Climatology*, 55, 145–161, <https://doi.org/10.1175/JAMC-D-15-0126.1>, 2016.