

# The Arctic Ocean marine carbon cycle: evaluation of air-sea CO<sub>2</sub> exchanges, ocean acidification impacts and potential feedbacks

N. R. Bates<sup>1</sup> and J. T. Mathis<sup>2</sup>

<sup>1</sup>Bermuda Institute of Ocean Sciences, Ferry Reach, Bermuda

<sup>2</sup>School of Fisheries and Ocean Sciences, University of Alaska Fairbanks, Fairbanks, Alaska, USA

Received: 8 June 2009 – Published in Biogeosciences Discuss.: 30 June 2009

Revised: 25 September 2009 – Accepted: 29 September 2009 – Published: 5 November 2009

**Abstract.** At present, although seasonal sea-ice cover mitigates atmosphere-ocean gas exchange, the Arctic Ocean takes up carbon dioxide (CO<sub>2</sub>) on the order of –66 to –199 Tg C year<sup>–1</sup> (10<sup>12</sup> g C), contributing 5–14% to the global balance of CO<sub>2</sub> sinks and sources. Because of this, the Arctic Ocean has an important influence on the global carbon cycle, with the marine carbon cycle and atmosphere-ocean CO<sub>2</sub> exchanges sensitive to Arctic Ocean and global climate change feedbacks. In the near-term, further sea-ice loss and increases in phytoplankton growth rates are expected to increase the uptake of CO<sub>2</sub> by Arctic Ocean surface waters, although mitigated somewhat by surface warming in the Arctic. Thus, the capacity of the Arctic Ocean to uptake CO<sub>2</sub> is expected to alter in response to environmental changes driven largely by climate. These changes are likely to continue to modify the physics, biogeochemistry, and ecology of the Arctic Ocean in ways that are not yet fully understood. In surface waters, sea-ice melt, river runoff, cooling and uptake of CO<sub>2</sub> through air-sea gas exchange combine to decrease the calcium carbonate (CaCO<sub>3</sub>) mineral saturation states ( $\Omega$ ) of seawater while seasonal phytoplankton primary production (PP) mitigates this effect. Biological amplification of ocean acidification effects in subsurface waters, due to the remineralization of organic matter, is likely to reduce the ability of many species to produce CaCO<sub>3</sub> shells or tests with profound implications for Arctic marine ecosystems

2002; Serreze and Francis, 2006; Overland et al., 2008), sea-ice loss (Cavaliere et al., 2003; Maslanik et al., 2007; Shimada et al., 2007; Giles et al., 2008; Comiso et al., 2008), and other physical changes (Wu et al., 2006; McGuire et al., 2006) as well as biology and ecosystem structure changes (Arrigo et al., 2008; Pabi et al., 2008). These changes and feedbacks could have profound impacts on the Arctic Ocean marine carbon cycle and the importance of the Arctic for the global carbon cycle and the balance of carbon dioxide (CO<sub>2</sub>) sinks and sources. In a recent review, Macdonald et al. (2009) compiled inorganic and organic carbon budgets of the Arctic Ocean. Here, we review the present understanding of the marine inorganic carbon cycle, air-sea CO<sub>2</sub> disequilibrium and rates of air-sea CO<sub>2</sub> gas exchange, the physical and biological processes that influence Arctic Ocean CO<sub>2</sub> sinks and sources, the impact of ocean acidification and the potential future drivers of change such as sea-ice loss, phytoplankton primary production (PP) and freshwater inputs. The geographic scope of this review encompasses the Arctic Ocean shelves (i.e. Barents, Kara, Laptev, East Siberian, Chukchi, Beaufort and Canadian Archipelago Seas) and central basin (i.e. Canada and Eurasian Basin), but does not extend to the gateways of the Arctic Ocean including the Greenland, Icelandic, Norwegian, and Bering Seas, and Hudson and Baffin Bays.


As context for this paper, a brief background on the physical and biogeochemical setting is presented in Sect. 2. Relevant information about historical sampling in the Arctic Ocean, methods for estimating air-sea CO<sub>2</sub> exchange rates, considerations about carbonate chemistry and relationships between shelf metabolism (net autotrophy versus net heterotrophy) and CO<sub>2</sub> sink/source issues are summarized in Sect. 3. Also included are considerations of present carbon pools and fluxes, and physical and biological processes within the Arctic Ocean that influence the inorganic carbon cycle (Sect. 3.5). With a synthesis of published results and data, the present status and magnitude of the Arctic Ocean

## 1 Introduction

The Arctic Ocean is currently experiencing rapid environmental change due to natural and anthropogenic factors that include warming (e.g. Serreze et al., 2000; Polyakov et al.,



Correspondence to: N. R. Bates  
(nick.bates@bios.edu)



**Fig. 1.** Schematic of the Arctic Ocean, central basin (Canada and Eurasian Basins) and Arctic continental shelves (with approximate boundaries of each Arctic shelf sea), and major rivers draining into the region. General surface circulation features are also shown. Inset: Characterization of Arctic Ocean continental shelves as “inflow”, “interior” or “outflow” shelves according to Carmack and Wassmann (2006).

CO<sub>2</sub> sink on the Arctic shelves and central basin is estimated, with predictions for near-term CO<sub>2</sub> flux rates in an era of rapid change (Sect. 4). The impacts of rapid change in the Arctic Ocean and the potential feedbacks on the marine inorganic carbon cycle and air-sea CO<sub>2</sub> exchange rate are also discussed in Sect. 5. Vulnerabilities of the Arctic marine carbon cycle relate to such factors as sea-ice loss and warming, biological and ecosystem changes, and hydrological/freshwater input changes. We also discuss the impact of ocean acidification, and biological and physical processes on the marine carbon cycle and calcium carbonate (CaCO<sub>3</sub>) mineral saturation states in the Arctic Ocean (Sect. 6).

## 2 The physical and biological setting of the Arctic Ocean

The relatively small Arctic Ocean ( $\sim 10.7 \times 10^6$  km<sup>2</sup>) is almost completely landlocked except for the Bering Strait, Canadian Archipelago, Fram Strait and Norwegian Sea gateways that allow exchanges with the Pacific and Atlantic Oceans (Fig. 1). The relatively broad, generally shallow (<200 m deep) and almost annular continental shelves surrounding the central basin comprise about 53% of the area of the Arctic Ocean, making it unique compared to other ocean basins. All of the Arctic Ocean continental shelves are highly varied, unique and difficult to characterize generically.

As a framework for this review, the Chukchi and Barents Seas can be characterized as “inflow” shelves (Carmack

and Wassmann, 2006) where the import of nutrient-rich water from the Pacific and Atlantic Oceans, respectively, sustains high rates of seasonal phytoplankton PP or net community production (NCP). Other shelves such as the Kara, Laptev, East Siberian and Beaufort Seas constitute “interior shelves”, which are highly influenced by exchanges with other shelves, while the Canadian Archipelago represents an “outflow shelf” where Arctic Ocean water is exported to the Atlantic Ocean via Hudson and Baffin Bays. In the central basin of the Arctic Ocean, waters of the Canada Basin or Beaufort Gyre are separated from the Eurasian Basin by the surface transpolar drift, with inputs and outputs of Atlantic Ocean water through the Fram Strait and shelf-basin exchanges of water across the entire Arctic Ocean. Central basin waters are highly stratified vertically with distinctly different physical and chemical properties of surface waters and subsurface halocline, Atlantic Ocean layer and deep-water layers, with deep waters having limited exchanges with the Atlantic Ocean.

The Arctic Ocean has an important role in the global freshwater cycle (e.g. Aagaard and Carmack, 1989; Wijffels et al., 1992; Woodgate and Aagaard, 2005) and Atlantic Ocean overturning circulation (e.g. Walsh and Chapman, 1990; Mysak et al., 1990; Hakkinen, 1993; Aagaard and Carmack, 1994; Wadley and Bigg, 2002). Surrounding the Arctic Ocean is an extensive land margin and watershed that contributes freshwater and material inputs such as inorganic nutrients and dissolved organic carbon (DOC) to the Arctic Ocean (Opsahl et al., 1999; Dittmar and Kattner,

2003; Benner et al., 2003; Guo et al., 2004). Siberian (Ob, Yenisey, Lena and Kolyma) and North American (Mackenzie, Yukon) rivers contribute significantly to land-ocean inputs (Peterson et al., 2002; McGuire et al., 2006; Serreze et al., 2006; Cooper et al., 2008). Finally, air-sea interaction controls the seasonal sea-ice advance and retreat, and inter-annual changes of sea-ice distributions and thickness, and sea-ice export from the Arctic Ocean (Carmack and Chapman, 2003). In wintertime, the Arctic Ocean is completely covered by sea-ice (except for polynyas and leads) with thick (3–7 m) multi-year ice occupying the interior of the central basin and thinner seasonal sea-ice (1–2 m) across the Arctic Ocean shelves and periphery (e.g. significant parts of the Bering Sea shelf; Hudson and Baffin Bays; Greenland, Iceland and Norwegian Sea) that melts and retreats towards the Pole during the summertime. While the seasonal production and melting of sea-ice contributes significantly to the physical structure and mixing of the Arctic, sea-ice melt and brine rejection during sea-ice formation also modifies the biogeochemical properties of surface and halocline layers (Steele and Boyd, 1998; Weingartner et al., 1998; Carmack and Chapman, 2003). The Arctic Ocean sea-ice cover is also particularly sensitive to interannual changes and variability of atmosphere-ocean-sea-ice forcing and feedbacks associated with climate variability such as the Arctic Oscillation (AO) and Arctic Sea-Ice Oscillation (ASIO) (e.g. Rigor et al., 2002; Overland and Wang, 2005; Wang et al., 2005).

### 3 Considerations about sampling, carbonate chemistry, air-sea CO<sub>2</sub> exchange calculations and continental shelf CO<sub>2</sub> sink-sources issues.

#### 3.1 Marine inorganic carbon cycle sampling in the Arctic Ocean


Compared to many other open-ocean and coastal environments, relatively few studies of the marine inorganic carbon cycle and air-sea CO<sub>2</sub> exchange rates have been conducted in the Arctic. The harsh polar climate and difficult logistical support have limited most studies to opportunistic icebreaker surveys conducted on the Arctic shelves during the summertime sea-ice retreat with a couple of transpolar surveys across the deep basin (e.g. Arctic Ocean Section; Tucker and Cate, 1996; Anderson et al., 1998a; Jutterstrom and Anderson, 2005; Jones et al., 2008; Tanhua et al., 2009; Carbon in the North Atlantic (CARINA) datasets at [http://cdiac.ornl.gov/oceans/CARINA/CARINA\\_table.html](http://cdiac.ornl.gov/oceans/CARINA/CARINA_table.html)). It is important to note that spring and summer observations of the Arctic inorganic carbon cycle are highly limited and virtually absent during winter sea-ice cover. The absence of long-term ocean time-series (e.g. Winn et al., 1994; Bates, 2007; Santana-Casiano et al., 2007), repeat hydrographic surveys (e.g. Sabine et al., 2004; Feely et al., 2004) and underway shipboard surface observations (Takahashi et al.,

2002; 2009) that have benefited understanding of seasonal and interannual change in other ocean basins and coastal ocean environments has considerably constrained our knowledge of the marine carbon cycle, and CO<sub>2</sub> sink and source terms in the Arctic Ocean.

#### 3.2 Inorganic carbon chemistry considerations

Studies of the marine inorganic carbon cycle in the Arctic Ocean have focused on measurements of one or more of the four directly observable inorganic carbon system parameters, that is either dissolved inorganic carbon (DIC), total alkalinity (TA), partial pressure of CO<sub>2</sub> (*p*CO<sub>2</sub>), or pH (Dickson et al., 2007). Early historical data of TA and pH were collected in the Arctic Ocean (Yamamoto-Kawai et al., 2005) but have relatively limited use today due to inherent uncertainties associated with the analytical techniques used. More recently, improvement of analytical techniques, the advent of standardized methods (DOE, 1994; Dickson et al., 2007) and use of seawater standards and calibration have allowed precise and accurate data to be collected in the Arctic Ocean, and subsequently used for assessments of stocks, distributions and transformations of the marine inorganic carbon cycle.

Seawater inorganic carbon chemistry is governed by a series of reactions, such that:



For example, DIC is defined as (DOE, 1994; Dickson et al., 2007):

$$\text{DIC} = [\text{CO}_2^*] + [\text{HCO}_3^-] + [\text{CO}_3^{2-}] \quad (\text{R2})$$

where [CO<sub>2</sub>\*] represents the concentration of all unionized carbon dioxide, present either as H<sub>2</sub>CO<sub>3</sub> or as CO<sub>2</sub>. The TA of seawater is defined as:

$$\begin{aligned} \text{TA} = &[\text{HCO}_3^-] + 2[\text{CO}_3^{2-}] + [\text{B}(\text{OH})_4^-] + [\text{OH}^-] + & (\text{R3}) \\ &[\text{HPO}_4^{2-}] + 2[\text{PO}_4^{3-}] + [\text{SiO}(\text{OH})_3^-] + [\text{HS}^-] + [\text{NH}_3] \\ &- [\text{H}^+] - [\text{HSO}_4^-] - [\text{HF}] - [\text{H}_3\text{PO}_4] - \text{minor species} \end{aligned}$$

where [HCO<sub>3</sub><sup>-</sup>] + 2[CO<sub>3</sub><sup>2-</sup>] + B(OH)<sub>4</sub><sup>-</sup> are the primary components of seawater TA. Other species have minor contribution to TA. DIC and TA are expressed as μmoles kg<sup>-1</sup>. *p*CO<sub>2</sub> or fugacity of CO<sub>2</sub> (*f*CO<sub>2</sub>) is expressed as μatm, while pH is negative log of [H<sup>+</sup>] generally expressed on the total seawater pH scale (please refer to Zeebe and Wolf-Gladrow (2001) and Dickson et al. (2007) for further information about the inorganic carbon system).

The complete seawater inorganic carbon system (i.e. [CO<sub>2</sub>\*], [HCO<sub>3</sub><sup>-</sup>], [CO<sub>3</sub><sup>2-</sup>], [H<sup>+</sup>] and CaCO<sub>3</sub> mineral

saturation states,  $\Omega$ ) can be calculated from a combination of any two measured parameters (i.e. DIC, TA,  $p\text{CO}_2$  and pH). In this paper, we used the thermodynamic model of Zeebe and Wolf-Gladrow (2001), temperature and salinity data, the  $\text{CO}_2$  solubility equations of Weiss (1974), and dissociation constants for carbonic acid (i.e.  $pK_1$  and  $pK_2$ ) of Mehrbach et al. (1973; as refit by Dickson and Millero, 1997), borate (Dickson, 1990) and phosphate (DOE, 1994). It should be noted that the experimental determinations of carbonic acid dissociation constants (i.e. Mehrbach et al., 1973; Goyet and Poisson, 1989; Roy et al., 1993) have lower limits of  $-1^\circ\text{C}$  or  $0^\circ\text{C}$ , and that the equations governing  $pK_1$  and  $pK_2$  have to be linearly extrapolated for seawater with temperatures less than  $-0^\circ\text{C}$ . The difference in calculated seawater  $p\text{CO}_2$  for different  $pK_1$  and  $pK_2$  determinations is relatively small ( $<5 \mu\text{atm}$ ) at temperatures less than  $0^\circ\text{C}$  (Bates, 2006). The equations governing  $pK_1$  and  $pK_2$  are also a function of salinity. On the Arctic shelves, salinity can be less than 10–15 in surface waters (upper few metres) locally influenced by sea-ice melt (e.g. Bates et al., 2009; Yamamoto-Kawai et al., 2009; Chierici and Fransson, 2009; Fransson et al., 2009) which can add, for example, uncertainty of  $\sim 10 \mu\text{atm}$  and  $\sim 0.1$  in the calculation of seawater  $p\text{CO}_2$  and  $\Omega$ , respectively.

$\text{CaCO}_3$  mineral production and dissolution is governed by the following chemical reaction:



The  $\text{CaCO}_3$  mineral saturation state ( $\Omega$ ) is defined as the ion product of carbonate and calcium concentrations, thus:

$$\Omega_{\text{aragonite}} = [\text{Ca}^{2+}][\text{CO}_3^{2-}]/K_{sp}^* \quad (\text{R5})$$

The experimental solubility product (i.e.  $K_{sp}^*$ ) of Mucci (1983) was used here. In general,  $\text{CaCO}_3$  production and dissolution rates vary as a function of saturation state for aragonite ( $\Omega_{\text{aragonite}}$ ) or calcite ( $\Omega_{\text{calcite}}$ ). For example, dissolution is generally favored when  $\Omega$  values are lower than 1, while aragonite formation generally occurs at  $\Omega$  values greater than 1.

### 3.3 Air-sea $\text{CO}_2$ exchange rate considerations

Different approaches have been used to estimate rates of air-sea  $\text{CO}_2$  exchange for the Arctic shelves or entire Arctic Ocean. These approaches include: (1) differences in atmospheric and seawater  $p\text{CO}_2$  (either by direct measurements of  $p\text{CO}_2$  or calculated from other carbonate parameters) that gives  $\Delta p\text{CO}_2$  or air-sea  $\text{CO}_2$  disequilibrium and, direction of air-sea  $\text{CO}_2$  gas exchange (e.g. Olsen et al., 2003; Omar et al., 2007; Bates et al., 2006; Fransson et al., 2009); (2) carbon mass balance assessments (e.g. Anderson et al., 1998a; Fransson et al., 2001; Kaltin and Anderson, 2005; Macdonald et al., 2009); and (3) direct measurements of air-sea  $\text{CO}_2$

exchange through eddy correlation techniques (Semiletov et al., 2007).

In the first approach,  $\Delta p\text{CO}_2$  data combined with knowledge of gas transfer velocities commonly parameterized as a function of contemporaneous windspeed data (e.g. Wanninkhof, 1992; Wanninkhof and McGillis, 1999) provide estimates of air-sea  $\text{CO}_2$  exchange rates. The direction of air-sea gas exchange is controlled by  $\Delta p\text{CO}_2$  with the magnitude dictated by gas transfer velocities. If surface water  $p\text{CO}_2$  values are undersaturated with respect to the atmosphere,  $\Delta p\text{CO}_2$  values are negative and the ocean has the potential to uptake  $\text{CO}_2$  (i.e. sink of  $\text{CO}_2$ ). If  $\Delta p\text{CO}_2$  values are positive (i.e. oversaturated) the ocean has the potential to release  $\text{CO}_2$  to the atmosphere (i.e. source of  $\text{CO}_2$ ).

In the second approach, air-sea  $\text{CO}_2$  exchange rates can be calculated by mass balance assessment of carbon stocks and inputs/outputs of carbon (both inorganic and organic), or temporal/depth changes of DIC that account for temperature and salinity variability and such processes as phytoplankton PP and respiration, vertical entrainment/detrainment and vertical diffusion. Both approaches have considerable uncertainties with caveats and assumptions used in the approaches. The third approach, using eddy correlation techniques, has only been used once in the Arctic Ocean (Semiletov et al., 2007). This method provides instantaneous fluxes and often yields different results to the other bulk methods. The results of these different approaches are discussed in more detail in Sect. 4 for individual shelves and the central basin of the Arctic Ocean.

### 3.4 Continental shelf net metabolism and $\text{CO}_2$ sink-sources issues

The  $\text{CO}_2$  source or sink status and net metabolism of the coastal ocean have been highly debated over the last few decades. In early papers, it was first suggested that continental shelves sequestered significant amounts of  $\text{CO}_2$  and that the global coastal ocean was net autotrophic (i.e. net consumption of  $\text{CO}_2$ ) and a significant sink for atmospheric  $\text{CO}_2$  (Walsh et al., 1985; Walsh, 1991). Smith and Hollibaugh (1993) subsequently argued that the nearshore coastal ocean was net heterotrophic (i.e. net production of  $\text{CO}_2$ ) and offshore coastal ocean neutral, and thus continental shelves were potential sources of  $\text{CO}_2$  to the atmosphere. Other studies have since suggested that the  $\text{CO}_2$  source or sink status (Cai and Dai, 2004; Borges, 2005; Borges et al., 2005; Cai et al., 2006; Chen and Borges, 2009), and the metabolic status of continental shelves (i.e. net autotrophy versus net heterotrophy) is highly variable in time and space (e.g. Wollast, 1998; Gattuso et al., 1998).

The net metabolism (either net autotrophy or heterotrophy) of a continental shelf is influenced by a balance of transports, transformation and exchanges, such as: the input of organic matter (OM) from terrestrial sources, OM production and respiration in the coastal ocean, retention or export

of OM from the shelf and, the net air-sea exchange of  $\text{CO}_2$ . The balance of these processes dictates the metabolic status (i.e. net autotrophic or heterotrophic), but can also influence the potential  $\text{CO}_2$  sink or source status of the continental shelf in question. The spatial and temporal variability of seawater  $p\text{CO}_2$  on continental shelves is acted upon by a complex interplay of connected and quasi-independent physical and biological factors. For example, the physicochemical process of air-sea  $\text{CO}_2$  exchange can proceed independently of the biological processes (e.g. production versus consumption of OM) that dictate the metabolism of the coastal ocean. The variability of  $p\text{CO}_2$  or  $\Delta p\text{CO}_2$  and separation of biological and physical processes can lead, for example, to a net heterotrophic continental shelf still being a sink for atmospheric  $\text{CO}_2$  (Ducklow and McAllister, 2005). Ducklow and McAllister (2005) suggested that the global coastal ocean was net autotrophic, and thus a potential sink for atmospheric  $\text{CO}_2$ .

Recent syntheses of coastal carbon cycle studies by Chen and Borges (2009) suggest that in general high-latitude and temperate continental shelf seas tend to be net annual sinks of atmospheric  $\text{CO}_2$ , while upwelling dominated, and subtropical to tropical coastal environments tend to release  $\text{CO}_2$  to the atmosphere (Cai and Dai, 2004). In addition, localized shallow areas influenced by rivers tend to be sources of  $\text{CO}_2$  to the atmosphere, while the deeper, more expansive areas of the continental shelves tend to be oceanic sinks for atmospheric  $\text{CO}_2$  (Cai et al., 2006; Chen and Borges, 2009). In summary, Chen and Borges (2009) suggest that the polar continental shelves, for example, are likely to be potential sinks for atmospheric  $\text{CO}_2$ . However, this potential is also influenced by seasonal sea-ice cover, which acts as a barrier to air-sea  $\text{CO}_2$  gas exchange compared to open water conditions (Bates, 2006).

### 3.5 Carbon pools, fluxes and physical/biological processes in the Arctic Ocean


There are multiple influences on the marine carbon cycle and air-sea exchange of  $\text{CO}_2$  in the Arctic Ocean. The inorganic carbon cycle is dominated by inter-ocean exchanges between the Pacific and Atlantic Oceans (Macdonald et al., 2009) and subsequent biogeochemical modifications and transformations of water while resident in the Arctic Ocean during transit between these oceans. In addition, river inputs of freshwater and materials, sea-ice production and melting, and atmosphere-ocean interaction and exchanges also have profound influence.

The upper waters of the Arctic Ocean contain approximately 25 Pg ( $10^{15}$  g) of inorganic carbon (i.e. in the form of  $[\text{HCO}_3^-]$ ,  $[\text{CO}_3^{2-}]$  and  $[\text{CO}_2^*]$ ; Reactions R1 and R2) and  $\sim 2$  Pg of organic carbon (in the form of living organisms, detritus, and other materials). The fluxes of carbon in and out of the Arctic Ocean are dominated by seawater exchanges with other oceans (Figs. 2 and 3; process 1). Seawater inflow of

Pacific Ocean water through Bering Strait into the Chukchi Sea and Atlantic Ocean water into the Barents Sea supplies a combined inflow of  $\sim 3$  Pg per year of inorganic carbon into the Arctic Ocean with a similar outflow of carbon from the Arctic through Fram Strait. As shown later in Sect. 4, ocean-atmosphere  $\text{CO}_2$  exchanges constitute at present a net air-to-sea flux of  $\sim 66$  to  $199 \text{ Tg C yr}^{-1}$  ( $10^{12}$  g) (Figs. 2 and 3; process 2). River inputs and coastal erosion constitute a net land to ocean inorganic and organic carbon flux of  $\sim 12 \text{ Tg yr}^{-1}$  (Macdonald et al., 2009; Fig. 3; process 3).

Factors influencing inorganic carbon cycling within the Arctic Ocean include physical and biological processes (see Figs. 2 and 3). Processes at the air-sea interface include: exposure of surface waters to gas exchange during seasonal sea-ice retreat (process 4); gas exchange in sea-ice melt (process 5), and river influenced surface waters (process 6); gas-exchange through sea-ice (process 7); exposure of surface waters to gas exchange in polynyas and leads (process 8), and; carbon export via brine rejection during sea-ice formation (process 9). Sea-ice cover has generally been thought of as a barrier to air-sea  $\text{CO}_2$  gas exchange compared to open water conditions (Bates, 2006). Whether sea-ice facilitates the uptake or release of  $\text{CO}_2$  (process 7) is an area of active research with no definitive answer (Gosink et al., 1978; Semiletov et al., 2004; Delille et al., 2007; Nagurnyi, 2008). Processes within the water column important for carbon balance and transformations in the mixed layer include: Cooling of waters during transit northward into the Arctic or temperature change within the Arctic Ocean (process 10); across-shelf transport of water and carbon (process 11); vertical diffusion, entrainment and detrainment through mixing (process 12), and; shelf-to-basin transport (process 13). Biologically mediated processes include: pelagic phytoplankton primary production process 14), and export of organic carbon (process 15) from surface waters to the subsurface. Rates of new and export production have been estimated at  $\sim 135 \text{ Tg C yr}^{-1}$  with large uncertainties (Macdonald et al., 2009). In the subsurface, remineralization of organic matter to  $\text{CO}_2$  in shelf and halocline waters (process 16) and sediments (process 17) can also contribute. There appears to be an active microbial community in the water-column both in summer and winter that contribute to heterotrophic production (i.e. remineralization of OM to  $\text{CO}_2$ ; e.g. Alonso-Saez et al., 2008; Garneau et al., 2008). An additional complication is the recent finding that anaerobic microbial production of, and subsequent release of alkalinity (process 18) can act to decrease seawater  $p\text{CO}_2$ , which in turn may enhance the uptake of atmospheric  $\text{CO}_2$  onto continental shelves where sediments significantly interact with surface waters (Thomas et al., 2009).

## Inflow Shelf (e.g., Chukchi and Barents Sea)



**Fig. 2.** Schematic of processes potentially influencing the inorganic carbon cycle and air-sea  $\text{CO}_2$  gas exchange on “inflow” shelves of the Arctic (e.g. Barents and Chukchi Seas). The two panels represent physical and biological processes likely operating during the summertime sea-ice free period (Panel a), and during the wintertime sea-ice covered period (Panel b). The processes are denoted by numeral with the caveat that the size of arrow does not necessarily reflect magnitude of flux, transport or transformation of  $\text{CO}_2$ . Blue arrows denote processes that likely decrease seawater  $p\text{CO}_2$ , while red arrows processes that likely increase seawater  $p\text{CO}_2$ . Black arrows denote processes that do not impact  $p\text{CO}_2$  directly or reflect uncertainty as to whether the process decreases or increases seawater  $p\text{CO}_2$ . The processes include: 1. northward transport of inorganic carbon; 2. air-sea gas exchange; 4. exposure of surface water to the atmosphere due to sea-ice retreat and melting; 5. localized air-sea gas exchange from surface water highly influenced by sea-ice melt; 7. air-sea gas exchange through sea-ice; 8. winter air-sea gas exchange in leads and polynyas, 9. inorganic carbon flux due to brine-rejection during deep-water formation in fall and winter. 10. cooling of surface waters during northward transport on Atlantic or Pacific Ocean waters into the Arctic Ocean; 11. between shelf transport of water and carbon; 12. redistribution of inorganic carbon between mixed layer and subsurface due to vertical diffusion and vertical entrainment/detrainment due to mixing; 13. shelf-basin exchanges of inorganic carbon (i.e. DIC) and organic carbon due to generalized circulation and eddy mediated transport; 14. net uptake of  $\text{CO}_2$  due to phytoplankton photosynthesis or new production; 15. export flux of organic matter (OM) or export production; 16. remineralization of organic matter back to  $\text{CO}_2$  either in subsurface waters or in sediments; 17. release of  $\text{CO}_2$  from sediments, and; 18. release of alkalinity from sediments due to anaerobic processes in sediments.

### 4 Present state of knowledge about the marine inorganic carbon cycle and air-sea $\text{CO}_2$ exchange rates of the Arctic Ocean


#### 4.1 Arctic Ocean shelves: inflow shelves

##### 4.1.1 Barents Sea shelf

The Barents Sea comprises a broad and shallow continental shelf that is dominated by inflowing Atlantic Ocean water with minimal freshwater inputs (Fig. 1). In the Barents

Sea, warm and salty surface Atlantic Ocean water inflows from the Norwegian Sea into the Barents Sea via the Norwegian Atlantic Current (Skelvan et al., 1999, 2005; Omar et al., 2007), with  $\sim 2\text{ Sv}$  exchanged between the Atlantic Ocean and Barents Sea (Gammelsrod et al., 2008). Cooling of surface water and brine rejection during sea-ice formation is subsequently thought to transform this surface water into dense subsurface water that is subsequently exported to the Kara Sea and central basin of the Arctic Ocean (Anderson et al., 1994; Fransson et al., 2001; Kaltin et al., 2002; Omar et

## Interior Shelf (e.g., Siberian and Beaufort Shelves)



**Fig. 3.** Schematic of processes potentially influencing the inorganic carbon cycle and air-sea  $\text{CO}_2$  gas exchange on “interior” shelves of the Arctic (e.g. Beaufort and Siberian shelf Seas such as the Kara, Laptev and East Siberian Sea). The two panels represent processes likely operating during the summertime sea-ice free period (Panel a), and during the wintertime sea-ice covered period (Panel b). The processes are denoted by numeral with the caveat that the size of arrow does not necessarily reflect magnitude of flux, transport or transformation of  $\text{CO}_2$ . Blue arrows denote processes that likely decrease seawater  $p\text{CO}_2$ , while red arrows processes that likely increase seawater  $p\text{CO}_2$ . Black arrows denote processes that do not impact  $p\text{CO}_2$  directly or reflect uncertainty as to whether the process decreases or increases seawater  $p\text{CO}_2$ . The processes include those described for Figure 2 but also include the following: 3. river input of organic carbon and DIC. This can also include inputs of carbon from coastal erosion of sediments, for example, and; 6. efflux of  $\text{CO}_2$  from river dominated surface waters. Process 10 includes warming of surface waters during sea-ice free period from solar input or lateral transport of warmer waters from upstream “inflow shelf”.

al., 2007) thereby contributing to the physical and chemical properties of the Eurasian Basin halocline and deeper waters (Jones and Anderson, 1986).

In the earliest study of the inorganic carbon cycle in the Arctic Ocean, Kelley (1970) directly observed undersaturated seawater  $p\text{CO}_2$  across the Barents Sea with  $\Delta p\text{CO}_2$  values ranging from  $\sim -50$  to  $-150 \mu\text{atm}$ . More recently, other studies have shown seawater  $p\text{CO}_2$  values of  $\sim 220$ – $350 \mu\text{atm}$  with a seasonal minimum occurring during the summertime (May to October) (Omar et al., 2003; Nakaoka et al., 2006; Omar et al., 2007). Given that the  $\Delta p\text{CO}_2$  values ranged from  $\sim -20$  to  $-110 \mu\text{atm}$ , the Barents Sea has a strong potential for significant uptake of  $\text{CO}_2$  from the atmosphere.

Nakaoka et al. (2006) and Omar et al. (2007) estimated an annual ocean  $\text{CO}_2$  uptake of  $46 \pm 18 \text{ g C m}^{-2} \text{ yr}^{-1}$  and  $51 \pm 8 \text{ g C m}^{-2} \text{ yr}^{-1}$ , respectively, in the Barents Sea (Table 1), accounting for sea-ice cover during the wintertime. Using mass balance approaches, and accounting for the influence of phytoplankton primary production, freshwater inputs and mixing on vertical profiles of DIC and inorganic nutrients, Fransson et al. (2001) computed an annual ocean  $\text{CO}_2$  uptake of  $44 \pm 10 \text{ g C m}^{-2} \text{ yr}^{-1}$ . Kaltin et al. (2002), using similar mass balance approaches applied to inorganic carbon data collected from a section across the Barents Sea estimated an ocean  $\text{CO}_2$  uptake of  $29 + 11 \text{ g C m}^{-2}$  for 3 months of the year (i.e. May to July). For comparison, Takahashi et al. (2009) estimated a climatological mean uptake of  $\sim 5$ – $30 \text{ g C m}^{-2} \text{ yr}^{-1}$  from a very limited dataset of surface

**Table 1.** Areas, depths, residence times, air-sea CO<sub>2</sub> exchange rates expressed in mmol C m<sup>-2</sup> d<sup>-1</sup> and g C m<sup>-2</sup> d<sup>-1</sup>, and annual air-sea CO<sub>2</sub> exchange rate expressed in Tg C (10<sup>12</sup> g C). Negative air-sea CO<sub>2</sub> exchange rates indicates ocean uptake of CO<sub>2</sub> (i.e., CO<sub>2</sub> sink).

	Area <sup>a</sup> (km <sup>2</sup> )	Depth <sup>a</sup> (m)	Residence <sup>a</sup> (years)	Air-Sea CO <sub>2</sub> flux (mmol C m <sup>-2</sup> d <sup>-1</sup> )	Air-Sea CO <sub>2</sub> flux (g C m <sup>-2</sup> yr <sup>-1</sup> )	Annual <sup>f</sup> CO <sub>2</sub> flux (Tg C yr <sup>-1</sup> )	Reference
Barents Sea	1 512 000	200	1 <sup>b</sup>	n/a	-46±18	-70±27	Nakaoka et al. (2006)
				n/a	-51±8	-77±12	Omar et al. 2007
				n/a	-44±10	-67±15	Fransson et al. (2001) <sup>g</sup>
				n/a	-29±11	-44±16	Kaltin et al. (2002) <sup>g</sup>
Kara Sea	926 000	131	2.5	n/a	-1	-1.0	Fransson et al. (2001) <sup>g</sup>
				n/a	n/a	-5.7	Anderson et al. (1998a, b) <sup>g</sup>
Laptev Sea	498 000	48	3.5±2	-2.1	-2.4	-1.2	Nitishinsky et al. (2007) <sup>g</sup>
				n/a	n/a	-4.3	Anderson et al. (1998a, b) <sup>g</sup>
East Siberian Sea	987 000	58	3.5±2	-1 to -10.9	n/a	-1.2/-13	Semiletov et al. (2007)
				+0.3	n/a	+0.3	Nitishinsky et al. (2007) <sup>g</sup>
				n/a	n/a	-5.9	Anderson et al. (1998a, b) <sup>g</sup>
Chukchi Sea	620 000	80	0.2-1.2	-12	-17 <sup>d</sup>	-11	Murata and Takizawa (2003)
				-40±22	-48±10 <sup>d</sup>	-36±6	Bates et al. (2006)
				n/a	-61±10 <sup>e</sup>	-46±6	Bates et al. (2006)
				n/a	-86±22	-53±14	Kaltin and Anderson (2001) <sup>g</sup>
Beaufort Sea	178 000	124	0.5-1.0	n/a	n/a	-2.9	Anderson et al. (1998a, b) <sup>g</sup>
				-12	-17 <sup>d</sup>	-2	Murata and Takizawa (2003)
Canadian Archipelago	1 490 000	290	0.5-12	?	?	-16/-24	scaled to Beaufort Sea shelf
Central basin	4 489 000	2748	2-30 <sup>c</sup>	<-1 to -3	-4 to -12	-6 to -19	Bates et al. (2006)
Arctic Ocean	10 700 000					-65 -175 -81 -199	low estimate from above high estimate from above low estimate from above <sup>k</sup> high estimate from above <sup>k</sup>
Arctic Ocean (other studies)						-129±65	Anderson et al. (1990) <sup>g</sup>
						-70±65	Anderson et al. (1994) <sup>g</sup>
						-110	Lundberg and Haugen (1996) <sup>g,h</sup>
						-24±17	Anderson et al. (1998b) <sup>g</sup>
						-41±18	Anderson et al. (1998b) <sup>g,j</sup>
						-31	Kaltin and Anderson (2005) <sup>g</sup>
						-66	Bates (2006)

Notes: <sup>a</sup> Macdonald et al. (2009) estimates of shelf areas, mean depth (m) and residence times; <sup>b</sup> Kaltin et al. (2002) estimate of Barents Sea shelf water residence time; <sup>c</sup> residence time of surface and halocline waters (upper 0-300 m only) from Macdonald et al. (2009); <sup>d</sup> summertime open water only (~100 d); <sup>e</sup> including wintertime air-sea CO<sub>2</sub> exchange; <sup>f</sup> annual flux estimated from annual uptake (g C yr<sup>-1</sup>) multiplied by area of Arctic shelf from Macdonald et al. (2009); <sup>g</sup> mass balance assessment; <sup>h</sup> includes Norwegian Sea (Lundberg and Haugen, 1996); <sup>j</sup> includes river contribution (Anderson et al., 1998b); <sup>k</sup> includes air-sea CO<sub>2</sub> exchange rates for the Canadian Archipelago scaled to estimates of the Beaufort Sea shelf.

underway *p*CO<sub>2</sub> data, with the oceanic CO<sub>2</sub> sink decreasing in a general trend across the Barents Sea from the Atlantic Ocean inflow to the Kara Sea. Similar spatial trends were also observed by Kelley (1970), with Δ*p*CO<sub>2</sub> values generally increasing from ~-50 μatm close to the coastline to approximately -150 μatm at the outer extent of the Barents Sea shelf.

The seasonal changes in surface *p*CO<sub>2</sub> and air-sea CO<sub>2</sub> exchange rates for the Barents Sea have been largely attributed to seasonal changes in phytoplankton primary production, warming and cooling, and air-sea CO<sub>2</sub> fluxes. In a general sense, cooling of surface waters during transit from the Atlantic Ocean to the Barents Sea throughout the year acts to decrease seawater *p*CO<sub>2</sub> at a thermodynamic rate of ~4.1% per °C (Takahashi et al., 1993; Millero, 1995). However, during the sea-ice free summertime, surface seawater *p*CO<sub>2</sub> decreases by ~100 μatm despite significant seasonal warm-

ing of the surface layer (~4-6°C) and ocean uptake of CO<sub>2</sub> through enhanced rates of air-sea CO<sub>2</sub> exchange. As Kaltin et al. (2002) and Omar et al. (2007) suggest, these processes are more than compensated for by summertime photosynthetic uptake of CO<sub>2</sub> by phytoplankton PP (or new production) in the euphotic zone that acts to decrease seawater DIC and *p*CO<sub>2</sub>. Indeed, the annual rate of net PP for the Barents Sea ranges from ~<40-70 g C m<sup>-2</sup> yr<sup>-1</sup> (Kaltin et al., 2002; Sakshaug, 2004; Macdonald et al., 2009), with rates of phytoplankton PP greater than other Arctic Ocean shelves with the exception of the Chukchi Sea.

The seasonal fall and winter rebound of seawater *p*CO<sub>2</sub> to values just below equilibrium with the atmosphere observed by Nakaoka et al. (2006) likely results from a continued uptake of CO<sub>2</sub> through gas exchange during sea-ice formation and brine rejection (Anderson et al., 2004; Omar et al., 2005) and vertical entrainment by mixing with CO<sub>2</sub> rich

subsurface waters. Despite the absence of supporting data, it is likely that subsurface  $p\text{CO}_2$  increases seasonally from fall and winter from the seasonal remineralization of organic matter, produced during earlier summertime enhanced rates of phytoplankton primary production, as observed in other Arctic Ocean marine environments (e.g. Yager et al., 1995; Bates et al., 2009).

#### 4.1.2 Chukchi Sea shelf

The wide and shallow Chukchi Sea shelf, with the shelf break extending hundreds of kilometers northward from the coast, represents a particularly extensive portion of the Arctic shelf system (Fig. 1). The Bering Strait acts as the gateway for Pacific Ocean waters to enter the Arctic Ocean via the Chukchi Sea (Coachman et al., 1975; Bjork, 1989). The mean annual inflow of Pacific Ocean water at Bering Strait is  $\sim 0.8$  Sv, with higher transport during the open-water sea-ice free season (Roach et al., 1975; Woodgate et al., 2005a, b; Woodgate and Aagaard, 2005), with minor inputs to the Chukchi Sea from the East Siberian Sea shelf via the intermittent Siberian Coastal Current through Long Strait (Weingartner et al., 1999). In general, there is a northward transport of water across the shallow Chukchi Sea shelf, with four major outflows ( $\sim 0.1$ – $0.3$  Sv each; Woodgate et al., 2005a) from the Chukchi Sea into the Canada Basin and Beaufort Sea shelf through Long Strait, Central Channel, Herald Valley and Barrow Canyon (Paquette and Bourke, 1974; Weingartner et al., 1998, 2005; Woodgate et al., 2005b). As with other Arctic Ocean shelves, physical transformations and seasonal sea-ice cover play a major role in shaping the Chukchi Sea shelf water-masses and ecosystem. During the sea-ice covered season, watermasses on the Chukchi Sea shelf are confined to a small range of temperature-salinity space due to vertical homogenization by ventilation, brine rejection and convective mixing (Woodgate et al., 2005a). During the sea-ice free season, local sea-ice melt contributes to Pacific Ocean water to produce a relatively warm, fresher surface mixed layer water (upper 0– $\sim 30$  m, salinity typically  $< 31$ ; temperature  $> -1.5$ ) with denser resident winter water remaining in the subsurface. Inflow of inorganic nutrient rich Pacific water supports high rates of phytoplankton PP or NCP on the Chukchi Sea shelf compared to other Arctic Ocean shelves ( $> 300$  g C m<sup>2</sup> y<sup>-1</sup> or 0.3–2.8 g C m<sup>2</sup> d<sup>-1</sup>; Hameedi, 1978; Cota et al., 1996; Sambrotto et al., 1993; Hansell et al., 1993; Wheeler et al., 1996; Gosselin et al., 1997; Hill and Cota, 2005; Bates et al., 2005a; Mathis et al., 2009; Macdonald et al., 2009).

In early studies of the Chukchi Sea, Semiletov et al. (1999) observed undersaturated seawater  $p\text{CO}_2$  ( $\sim 200$ – $350$   $\mu\text{atm}$ ) during the sea-ice free period across the shelf with lowest values ( $\sim 150$ – $200$   $\mu\text{atm}$ ) observed close to the ice edge at the northern shelf break. More recently, other studies have observed similar undersaturated seawater  $p\text{CO}_2$  conditions from May to September ( $\sim 150$ – $350$   $\mu\text{atm}$ ; Pipko et al., 2002;

Murata and Takizawa, 2003; Bates et al., 2005a, 2006; Bates, 2006; Chen and Gao, 2007; Fransson et al., 2009). Summertime  $\Delta p\text{CO}_2$  values are typically in the range of  $-50$  to  $200$   $\mu\text{atm}$ . Extremely low summertime  $p\text{CO}_2$  conditions ( $< 100$   $\mu\text{atm}$ ) were observed in the surface layer in the vicinity of Barrow Canyon at the northern edge of the Chukchi Sea shelf in a location with very high rates of phytoplankton primary production (Bates et al., 2005a; Hill and Cota, 2005) and north of Wrangel Island (Fransson et al., 2009). In the western Chukchi Sea near Long Strait, summertime seawater  $p\text{CO}_2$  conditions are weakly undersaturated (Fransson et al., 2009) perhaps reflecting outflow of higher  $p\text{CO}_2$  surface waters from the East Siberian Sea with the intermittent Siberian Coastal Current (Weingartner et al., 1999). Early season observations under near complete sea-ice cover also indicate that Chukchi Sea shelf “winter” surface waters were not as undersaturated with respect to the atmosphere ( $\Delta p\text{CO}_2$  values of  $\sim -20$ – $60$   $\mu\text{atm}$ ) compared to the summertime sea-ice free period (Bates, 2006). In contrast to strongly undersaturated surface waters in the Chukchi Sea, subsurface waters seawater  $p\text{CO}_2$  values were highly oversaturated with respect to the atmosphere (Bates, 2006; Bates et al., 2009).

Estimates of the rates of air-sea  $\text{CO}_2$  exchange during the sea-ice free period in the summertime have ranged from  $\sim 20$ – $90$  mmol  $\text{CO}_2$  m<sup>-2</sup> d<sup>-1</sup> (Wang et al., 2003; Murata and Takizawa, 2003; Bates, 2006; Fransson et al., 2009) with the Chukchi Sea shelf a strong sink of  $\text{CO}_2$ . These studies indicate that the ocean uptake of  $\text{CO}_2$  from the atmosphere is greater for the Chukchi Sea shelf than other Arctic Ocean shelves. However, in regions of the Chukchi Sea shelf where sea-ice cover remained high ( $> 80\%$ ) during the summertime, air-sea  $\text{CO}_2$  exchange rates were estimated to be generally low ( $< 1$  mmol  $\text{CO}_2$  m<sup>-2</sup> d<sup>-1</sup>; Bates, 2006). Similarly, wintertime air-sea  $\text{CO}_2$  exchange rates during complete sea-ice cover were estimated to be minor ( $< 1$  mmol  $\text{CO}_2$  m<sup>-2</sup> d<sup>-1</sup>; Bates, 2006). The annual ocean  $\text{CO}_2$  uptake for the Chukchi Sea shelf has been estimated at  $24$ – $108$  g C m<sup>-2</sup> yr<sup>-1</sup> (Table 1; Bates, 2006). Similarly, using mass balance approaches, Kaltin and Anderson (2005) estimated that the annual ocean  $\text{CO}_2$  uptake for the Chukchi Sea to be  $86 \pm 22$  g C m<sup>-2</sup> yr<sup>-1</sup>.

The seasonal changes in surface  $p\text{CO}_2$  and strong sink for atmospheric  $\text{CO}_2$  on the Chukchi Sea shelf have been largely attributed to cooling of surface waters during the northward transit of waters across the Chukchi Sea shelf (Murata and Takizawa, 2003) and high rates of summertime phytoplankton PP that acts to decrease seawater DIC and  $p\text{CO}_2$  (Bates, 2006). As in the Barents Sea, the fall and winter rebound of seawater  $p\text{CO}_2$  to values just below equilibrium with the atmosphere likely results from a continued uptake of  $\text{CO}_2$  through gas exchange during sea-ice formation and brine rejection (Anderson et al., 2004; Omar et al., 2005) and vertical entrainment by mixing with  $\text{CO}_2$  rich subsurface waters.

## 4.2 Arctic Ocean shelves: interior shelves

### 4.2.1 Siberian Sea shelves (i.e. Kara, Laptev and East Siberian Seas)

The Kara Sea shelf is controlled by exchanges with the Barents Sea and adjoining Eurasian Basin (Fig. 1). Freshwater runoff from the Ob and Yenisey Rivers mix with more saline water of Atlantic Ocean origin, before entering the Laptev Sea or exported into the central basin of the Arctic Ocean (e.g. Anderson et al., 1994; Jones et al., 1991; Fransson et al., 2001). There is a general flow from the Laptev Sea across the East Siberian Sea and into the Chukchi Sea through the Siberian Coastal current (e.g. Weingartner et al., 1999; Nitishinsky et al., 2008), but circulation patterns can reverse with considerable transport from the East Siberian Sea shelf across the northeastern Laptev Sea shelf (e.g. Dmitrenko et al., 2008). Major freshwater inputs to the Laptev Sea occur primarily through the Lena River (e.g. Gordeev et al., 1996; Olsson and Anderson, 1997). On all of these shelves, seasonal sea-ice melt and formation, and brine rejection in coastal polynyas are important processes (e.g. Dmitrenko et al., 2008) with sea-ice production in this region occurring at the highest rate compared to other Arctic Ocean shelves.

There have been few studies of inorganic carbon on the Siberian Sea shelves, with surface seawater  $p\text{CO}_2$  and  $\Delta p\text{CO}_2$  values highly variable over time and space. During the sea-ice free period, surface waters of the Kara Sea appear to be undersaturated with respect to the atmosphere. Semiletov et al. (1999) report a range of 273–405  $\mu\text{atm}$  for surface seawater  $p\text{CO}_2$  while Kelley (1970) reported  $\Delta p\text{CO}_2$  values ranging from –30 to –100  $\mu\text{atm}$  across much of the Kara Sea in 1967. In contrast, supersaturated surface seawater  $p\text{CO}_2$  were observed nearshore and close to the outflow of the Ob and Yenisey Rivers (Kelley, 1970; Makkaveev, 1994; Semiletov et al., 1999). In the other Siberian Sea shelves, generally undersaturated surface seawater  $p\text{CO}_2$  (216–~400  $\mu\text{atm}$ ) for the Laptev Sea and highly variable seawater  $p\text{CO}_2$  conditions in the East Siberian Sea (~300–500  $\mu\text{atm}$ ) have been reported (Semiletov et al. 1999, 2007). High values for surface seawater  $p\text{CO}_2$  were observed nearshore on the shelf within the river plume of the Lena River (~850  $\mu\text{atm}$ ) that drains into the Laptev Sea, and the Kolyma River (~500  $\mu\text{atm}$ ) that drains into the East Siberian Sea shelf (Semiletov et al., 1999, 2007). In addition, very high values (~500 to +1500  $\mu\text{atm}$ ) have been observed in bottom waters of the inner shelf and also in the nearshore bays (e.g. Tiksi Bay) and estuaries of the Laptev and East Siberian Seas (Semiletov et al. 1999, 2007).

Fransson et al. (2001), using carbon mass balance approaches, computed a small annual ocean  $\text{CO}_2$  uptake of  $1 \text{ g C m}^{-2} \text{ yr}^{-1}$  for the Kara Sea (Table 1). For the Laptev Sea, mass balance assessments integrated across the shelf indicate that this sea is a sink for atmospheric  $\text{CO}_2$  during the sea-ice free summertime (~2.1  $\text{mmoles C m}^{-2} \text{ d}^{-1}$ ;

July–September; Nitishinsky et al., 2008) despite supersaturated seawater  $p\text{CO}_2$  conditions (i.e. sources of  $\text{CO}_2$  to the atmosphere) in nearshore bays (Olsson and Anderson, 1997) and Lena River outflow (Semiletov et al., 1999, 2007). Direct eddy correlation measurements of air-sea  $\text{CO}_2$  exchange in late summer (September) close to the multiyear sea-ice at the shelf break of the Laptev Sea yielded more contradictory results with air-sea  $\text{CO}_2$  exchange rates varying from invasion of  $1.7 \text{ mmoles C m}^{-2} \text{ d}^{-1}$  to evasion of  $1.2 \text{ mmoles C m}^{-2} \text{ d}^{-1}$  (Semiletov et al., 2007). If we extrapolate the results of Nitishinsky et al. (2008) and assume a 100 d sea-ice free period, and negligible air-sea gas exchange during wintertime sea-ice cover, an annual ocean  $\text{CO}_2$  uptake of ~2.4  $\text{g C m}^{-2} \text{ yr}^{-1}$  is computed (Table 1). For the East Siberian Sea shelf, the sink or source terms are uncertain. Nitishinsky et al. (2008) estimated that the East Siberian Sea shelf is a small source of  $\text{CO}_2$  to the atmosphere (~0.3  $\text{mmoles C d}^{-1} \text{ m}^{-2}$ ). However, Semiletov et al. (2007) suggest that the western area of the shelf is a source of  $\text{CO}_2$  to the atmosphere ( $1 \pm 1.6 \text{ mmoles C d}^{-1} \text{ m}^{-2}$ , 2003; and  $10.9 \pm 12.6 \text{ mmoles C m}^{-2} \text{ d}^{-1}$ , 2004) while the eastern area, influenced by mixing with Pacific Ocean water from the Chukchi Sea shelf, is a sink for atmospheric  $\text{CO}_2$ .

The above studies suggest that the Siberian Sea shelves range from minor sinks to minor sources of  $\text{CO}_2$  to the atmosphere, with the causes uncertain at present. Nitishinsky et al. (2008) suggest that the drawdown of  $p\text{CO}_2$  from summertime photosynthetic uptake more than compensates for seasonal warming, and mixing with river freshwater and high- $\text{CO}_2$  subsurface waters, all of which act to increase seawater  $p\text{CO}_2$ . Semiletov et al. (2007) showed highly supersaturated values for subsurface waters on all of the Siberian Sea shelves, presumably resulting from remineralization of organic matter introduced from the Siberian Rivers (e.g. Anderson et al., 1990; Cauwet and Sidorov, 1996; Kattner et al., 1999) and from vertical export of organic carbon despite relatively low rates of summertime phytoplankton primary production (compared to other Arctic Ocean shelves) in the euphotic zone (~6–12  $\text{g C m}^{-2} \text{ yr}^{-1}$ ; Macdonald et al., 2009). The East Siberian Sea shelf may be more highly influenced by these processes and hence more likely to be net heterotrophic and a source of  $\text{CO}_2$  to the atmosphere. The Laptev Sea also has a permanent flaw lead polynya (Dmitrenko et al., 2005; Nitishinsky et al., 2008) that may contribute to ocean  $\text{CO}_2$  uptake from the atmosphere during wintertime brine rejection and sea-ice formation as shown elsewhere (Anderson et al., 2004; Omar et al., 2005), but also winter outgassing of  $\text{CO}_2$  from the polynya.

### 4.2.2 Beaufort Sea shelf

The Beaufort Sea shelf has a relatively narrow (~50 km) continental shelf in contrast to the adjacent Chukchi Sea shelf (Fig. 1). The eastern Beaufort Sea shelf is heavily

influenced by outflow from the Mackenzie River, and this input of freshwater and terrigenous input dominates the physical and chemical properties of the Beaufort Sea shelf (Carmack and Macdonald, 2002; Macdonald et al., 2009). Compared to the adjacent Chukchi Sea shelf, annual rates of phytoplankton PP for the Beaufort Sea shelf were relatively low ( $\sim 7\text{--}17\text{ g C m}^{-2}\text{ yr}^{-1}$ ; Macdonald et al., 2009), similar to other “interior” type shelves (Carmack and Wassmann, 2006) such as the Siberian Sea shelves. In the adjoining Canada Basin, wind-driven, clockwise rotation of the Beaufort Gyre controls regional scale sea-ice and upper ocean circulation (Carmack and Chapman, 2003). Along the northern edge of the Chukchi and Beaufort Seas, a narrow eastward boundary current is present (Nikolopoulos et al., 2009) which is constrained to follow the isobaths along the shelf break due to strong dynamic controls (Winsor and Chapman, 2004; Spall, 2007). The generalized regional water circulation along the Beaufort Sea shelf is directed eastward with shelf waters renewed principally by Pacific Ocean waters that enter through Bering Strait and exit the Chukchi Sea off the Alaskan coast. Shelf-basin exchange facilitated by the generation of mesoscale eddies (e.g. Manley and Hunkins, 1985; Muench et al., 2000; Pickart et al., 2005; Mathis et al., 2007a), and winter brine rejection during sea-ice formation are also important for these shelves.

As with the Siberian Sea shelves, few studies have been conducted on the Beaufort Sea shelf. Seawater  $p\text{CO}_2$  conditions appear to be highly variable ( $\sim 150\text{--}350\text{ }\mu\text{atm}$ ) in the western part of the Beaufort Sea shelf (Murata and Takizawa, 2003; Bates, 2006). Extremely low seawater  $p\text{CO}_2$  conditions ( $<100\text{ }\mu\text{atm}$ ) have also been observed in the surface layer of the Beaufort Sea shelf east of Point Barrow in a region of characterized by high proportions of freshwater (10–20%) and sea-ice melt (10–25%) (Cooper et al., 2005; Bates, 2006). Murata and Takizawa (2003) have estimated that rate of ocean  $\text{CO}_2$  uptake was relatively low ( $-12\text{ mmol CO}_2\text{ m}^{-2}\text{ d}^{-1}$ ) during the summertime compared to the adjacent Chukchi Sea. In areas of the western Beaufort Sea shelf where sea-ice cover remained high ( $>80\%$ ), air-to-sea  $\text{CO}_2$  fluxes were generally low ( $<1\text{--}10\text{ mmol CO}_2\text{ m}^{-2}\text{ d}^{-1}$ ; Bates, 2006; Fransson et al., 2009). In the eastern Beaufort Sea shelf, summertime surface seawater  $p\text{CO}_2$  values were generally weakly undersaturated (Mucci et al., 2008) or close to equilibrium with the atmosphere particularly in the Banks Island polynya (Fransson et al., 2009). Fransson et al. (2009) estimated that the summertime ocean  $\text{CO}_2$  uptake was  $\sim 6\text{ mmol CO}_2\text{ m}^{-2}\text{ d}^{-1}$  for the eastern Beaufort Sea shelf. The annual ocean  $\text{CO}_2$  uptake for the Beaufort Sea shelf has been estimated at  $1.2\text{ g C m}^{-2}\text{ yr}^{-1}$ , low compared to other Arctic Ocean shelves (Table 1; Murata and Takizawa, 2003).

### 4.3 Arctic Ocean shelves: outflow shelves


#### 4.3.1 Canadian Archipelago shelf

The shelf of the Canadian Archipelago, east of the Beaufort Sea shelf (Fig. 1), has been poorly sampled for marine inorganic carbon cycle properties. There is little information to estimate rates of air-sea  $\text{CO}_2$  gas exchange and the processes that control the inorganic carbon cycle. In the adjacent Hudson Bay, Else et al. (2008) observed mean rates of air-sea  $\text{CO}_2$  exchange of  $0.7\text{ mmol CO}_2\text{ m}^{-2}\text{ d}^{-1}$  directed towards the ocean during the sea-ice free period. On an annual timescale, a small ocean  $\text{CO}_2$  uptake for the Hudson Bay shelf is estimated at  $0.07\text{ g C m}^{-2}\text{ yr}^{-1}$  (Else et al., 2008).

In the CARINA dataset (Fig. 4), computation of  $p\text{CO}_2$  from DIC and alkalinity datasets indicate that surface  $p\text{CO}_2$  values ranged from  $\sim 300\text{--}400\text{ }\mu\text{atm}$  on the Canadian Archipelago shelf and generally undersaturated with respect to the atmosphere. During the CASES project in the early 2000's, surface  $p\text{CO}_2$  values in the eastern Beaufort Sea shelf ranged from  $\sim 300\text{--}430\text{ }\mu\text{atm}$  with significant spatial variability (Mucci et al., 2008). More recently, Fransson et al. (2009) observed summertime surface  $p\text{CO}_2$  values of  $\sim 150\text{--}300\text{ }\mu\text{atm}$  in the Canadian Archipelago shelf. Surface waters were strongly undersaturated ( $\Delta p\text{CO}_2$  values of  $-100$  to  $-200\text{ }\mu\text{atm}$ ) and a very strong ocean sink of  $140\text{ mmol m}^{-2}\text{ d}^{-1}$  was shown for Lancaster Sound in particular (Fransson et al., 2009). If we conservatively assume that air-sea  $\text{CO}_2$  exchange rates are similar for the Canadian Archipelago as the “upstream” Beaufort Sea shelf, the annual ocean uptake of  $\text{CO}_2$  for the Canadian Archipelago shelf is computed at 16 to  $24\text{ g C m}^{-2}\text{ yr}^{-1}$  (Table 1).

#### 4.4 Arctic Ocean central basin: Canada and Eurasian Basins

The central basin of the Arctic Ocean is strongly stratified (Kinney et al., 1970; Aagaard et al., 1981; Jones and Anderson, 1986; Anderson et al., 1989, 1999; Aagaard and Carmack, 1994; Schlosser et al., 1995; Codispoti et al., 2005). In the Eurasian and Canada Basin (Fig. 1), surface waters of the central basin are derived from a mixture of local sea-ice melt, fresh water inputs, exchanges with shelf waters of Pacific Ocean and Atlantic Ocean origin (Jones et al., 2008). In the Canada Basin, a strong vertical density gradient separates nutrient poor surface waters or polar mixed layer (PML) from underlying nutrient rich halocline waters that can be subdivided into Upper Halocline Layer (UHL) and Lower Halocline Layer (LHL). The shelf surface waters and underlying upper halocline waters have a predominantly Pacific Ocean origin based on characteristic temperature, salinity (Aagaard et al., 1981), inorganic nutrient, dissolved oxygen, barium and alkalinity distributions (e.g. Wallace et al., 1987; Wilson and Wallace, 1990; Jones et al., 1991; Salmon and McRoy, 1994; Falkner et al., 1994; Anderson, 1995; Anderson et al.,



**Fig. 4.** Surface seawater  $p\text{CO}_2$  values computed from DIC and TA data collected in the Arctic Ocean and part of the CARINA dataset (data obtained at <http://cdiac.ornl.gov/>). The following cruise data were used to generate the surface  $p\text{CO}_2$  map. The metadata is given as CDIAc cruise number, Cruise Name, CARINA Table ID, ship, cruise dates, Chief Scientist (in brackets) and scientist responsible for inorganic carbon measurements. 3, 06AQ19960712, 3, Polarstern, 7/12–9/6/1996 (E. Augstein), L. Anderson; 17, 06MT19970707, 18, Meteor, 7/7–8/9/1997 (F. Schott), A. Koertzing/L. Mintrop; 37, 18HU19970509, 44, Hudson, 5/9–6/11/1997 (R. A. Clarke), P. Jones/R. Gershey; 38, 18SN19940724, 47, St. Laurent, 7/24–9/1/1994 (K. Aagaard/E. Carmack/J. Swift); P. Jones/K. Azetsu-Scott; 39, 18SN19970803, 48, St. Laurent, 8/3–8/18/1997 (F. McLaughlin/K. Falkner), P. Jones/K. Azetsu-Scott; 40, 18SN19970831, 49, St. Laurent, 8/31–9/16/1997 (F. McLaughlin/K. Falkner), P. Jones/K. Azetsu-Scott; 41, 18SN19970924, 50, St. Laurent, 9/24–10/15/1997 (F. McLaughlin/K. Falkner), P. Jones/K. Azetsu-Scott; 108, 58AA20000923, 124, H. Mosby, 9/23–10/03/2000 (S. Osterhus) A. Omar; 128, 58JM20030710, 145, Jan Mayen, 7/10–7/18/2003 (P. Wassmann), R. Bellerby/C. Kivimae; 129, 58JM20040724, 146, Jan Mayen, 7/24–7/31/2004 (P. Wassmann), R. Bellerby/C. Kivimae; 130, 58JM20050520, 147, Jan Mayen, 5/20–6/2/2005 (P. Wassmann), R. Bellerby/C. Kivimae; 131, 58LA19860719, 148, Lance, 7/19–7/26/1986 (chief scientist not listed), L. Anderson; 160, 77DN19910726, 177, Oden, 7/26–9/3/1991 (L. Anderson/J.-E. Hellsvik), L. Anderson; 162, 77DN20020420, 179, Oden, 4/20–6/6/2002, L. Anderson; L. Anderson; 164, 90AQ19940706, 181, A. Fedorov, 7/5–8/8/1994 (chief scientist not listed), L. Anderson.

1999; Rudels et al., 1996; Guay and Falkner, 1997; Jones et al., 2003). In surface waters of the Canada Basin, the river fraction can be up to 20% (Macdonald et al., 2002; Cooper et al., 2005), while localized to the upper few metres, the sea-ice melt fraction can also exceed 25% (Cooper et al., 2005) during the summertime retreat of sea-ice. In the Eurasian Basin, surface waters and halocline waters are predominantly of Atlantic Ocean origin that have been modified on the Barents Sea and Siberian Seas shelves adjacent to the

Eurasian Basin (Anderson et al., 1994). The upper halocline layer is absent from most of the Eurasian Basin (Anderson et al., 1994a). At greater depths of the central basin is the Atlantic Water Layer (AWL;  $\sim 200$ – $800$  m deep) and the Arctic Ocean Deep Water (AODW), both of which have Atlantic Ocean origins (e.g. Jones et al., 1991, 1995).

The central basin of the Arctic has been poorly sampled for marine inorganic carbon cycle properties. The Arctic Ocean Section (AOS) expedition in 1994 sampled across the Eurasian Basin and Canada Basin, west of the Chukchi Cap (Anderson et al., 1998a). Calculation of  $p\text{CO}_2$  from AOS DIC and TA data (obtained at <http://cdiac.ornl.gov/oceans/CARINA/>) indicate that the surface waters had seawater  $p\text{CO}_2$  values of  $\sim 300$ – $330$   $\mu\text{atm}$  and  $\Delta p\text{CO}_2$  values of  $\sim -40$ – $70$   $\mu\text{atm}$ , and hence undersaturated with respect to the atmosphere. In the Canada Basin, adjacent to the Chukchi Sea shelf, lower seawater  $p\text{CO}_2$  values of  $\sim 240$ – $280$   $\mu\text{atm}$  and  $\Delta p\text{CO}_2$  values of  $\sim -150$ – $25$   $\mu\text{atm}$  have also been observed (Bates, 2006; Bates et al., 2006). More recently, in the Makarov Basin of the Canada Basin, low surface seawater  $p\text{CO}_2$  values of  $\sim 150$ – $250$   $\mu\text{atm}$  and  $\Delta p\text{CO}_2$  values of  $\sim -100$  to  $200$   $\mu\text{atm}$  have been observed under sea-ice (Fransson et al., 2009). These observations suggest that the central basin of the Arctic Ocean has the potential to absorb significant amounts of  $\text{CO}_2$  (Anderson and Kaltin, 2001; Bates et al., 2006). The capacity for sea-ice to allow direct gas exchange has been debated (e.g. Gosink et al., 1976; Semiletov et al., 2004; Delille et al., 2007; Nagurnyi, 2008), and here, we assume that sea-ice is an effective barrier to gas exchange. Wintertime area of open water in leads and polynyas has been estimated at 1% in the central basin (Gow and Tucker, 1990). In regions of the Canada Basin with 100% sea-ice cover, Bates et al. (2006) used an effective sea-ice cover of 99% to compute air-sea  $\text{CO}_2$  influx of  $< 3$   $\text{mmoles CO}_2 \text{ m}^{-2} \text{ d}^{-1}$  to account for exchange through leads, polynyas, fractures and brine channels (i.e. 1% area of the central basin), with much higher rates of exchange in areas without complete sea-ice cover.

#### 4.5 Arctic Ocean (shelves and central basin): estimating annual air-sea $\text{CO}_2$ exchange rates and the Arctic Ocean $\text{CO}_2$ sink

Early carbon mass balance estimates of the rate of air-sea  $\text{CO}_2$  exchange suggested that the entire Arctic Ocean was a sink for  $\text{CO}_2$  in the range of 70 to 129  $\text{Tg C year}^{-1}$  (Table 1: Anderson et al., 1990, 1994; Lundberg and Haugen, 1996). Subsequently, Anderson et al. (1998b) and Kaltin and Anderson (2005) used similar carbon mass balance approaches but revised their estimates of the Arctic Ocean  $\text{CO}_2$  sink downward to 24 and 31  $\text{Tg C year}^{-1}$ . More recently, direct  $\Delta p\text{CO}_2$  observations and air-sea  $\text{CO}_2$  exchange rate estimates revised the Chukchi Sea shelf  $\text{CO}_2$  sink upward and the Arctic Ocean  $\text{CO}_2$  sink was estimated at 66  $\text{Tg C year}^{-1}$  (Bates, 2006; Bates et al., 2006). A review of published

data suggests that the Arctic Ocean CO<sub>2</sub> sink ranges from approximately 66 to 175 Tg C yr<sup>-1</sup> depending on lowest to highest flux rates estimates from available studies (Table 1). In light of recent observations in the Canadian Archipelago shelf (Fransson et al., 2009), the addition of a Canadian Archipelago shelf ocean CO<sub>2</sub> sink would increase the Arctic Ocean CO<sub>2</sub> sink to 81 to 199 Tg C yr<sup>-1</sup> (Table 1). In comparison to the mean annual global ocean CO<sub>2</sub> uptake of approximately 1400 Tg C yr<sup>-1</sup> (Takahashi et al., 2009), the Arctic Ocean CO<sub>2</sub> sink potentially contributes ~5–14% to the global balance of CO<sub>2</sub> sinks and sources.

#### 4.5.1 The present day annual Arctic Ocean CO<sub>2</sub> sink: caveats and considerations

Our review of published results from both carbon mass balance and observational studies indicates that the Arctic Ocean is a CO<sub>2</sub> sink of approximately 66 to 199 Tg C yr<sup>-1</sup>, at present. However, it is important to note that these estimates are based on very limited spring and summer data, and near absence of wintertime data.

On the Arctic Ocean shelves, there is a potential for wintertime outgassing of CO<sub>2</sub>. For example, in the nearshore regions of the shelves, surface summer  $\Delta p\text{CO}_2$  values are near neutral (near the Mackenzie River outflow and Banks Island polynya; Fransson et al., 2009) or positive (highly positive in places like Tiksi Bay and in the river outflows onto the Siberian and Beaufort shelves; e.g. Semiletov, 1999; Semiletov et al., 2007). There are also indications that the highly river-dominated surface waters on the Laptev and East Siberian Sea shelves have positive  $\Delta p\text{CO}_2$  values (inferred from the very low  $\Omega$  values for river end-members shown by Salisbury et al., 2008). Winter mixing with CO<sub>2</sub> rich subsurface waters are highly likely to create conditions favoring the potential for surface waters to outgas CO<sub>2</sub> during winter through polynyas and leads. In the Laptev Sea, for example, the wintertime flaw-lead polynya occurs over the outer shelf at a distance of ~50–150 km from the nearshore areas with positive summertime  $\Delta p\text{CO}_2$  values (Dmitrenko et al., 2005). However, the prevalent offshore circulation pathways on the Laptev Sea are likely to transport positive  $\Delta p\text{CO}_2$  surface waters to the region of the law-lead polynya. In total, the area of polynyas and flaw-leads on the Arctic shelves has been reported at  $3.5 \times 10^{10} \text{ m}^2$  (e.g. Winsor and Bjork, 2000). Assuming a 100 d period for gas exchange from polynyas and leads, gas exchange rates of 10 mmol CO<sub>2</sub> m<sup>-2</sup> d<sup>-1</sup> would result in an efflux of 0.4 Tg C. Compared to influxes in the Chukchi and Barents Seas,  $\Delta p\text{CO}_2$  values would have to be in the order of +150  $\mu\text{atm}$  for an efflux of 0.4 Tg, and +1500  $\mu\text{atm}$ , for a wintertime polynya/lead efflux of 4 Tg.

A counterbalance to wintertime outgassing from polynyas and leads, may be wintertime net influx of carbon due to brine rejection during deep water formation (Anderson et al., 2004; Omar et al., 2005). Omar et al. (2005), scaling results from Storfjorden, Svalbard, reported a wintertime influx of

2.3, 6.8 and 33 Tg C for coastal shelf polynyas, central basin polynyas, and brine-rejection during ice formation in the seasonally sea-ice free areas of the Arctic, respectively. The caveat for this study is that the results were extrapolated from Storfjorden to the entire Arctic Ocean.

Under-sea ice observations across the central basin (Anderson et al., 1994b; Jutterstrom and Anderson, 2005; Tanhua et al., 2009; Fransson et al., 2009) indicate that  $\Delta p\text{CO}_2$  values are negative with the potential to absorb CO<sub>2</sub> (although blocked by sea-ice). Given the 2–12 year residence times of surface waters in the central basin, we expect that the under-ice trans-Arctic sections are likely to reflect wintertime conditions as well. In addition, the Revelle factors (Sabine et al., 2004) and pre-conditions of Atlantic and Pacific surface waters entering the Arctic and cooling during transit into the Arctic are likely to favor negative  $\Delta p\text{CO}_2$  values and thereby the potential for Arctic Ocean surface waters to absorb CO<sub>2</sub>.

## 5 Feedbacks and vulnerabilities

The Arctic Ocean is currently in the forefront of climate change caused by both natural and anthropogenic factors that influence the cryosphere, hydrological and biogeochemical cycles of the Arctic. However, due to undersampling of the Arctic Ocean, many uncertainties remain about the marine carbon cycle and, as shown earlier in this review, the uptake of CO<sub>2</sub> and the physical and biological processes that control the inorganic carbon cycle on the shelves and in the central basin. It is thus difficult to quantitatively predict changes in processes and feedbacks in the Arctic Ocean. However, here we discuss the potential vulnerabilities of the inorganic carbon cycle due to natural and anthropogenic factors, including: (1) sea-ice loss; (2) warming, circulation and other physical changes; (3) changes in biology and ecosystem structure of the surface and halocline water, and; (4) changes in the hydrological cycle and freshwater inputs to the Arctic Ocean. We use Figs. 2 and 3 as a framework for this discussion and Table 2 provides likely impact of process, confidence level and potential time-scale of impact.

### 5.1 Sea-ice loss

Mean air temperatures over the Arctic have increased by ~2–3°C in summertime and by ~4°C in winter since the 1950's (Chapman and Walsh 2003) with projections for a further 4–5°C increase by the end of the 21st century (ACIA, 2005). In conjunction with these higher temperatures, sea ice cover in the Arctic Ocean has contracted over the past three decades (Levi 2000; Parkinson 2000; Cavalieri et al., 2003; Wang et al., 2005; Overland and Wang, 2005; Rothrock and Zhang, 2005; Stroeve et al., 2005, 2007), with the pace of decline accelerating beyond model predictions over the last several years (Winton, 2006; Holland et al., 2006; Overland and Wang, 2007; Stroeve et al., 2007; Maslanik et al.,

**Table 2.** Summary of physical and biological processes that can influence the inorganic carbon cycle and increase/decrease the Arctic CO<sub>2</sub> sink. The numeral identification of each process is taken from Figs. 2 and 3. Confidence level is also given, as well as time-scale (over the next decade or century) and impact (if quantified).

Process	Process ID (Refer to Fig. 2 and 3)	Increase/ decrease CO <sub>2</sub> sink	Confidence	Time-scale and Impact Decade	Century
<i>Sea-ice changes and ocean-atmosphere interaction</i>					
Sea-ice loss	7	Increase	Medium	+ 2 Tg C yr <sup>-1</sup>	?
Air-sea gas exchange	3	Uncertain	Low	?	?
Sea-ice gas exchange	14	Uncertain	Low	?	?
Polynya/lead gas exchange	13	Decrease	Low	?	?
Brine rejection	15	Increase	Low	?	?
Sea-ice melt gas exchange	8	Decrease	Low	?	?
River-water gas exchange	17	Decrease	Low	?	?
<i>Warming, circulation and other physical changes</i>					
Northward transport	1	n/a	Medium	?	?
Less cooling during northward transport of surface water	2	Decrease	Medium	?	?
Shelf-basin transport	9	Uncertain	Low	?	?
Vertical mixing and diffusion	5	Decrease	Low	?	?
River input and coastal erosion	16	Decrease	Medium	?	?
<i>Biological processes</i>					
Primary production (new)	4	Increase	Medium	?	?
Export production	6	Increase	Low	?	?
Subsurface OM remineralization	10	Uncertain	Low	?	?
Sediment respiration of CO <sub>2</sub>	11	Uncertain	Low	?	?
Sediment release of TA	12	Uncertain	Low	?	?

2007; Shimada et al., 2006). The length of the seasonal sea-ice free period has increased (e.g. Smith 1998; Rigor et al., 2002; Rigor and Wallace, 2004; Arrigo et al., 2008; Comiso et al., 2008), and sea-ice thickness observed over the central Arctic Ocean has decreased as thick multi-year sea-ice is replaced by thinner first-year sea-ice (e.g. Rothrock et al., 2003; Giles et al., 2008). The decline in Arctic sea-ice cover has been attributed to a combination of factors, including: (1) increased Arctic temperatures (Rothrock and Zhang 2005; Lindsay and Zhang 2005); (2) increased advection of warmer water into the Arctic Ocean from the Atlantic Ocean through the Fram Strait and Barents Sea gateways (Steele and Boyd 1998; Dickson et al., 2000) and from the Pacific Ocean through Bering Strait (Maslowski et al., 2001; Shimada et al., 2006), and; (3) increased advection of sea-ice out of the Arctic Ocean through Fram Strait due to recent changes in atmospheric circulation patterns (Rigor and Wallace, 2004; Maslanik et al., 2007; Serreze et al., 2007). In combination, these factors contribute to a positive feedback where sea-ice loss reinforces surface warming due to reduced surface albedo and increased shortwave penetration, which in turn inhibits sea-ice formation in wintertime and allows for

acceleration of sea-ice loss during the summertime (Perovich et al., 2007).

### 5.1.1 Sea-ice loss and exposure of surface waters

The loss of sea-ice is expected to both reduce the % sea-ice cover and expose undersaturated surface waters of the Arctic shelves and central basin (i.e. Canada and Eurasian Basins) (Figs. 2 and 3, process 4). Prior to 2007, the rate of sea-ice loss from the central Arctic Ocean was estimated at  $\sim 36\,000\text{ km}^2\text{ yr}^{-1}$  (Cavalieri et al., 2003). In surface waters of the Canada Basin, a mean ocean CO<sub>2</sub> uptake of  $46\pm 8\text{ mmol CO}_2\text{ m}^{-2}\text{ d}^{-1}$  was observed in 2002 and 2004 in sea-ice free areas during summertime. These air-sea CO<sub>2</sub> exchange rates were scaled to the entire central basin and annual reduction of sea-ice extent to estimate an increase of the ocean CO<sub>2</sub> sink of  $2.0\pm 0.3\text{ Tg C year}^{-1}$  due to sea-ice loss with the effect compounding over time (Bates et al., 2006). In 2007 and 2008, sea-ice extent reached a seasonal minimum 25% lower than any previously observed in the satellite record (Maslanik et al., 2007; Comiso et al., 2008) constituting an additional exposure of  $\sim 600\,000\text{ km}^2$  of surface

waters to air-sea gas exchange. Assuming a status quo of inorganic carbon distributions in surface waters of the Arctic, this recent loss of summertime sea-ice may have increased the ocean uptake of CO<sub>2</sub> in the Arctic Ocean by an additional 33±10 Tg C year<sup>-1</sup>. If sea-ice cover was lost completely over the central basin during summertime, surface waters of the Arctic Ocean might absorb an additional ~280 to 1200 Tg of CO<sub>2</sub> (Anderson and Kaltin, 2001; Bates et al., 2006). The range of estimates depends on differences in surface water carbonate chemistry observed between the Canada Basin and Eurasian Basin. For example, Anderson and Kaltin (2001) computed how much CO<sub>2</sub> was needed to equilibrate the surface mixed layer with the atmosphere using vertical profiles of DIC data from the Eurasian Basin.

The above estimates of CO<sub>2</sub> uptake changes due to sea-ice loss have many uncertainties and associated caveats. These studies are only applicable to the near-term future (<decade) since it was assumed that the driving force of gas exchange (e.g.  $\Delta p\text{CO}_2$  and DIC distributions of surface and underlying halocline layer waters) would not change significantly over the residence time of surface waters in the Arctic Ocean (e.g. 2–30 years; Macdonald et al., 2009). However, in the era of rapid change in the Arctic, inorganic carbon distributions and air-sea CO<sub>2</sub> exchange rates are highly likely to change due to a host of other factors. Since the effect of these other factors on the marine inorganic carbon cycle can be opposing or amplifying in nature, future predictions of the trajectory of the Arctic Ocean CO<sub>2</sub> sink/source terms are difficult to make.

### 5.1.2 Other sea-ice and gas exchange issues

The loss of sea-ice in the Arctic Ocean should result in greater open water area, and increased air-sea interaction, with a variety of consequences. At the same time, increased latitudinal atmospheric pressure gradients could result in increased windspeed and storm events over the Arctic. Although the direction of air-sea gas exchange is forced by  $\Delta p\text{CO}_2$  values, the rate of gas exchange of CO<sub>2</sub> (Figs. 2 and 3, process 2) and other gases is primarily driven by windspeed and atmospheric boundary layer-surface water interactions. Given these dictates, even though  $\Delta p\text{CO}_2$  values might reduce, air-sea CO<sub>2</sub> gas exchange rates can increase since gas transfer velocities increase exponentially relative to windspeed (Wanninkhof, 1992; Wanninkhof and McGillis, 1999) and CO<sub>2</sub> flux in other regions can be dominated by storms and high-wind events (Bates and Merlivat, 2001).

Wintertime sea-ice has thinned over the last few decades and there may be potentially greater air-sea gas exchange through sea-ice (Gosink et al., 1978; Semiletov et al., 2004; Delille et al., 2007) with sea-ice a potentially weaker barrier to gas exchange (Figs. 2 and 3, process 7). Sea-ice can both release and absorb CO<sub>2</sub> (Delille et al., 2007) depending on changes in the net metabolism of the sea-ice biological community, and the net impact on air-sea CO<sub>2</sub> exchange is uncer-

tain. Brine rejection during sea ice formation (Figs. 2 and 3, process 9) is thought to transfer carbon from sea-ice to deep water (Anderson et al., 2004; Rysgaard et al., 2007) with estimates of its impact in the Arctic Ocean at ~43 Tg C yr<sup>-1</sup> (Omar et al., 2005). Calcium carbonate minerals such as Ikaite also form in sea-ice (Dieckmann et al., 2008) indicative of high  $\Omega$  and low  $p\text{CO}_2$ . However, localized surface water with high percentages of sea-ice melt or river freshwater have been observed with low  $\Omega$  values (Yamamoto-Kawai et al., 2009; Chierici and Fransson, 2009; Bates et al., 2009) and high  $p\text{CO}_2$  (Figs. 2 and 3, process 5 and 6), suggesting that these summertime surface areas could outgass CO<sub>2</sub>.

In wintertime, wind-driven polynyas open up on the Chukchi and Laptev Sea shelves in particular (Cavaliere and Martin, 1994; Dmitrenko et al., 2005; Nitishinsky et al., 2007), resulting in the formation of dense water via brine-driven convection (Weingartner et al. 1998). Since brine rejection and formation of dense water appears to facilitate uptake of CO<sub>2</sub> from the atmosphere (Anderson et al., 2004; Omar et al., 2005), enhanced opening of polynyas and leads may increase the sink of CO<sub>2</sub> in the Arctic Ocean in the near-future depending on inorganic carbon distributions and future air-sea CO<sub>2</sub> disequilibrium. In counterbalance, polynyas and leads may outgass CO<sub>2</sub> in shelf areas with positive  $\Delta p\text{CO}_2$  values, especially in the Siberian shelves (Sect. 4.2; Figs. 2 and 3, process 8).

The cumulative effect of the above processes (i.e. 5–9) on the balance of CO<sub>2</sub> sinks and sources in the Arctic Ocean is highly uncertain and in the absence of definitive data their impact in the near-term has to be considered minor at present (Table 2).

### 5.2 Warming, circulation and other physical changes in the Arctic

In the recent past (1998–2006), Pabi et al. (2008) observed positive anomalies in surface temperatures of up to 2°C in the regions of significant sea-ice loss (mainly on the “inflow” Arctic shelves). This suggests that the cooling of surface waters during transit northward into the Arctic has reduced (Figs. 2 and 3, process 10), and that summertime surface waters have become warmer over the last few decades (ACIA, 2005). Due to simple thermodynamics, cooling/warming should decrease/increase seawater  $p\text{CO}_2$  at a rate of ~4.1% per °C (Takahashi et al., 1993; Millero, 1995). In inflow shelf and central basin surface waters, if we assume present day surface water  $p\text{CO}_2$  values of 250  $\mu\text{atm}$ , a 1°C warming (or 1°C less cooling!) should increase seawater  $p\text{CO}_2$  by ~8–12  $\mu\text{atm}$  per 1°C and reduce  $\Delta p\text{CO}_2$  values and rates of ocean CO<sub>2</sub> uptake by ~10%. If surface waters equilibrate with the predicted 4–5°C warming of the atmosphere over the Arctic by the end of the 21st century (ACIA 2005), and if present-day inorganic carbon distributions in the upper layers of the Arctic remain unchanged, the Arctic Ocean CO<sub>2</sub> sink might reduce by ~40–50% due to warming.

The influence of change in the northward transport of water and carbon (Figs. 2 and 3, process 1) into the Arctic through the gateways of the Arctic (e.g. Bering Strait) is also uncertain at present. Water mass transport through Bering Strait appears to have increased over the last decade (Woodgate et al., 2005). Given that the inflow of Pacific and Atlantic Ocean waters transports  $\sim 3 \text{ Pg C yr}^{-1}$  into the Arctic, small changes in the transport of carbon into and out of the Arctic, and changes in the  $\text{CO}_2$  preconditions of the Bering Sea and North Atlantic Ocean may have significant and rapid impact on the balance of Arctic Ocean  $\text{CO}_2$  sinks and sources.

Wind-driven air-sea interaction and eddy formation may also change, including wind-induced upwelling/downwelling (Melling, 1993; Pickart et al. 2006, 2009), and formation of eddies that contribute to exchanges of water and materials between shelves (Figs. 2 and 3, process 11) the shelf and basin (process 13; Manley and Hunkins, 1985; Muench et al., 2000; Pickart et al., 2005), particularly in the western Arctic between the Beaufort Sea shelf and Canada Basin (Carmack and Macdonald, 2002; Mathis et al., 2007a; Spall et al., 2008). Deeper mixing may also entrain  $\text{CO}_2$  from subsurface waters (Figs. 2 and 3, process 12), but if nutrients are also entrained to support enhanced rates of new and export production (Figs. 2 and 3, process 14 and 15), there may be no net impact on the balance of  $\text{CO}_2$  sinks and sources.

The cumulative effect of changes to these processes (i.e. 1, 10–13) also remains uncertain at present (Table 2).

### 5.3 Changes in biology and ecosystem structure in surface and halocline waters of the Arctic Ocean

The loss of sea-ice in the Arctic Ocean and greater open water area should also enhance upwelling at the shelf-break and potentially increase the input of nutrients from subsurface waters to the Arctic shelves (ACIA 2005). The growing season for phytoplankton should increase with reduced sea-ice extent and longer open-water conditions. As a consequence of increased phytoplankton PP, new production or NCP (Figs. 2 and 3, process 14), the drawdown of  $p\text{CO}_2$  and DIC should increase the air-sea  $\text{CO}_2$  disequilibrium (i.e.  $\Delta p\text{CO}_2$ ) and increase the net oceanic uptake of  $\text{CO}_2$  (Anderson and Kaltin, 2001; Bates et al., 2006). Recent reductions of sea-ice cover, particularly in the western Arctic Ocean, have resulted in a longer phytoplankton growing season and a  $\sim 30$ – $60\%$  increase in primary production over the last decade, with a  $\sim 10$ – $40\%$  increase in phytoplankton primary production over the last several years (Arrigo et al., 2008). However, enhanced stability of the water column observed in the Beaufort Sea shelf (Trembley et al., 2008), if widespread across the Arctic, could reduce the upward flux of nutrients and rates of new production with reduced net impact on surface  $p\text{CO}_2$ . Enhanced export of organic matter (Figs. 2 and 3, process 15) may also enhance the rates of

rem mineralization of OM to  $\text{CO}_2$  in subsurface waters (Figs. 2 and 3, process 16) and sediments (Figs. 2 and 3, process 17), increasing the potential for subsurface waters to reduce air-sea disequilibrium in the surface through processes such as upwelling, mixing and vertical diffusion.

For sea-ice communities that generally account for a relatively small fraction of primary production, the reduction of sea-ice extent and thickness will reduce the contribution of sea-ice algae to primary production (Subba Rao and Platt, 1984; Legendre et al., 1992; Gosselin et al., 1997) and their influence on surface water inorganic carbon distributions and air-sea  $\text{CO}_2$  disequilibrium.

Other factors may also influence the marine inorganic carbon cycle and present-day  $\text{CO}_2$  sink in the Arctic Ocean. For example, reduced sea-ice cover has been proposed to favor a “phytoplankton–zooplankton” dominated ecosystem rather than a “sea-ice algae–benthos” dominated ecosystem over the Arctic shelves (Piepenburg, 2005). At present on the highly productive Chukchi Sea shelf,  $\sim 10\%$  of NCP is converted to dissolved organic carbon (DOC) and  $\sim 15\%$  of NCP is converted to suspended particulate organic carbon (POC) (Mathis et al., 2007b, 2009) that gets exported from the shelf into the Canada Basin beneath the mixed layer (Bates et al., 2005b). The remaining 75% of NCP is exported from the mixed layer to the sea floor as sinking particles (Figs. 2 and 3, process 15), that sustain the rich benthos on the sea floor of the Chukchi Sea shelf (Grebmeier et al., 2008). In the Bering Sea, earlier sea-ice loss has led to ecosystem changes and altered pelagic–benthic coupling (e.g. Stabeno et al., 2001, 2002; Hunt et al., 2002; Overland et al., 2001; Macklin et al., 2002; Napp et al., 2002; Bond et al., 2003; Merico et al., 2004). If there are ecosystem shifts in the future, for example on the Chukchi and Beaufort Seas shelves, the export of organic carbon and pelagic–benthic coupling might decrease, despite concurrent increases in phytoplankton PP.

There may also be changes in the inorganic carbon distributions of subsurface waters on the Arctic shelves and halocline layer waters of the central basin. In the central basin, density stratification generally acts as a barrier to mixing between nutrient-poor surface waters and nutrient-rich deep waters (Aagaard et al., 1981; Jones and Anderson, 1986). Halocline waters generally have much higher  $p\text{CO}_2$  and DIC content than surface waters (Jutterstrom and Anderson, 2004), with low rates of upward vertical diffusion between these waters and the surface mixed layer (Wallace et al., 1987; Wilson and Wallace, 1990; Bates, 2006). The mixed layer typically extends to 10–50 m, with depth heavily dependent on seasonality of winds and sea-ice cover (Carmack and Macdonald, 2002). The reduction of sea-ice extent may facilitate deeper mixing and erosion into inorganic nutrient and  $\text{CO}_2$  rich halocline waters. Depending on the stoichiometry of nutrients and carbon entrained into the mixed layer, enhancement of phytoplankton PP in typically highly oligotrophic surface waters of the Arctic (e.g. English, 1961) and subsequent vertical export of organic carbon, this mech-

anism could either enhance or suppress air-sea CO<sub>2</sub> disequilibrium and CO<sub>2</sub> exchange between atmosphere and ocean (Figs. 2 and 3, process 12). The halocline is believed to ventilate predominantly through lateral processes and the transfer of water from the shelves into the interior formation of dense water on the shelves via brine-driven convection (Weingartner et al., 1998).

If the marine inorganic carbon cycle on the Arctic shelves is altered, the transfer of inorganic and organic carbon into the halocline may also change. For example, if there is increased transport of DOC from the shelves into the subsurface halocline of the central basin, remineralization of DOC by the microbial community may enhance *p*CO<sub>2</sub> and DIC contents of surface waters, and in turn, the upward supply of CO<sub>2</sub> to the surface layer through vertical diffusion and mixing (Figs. 2 and 3, process 12). The release of alkalinity from the sediments, produced by anaerobic microbial decomposition, may also be important for suppressing seawater *p*CO<sub>2</sub> contents in shelf waters that could in turn enhance air-sea CO<sub>2</sub> disequilibrium (Thomas et al., 2009). However, no evidence has yet been forthcoming about the potential influence of this process for Arctic shelves (Figs. 2 and 3, process 18). In summary, ecosystem changes in the Arctic Ocean shelves and in the central basin, both in the surface and underlying halocline layer, may alter inorganic carbon distributions and the CO<sub>2</sub> sink or source terms of the Arctic as a whole.

#### 5.4 Changes in the hydrological cycle and freshwater inputs to the Arctic Ocean

Significant changes in the hydrological cycle of the Arctic are also predicted by the end of the 21st century (ACIA 2005). Freshwater inputs from Arctic rivers, transport of sediment and dissolved materials and coastal erosion (Figs. 2 and 3, process 3) constitute a flux of  $\sim 12 \text{ Tg C yr}^{-1}$  at present (Macdonald et al., 2009) and expected to increase. At present, Arctic rivers influence the hydrographic properties of the Siberian Sea and Beaufort Sea shelves in particular. The eastern Beaufort Sea shelf is heavily impacted by Mackenzie River inflow of freshwater (e.g. Macdonald et al., 1995, 1999, 2002, 2009; Kadko and Swart, 2005) and plumes of elevated sediment loads and chromophoric dissolved organic matter (CDOM) can be followed from the Mackenzie outflow into the Canada Basin using satellite remote sensing. However, the impact of increased river flow on the chemical characteristics of the Arctic is uncertain at present. The freshwater input of DOC will presumably increase, but its contribution to inorganic carbon distributions on the shelves from bacterial remineralization of DOC to CO<sub>2</sub> is highly dependent on the lability of DOC and its decomposition relative to residence times in the Arctic. While most riverine DOC is thought to be highly refractory (Hansell et al., 2004), enhanced open-water may increase the photochemical breakdown of refractory DOC to CO<sub>2</sub> in surface waters. Given the relatively small flux of carbon from freshwater and coastal

erosion, this process is likely to influence the CO<sub>2</sub> sinks and sources of the Arctic over the longer-term rather than the next decade or so.

#### 5.5 Summary of near-term impacts on Arctic Ocean air-sea CO<sub>2</sub> fluxes

In the near-term, summertime sea-ice loss is expected to increase the uptake of CO<sub>2</sub> by surface waters because the exposed surface waters have a lower *p*CO<sub>2</sub> than the atmosphere. The loss of sea-ice in the Arctic Ocean also is likely to result in greater open water area, and increased air-sea interaction, with the likelihood of increased rates of air-sea gas exchange. Wintertime sea-ice is now thinner than previous decades and there may be potentially greater atmosphere-ocean gas exchange directly through sea-ice, with sea-ice a potentially weaker barrier to gas exchange. As a consequence of increased phytoplankton growth due to sea-ice loss and expansion of open water in the Arctic, the potential for the Arctic Ocean to uptake CO<sub>2</sub> should also increase.

In mitigation, reduced cooling of water during transit poleward and increased absorption of solar radiation resulting in warming of surface water relative to previous decades should act to increase the *p*CO<sub>2</sub> of seawater. This process may somewhat counteract the impacts of sea-ice loss and increased phytoplankton growth on the atmosphere-ocean exchange of CO<sub>2</sub>.

#### 5.6 Ocean acidification effects and changes in carbonate mineral saturation states

As a consequence of the ocean uptake of anthropogenic CO<sub>2</sub>, surface *p*CO<sub>2</sub> and DIC concentration have increased while pH has decreased in the upper ocean over the last few decades (e.g. Winn et al., 1994; Bates et al., 1996; Bates, 2007; Bates and Peters, 2007; Takahashi et al., 2009). This gradual process, termed *ocean acidification*, has long been recognized (e.g. Broecker and Takahashi, 1966; Broecker et al., 1971; Bacastow and Keeling, 1973). Estimates based on the Intergovernmental Panel on Climate Change (IPCC) projections of future CO<sub>2</sub> emission and subsequent ocean absorption of anthropogenic CO<sub>2</sub> (Bindoff et al., 2007; IPCC, 2007) predict that upper ocean pH will decrease by 0.3–0.5 units over the next century and beyond (Caldeira and Wickett, 2003, 2005; Doney, 2006). The effects of ocean acidification are potentially far-reaching in the global ocean, particularly for calcifying fauna (e.g. Buddemeier et al., 2004; Royal Society, 2005; Fabry et al., 2008). Decreased pH reduces the saturation states ( $\Omega$ ) of calcium carbonate (CaCO<sub>3</sub>) minerals such as aragonite and calcite, and increases the potential for reduced production of CaCO<sub>3</sub> by calcifying fauna (e.g. Buddemeier et al., 2004; Royal Society, 2005; Fabry et al., 2008; Doney et al., 2009) and dissolution of CaCO<sub>3</sub> in the water-column and sediments. Model studies suggest that CaCO<sub>3</sub> mineral undersaturation due to

ocean acidification should impact the Arctic before other regions (Orr et al., 2005, 2006; Steinacher et al., 2009).

In the Arctic Ocean, sea-ice melt, river runoff and phytoplankton photosynthesis and respiration appear to be the dominant processes that induce divergent directions for  $\text{CaCO}_3$  saturation states for surface and subsurface waters. As such, these processes may suppress or enhance  $\Omega$  values that in turn amplify or mitigate the impacts of ocean acidification in the Arctic.

### 5.7 Suppression and enhancement of $\Omega$ in surface waters

In surface waters, freshwater contributions from sea-ice melt and river runoff appear to be the primary processes that act to decrease  $\text{CaCO}_3$  saturation states. Salisbury et al. (2008) have shown that the  $\Omega$  values of major Arctic rivers are close to zero for the run-off end-member. In the Chukchi Sea and Canada Basin, surface mixed layer waters were generally saturated with respect to aragonite ( $\Omega_{\text{aragonite}}$ ) but  $\Omega_{\text{aragonite}}$  values decreased as runoff fraction increased (Bates et al., 2009). These findings are consistent with observations that a wide variety of Arctic rivers are net heterotrophic (e.g. Chen and Borges, 2009) with high  $p\text{CO}_2$  contents (Kelley, 1970; Makkayeev, 1994; Semiletov, 1999; Semiletov et al., 2007; Nitishinsky et al., 2007) and corresponding low  $\Omega$  values. River DOC remineralization and respiration in combination with many Arctic rivers having high DIC:TA ratios (Anderson et al., 1988; Olsson and Anderson, 1997; Cooper et al., 2008; Bates et al., 2009) act to decrease  $\Omega$  values.

$\Omega$  values have also been observed to decrease with increasing sea-ice melt fraction (>15%; Bates et al., 2009), and in a few localized surface areas in the Canada Basin, both aragonite and calcite undersaturation have been observed. Similarly, Chierici and Fransson (2009) have observed localized areas where surface waters were undersaturated with respect to aragonite ( $\Omega_{\text{aragonite}}$ ) on the Canadian Archipelago and Beaufort Sea shelves which they attribute to the influence of freshwater contributions from sea-ice melt and river runoff. Cooling and air-sea uptake of  $\text{CO}_2$  (which increases  $p\text{CO}_2$  and DIC) will also act to decrease  $\Omega$  values, but are relatively minor impacts compared to sea-ice melt and river runoff inputs to the Arctic. Low  $\Omega$  values have also been attributed to upwelling of  $\text{CO}_2$ -rich subsurface waters off the Chukchi Sea shelf (Chierici and Fransson, 2009).

Oversaturated  $\Omega$  values are been observed for surface waters across much of the Chukchi Sea shelf and adjacent Canada Basin (Jutterstrom and Anderson, 2005; Bates et al., 2009; Chierici and Fransson, 2009). The oversaturated conditions for both aragonite and calcite have been attributed to the high rates of phytoplankton PP that decrease  $p\text{CO}_2$  and DIC, and increase  $\Omega$  values on the Chukchi Sea shelf (Bates et al., 2009), as well as low DIC:TA ratios and Revelle Factors present in the Canada Basin (Bates, 2006) that facilitates oversaturated  $\Omega$  conditions.

### 5.8 Suppression of $\Omega$ in subsurface waters

In the North Pacific Ocean, Feely et al. (1988) observed seasonal enhancement of surface  $\Omega$  values and suppression of subsurface  $\Omega$  values due to seasonal phytoplankton PP, export of organic matter and subsequent remineralization to  $\text{CO}_2$ . A similar mechanism occurs in the Chukchi Sea where high rates of summertime phytoplankton PP (Hill and Cota, 2005; Bates et al., 2005; Mathis et al., 2009), vertical export of organic carbon (Lepore et al., 2007) and buildup of  $\text{CO}_2$  in subsurface waters (primarily through subsurface remineralization of OM to  $\text{CO}_2$ , and benthic respiration) suppresses  $\Omega$  values below saturation. Subsurface waters were observed to be undersaturated with respect to aragonite and calcite, and thus potentially corrosive to  $\text{CaCO}_3$  for the Chukchi Sea shelf sediments and benthic ecosystem.

### 5.9 Amplification and mitigation of on acidification impacts in the Arctic Ocean

In many oceanic basins, the oceanic uptake of anthropogenic  $\text{CO}_2$  has resulted in the shoaling of the aragonite saturation horizon (i.e.  $\Omega_{\text{aragonite}} = 1$ ) by 40 to 200 m (Feely et al., 2008) over the past century. Given the scenarios for pH changes in the Arctic Ocean (Orr et al., 2005; Steinacher et al., 2009), the Arctic shelves will be increasingly impacted by ocean acidification and presence of  $\text{CaCO}_3$  mineral undersaturated waters. Superimposed on the ocean acidification effects induced by ocean uptake of anthropogenic  $\text{CO}_2$ , seasonal biological processes are likely to mitigate (enhance  $\Omega$ ) and amplify (suppress  $\Omega$ ) the impacts of ocean acidification on the Arctic Ocean shelves.

The seasonal changes in  $\Omega$  induced by biology primarily have been described as a seasonal “*Phytoplankton-Carbonate Saturation State*” (PhyCASS) interaction that drives divergent trajectories for carbonate chemistry in surface and subsurface waters of Arctic Ocean shelves like the Chukchi Sea (Bates et al., 2009). For inflow shelves (Carmack and Wassmann, 2006) like the Chukchi Sea and Barents Sea, high rates of summertime phytoplankton PP or NCP and vertical export of OM during seasonal sea-ice loss may drive a strong seasonal divergence in surface and subsurface  $\Omega$ . The seasonal divergence in  $\Omega$  is perhaps attenuated in the Barents Sea compared to the Chukchi Sea since the biogeochemical modification of  $\Omega$  is probably dispersed within a deeper mixed layer and thicker subsurface layer. For interior shelves such as the Siberian and Beaufort Sea shelves, much lower rates of summertime phytoplankton PP, and much greater influence of river runoff (with lower  $\Omega$ ) is also likely to result in attenuated divergence in  $\Omega$  for surface and subsurface waters compared to the Chukchi Sea. On all shelves, a combination of Pacific or Atlantic Ocean water inflow and wintertime homogenization of shelf waters through wind-mixing and brine rejection processes are likely

to “rectify” the seasonal divergences in  $\Omega$  between surface and subsurface waters.

In an era of rapid change in the Arctic, sea-ice loss, surface warming and increased phytoplankton PP on the Arctic shelves may act to enhance surface  $\Omega$  and suppress subsurface  $\Omega$  amplifying the existing impact of ocean acidification. This may be mitigated somewhat by increased freshwater contributions through sea-ice melt and river runoff that suppress surface water  $\Omega$ . In the central basin of the Arctic Ocean, potentially corrosive waters to aragonite and calcite are found in the upper halocline layer (Jutterstrom and Anderson, 2005; Bates et al., 2009), with highly saturated  $\Omega$  values in surface water. Increased air-sea interaction in regions experiencing continued seasonal sea-ice loss will likely exhibit increased vertical mixing of  $\text{CO}_2$ -rich subsurface waters, increased shelf-basin exchanges and ocean uptake of  $\text{CO}_2$  through gas exchange. Combined, these factors will reduce the vertical gradients of  $\Omega$  between surface and subsurface with the net result of suppressing surface  $\Omega$  over the residence time of surface water in the central basin (2–30 years; Macdonald et al., 2009).

## 6 Conclusions

At present, the Arctic Ocean continental shelves and central basin have lower  $\text{CO}_2$  content than the atmosphere, and these areas are potential sinks of  $\text{CO}_2$  from the atmosphere. There are however localized areas of sea-ice melt and river inputs where the opposite is observed, and these areas are potential sources of  $\text{CO}_2$  to the atmosphere. The  $\text{CO}_2$  chemistry of the Arctic Ocean is highly influenced by physical and biological processes such as seasonal phytoplankton PP during summertime sea-ice retreat towards the pole, as well as temperature effects (both cooling and warming), shelf-basin exchanges and formation of dense winter waters, and river inputs of freshwater and carbon.

Although seasonal sea-ice cover provides a barrier to atmosphere-ocean gas exchange, the Arctic Ocean is a sink for  $\text{CO}_2$ , on the order of  $-65$  to  $-199 \text{ Tg C year}^{-1}$ , contributing 5–14% to the global balance of  $\text{CO}_2$  sinks and sources. The Arctic Ocean has become an important influence on the global carbon cycle, with the marine carbon cycle and atmosphere-ocean  $\text{CO}_2$  exchanges sensitive to Arctic Ocean and global climate change feedbacks. In the near-term, further sea-ice loss and increases in phytoplankton PP rates are expected to increase the uptake of  $\text{CO}_2$  by Arctic surface waters, although mitigated somewhat by warming in the Arctic. Thus, the capacity of the Arctic Ocean to uptake  $\text{CO}_2$  is expected to increase and alter in response to environmental changes driven largely by climate. These changes are likely to continue to modify the physics, biogeochemistry, and ecology of the Arctic Ocean in ways that are not yet fully understood. Finally, in response to increased phytoplankton PP and uptake of

human-produced  $\text{CO}_2$ , the benthic ecosystem of the Arctic shelves are expected to be negatively impacted by ocean acidification which reduces the ability of many species to produce calcium carbonate shells or tests with profound implications for Arctic marine ecosystems.

Edited by: A. V. Borges

## References

- Aagaard, K., Coachman, L. K., and Carmack, E. C.: On the halocline of the Arctic Ocean, *Deep-Sea Res.*, 28A(6), 529–545, 1981.
- Aagaard, K. and Carmack, E. C.: The role of sea ice and other fresh water in the Arctic circulation, *J. Geophys. Res.*, 94(C10), 14485–14498, 1989.
- Aagaard, K. and Carmack, E. C.: The Arctic Ocean and Climate: A perspective, in: *The Polar Oceans and Their Role in Shaping the Global Environment*, edited by: Johannessen, O. M., Muench, R. D., and Overland, J. E., American Geophysical Union, Geophys. Monogr., 85, 5–20, 1994.
- Alonso-Saez, L., Sanchez, O., Gasol, J. M., Balague, V., and Pedros-Alio, C.: Winter-to-summer changes in the composition and single-cell activity of near-surface Arctic prokaryotes, *Environ. Microbiol.*, 10(9), 2444–2454, doi:10.1111/j.1462-2920.2008.01674.x, 2008.
- Anderson, L. G.: Chemical oceanography of the Arctic and its shelf sea, in: *Arctic Oceanography: Marginal Ice Zones and Continental Shelves*, edited by: Smith Jr., W. O. and Grebmeier, J. M., Coastal and Estuarine Studies, American Geophysical Union, Washington, DC, USA, 183–202, 1995.
- Anderson, L. G. and Kaltin, S.: Carbon fluxes in the Arctic Ocean – potential impact by climate change, *Polar Res.*, 20(2), 225–232, 2001.
- Anderson, L. G., Jones, E. P., Lindegren, R., Rudels, B., and Sehlstedt, P. I.: Nutrient regeneration in cold, high salinity bottom water of the Arctic shelves, *Cont. Shelf. Res.*, 8, 1345–1355, 1988.
- Anderson, L. G., Dyrssen, D., and Jones, E. P.: An assessment of the transport of atmospheric  $\text{CO}_2$  into the Arctic ocean, *J. Geophys. Res.*, 95(C2), 1703–1711, 1990.
- Anderson, L. G., Olsson, K., and Skoog, A.: Distribution of dissolved inorganic and organic carbon in the Eurasian Basin of the Arctic Ocean, in: *The Polar Oceans and Their Role in Shaping the Global Environment*, edited by: Johannessen, O. M., Muench, R. D., and Overland, J. E., American Geophysical Union, Geophysical Monograph, 85, 252–262, 1994a.
- Anderson, L. G., Bjork, G., Holby, O., Jones, E. P., Kattner, G., Koltermann, K. P., Liujeband, B., Lindegren, R., Rudels, R., and Swift, J.: Water masses and circulation in the Eurasian Basin: Results from the Oden 91 Expedition, *J. Geophys. Res.-Oceans*, 99(C2), 3273–3283, 1994b.
- Anderson, L. G., Olsson, K., and Chierici, M.: A carbon budget for the Arctic Ocean, *Global Biogeochem. Cy.*, 12(3), 455–465, 1998a.
- Anderson, L. G., Olsson, K., Jones, E. P., Chierici, M., and Fransson, A.: Anthropogenic carbon dioxide in the Arctic Ocean: Inventory and sinks, *J. Geophys. Res.*, 10(C12), 27707–27716, 1998b.

- Anderson, L. G., Jones, E. P., and Rudels, B.: Ventilation of the Arctic Ocean estimated by a plume entrainment model constrained by CFCs, *J. Geophys. Res.*, 104(C6), 13423–13429, 1999.
- Anderson, L. G., Falck, E., Jones, E. P., Jutterstrom, S., and Swift, J. H.: Enhanced uptake of atmospheric CO<sub>2</sub> during freezing of seawater: A field study in Storfjorden, Svalbard, *J. Geophys. Res.*, 109(C6), C06004, doi:10.1029/2003JC002120, 2004.
- Arctic Climate Impact Assessment (ACIA): Cambridge University Press, Cambridge, UK and NY, NY, USA, 2005.
- Arrigo, K. R., van Dijken, G. L., and Pabi, S.: The impact of a shrinking Arctic ice cover on marine primary production, *Geophys. Res. Lett.*, 35, L19603, doi:10.1029/2008GL035028, 2008.
- Bacastow, R. D. and Keeling, C. D.: Atmospheric carbon dioxide and radiocarbon in the natural carbon cycle II. Changes from A. D. 1700 to 2070 as deduced from a geochemical model, in: *Carbon and the biosphere*, edited by: Woodwell, G. M. and Pecan, E. V., US Atomic Energy Commission, 86–135, 1973.
- Bates, N. R.: Air-sea CO<sub>2</sub> fluxes and the continental shelf pump of carbon in the Chukchi Sea adjacent to the Arctic Ocean, *J. Geophys. Res. Oceans*, 111, C10013, doi:10.129/2005JC003083, 2006.
- Bates, N. R.: Interannual variability of the oceanic CO<sub>2</sub> sink in the subtropical gyre of the North Atlantic Ocean over the last two decades, *J. Geophys. Res. Oceans*, 112, C09013, doi:2006JC003759, 2007.
- Bates, N. R. and Merlivat, L.: The influence of short-term wind variability on air-sea CO<sub>2</sub> exchange, *Geophys. Res. Lett.*, 28(17), 3281–3284, 2001.
- Bates, N. R. and Peters, A. J.: The contribution of atmospheric acid deposition to ocean acidification in the subtropical North Atlantic Ocean, *Mar. Chem.*, 107, 547–558, doi:10.1016/j.marchem.2007.08.002, 2007.
- Bates, N. R., Michaels, A. F., and Knap, A. H.: Seasonal and interannual variability of the oceanic carbon dioxide system at the US, JGOFS Bermuda Atlantic Time-series Site, Deep-Sea Res. II, 43(2–3), 347–383, doi:10.1016/0967-0645(95)00093-3, 1996.
- Bates, N. R., Best, M. H. P., and Hansell, D. A.: Spatio-temporal distribution of dissolved inorganic carbon and net community production in the Chukchi and Beaufort Seas, *Deep-Sea Res. II*, 52(22–24), 3303–3323, doi:10.1016/j.dsr2.2005.10.005, 2005a.
- Bates, N. R., Hansell, D. A., Moran, S. B., and Codispoti, L. A.: Seasonal and spatial distributions of particulate organic matter (POM) in the Chukchi and Beaufort Seas, *Deep-Sea Res. II*, 52(22–24), 3324–3343, doi:10.1016/j.dsr2.2005.10.003, 2005b.
- Bates, N. R., Moran, S. B., Hansell, D. A., and Mathis, J. T.: An increasing CO<sub>2</sub> sink in the Arctic Ocean due to sea-ice loss, *Geophys. Res. Lett.*, 33(23), L23609, doi:10.1029/2006GL027028, 2006.
- Bates, N. R., Mathis, J. T., and Cooper, L.: The effect of ocean acidification on biologically induced seasonality of carbonate mineral saturation states in the Western Arctic Ocean, *J. Geophys. Res.-Oceans*, doi:10.1029/2008JC004862, in press, 2009.
- Benner, R., Benitez-Nelson, B., Kaiser, K., and Amon, R. M. W.: Export of young terrigenous dissolved organic carbon from rivers to the Arctic Ocean, *Geophys. Res. Lett.*, 31, doi:10.1029/2003GL019251, 2004.
- Bindoff, N. L., Willebrand, J., Artale, V., Cazenave, A., Gregory, J., Gulev, S., Hanawa, K., Le Quéré, C., Levitus, S., Nojiri, Y., Shum, C. K., Talley, L. D., and Unnikrishnan, A.: Observations: Oceanic Climate Change and Sea Level, in: *Climate Change 2007: The Physical Science Basis, Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L.: Cambridge University Press, Cambridge, UK and NY, NY, USA, 2007.
- Björk, G.: A one-dimensional time-dependent model for the vertical stratification of the upper Arctic Ocean, *J. Phys. Oceanogr.*, 19, 52–67, 1989.
- Bond, N. A., Overland, J. E., Spillane, M., and Stabeno, P.: Recent shifts in the state of the North Pacific, *Geophys. Res. Lett.*, 30(23), 2183, doi:10.1029/2003GL018597, 2003.
- Borges, A. V.: Do we have enough pieces of the jigsaw to integrate CO<sub>2</sub> fluxes in the coastal ocean?, *Estuaries*, 28(1), 3–27, 2005.
- Borges, A. V., Delille, B., and Frankignoulle, M.: Budgeting sinks and sources of CO<sub>2</sub> in the coastal ocean: Diversity of ecosystems counts, *Geophys. Res. Lett.*, 32, L14601, doi:10.1029/2005GL023053, 2005.
- Broecker, W. S. and Takahashi, T.: Calcium carbonate precipitation on the Bahama Banks, *J. Geophys. Res.*, 71, 1575–1602, 1966.
- Broecker, W. S., Li, Y.-H., Peng, T.-H.: Carbon dioxide – man’s unseen artifact, in: *Impingement of man on the oceans*, edited by: Hood, D. W., John Wiley and Sons, Inc, 287–324, 1971.
- Buddemeier, R. W., Kleypas, J. A., and Aronson, R. B.: Coral Reefs and Global Climate Change: Potential Contributions of Climate Change to Stresses on Coral Reef Ecosystems, 44, (download report at: [http://www.pewclimate.org/global-warming/indepth/all-reports/coral\\_reefs/index.cfm](http://www.pewclimate.org/global-warming/indepth/all-reports/coral_reefs/index.cfm)), Pew Center on Climate Change, 2004.
- Cai, W.-J. and Dai, M.: A Comment on “Enhanced open ocean storage of CO<sub>2</sub> from shelf sea pumping”, *Science*, 306, 1477c, 2004.
- Cai, W.-J., Dai, M., and Wang, Y.: Air-sea exchange of carbon dioxide in ocean margins: A province-based synthesis, *Geophys. Res. Lett.*, 33, L12603, doi:10.1029/2006GL026219, 2006.
- Caldeira, K. and Wickett, M. E.: Anthropogenic carbon and ocean pH, *Nature*, 425(6956), 365–365, doi:10.1038/425365a, 2003.
- Caldeira, K. and Wickett, M. E.: Ocean model predictions of chemistry changes from carbon dioxide emissions to the atmosphere and ocean, *J. Geophys. Res., Oceans*, 110(C9), C09S04, doi:10.1029/2004JC002671, 2005.
- Carmack, E. C. and Chapman, D. C.: Wind-driven shelf/basin exchange on an Arctic shelf: The joint roles of ice cover extent and shelf-break bathymetry, *Geophys. Res. Lett.*, 30(14), 1778, doi:10.1029/2003GL017526, 2003.
- Carmack, E. C. and Macdonald, R. W.: Oceanography of the Canadian shelf of the Beaufort Sea: A setting for marine life, *Arctic*, 55, Supplement 1, 29–45, 2002.
- Carmack, E. and Wassmann, P.: Food webs and physical-biological coupling on pan-Arctic shelves: Unifying concepts and comprehensive perspectives, *Prog. Oceanogr.*, 71, 446–477, doi:10.1016/j.pocean.2006.10.004, 2006.
- Cavaliere, D. J. and Martin, S.: The contribution of Alaskan, Siberian, and Canadian coastal polynyas to the cold halocline layer of the Arctic Ocean, *J. Geophys. Res.*, 99(C9), 18343–18362, 1994.
- Cavaliere, D. J., Parkinson, C. L., and Vinnikov, K. Y.: 30-year satellite record reveals contrasting Arctic and Antarctic

- decadal sea ice variability, *Geophys. Res. Lett.*, 30(18), 1970, doi:10.1029/2003GL018031, 2003.
- Cauwet, G. and Sidorov, I.: The biogeochemistry of Lena River: organic carbon and nutrients distribution, *Marine Chemistry*, 53, 211–227, 1996.
- Chapman, W. L. and Walsh, J. E.: Observed climate change in the Arctic: updated from Chapman and Walsh, 1993: Recent variations of sea ice and air temperatures in high latitudes, *Bull. Am. Met. Soc.*, 74(1), 33–47, 2003.
- Chen, C. T. A. and Borges, A. V.: Reconciling opposing views on carbon cycling in the coastal ocean: Continental shelves as sinks and near-shore ecosystems as sources of atmospheric CO<sub>2</sub>. *Deep-Sea Res. II*, 56(8–10), 578–581, doi:10.1016/j.dsr2.2009.01.001, 2009.
- Chen, L. Q. and Gao, Z. Y.: Spatial variability in the partial pressures of CO<sub>2</sub> in the northern Bering and Chukchi seas, *Deep-Sea Res. II*, 54(23–26), 2619–2629, doi:10.1016/j.dsr2.2007.08.010, 2007.
- Chierici, M. and Fransson, A.: Calcium carbonate saturation in the surface water of the Arctic Ocean: undersaturation in freshwater influenced shelves, *Biogeosci. Discuss.*, 6, 4963–4991, 2009.
- Coachman, L. K., Aagaard, K., and Tripp, R. B.: *Bering Strait: The Regional Physical Oceanography*, University of Washington Press, Seattle, 169 pp., 1975.
- Codispoti, L., Flagg, C., and Kelly, V.: Hydrographic conditions during the 2002 SBI process experiments. *Deep Sea Res., Part II*, 52, 3199–3226, 2005.
- Comiso, J. C., Parkinson, C. L., Gersten, R., and Stock, L.: Accelerated decline in the Arctic sea ice cover, *Geophys. Res. Lett.*, 35, L01703, doi:10.1029/2007GL031972, 2008.
- Cooper, L. W., Benner, R., McClelland, J. R., Peterson, B. J., Holmes, R. M., Raymond, P. A., Hansell, D. A., Grebmeier, J. M., and Codispoti, L. A.: Linkages among runoff, dissolved organic carbon, and the stable oxygen isotope composition of seawater and other water mass indicators in the Arctic Ocean, *J. Geophys. Res.*, 110, G02013, doi:10.1029/2005JG000031, 2005.
- Cooper, L. W., McClelland, J. W., Holmes, R. M., Raymond, P. A., Gibson, J. J., Guay, C. K., and Peterson, B. J.: Flow-weighted values of runoff tracers ( $\delta^{18}\text{O}$ , DOC, Ba, alkalinity) from the six largest Arctic rivers. *Geophys. Res. Lett.*, 35, L18606, doi:10.1029/2008GL035007, 2008.
- Cota, G. F., Pomeroy, L. R., Harrison, W. G., Jones, E. P., Peters, F., Sheldon W. M., and Weingartner T. R.: Nutrients, primary production and microbial heterotrophy in the southeastern Chukchi Sea: Arctic summer nutrient depletion and heterotrophy, *Marine Ecol., Prog. Ser.*, 135(1–3), 247–258, 1996.
- Delille, B., Jourdain, B., Borges, A. V., Tison, J.-L., and Delille, D.: Biogas (CO<sub>2</sub>, O<sub>2</sub>, dimethylsulfide) dynamics in spring Antarctic fast ice, *Limnol. Oceanogr.*, 52, 1367–1379, 2007.
- Dickson, A. G.: Thermodynamics of the dissolution of boric acid in synthetic seawater from 273.15°K to 318.15°K, *Deep-Sea Res.*, 37A(5), 755–766, 1990.
- Dickson, A. G. and Millero, F. J.: A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media, *Deep-Sea Res.*, 34, 1733–1743, 1987.
- Dickson, A. G., Sabine, C. L., and Christian, J. R.: *Guide to best practices for ocean CO<sub>2</sub> measurements*, Sidney, British Columbia, North Pacific Marine Science Organization, PICES Special Publication 3, 2007.
- Dickson, R. R., Osborn, T. J., Hurrell, J. W., Meincke, J., Blindheim, J., Adlandsvik, B., Vinje, T., Alekseev, G., and Maslowski, W.: Arctic ocean response to the North Atlantic oscillation, *J. Climate*, 13, 2671–2696, 2000.
- Dieckmann, G. S., Nehrke, G., Papadimitriou, S., Göttlicher, J., Steininger, R., Kennedy, H., Wolf-Gladrow, D., and Thomas, D. N.: Calcium carbonate as ikaite crystals in Antarctic sea ice, *Geophys. Res. Lett.*, 35, L08501, doi:10.1029/2008GL033540, 2008.
- Dittmar, T. and Kattner, G.: The biogeochemistry of the river and shelf ecosystem of the Arctic Ocean: a review, *Mar. Chem.*, 83, 103–120, doi:10.1016/S0304-4203(03)00105-1, 2003.
- Dmitrenko, I. A., Tyshko, K. N., Kirillov, S. A., Eicken, H., Hölemann, J. A., and Kassense, H.-M.: Impact of flaw polynyas on the hydrography of the Laptev Sea, *Global Planet. Change*, 48(1–3), 9–27, doi:10.1016/j.gloplacha.2004.12.016, 2005.
- Dmitrenko, I. A., Kirillov, S. A., and Tremblay, B. L.: The long-term and interannual variability of summer fresh water storage over the eastern Siberian shelf: Implications for climate change, *J. Geophys. Res.*, 113, C03007, doi:10.1029/2007JC004304, 2008.
- DOE: *Handbook of Methods for the Analysis of the Various Parameters of the Carbon Dioxide System in Seawater*; version 2.0, (Dickson, A. G. and Goyet, C.), US Department of Energy CO<sub>2</sub> Science Team Report, 1994.
- Doney, S. C.: The dangers of ocean acidification, *Sci. Am.*, 294(3), 58–65, 2006.
- Doney, S. C., Fabry, V. J., Feely, R. A., and Kleypas, J. A.: Ocean acidification: The other CO<sub>2</sub> problem, *Annual Reviews of Marine Science*, 1, 169–192, 2009.
- Ducklow, H. W. and McAllister, S. L.: Biogeochemistry of carbon dioxide in the coastal oceans, in: *The Sea, Volume 13, The Global Coastal Ocean-Multiscale Interdisciplinary Processes*, edited by: Robinson, A. R. and Brink, K., J. Wiley and Sons, NY, USA, 2005.
- Else, B. G. T., Papakyriakou, T. N., Granskog, M. A., and Yackel, J. J.: Observations of sea surface *f*CO<sub>2</sub> distributions and estimated air-sea CO<sub>2</sub> fluxes in the Hudson Bay region (Canada) during the open water season, *J. Geophys. Res. Oceans*, 113, C08026, doi:10.1029/2007JC004389, 2008.
- English, T. S.: Some biological oceanographic observations in the central north Polar Sea, Drift Station Alpha, 1957–1958. *Arctic Institute of North America Scientific Report 15*, 1961.
- Fabry, V. J., Seibel, B. A., Feely, R. A., and Orr, J. C.: Impacts of ocean acidification on marine fauna and ecosystem processes, *ICES Journal of Marine Science*, 65, 414–432, doi:10.1093/icesjms/fsn048, 2008.
- Falkner, K. K., MacDonald, R. W., Carmack, E. C., and Weingartner, T.: The potential of barium as a tracer of Arctic water masses, in: *Polar Oceans and Their Role in Shaping the Global Environment*, edited by: Johannessen, O. M., Muench, R. D., and Overland, J. E., American Geophysical Unions, *Geophys. Monogr.*, 85, 63–76, 1994.
- Feely, R. A., Byrne, R. H., Acker, J. G., Betzer, P. R., Chen, C. T.-A., Gendron, J. R., and Lamb, M. F.: Winter summer variations of calcite and aragonite saturation in the northeast Pacific, *Mar. Chem.*, 25(3), 227–241, 1988.
- Feely, R. A., Sabine, C. L., Lee, K., Berelson, W., Kleypas, J., Fabry, V. J., and Millero, F. J.: Impact of anthropogenic CO<sub>2</sub> on

- the CaCO<sub>3</sub> system in the oceans, *Science*, 305(5682), 362–366, doi:10.1126/science.1097329, 2004.
- Feely, R. A., Sabine, C. L., Hernandez-Ayon, J. M., Ianson, D., and Hales, B.: Evidence for upwelling of corrosive “acidified” water onto the continental shelf, *Science*, 320, 1490–1492, doi:10.1126/science.1155676, 2008.
- Fransson, A., Chierici, M., Anderson, L. C., Bussmann, I., Kattner, G., Jones, E. P., and Swift, J. H.: The importance of shelf processes for the modification of chemical constituents in the waters of the Eurasian Arctic Ocean: implication for carbon fluxes, *Cont. Shelf Res.* 21(3), 225–242, doi:10.1016/S0278-4343(00)00088-1, 2001.
- Fransson, A., Chierici, M., and Nojiri, Y.: New insights into the spatial variability of the surface water carbon dioxide in varying sea ice conditions in the Arctic Ocean., *Cont. Shelf Res.*, 29(10), 1317–1328, doi:10.1016/j.csr.2009.03.008, 2009.
- Gammelsrod, T., Leikvin, O., Lien, V., Budgell, W. P., Loeng, H., and Maslowski, W.: Mass and heat transports in the NE Barents Sea: Observations and models, *J. Marine Syst.*, 75(1–2), 56–69, doi:10.1016/j.jmarsys.2008.07.010, 2009.
- Garneau, M. E., Vincent, W. F., Terrado, R., and Lovejoy, C.: Importance of particle-associated bacterial heterotrophy in a coastal Arctic ecosystem, *J. Marine Syst.*, 75(1–2), 185–197, doi:10.1016/j.jmarsys.2008.09.002, 2008.
- Gattuso, J. P., Frankignoulle, M., and Wollast, R.: Carbon and carbonate metabolism in coastal aquatic ecosystems, *Ann. Rev. Ecol. Syst.*, 29, 405–434, 1998.
- Giles, K. A., Laxon, S. W., and Ridout, A. L.: Circumpolar thinning of Arctic sea ice following the 2007 record ice extent minimum, *Geophys. Res. Lett.*, 35(22), L22502, doi:10.1029/2008GL035710, 2008.
- Gordeev, V. V., Martin, J. M., Sidorov, I. S., Sidorova, M. V.: A reassessment of the Eurasian River input of water, sediment, major elements, and nutrients to the Arctic Ocean, *J. Am. Sci.*, 296, 664–691, 1996.
- Gosink, T. A., Pearson, J. G., and Kelley, J. J.: Gas movement through sea ice, *Nature*, 263, 41–42, doi:10.1038/263041a1, 1976.
- Gosselin, M., Lavoie, M., Wheeler, P. A., Horner, R. A., and Booth, B. C.: New measurements of phytoplankton and ice algal production in the Arctic Ocean, *Deep-Sea Res. II*, 44, 1623–1644, doi:10.1016/S0967-0645(97)00054-4, 1997.
- Gow, A. J. and Tucker, W. B.: Sea ice in the polar regions, in: *Polar Oceanography*, edited by: Smith, W. O., Academic Press, London, UK, 47–122, 1990.
- Goyet, C., and Poisson, A. P.: New determination of carbonic acid dissociation constants in seawater as a function of temperature and salinity, *Deep-Sea Res.*, 36, 1635–1654, 1989.
- Grebmeier, J. M., Bates, N. R., and Devol, A.: Continental Margins of the Arctic Ocean and Bering Sea, in: *North American Continental Margins: A Synthesis and Planning Workshop*, US Carbon Cycle Science Program, Washington DC, edited by: Hales, B., Cai, W.-J., Mitchell, B. G., Sabine, C. L., and Schofield, O., 120 pp., 61–72, 2008.
- Guay, C. K. and Falkner, K. K.: Barium as a tracer of Arctic halocline and river waters, *Deep-Sea Res. II*, 44(8), 1543–1570, doi:10.1016/S0967-0645(97)00066-0, 1997.
- Guo, L., Semiletov, I., Gustafsson, O., Ingri, J., Andersson, P., Dudarev, O., and White, D.: Characterization of Siberian Arctic estuarine sediments: implications for terrestrial organic carbon export, *Global Biogeochem. Cy.*, 18, GB1036, doi:10.1029/2003GB002087, 2004.
- Häkkinen, S.: An Arctic source for the Great Salinity Anomaly: A simulation of the Arctic ice-ocean system for 1955–1975, *J. Geophys. Res.*, 98(C9), 16397–16410, 1993.
- Hameedi, M. J.: Aspects of water column primary productivity in Chukchi Sea during summer, *Mar. Biol.*, 48(1), 37–46, 1978.
- Hansell, D. A., Whitledge, T. E., and Goering, J. J.: Patterns of nitrate utilization and new production over the Bering-Chukchi shelf, *Cont. Shelf Res.*, 13, 601–628, 1993.
- Hansell, D. A., Kadko, D., and Bates, N. R.: Degradation of terrigenous dissolved organic carbon in the western Arctic Ocean. *Science*, 304(5672), 858–861, doi:10.1126/science.1096175, 2004.
- Hill, V. and Cota, G.: Spatial patterns of primary production on the shelf, slope and basin of the Western Arctic in 2002, *Deep-Sea Res. II*, 52(22–24), 3344–3354, doi:10.1016/j.dsr.2005.10.001, 2005.
- Holland, M. M., Bitz, C. M., and Tremblay, B.: Future abrupt reductions in the summer Arctic sea ice, *Geophys. Res. Lett.*, 33(23), L23503, doi:10.1029/2006GL028024, 2006.
- Hunt, G. L., Stabeno, P., Walters, G., Sinclair, E., Brodeur, R. D., Napp, J. M., and Bond, N. A.: Climate change and control of the southeastern Bering Sea pelagic ecosystem, *Deep-Sea Res. II*, 49(26), 5821–5853, doi:10.1016/S0967-0645(02)00321-1, 2002.
- IPCC, 2007: *The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Avery, K. B., Tignor, M., and Miller, H. L., Cambridge University Press, Cambridge, UK and NY, NY, USA, Climate Change, 996 pp., 2007.
- Jones, E. P. and Anderson, L. G.: On the origin of chemical properties of the Arctic Ocean halocline, *J. Geophys. Res.*, 91(C9), 10759–10767, 1986.
- Jones, E. P., Anderson, L. G., and Wallace, D. W. R.: Tracers of near-surface, halocline and deep waters in the Arctic Ocean: Implications for circulation, *J. Mar. Syst.*, 2, 241–255, 1991.
- Jones, E. P., Rudels, B., and Anderson, L. G.: Deep waters of the Arctic ocean – origin and circulation, *Deep-Sea Res. I*, 42, 737–760, 1995.
- Jones, E. P., Swift, J. H., Anderson, L. G., Lipzer, M., Civatarese, G., Falkner, K. K., Kattner, G., and McLaughlin, F.: Tracing Pacific water in the North Atlantic Ocean. *J. Geophys. Res.*, 108(C4), 3116, doi:10.1029/2001JC001141, 2003.
- Jones, E. P., Anderson, L. G., Jutterstrom, S., Mintrop, L., and Swift, J. H.: Pacific freshwater, river water and sea-ice meltwater across the Arctic Ocean basins: Results from the 2005 Beringia Expedition, *J. Geophys. Res.*, 113(C8), C08012, doi:10.1029/2007JC004124, 2008.
- Jutterström, S. and Anderson, L. G.: The saturation of calcite and aragonite in the Arctic Ocean, *Marine Chem.*, 94(1–4), 101–110, doi:10.1016/j.marchem.2004.08.010, 2005.
- Kadko, D. and Swart, P.: The source of the high heat and freshwater content of the upper ocean at the SHEBA site in the Beaufort Sea in 1997, *J. Geophys. Res.*, 109, C01022, doi:10.1029/2002JC001734, 2004.
- Kaltin, S. and Anderson, L. G.: Uptake of atmospheric carbon dioxide in Arctic shelf seas: Evaluation of the relative importance of processes that influence pCO<sub>2</sub> in water transported

- over the Bering–Chukchi Sea shelf, *Mar. Chem.*, 94(1–4), 67–79, doi:10.1016/j.marchem.2004.07.010, 2005.
- Kaltin, S., Anderson, L. G., Olsson, K., Fransson, A., and Chierici, M.: Uptake of atmospheric carbon dioxide in the Barents Sea, *J. Marine Syst.*, 38(1–2), 31–45, doi:10.1016/S0924-7963(02)00168-9, 2002.
- Kattner, G., Lobbes, J. M., Fitznar, H. P., Engbrodt, R., Nothig, E.-M., and Lara, R. J.: Tracing dissolved organic substances and nutrients from the Lena River through Laptev Sea (Arctic), *Marine Chemistry*, 65(1–2), 25–39, doi:10.1016/S0304-4203(99)00008-0, 1999.
- Kelley Jr., J. J.: Carbon dioxide in the surface waters of the North Atlantic and the Barents and Kara Sea, *Limnol. Oceanogr.*, 15(1), 80–87, 1970.
- Kinney, P., Arhelger, M. E., and Burrell, D. C.: Chemical characteristics of water masses in the Amerasian Basin of the Arctic Ocean, *J. Geophys. Res.*, 75, 4097–4104, 1970.
- Legendre, L., Ackley, S. F., Dieckmann, G. S., Gullicksen, B., Horner, R., Hoshiai, T., Melnikov, I. A., Reeburgh, W. S., Spindler, M., and Sullivan, C. W.: Ecology of sea ice biota: 2. Global significance, *Polar Biol.*, 12, 429–444, 1992.
- Lepore, K., Moran, S. B., Grebmeier, J. M., Cooper, L. W., Lalande, C., Maslowski, W., Hill, V., Bates, N. R., Hansell, D. A., Mathis, J. T., and Kelly, R. P.: Seasonal and interannual changes in particulate organic carbon export and deposition in the Chukchi Sea. *Journal of Geophysical Research*, 112, C10024, doi:10.1029/2006JC003555, 2007.
- Levi, B. G.: The decreasing Arctic ice cover. *Physics Today*, 53(1), 19–20, 2000.
- Lindsay, R. W. and Zhang, J.: The thinning of Arctic sea ice, 1988–2003: Have we passed a tipping point?, *J. Climate*, 18(22), 4879–4894, doi:10.1175/JCLI3587.1, 2005.
- Lundberg, L. and Haugen, P. M.: A Nordic Seas – Arctic Ocean carbon budget from volume flows and inorganic carbon data, *Global Biogeochem. Cy.*, 10, 493–510, 1996.
- Macdonald, R. W., Paton, D. W., Carmack, E. C., and Omstedt, A.: The fresh-water budget and under-ice spreading of Mackenzie River water in the Canadian Beaufort Sea based on salinity and  $O^{18}/O^{16}$  measurements in water and ice, *J. Geophys. Res.*, 100(C1), 895–919, 1995.
- Macdonald, R. W., Carmack, E. C., and Paton, D. W.: Using the delta  $O^{18}$  composition in landfast ice as a record of Arctic estuarine processes, *Marine Chemistry*, 65(1–2), 3–24, doi:10.1016/S0304-4203(99)00007-9, 1999.
- Macdonald, R. W., McLaughlin, F. A., and Carmack, E. C.: Fresh water and its sources during the SHEBA drift in the Canada Basin of the Arctic Ocean, *Deep Sea Res., Part I*, 49(10), 1769–1785, doi:10.1016/S0967-0637(02)00097-3, 2002.
- Macdonald, R. W., Anderson, L. G., Christensen, J. P., Miller, L. A., Semiletov, I. P., and Stein, R.: The Arctic Ocean, in: *Carbon and Nutrient Fluxes in Continental Margins: A Global Synthesis*, edited by: Liu, K. K., Atkinson, L., Quinones, R., and Talue-McManus, L., *Global Change – The IGBP Series*, Springer, New York, USA, 291–303, 2009.
- Macklin, S. A., Radchenko, V. I., Saitoh, S., and Stabeno, P. J.: Variability in the Bering Sea ecosystem, *Prog. Oceanogr.*, 55(1–2), 1–4, 2002.
- Makkaveev, P. N.: The dissolved inorganic carbon in the Kara Sea and in the mouths of the Ob and Yenisei Rivers, *Okeanologiya*, 34, 668–672, 1994.
- Manley, T. O. and Hunkins, K.: Mesoscale eddies of the Arctic Ocean. *J. Geophys. Res.*, 90(NC3), 4911–4930, 1985.
- Maslanik, J., Drobot, S., Fowler, C., Emery, W., and Barry, R.: On the Arctic climate paradox and the continuing role of atmospheric circulation in affecting sea ice conditions, *Geophys. Res. Lett.*, 34, L03711, doi:10.1029/2006GL028269, 2007.
- Maslowski, W., Marble, D. C., Walczowski, W., and Semtner, A. J.: On large-scale shifts in the Arctic Ocean and sea-ice conditions during 1979–98, *Ann. Glaciol.*, 33, 545–550, 2001.
- Mathis, J. T., Pickart, R. S., Hansell, D. A., Kadko, D., and Bates, N. R.: Eddy transport of organic carbon and nutrients from the Chukchi Shelf: Impact on the upper halocline of the western Arctic Ocean, *J. Geophys. Res.*, 112(C5), C05011, doi:10.1029/2006JC003899, 2007a.
- Mathis, J. T., Hansell, D. A., Kadko, D., Bates, N. R., and Cooper, L. W.: Determining net dissolved organic carbon production in the hydrographically complex western Arctic Ocean, *Limnol. Oceanogr.*, 52(5), 1789–1799, 2007b.
- Mathis, J. T., Hansell, D. A., and Bates, N. R.: Interannual variability of dissolved inorganic carbon distribution and net community production during the Western Arctic Shelf-Basin Interactions Project, *Deep-Sea Res. II*, 56(17), 1213–1222, doi:10.1016/j.dsr2.2008.10.017, 2009.
- McGuire, A. D., Chapin, F. S., Walsh, J. E., and Wirth, C.: Integrated regional changes in arctic climate feedbacks: implications for the global climate system. *Ann. Rev. Environ. Res.*, 31, 61–91, doi:10.1146/annurev.energy.31.020105.100253, 2006.
- Mehrbach, C., Culberson, C. H., Hawley, J. E., and Pytkowicz, R. M.: Measurement of the apparent dissociation constants of carbonic acid in seawater at atmospheric pressure, *Limnol. Oceanogr.*, 18, 897–907, 1973.
- Melling, H.: The formation of a haline shelf front in wintertime in an ice-covered arctic sea, *Cont. Shelf Res.*, 13, 1123–1147, 1993.
- Merico, A., Tyrrell, T., Lessard, E. J., Oguz, T., Stabeno, P. J., Zeeman, S. I., and Whitley, T. E.: Modelling phytoplankton succession on the Bering Sea shelf: role of climate influences and trophic interactions in generating *Emiliania huxleyi* blooms 1997–2000, *Deep-Sea Res. I*, 51(12), 1803–1826, doi:10.1016/j.dsr.2004.07.003, 2004.
- Millero, F. J.: Thermodynamics of the carbon dioxide system in the oceans, *Geochim. Cosmochim. Acta*, 59, 661–677, 1995.
- Mucci, A.: The solubility of calcite and aragonite in seawater at various salinities, temperatures, and one atmosphere total pressure. *Am. J. Sci.*, 283, 780–799, 1983.
- Mucci, A., Forest, A., Fortier, L., Fukuchi, M., Grant, J., Hattori, H., Hill, P., Lintern, G., Makbe, R., Magen, C., Miller, L., Sampei, M., Sasaki, H., Sundby, B., Walker, T., and Wassman, P.: Chapter 7: Organic and inorganic fluxes, in: *On Thin Ice: A Synthesis of the Canadian Arctic Shelf Exchange Study (CASES)*, edited by: Fortier, L., Barber, D., and Michaud, J., University of Manitoba Publication, 113–141, 2008.
- Muench, R. D., Gunn, J. T., Whitley, T. E., Schlosser, P., and Smethie, W.: An Arctic Ocean cold core eddy, *J. Geophys. Res.*, 105(C10), 23997–24006, 2000.
- Murata, A. and Takizawa, T.: Summertime  $CO_2$  sinks in shelf and slope waters of the western Arctic Ocean, *Cont. Shelf Res.*, 23(8), 753–776, doi:10.1016/S0278-4343(03)00046-3, 2003.
- Mysak, L. A., Manak, D. K., and Marsden, R. F.: Sea ice anomalies

- observed in the Greenland and Labrador Seas during 1901–1984 and their relation to an interdecadal Arctic climate cycle, *Clim. Dynam.*, 5(2), 111–133, 1990.
- Nagurnyi, A. P.: On the role of Arctic sea-ice in seasonal variability of carbon dioxide concentration in Northern Latitudes, *Russ. Meteorol. Hydrol.*, 33(1), 43–47, 2008.
- Nakaoka, S., Aiki, A., Nakazawa, T., Hashida, G., Morimoto, S., Yamanouchi, T., and Yoshikawa-Inoue, H.: Temporal and spatial variations of oceanic  $p\text{CO}_2$  and air-sea  $\text{CO}_2$  flux in the Greenland Sea and the Barents Sea, *Tellus*, 58(2), 148–161, 2006.
- Napp, J. M., Baier, C. T., Brodeur, R. D., Coyle, K. O., Shiga, N., and Mier, K.: Interannual and decadal variability in zooplankton communities of the southeast Bering Sea shelf, *Deep-Sea Res. II*, 49(26), 5991–6008, doi:10.1016/S0967-0645(02)00330-2, 2002.
- Nikolopoulos, A., Pickart, R. S., Fratantoni, P. S., Shimada, K., Torres, D. J., and Jones, E. P.: The western Arctic boundary current at 152° W: Structure, variability, and transport, *Deep-Sea Res. II*, 56(17), 1164–1181, doi:10.1016/j.dsr2.2008.10.014, 2009.
- Nitishinsky, M., Anderson, L. G., and Hölemann, J. A.: Inorganic carbon and nutrient fluxes on the Arctic Shelf, *Cont. Shelf Res.*, 27(10–11), 1584–1599, doi:10.1016/j.csr.2007.01.019, 2007.
- Olsen, A., Bellerby, R. G. J., Johannessen, T., Omar, A. M., and Skjelvan, I.: Interannual variability in the wintertime air-sea flux of carbon dioxide in the northern North Atlantic, 1981–2001, *Deep-Sea Res. I*, 50(10–11), 1323–1338, doi:10.1016/S0967-0637(03)00144-4, 2003.
- Olsson, K. and Anderson, L. G.: Input and biogeochemical transformation of dissolved carbon in the Siberian shelf seas, *Cont. Shelf Res.*, 17(7), 819–833, 1997.
- Omar, A. M., Johannessen, T., Kaltin, S., and Olsen, A.: Anthropogenic increase of oceanic  $p\text{CO}_2$  in the Barents Sea surface water, *J. Geophys. Res.*, 108(C12), 3388, doi:10.1029/2002JC001628, 2003.
- Omar, A., Johannessen, T., Bellerby, R. G. J., Olsen, A., Anderson, L. G., and Kivimäe, C.: Sea ice and brine formation in Storfjorden: Implications for the Arctic wintertime air-sea  $\text{CO}_2$  flux, in: *The Nordic Seas: An Integrated Perspective: Oceanography, climatology, biogeochemistry, and modeling*, edited by: Drange, H., Dokken, T., Furevik, T., Gerdes, R., and Berger, W., *Geoph. Monogr.*, 158, 177–187, 2005.
- Omar, A. M., Johannessen, T., Olsen, A., Kaltin, S., and Rey, F.: Seasonal and interannual variability of the air-sea  $\text{CO}_2$  flux in the Atlantic sector of the Barents Sea, *Mar. Chem.*, 104(3–4), 203–213, doi:10.1016/j.marchem.2006.11.002, 2007.
- Opsahl, S., Benner, R., and Amon, R. W.: Major flux of terrigenous dissolved organic matter through the Arctic Ocean, *Limnol. Oceanogr.*, 44(8), 2017–2023, 1999.
- Orr, J. C., Fabry, V. J., Aumont, O., Bopp, L., Doney, S. C., Feely, R. A., Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F., Key, R. M., Lindsay, K., Maier-Reimer, E., Matear, R., Monfray, P., Mouchet, A., Najjar, R. G., Plattner, G. K., Rodgers, K. B., Sabine, C. L., Sarmiento, J. L., Schlitzer, R., Slater, R. D., Totterdell, I. J., Weirig, M. F., Yamanaka, Y., and Yool, A.: Anthropogenic ocean acidification over the twenty-first century and its impacts on calcifying organisms, *Nature*, 437, 681–686, 2005.
- Orr, J., Anderson, L. G., Bates, N. R., Bopp, L., Fabry, V. J., Jones, E. P., and Swingedouw, D.: Arctic Ocean acidification, AGU/TOS/ASLO meeting, Hawaii, February 2006.
- Overland, J. E. and Wang, M.: The Arctic climate paradox: The recent decrease of the Arctic Oscillation, *Geophys. Res. Lett.*, 32(6), L06701, doi:10.1029/2004GL021752, 2005.
- Overland, J. E. and Wang, M.: Future regional Arctic sea ice declines, *Geophys. Res. Lett.*, 34(17), L17705, doi:10.1029/2007GL030808, 2007.
- Overland, J. E., Wang, M., and Salo, S.: The recent Arctic warm period, *Tellus A*, 60(4), 589–597, doi:10.1111/j.1600-0870.2008.00327.x, 2008.
- Pabi, S., van Dijken, G. L., and Arrigo, K. R.: Primary production in the Arctic Ocean, 1998–2006, *J. Geophys. Res.*, 113(C8), C08005, doi:10.1029/2007JC004578, 2008.
- Paquette, R. G. and Bourke, R. H.: Observations on the coastal current of arctic Alaska, *J. Mar. Res.*, 32, 195–207, 1974.
- Parkinson, C. L.: Variability of Arctic sea ice: The view from space, and 18-year record, *Arctic*, 53, 341–358, 2000.
- Perovich, D. K., Light, B., Eicken, H., Jones, K. F., Runciman, K., and Nghiem, S. V.: Increasing solar heating of the Arctic Ocean and adjacent seas, 1979–2005: Attribution and role in the ice-albedo feedback, *Geophys. Res. Lett.*, 34, L19505, doi:10.1029/2007GL031480, 2007.
- Peterson, B. J., Holmes, R. M., McClelland, J. W., Vorosmarty, C. J., Lammers, R. B., Shiklomanov, A. I., Shiklomanov, I. A., and Rahmstorf, S.: Increasing river discharge to the Arctic Ocean, *Science*, 298, 2171–2173, doi:10.1126/science.1077445, 2002.
- Pickart, R. S., Weingartner, T. J., Zimmermann, S., Torres, D. J., and Pratt, L. J.: Flow of winter-transformed Pacific water into the Western Arctic, *Deep-Sea Res. II*, 52(24–26), 3175–3198, doi:10.1016/j.dsr2.2005.10.009, 2005.
- Pickart, R. S., Fratantoni, P. S., Goldsmith, R. A., Moore, G. W. K., Agnew, T., and Vandeweghe, J.: Upwelling in the western Arctic boundary current north of Alaska, *EOS Trans. AGU*, 87(36), Ocean Sci. Meet. Suppl., Abstract OS350–07, 2006.
- Pickart, R. S., Moore, G. W. K., Torres, D. J., Fratantoni, P. S., Goldsmith, R. A., and Yang, J.: Upwelling on the continental slope of the Alaskan Beaufort Sea: Storms, ice, and oceanographic response, *J. Geophys. Res.*, 114, C00A13, doi:10.1029/2208JC005009, 2009.
- Piepenburg, D.: Recent research on Arctic benthos: common notions need to be revised, *Polar Biol.*, 28(10), 733–755, doi:10.1007/s00300-005-0013-5, 2005.
- Pipko I. I., Semiletov I. P., Tishchenko P. Y., Pugach S. P., and Christensen J. P.: Carbonate chemistry dynamics in Bering Strait and the Chukchi Sea, *Prog. Oceanogr.*, 55(1–2), 77–94, doi:10.1016/S0079-6611(02)00071-X, 2002.
- Polyakov, I. V., Alekseev, G. V., Timokhov, L. A., Bhatt, U. S., Colony, R. L., Simmons, H. L., Walsh, D., Walsh, J. E., and Zakharov, V. F.: Variability of the intermediate Atlantic water of the Arctic Ocean over the last 100 years, *J. Clim.*, 17(23), 4485–4497, doi:10.1175/JCLI-3224.1, 2004.
- Rigor, I. G. and Wallace, J. M.: Variations in the age of Arctic sea-ice extent, *Geophys. Res. Lett.*, 31(9), L09401, doi:10.1029/2004GL019492, 2004.
- Rigor, I. G., Wallace, J. M., and Colony, R. L.: Response of sea ice to the Arctic Oscillation, *J. Climate*, 15(18), 2648–2663, doi:10.1175/1520-0442(2002)015, 2002.
- Roach, A. T., Aagaard, K., Pease, C. H., Salo, S. A., Weingartner, T. J., Pavlov, V., and Kulakov, M.: Direct measurements of transport and water properties through Bering Strait, *J. Geophys. Res.*, 100(C9), 18443–18457, 1995.

- Rothrock, D. A. and Zhang, J.: Arctic Ocean sea ice volume: What explains its recent depletion?, *J. Geophys. Res.*, 110(C1), C01002, doi:10.1029/2004JC002282, 2005.
- Rothrock, D. A., Zhang, J., and Yu, Y.: The arctic ice thickness anomaly of the 1990s: A consistent view from observations and models, *J. Geophys. Res.*, 108(C3), 3083, doi:10.1029/2001JC001208, 2003.
- Roy, R. N., Roy, L. N., Vogel, R., Moore, C. P., Pearson, T., Good, C. E., Millero, F. J., and Campbell, D.: Determination of the ionization constants of carbonic acid in seawater, *Marine Chem.*, 44, 249–268, 1993.
- Royal Society: Ocean acidification due to increasing atmospheric carbon dioxide, The Clyvedon Press, Ltd, Cardiff, UK, 2005.
- Rudels, B., Anderson, L. G., and Jones, E. P.: Formation and evolution of the surface mixed layer and halocline of the Arctic Ocean, *J. Geophys. Res.-Ocean.*, 101(C4), 8807–8821, 1996.
- Rysgaard, S., Glud, R. N., Sejr, M. K., Bendtsen, J., and Christensen, P. B.: Inorganic carbon transport during sea ice growth and decay: A carbon pump in polar seas, *J. Geophys. Res.*, 112, C03016, doi:10.1029/2006JC003572, 2007.
- Sabine, C. L., Feely, R. A., Gruber, N., Key, R. M., Lee, K., Bullister, J. L., Wanninkhof, R., Wong, C. S., Wallace, D. W. R., Tilbrook, B., Millero, F. J., Peng, T.-H., Kozyr, A., Ono, T., and Rios, A. F.: The oceanic sink for anthropogenic CO<sub>2</sub>, *Science*, 305, 367–371, doi:10.1126/science.1097403, 2004.
- Sakshaug, E.: Primary and secondary production in the Arctic seas, in: *The Organic Carbon Cycle in the Arctic Ocean*, edited by: Stein, R. and Macdonald, R. W., Springer Publishing Company, Heidelberg, Germany, 57–81, 2004.
- Salisbury, J. E., Green, M., Hunt, C., and Campbell, J.: Coastal acidification by rivers: A threat to shellfish?, *EOS, Transactions*, 89(50), 513–528, 2008.
- Salmon, D. K. and McRoy, C. P.: Nutrient-based tracers in the Western Arctic: A new lower halocline water defined, in: *The Polar Oceans and Their Role in Shaping the Global Environment*, edited by: Johannessen, O. M., Muench, R. D., and Overland, J. E., American Geophysical Union, Geoph. Monogr., 85, 47–61, 1994.
- Sambrotto, R. N., Goering, J. J., and McRoy, C. P.: Large yearly production of phytoplankton in the western Bering Sea, *Science*, 225, 1147–1155, 1984.
- Santana-Casiano, J. M., Gonzalez-Davila, M., Rueda, M.-J., Llinas, O., and Gonzalez-Davila, E.-F.: The interannual variability of oceanic CO<sub>2</sub> parameters in the western Atlantic subtropical gyre at the ESTOC site, *Global Biogeochem. Cy.*, 21, GB1015, doi:10.1029/2006GB002788, 2007.
- Schlosser, P., Swift, J. H., Lewis, D., and Pfirman, S.: The role of the large-scale Arctic Ocean circulation in the transport of contaminant, *Deep-Sea Res. II*, 42, 1341–1368, 1995.
- Semiletov, I. P.: Aquatic sources of CO<sub>2</sub> and CH<sub>4</sub> in the Polar regions, *J. Atmos. Sci.*, 56(2), 286–306, 1999.
- Semiletov, I., Makshatas, A. I., Akasofu, S.-I., and Andreas, E. L.: Atmospheric CO<sub>2</sub> balance: The role of arctic sea ice, *Geophys. Res. Lett.*, 31(5), L05121, doi:10.1029/2003GL017996, 2004.
- Semiletov, I. P., Pipko, I. I., Repina, I., and Shakhova, N. E.: Carbonate chemistry dynamics and carbon dioxide fluxes across the atmosphere-ice-water interface in the Arctic Ocean, *J. Marine Syst.*, 66(1–4), 204–226, doi:10.1016/j.jmarsys.2006.05.012, 2007.
- Serreze, M. C. and Francis, J. A.: The Arctic amplification debate, *Clim. Change*, 76, 241–264, 2006.
- Serreze, M. C., Walsh, J. E., Chapin, F. S., Osterkamp, T., Dyurgerov, M., Romanovsky, V., Oechel, W. C., Morison, J., Zhang, T., and Barry, R. G.: Observational evidence of recent change in the northern high-latitude environment, *Clim. Change*, 46, 159–207, 2000.
- Serreze, M. C., Barrett, A. P., Slater, A. G., Woodgate, R. A., Aagaard, K., Lammers, R. B., Steele, M., Moritz, R., Meredith, M., and Lee, C. M.: The large-scale freshwater cycle of the Arctic, *Journal of Geophysical Research*, 111, C11010, doi:10.1029/2005JC003424, 2006.
- Serreze, M. C., Holland, M. M., and Stroeve, J.: Perspectives on the Arctic's shrinking sea-ice cover, *Science*, 315(5818), 1533–1536, doi:10.1126/science.1139426, 2007.
- Shimada, K., Kamoshida, T., Itoh, M., Nishino, S., Carmack, E., McLaughlin, F., Zimmermann, S., and Proshutinsky A.: Pacific Ocean inflow: influence on catastrophic reduction of sea ice cover in the Arctic Ocean, *Geophys. Res. Lett.*, 33(8), L08605, doi:10.1029/2005GL025624, 2006.
- Skjelvan, I., Olsen, A., Anderson, L. G., Bellerby, R. G. J., Falck, E., Kasajima, Y., Kivimäe, C., Omar, A., Rey, F., Olsson, K. A., Johannessen, T., and Heinze, C.: A review of the biogeochemistry of the Nordic Seas and Barents Sea, in: *Drange, edited by: Drange, H., Dokken, T., Furevik, T., Gerdes, R., and Berger, W., The Nordic Seas: An Integrated Perspective Oceanography, Climatology and Modelling*, American Geophysical Union, Washington DC, USA, 157–175, 2005.
- Skjelvan, I. M., Johannessen, T., and Miller, L.: Interannual variability of *f*CO<sub>2</sub> in the Greenland and Norwegian Seas, *Tellus B*, 51, 477–489, 1999b.
- Smith, D. M.: Recent increase in the length of the melt season of perennial Arctic sea ice, *Geophys. Res. Lett.*, 25(5), 655–658, 1998.
- Smith, S. V. and Hollibaugh, J. T.: Coastal metabolism and the oceanic organic carbon balance, *Rev. Geophys.*, 31, 75–89, 1985.
- Spall, M. A., Pickart, R. S., Fratantoni, P. S., and Plueddemann, A. J.: Western Arctic shelfbreak eddies: Formation and transport, *J. Phys. Oceanogr.*, 38, 1644–1668, doi:10.1175/2007JPO3829.1, 2008.
- Stabeno, P. J., Bond, N. A., Kachel, N. B., Salo, S. A., and Schumacher, J. D.: On the temporal variability of the physical environment over the south-eastern Bering Sea, *Fish. Oceanogr.*, 10(1), 81–98, 2001.
- Stabeno P. J., Kachel N. B., Sullivan, M., and Whitley, T. E.: Variability of physical and chemical characteristics along the 70 m isobath of the southeastern Bering Sea, *Deep-Sea Res. II*, 49(26), 5931–5943, doi:10.1016/S0967-0645(02)00327-2, 2002.
- Steele, M. and Boyd, T.: Retreat of the cold halocline layer in the Arctic Ocean, *J. Geophys. Res.*, 103(C5), 10419–10435, 1998.
- Steinacher, M., Joos, F., Frolicher, T. L., Plattner, G.-K., and Doney, S. C.: Imminent ocean acidification of the Arctic projected with the NCAR global coupled carbon-cycle climate model, *Biogeosciences*, 6(4), 515–533, 2009.
- Stroeve, J. C., Serreze, M. C., Fetterer, F., Arbetter, T., Meier, W., Maslanik, J., and Knowles, K.: Tracking the Arctic's shrinking ice cover: Another extreme September minimum in 2004, *Geophys. Res. Lett.*, 32(4), L04501, doi:10.1029/2004GL021810, 2005.

- Stroeve, J., Holland, M. M., Meier, W., Scambos, T., and Serreze, M.: Arctic sea ice decline: Faster than forecast, *Geophys. Res. Lett.*, 34(9), L09501, doi:10.1029/2007GL029703, 2007.
- Subba Rao, D. V. and Platt, T.: Primary production of Arctic waters. *Polar Biol.*, 3(4), 191–201, 1984.
- Takahashi, T., Olafsson, J., Goddard, J. G., Chipman, D. W., and Sutherland, S. G.: Seasonal variation of CO<sub>2</sub> and nutrients in the high-latitude surface oceans: a comparative study, *Global Biogeochem. Cy.*, 7(4), 843–878, 1993.
- Takahashi, T., Sutherland, S. C., Sweeney, C., Poisson, A., Metzl, N., Tilbrook, B., Bates, N., Wanninkhof, R., Feely, R. A., Sabine, C., Olafsson, J., and Nojiri, Y.: Global sea-air CO<sub>2</sub> flux based on climatological surface ocean pCO<sub>2</sub>, and seasonal biological and temperature effects, *Deep-Sea Res. II*, 49(9–10), 1601–1622, doi:10.1016/S0967-0645(02)00003-6, 2002.
- Takahashi, T., Sutherland, S. C., Wanninkhof, R., Sweeney, C., Feely, R. A., Chipman, D. W., Hales, B., Friederich, G., Chavez, F., Sabine, C., Watson, A., Bakker, D. C. E., Schuster, U., Metzl, N., Yoshikawa-Inoue, H., Ishii, M., Midorikawa, Nojiri, Y., Kortzinger, A., Steinhoff, T., Hoppema, M., Olafsson, J., Arnason, T. S., Tilbrook, B., Johannessen, T., Olsen, A., Bellerby, R., Wong, C. S., Delille, B., Bates, N. R., and de Baar, H. J. W.: Climatological mean and decadal change in surface ocean pCO<sub>2</sub>, and net sea-air CO<sub>2</sub> flux over the global oceans, *Deep-Sea Res. II*, 56(8–10), 554–577, doi:10.1016/j.dsr2.2008.12.009, 2009.
- Tanhua, T., Jones, E. P., Jeansson, E., Jutterstrom, S., Smethie, W. M., Wallace, D. W. R., and Anderson, L. G.: Ventilation of the Arctic Ocean: Mean ages and inventories of anthropogenic CO<sub>2</sub> and CFC-11, *J. Geophys. Res. Oceans*, 114(C1), C01002, doi:10.1029/2008JC004868, 2009.
- Thomas, H., Schiettecatte, L.-S., Suykens, K., Kone, Y. J. M., Shadwick, E. H., Prowe, A. E. F., Bozec, Y., de Baar, H. J. W., and Borges, A. V.: Enhanced ocean carbon storage from anaerobic alkalinity generation in coastal sediments, *Biogeosciences*, 6, 267–274, 2009, <http://www.biogeosciences.net/6/267/2009/>.
- Tremblay, J.-É., Simpson, K., Martin, J., Miller, L., Gratton, Y., Barber, D., and Price, N. M.: Vertical stability and the annual dynamics of nutrients and chlorophyll fluorescence in the coastal, southeast Beaufort Sea, *J. Geophys. Res.*, 113, C07S90, doi:10.1029/2007JC004547, 2008.
- Tucker, W. and Cate, D.: The 1994 Arctic ocean Section: The first major scientific crossing of the Arctic Ocean, US Army Cold Regions Research and Engineering Laboratory, Special Report, 117, 96–23, 1996.
- Wadley, M. R. and Bigg, G. R.: Impact of flow through the Canadian Archipelago and Bering Strait on the North Atlantic and Arctic circulation: An ocean modelling study, *Q. J. Roy. Meteor. Soc.*, 128, 2187–2203, 2002.
- Walsh, J. E. and Chapman, W. L.: Arctic contribution to upper-ocean variability in the North Atlantic, *J. Climate*, 3(12), 1462–1473, 1990.
- Walsh, J. J.: Importance of continental margins in the marine biogeochemical cycling of carbon and nitrogen, *Nature*, 350, 53–55, 1991.
- Walsh, J. J., Premuzic, E. T., Gaffney, J. S., Rowe, G. T., Harbottle, G., Stoenner, R. W., Balsam, W. L., Betzer, P. R., and Macko, S. A.: Organic storage of CO<sub>2</sub> on the continental slope off the Mid-Atlantic bight, the southeastern Bering Sea, and the Peru coast, *Deep-Sea Res.*, 32, 853–883, 1985.
- Wallace, D. W. R., Moore, R. M., and Jones, E. P.: Ventilation of the Arctic Ocean cold halocline: rates of diapycnal and isopycnal transport, oxygen utilization and primary production inferred using chlorofluoromethane distributions, *Deep-Sea Res.*, 34, 1957–1979, 1987.
- Wang, J., Ikeda, M., Zhang, S., and Gerdes, R.: Linking the northern hemisphere sea-ice reduction trend and the quasi-decadal arctic sea-ice oscillation, *Clim. Dynam.*, 24(2–3), 115–130, 2005.
- Wang, W. Q., Yang, X. T., Huang, H. B., and Chen, L. Q.: Investigation on distribution and fluxes of sea–air CO<sub>2</sub> of the expedition areas in the Arctic Ocean, *Science in China Series D – Earth Sciences*, 46, 569–580, 2003.
- Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean, *J. Geophys. Res. Oceans*, 97, 7373–7382, 1992.
- Wanninkhof, R. and McGillis, W. R.: A cubic relationship between air-sea CO<sub>2</sub> exchange and windspeed, *Geophys. Res. Lett.*, 26(13), 1889–1892, 1999.
- Weingartner, T. J., Cavalieri, D. J., Aagaard, K., and Sasaki, Y.: Circulation, dense water formation, and outflow on the northeast Chukchi shelf, *J. Geophys. Res.*, 103(C4), 7647–7661, 1998.
- Weingartner, T. J., Danielson, S., Sasaki, Y., Pavlov, V., and Kulakov, M.: The Siberian Coastal Current: A wind- and buoyancy-forced Arctic coastal current, *J. Geophys. Res.*, 104(C12), 29697–29713, 1999.
- Weingartner, T., Aagaard, K., Woodgate, R., Danielson, S., Sasaki, Y., and Cavalieri, D.: Circulation on the north central Chukchi Sea shelf, *Deep-Sea Res. II*, 52(24–26), 3150–3174, doi:10.1016/j.dsr2.2005.10.015, 2005.
- Wheeler, P. A., Gosselin, M., Sherr, E., Thibault, D., Kirchman, D. L., Benner, R., and Whitledge, T. E.: Active cycling of organic carbon in the central Arctic Ocean, *Nature*, 380(6576), 697–699, 1996.
- Wijffels, S. E., Schmitt, R. W., Bryden, H. L., and Stigebrandt, A.: Transport of freshwater by the oceans, *J. Geophys. Res.*, 22, 155–162, 1992.
- Wilson, C. and Wallace, D. W. R.: Using the nutrient ratio NO/PO as a tracer of continental shelf waters in the central Arctic Ocean, *J. Geophys. Res.*, 95(C12), 22193–22208, 1990.
- Winn, C. D., Mackenzie, F. T., Carillo, C. J., Sabine, C. L., and Karl, D. M.: Air-sea carbon dioxide exchange in the North Pacific Subtropical Gyre: Implications for the global carbon budget, *Global Biogeochem. Cy.*, 8(2), 157–163, 1994.
- Winsor, P. and Bjork, G.: Polynya activity in the Arctic Ocean from 1958 to 1997, *J. Geophys. Res.*, 105(C4), 8789–8803, 2000.
- Winsor, P. and Chapman, D. C.: Pathways of Pacific water across the Chukchi Sea: A numerical model study, *J. Geophys. Res.*, 109, C03002, doi:10.1029/2003JC001962, 2004.
- Winton, M.: Does the Arctic sea-ice have a tipping point?, *Geophys. Res. Lett.*, 33(23), L23504, doi:10.1029/2006GL028017, 2006.
- Wollast, R.: Evaluation and comparison of the global carbon cycle in the coastal zone and in the open ocean, in: *The Sea 10, The Global Coastal Ocean*, edited by: Brinkm K. H. and Robinson, A. R., Wiley, New York, USA, 213–252, 1998.
- Woodgate, R. A. and Aagaard, K.: Revising the Bering Strait freshwater flux into the Arctic Ocean, *Geophys. Res. Lett.*, 32(2), L02602, doi:10.1029/2004GL021747, 2005.
- Woodgate, R. A., Aagaard, K., and Weingartner, T. J.: A year in the

- physical oceanography of the Chukchi Sea: Moored measurements from autumn 1990–1991, *Deep-Sea Res. II*, 52(24–26), 3116–3149, doi:10.1016/j.dsr2.2005.10.016, 2005.
- Woodgate, R. A., Aagaard, K., and Weingartner, T. J.: Monthly temperature, salinity and transport variability of the Bering Strait through flow, *Geophys. Res. Lett.*, 32(4), L04601, doi:10.1029/2004GL021880, 2005.
- Wu, B. Y., Wang, J., and Walsh, J. E.: Dipole anomaly in the winter Arctic atmosphere and its association with sea-ice motion. *Journal of Climate*, 19(2), 210–225, doi:10.1175/JCLI3619.1, 2006.
- Yager, P. L., Wallace, D. W. R., Johnson, K. M., Smith, W. O., Minnett, P. J., and Deming, J. W.: The Northeast Water Polynya as an atmospheric CO<sub>2</sub> sink: A seasonal rectification hypothesis, *J. Geophys. Res.*, 100, 4389–4398, 1995.
- Yamamoto-Kawai, M., Tanaka, N., and Pivovarov, S.: Freshwater and brine behaviors in the Arctic Ocean deduced from historical data of delta O-18 and alkalinity (1929–2002 AD), *J. Geophys. Res.*, 110(C10), C10003, doi:10.1029/2004JC002793, 2005.
- Yamamoto-Kawai, M., McLaughlin, F. A., Carmack, E. C., Nishino, S., and Shimada, K.: Aragonite undersaturation in the Arctic Ocean: effects of ocean acidification and sea ice melt, *Science*, accepted, 2009.
- Zeebe, R., and Wolf-Gladrow, D., 2001. CO<sub>2</sub> in Seawater: Equilibrium, Kinetics, Isotopes. Elsevier Oceanography Series, 65.