



This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Exploring local adaptation and the ocean acidification seascape – studies in the California Current Large Marine Ecosystem

G. E. Hofmann^{1,3}, T. G. Evans², M. W. Kelly¹, J. L. Padilla-Gamiño¹,
C. A. Blanchette³, L. Washburn^{3,4}, F. Chan⁵, M. A. McManus⁶, B. A. Menge⁵,
B. Gaylord^{7,8}, T. M. Hill^{7,9}, E. Sanford^{7,8}, M. LaVigne^{7,9,10}, J. M. Rose⁵,
L. Kapsenberg¹, and J. M. Dutton¹

¹Department of Ecology, Evolution and Marine Biology, University of California Santa Barbara, Santa Barbara, CA 93106-9620, USA

²Department of Biological Sciences, California State University East Bay, Hayward, CA 94542, USA

³Marine Science Institute, University of California Santa Barbara, Santa Barbara, CA 93106-6150, USA

⁴Department of Geography, University of California Santa Barbara, Santa Barbara, CA 93106-4060, USA

⁵Department of Zoology, Oregon State University, Corvallis, OR 97331-2914, USA

⁶Department of Oceanography, University of Hawaii at Manoa, Honolulu, Hawaii 96822, USA

⁷Bodega Marine Laboratory, University of California Davis, Bodega Bay, CA 94923, USA

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
⏪	⏩
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



⁸Department of Evolution and Ecology, University of California Davis, Davis, CA 95616, USA

⁹Department of Geology, University of California Davis, Davis, CA 95616, USA

¹⁰Department of Earth and Oceanographic Science, Bowdoin College, Brunswick ME, 04011, USA

Received: 31 March 2013 – Accepted: 19 June 2013 – Published: 16 July 2013

Correspondence to: G. E. Hofmann (hofmann@lifesci.ucsb.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

BGD

10, 11825–11856, 2013

Exploring local adaptation and the ocean acidification seascape

G. E. Hofmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

The California Current Large Marine Ecosystem (CCLME), a temperate marine region dominated by episodic upwelling, is predicted to experience rapid environmental change in the future due to ocean acidification. Aragonite saturation state within the California Current System is predicted to decrease in the future, with near-permanent undersaturation conditions expected by the year 2050. Thus, the CCLME is a critical region to study due to the rapid rate of environmental change that resident organisms will experience and because of the economic and societal value of this coastal region. Recent efforts by a research consortium – the Ocean Margin Ecosystems Group for Acidification Studies (OMEGAS) – has begun to characterize a portion of the CCLME; both describing the mosaic of pH in coastal waters and examining the responses of key calcification-dependent benthic marine organisms to natural variation in pH and to changes in carbonate chemistry that are expected in the coming decades. In this review, we present the OMEGAS strategy of co-locating sensors and oceanographic observations with biological studies on benthic marine invertebrates, specifically measurements of functional traits such as calcification-related processes and genetic variation in populations that are locally adapted to conditions in a particular region of the coast. Highlighted in this contribution are (1) the OMEGAS sensor network that spans the west coast of the US from central Oregon to southern California, (2) initial findings of the carbonate chemistry amongst the OMEGAS study sites, (3) an overview of the biological data that describes the acclimatization and the adaptation capacity of key benthic marine invertebrates within the CCLME.

1 Introduction

A leading imperative in global change biology is forecasting the impact of environmental change on key species and ecosystems (Buckley and Kingsolver, 2012; Hoffmann and Sgrò, 2011; Williams et al., 2008; Dawson et al., 2011), and on critical natural

BGD

10, 11825–11856, 2013

Exploring local adaptation and the ocean acidification seascape

G. E. Hofmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Exploring local adaptation and the ocean acidification seascape

G. E. Hofmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

resources (Gillson et al., 2013; Ibáñez et al., 2013; Pettoelli, 2012). As exemplified by the study of ocean acidification (OA) (Boyd, 2011; Hofmann et al., 2011; Kroeker et al., 2013), the issues in this scientific endeavor are complex. In addition to the natural complexity in ecosystems emanating from biotic interactions and variability in biological responses to the environment, physical environmental parameters can also vary dramatically, as is evident when comparing disparate ecosystems such as tropical reefs and polar seas. From a biological perspective, one way to better predict the consequences of ocean change is to examine ways in which populations could potentially respond to environmental change. One mechanism is inherent physiological plasticity that confers tolerance to a changing environment (Chown, 2012; Chown and Gaston, 2008; Helmuth, 2009; Huey et al., 2012); a second is outright evolutionary adaptation to rapidly changing conditions (Hoffmann and Sgrò, 2011; Kelly and Hofmann, 2012; Visser, 2008). Of the two responses, evolutionary adaptation may be the most difficult to study. The absence of extensive long-term datasets and complexity of biological processes combined with vast spatial expanses to cover have made studying evolution in the ocean a significant challenge.

In the oceans, major changes in temperature and carbonate chemistry are predicted in the coming decades (Feely et al., 2009; Gruber et al., 2012; Hauri et al., 2013; Orr et al., 2005) and in some ecosystems the rate of change will be rapid. For example, Australia is experiencing warming and extreme heat events (Wernberg et al., 2013) and the Western Antarctic Peninsula has warmed significantly in the last two decades with significant decreases in sea ice that have already altered this polar marine ecosystem (Steinberg et al., 2012). In the California Current System, the focus of this article, aragonite saturation state in the upwelling zone is expected to decrease rapidly in the future (Gruber et al., 2012; Hauri et al., 2013). Here, within the next 30 yr, summer-long undersaturation conditions are expected in nearshore regions in the top 60 m. Thus, the California Current Large Marine Ecosystem (CCLME) is a critical region to study, both in terms of the pace at which associated biota might experience rapid environmental change and because of the economic and societal value of this coastal

region (Costanza et al., 1997). From a biological perspective, we have limited information on how critical species might respond to this predicted shift in ocean chemistry. Major questions remain regarding whether marine species currently possess functional traits that would allow the tolerance of a changing environment, or whether they will be able to adapt to rapidly changing ocean conditions into the future.

Recently, a collaborative group of investigators, who have been studying biological, ecological and oceanographic conditions along in the US West coast for decades, has begun to tackle these questions in the CCLME using a strategy that involves co-locating oceanographic sensors with biological observations. This group, called OMEGAS (Ocean Margin Ecosystems Group for Acidification Studies), has worked as a collaborative body since 2010 studying an area spanning from central Oregon to southern California (Fig. 1). Research activities include gathering oceanographic observations (e.g., pH, temperature, salinity, $p\text{CO}_2$, alkalinity, current magnitude and direction) that are matched with biological data (physiological measurements, organismal-level responses to low pH conditions such as growth rates and calcification, and genetic data) for key species of calcifying benthic marine invertebrates. The central goal of the OMEGAS project is to consider physiological and ecological performance of organisms, and genetic variation within and between populations, within the broad mosaic of pH and saturation states found across the CCLME. Changes in pH and saturation state are driven by absorption of anthropogenic CO_2 into surface waters, and the commensurate change in ocean chemistry, both the reduction in pH and the decline in the concentration of carbonate ions, are physiologically challenging to calcifying marine organisms (Doney et al., 2009). Further, because pH and saturation state regimes vary widely across the CCLME, our research allows a “substitution of space for time” (Pickett, 1989) approach in an exploration of local adaptation of benthic marine invertebrates to carbonate chemistry (Sanford and Kelly, 2011). The combination of simultaneous oceanographic and biological research across a large latitudinal range will help identify areas on the coast that express a range of acidification conditions. This information may indicate areas that might be refuges from acidification in the future, and

Exploring local adaptation and the ocean acidification seascape

G. E. Hofmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Exploring local adaptation and the ocean acidification seascape

G. E. Hofmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

could reveal regions that are adaptation “hot spots;” i.e., places where selection for undersaturation-tolerant genotypes has been underway for long periods of time (Jacobs et al., 2004). Finally, discovery of tolerant populations within the environmental mosaic of the CCLME can provide an opportunity to identify mechanisms that underlie tolerance (Evans et al., 2013a; Evans and Hofmann, 2012). We also hope to address whether some species possess sufficient physiological and genetic variation to adapt to future change. Species that lack physiological plasticity may quickly become excluded from an environment when change occurs. Alternatively, organisms with the capacity to adapt should do so, and exploring the degree of standing genetic variation for pH tolerance may help us to forecast responses to environmental change (De Wit and Palumbi, 2012; Kelly et al., 2013; Pespeni et al., 2013a).

From a broad perspective, the OMEGAS strategy of co-locating sensors with biology allows one to explore the balance of physiological plasticity vs. capacity for adaptation in the light of present-day environmental conditions (Kelly and Hofmann, 2012). Such an assessment is critically important in allowing prediction of outcomes that rely on both species- and community-level responses (Kroeker et al., 2013; Wootton et al., 2008; Hall-Spencer et al., 2008). The goal of this article is to overview recent efforts of the OMEGAS consortium (<http://omegas.science.oregonstate.edu>) in the study of physiological plasticity and the capacity for adaptation in populations of benthic marine invertebrates across the CCLME. Here, we outline three lines of investigation that are integrated across the group and that frame our studies: (1) an overview of a spatially distributed sensor network for quantifying pH dynamics, (2) data on functional traits that may vary in populations and support physiological plasticity, and (3) data on genetic structure of the study populations that link to local adaptation in natural populations and the capacity to adapt to future environmental change in the CCLME.

2 The OMEGAS sensor network: identifying a pH mosaic in the CCLME

A central goal of researchers within the OMEGAS consortium is to link biological performance with environmental variability in ocean carbonate chemistry along what we hypothesized would be a mosaic or gradient of conditions that might foster local adaptation. Long-term observations have been invaluable in defining the rate of ocean acidification (OA) progression in low-latitude, open ocean biomes and records from sub-tropical gyre time-series stations (e.g., Bermuda Atlantic Time Series “BATS”, <http://bats.bios.edu/>; Hawaiian Ocean Times Series “HOTS”, http://hahana.soest.hawaii.edu/hot/hot_jgofs.html; European Station for Time Series in the Ocean “ESTOC” <http://www.eurosites.info/estoc.php>) show a decline of ocean pH -0.02 to 0.04 pH units over a 20 yr period against low-frequency seasonal oscillations of similar magnitude (Bates et al., 2012; Dore et al., 2009). For coastal regions, the scientific community is just now assessing the longer-term variability in pH. Recent analyses of long-term data sets indicate that pH is changing rapidly in coastal Washington (Wootton and Pfister, 2012), in coastal upwelling zones along the US Pacific coast (Harris et al., 2013; Chan et al., 2013), at a coastal region in the Netherlands (Provoost et al., 2010), and in the Monterey Bay area where low pH water is associated with low oxygen water masses that reach the shallow, nearshore regions (Booth et al., 2012). Cruise data have provided snapshots of carbonate chemistry along the coast of the CCLME (Feely et al., 2008), and suggested that at some locations in northern California, undersaturated waters shoaled in the inner shelf. Prior to OMEGAS, however, no coordinated inner-shelf time series were available that would allow evaluation of the frequency, intensity and spatial expanse with which coastal ecosystems experience rapid acidification.

The recent development of autonomously recording pH sensors (Martz et al., 2010) has helped to bridge this data gap. Easily deployed on either moorings or benthic (e.g., rocky intertidal) locations, these sensors facilitate the collection of environmental pH data in a variety of habitats and support the collection of long-term datasets that

BGD

10, 11825–11856, 2013

Exploring local adaptation and the ocean acidification seascape

G. E. Hofmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Exploring local adaptation and the ocean acidification seascape

G. E. Hofmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

chemistry. Although it is difficult to pin any one phenotypic element on a single selection factor in the environment, (e.g., temperature or pH alone), increasingly the importance of the phenotype and tolerance traits has been increasingly recognized as a key mechanism by which a species might respond to environmental change in a number of systems (Buckley and Kingsolver, 2012; Chown, 2012; Chown and Gaston, 2008; Hel-
muth, 2009). Recently within the OMEGAS research community, functional traits and physiological tolerances have been measured in an environmentally relevant context (Evans et al., 2013a, b; Kelly et al., 2013; Padilla-Gamiño et al., 2013; Pespeni et al., 2013a; Yu et al., 2011). The OMEGAS group hypothesized that such phenotypic traits would show variation across space if benthic invertebrates are acclimatized to the local conditions (Chown, 2012) such that they might also contribute to how species respond to environmental change.

Studies initiated by the OMEGAS research teams first focused on populations of ecologically important calcification-dependent benthic marine invertebrates, the purple sea urchin, *Strongylocentrotus purpuratus*, and the mussel, *Mytilus californianus*, across the OMEGAS study sites. Experiments conducted by the group included field experiments at different OMEGAS sites and laboratory mesocosms that tested the response of organisms from various populations to variation in pH. The overall results of the laboratory component demonstrated that the two study organisms displayed different degrees of sensitivity to shifts in ocean chemistry with sea urchins appearing to be more resilient to variation in partial pressure of carbon dioxide ($p\text{CO}_2$) than mussels. Specifically, larval culturing of purple sea urchins in CO_2 mesocosms showed sea urchin larvae were tolerant of $p\text{CO}_2$ levels that are representative of present-day $p\text{CO}_2$ levels documented by OMEGAS field sensors. That is, growth, morphology, early development to echinopluteus and development to metamorphosis were rarely affected (Chown, 2012; Kelly et al., 2013; Padilla-Gamiño et al., 2013; Pespeni et al., 2013a), although minor decreases in the size of the larval skeleton were seen at elevated $p\text{CO}_2$ s near $1000 \mu\text{atm } p\text{CO}_2$ (Padilla-Gamiño et al., 2013). The mechanistic underpinning of this tolerance and physiological plasticity is currently being examined in greater detail.

Exploring local adaptation and the ocean acidification seascape

G. E. Hofmann et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

However, in the analyses of urchin larvae from the CO₂ mesocosm experiments, the transcriptome displayed variable gene expression during development that likely compensates for carbonate under-saturation and challenges to calcification and homeostasis of intracellular pH (Evans et al., 2013a). In contrast, laboratory studies on mussel larvae (*M. californianus*) showed that low pH conditions reduced growth and shell strength (Gaylord et al., 2011). In light of the pH mosaic data from the sensors, these results suggest that mussel recruits will vary across the CCLME in terms of their susceptibility to drilling and crushing predators, and that susceptibility will increase through time.

When functional traits were compared from animals in populations across the OMEGAS study sites, we found a more complicated story. However, for the most part, results thus far highlight a resilience of function in the marine invertebrate study organisms. First, in line with the robustness of the early life embryonic stages of sea urchins in mesocosm studies, analysis of fertilization kinetics using adult purple sea urchins collected from OMEGAS sites showed that fertilization success was not affected by elevated $p\text{CO}_2$ (Fig. 3), with populations from both Fogarty Creek (FC) and further south at Bodega Marine Reserve (BMR) showing the same insensitivity to elevated $p\text{CO}_2$. Similarly, larval culturing of sea urchins demonstrated that larvae from northern and southern populations have similar and more resilient metabolic responses to $p\text{CO}_2$ (Kelly et al., 2013). Specifically, oxygen consumption rates in early pluteus did not change between larvae reared under different $p\text{CO}_2$ conditions (Fig. 4a), and this response was consistent between populations exposed to different upwelling regimes (Kelly et al., 2013). Examining a feature of juvenile and adult sea urchins – the mineralogy of the calcium carbonate skeleton – LaVigne et al. (2013) found that for the most part the composition of the skeleton was not different in urchins collected across the latitudinal gradient of the OMEGAS study sites. Adult spine composition (Mg/Ca and Sr/Ca ratios) was not different between populations from Fogarty Creek, Oregon in the north to near Lompoc Landing in southern California (LaVigne et al., 2013) (Fig. 4b).

pH) implies the existence of variation that will allow for adaptation to changes in that gradient in the future. In addition, the capacity to adapt to an environmental variable in *space*, suggests the capacity to adapt to changes in that variable through *time*. And conversely, limits to adaptation in space (for example failure to adapt to maximum temperatures above 20 °C at a southern range limit) suggest possible limits to adaptation in the face of future changes.

There are several approaches to describing local adaptation. The gold standard for demonstrating local adaptation is a set of reciprocal transplants among sites, followed by a comparison of some aspect of performance for “local” vs. “foreign” genotypes in each environment. However, when local genotypes can be shown to have the highest performance in a particular environment it will not always be clear which aspect of the environment they are responding to, as aspects of the environment (e.g., temperature, pH, biotic interactions) often co-vary. To test for adaptation to a specific environmental variable of interest, it will often be necessary to bring organisms from different populations into the laboratory, and examine variation in performance along a particular axis of environmental variation.

This type of laboratory experiment was done by OMEGAS investigators with sea urchins. Kelly et al. (2013) performed quantitative genetic crosses among purple sea urchins (*Strongylocentrotus purpuratus*) from two different sites with different upwelling regimes, and then reared the offspring of these crosses under high and low $p\text{CO}_2$. They found abundant genetic variation for the response to high $p\text{CO}_2$, but only minor differences between the offspring of males from the two sites. However these small differences were consistent with local adaptation to pH, with offspring from the site with more exposure to extreme pH levels showing a lesser sensitivity to low pH (Fig. 5). The lack of large differences among populations is also consistent with the biology of purple sea urchins, which have long-lived planktonic larvae, and therefore relatively homogenized populations (Edmands et al., 1996).

Another way to demonstrate genetic differences among populations is to identify differences in actual gene sequences. Pespeni et al. (2012) performed a genome-wide

Exploring local adaptation and the ocean acidification seascape

G. E. Hofmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



scan of polymorphisms for two distant populations of the purple urchin *S. purpuratus*. They found many polymorphisms whose frequencies differed strongly among populations, indicating that they were targets of selective forces that differed among these populations. Similarly, De Wit and Palumbi (2013) used transcriptome sequencing of three populations of the red abalone (*Haliotis rufescens*) to identify single nucleotide polymorphisms (SNPS) in genes that appeared to be under spatially varying selection. A limitation of the approach used in both of these studies is that variation in gene sequences often cannot be directly tied to variation in organismal performance. Furthermore, because of co-variation in environmental variables it is often difficult to know which aspect of the environment is selecting for changes in allele frequencies among populations. However studies like these provide a rich foundation for ongoing work attempting to identify the genetic basis of local adaptation. Notably, in a mesocosm experiment where urchin larvae from OMEGAS study sites were cultured under ecologically relevant $p\text{CO}_2$ conditions for the CCLME, Pespeni et al. (2013a) found significant allelic frequencies in 40 classes of functional proteins in gene classes for ion homeostasis and biomineralization, suggesting that standing genetic variation does exist within these study populations at the OMEGAS sites.

We are currently testing for local adaptation to carbonate chemistry in another species from the CCLME with a very different life history and dispersal mode from purple urchins. Coralline algae are an ecologically important group of marine calcifiers that live in almost every habitat in the world's oceans, from the tropics to the polar regions (Johansen, 1981). Multiple species can be found in abundance in the CCLME (Abbott and Hollenberg, 1976) providing substrata and settlement cues to invertebrate larvae and serving as habitat for many marine species (Johansen, 1981). These calcifying algae are highly vulnerable to ocean acidification (Koch et al., 2013) and to date, it is unclear how more acidified waters will affect algal physiological mechanisms dependent on HCO_3^- and CO_3^{2-} -availability such as photosynthesis and calcification. Coralline algae reproduce by releasing spores (Johansen, 1981) that can fully attach to the bottom within hours of release (Miklasz, 2012) and recruit near the parental alga.

Exploring local adaptation and the ocean acidification seascape

G. E. Hofmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



This feature of coralline algae life history could limit dispersal distance and increase the potential for local adaptation in this group. We are currently examining how ocean acidification can affect the growth of spores from the articulated coralline algae *Corallina vancouveriensis*, which is an abundant species in the intertidal zone within the CCLME (Abbott and Hollenberg, 1976). Using populations exposed to different oceanographic conditions we are examining whether there is genetic variation among individuals and how this genetic variation is distributed among populations. Differences in the potential for adaptation could have important implications for the distribution of *C. vancouveriensis* and the organisms in the CCLME that depend on them.

5 Summary and future directions

The OMEGAS consortium has used a strategy of co-locating sensors and biological observations to begin to describe patterns of acclimatization and local adaptation in the CCLME. Ultimately, the goal of this research consortium is use this information to forecast the impacts of future acidification in coastal regions around the globe. Co-locating sensors with measures of physiological performance makes sense, but should be more deeply appreciated in a biological context. Since it is the physical environment to make a large contribution to maintaining genetic variation in populations, we would predict that regions characterized by large variation in pH and changes in saturation state may have organisms with different physiological tolerances, and with genetic differences in OA tolerance (Kelly and Hofmann, 2012; Kelly et al., 2013; Pespeni et al., 2013a). Thus, from a global change biology perspective, this strategy also allows us to explore the extent to which physiological plasticity vs. the capacity for adaptation might play a role in species' responses to environmental change.

The OMEGAS sensor network has identified a mosaic of complex oceanographic conditions and we anticipate that the complexity of pH dynamics will be matched by the dynamics of other environmental factors such as oxygen levels in coastal oceans. Thus, future experimentation in the laboratory will increasingly begin to combine these factors

Exploring local adaptation and the ocean acidification seascape

G. E. Hofmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Exploring local adaptation and the ocean acidification seascape

G. E. Hofmann et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

ary at point conception, California, USA, J. Exp. Mar. Biol. Ecol., 340, 126–148, doi:10.1016/j.jembe.2006.09.022, 2007.

Bockmon, E. E., Frieder, C. A., Navarro, M. O., White-Kershek, L. A., and Dickson, A. G.: Controlled experimental aquarium system for multi-stressor investigation: carbonate chemistry, oxygen saturation, and temperature, Biogeosciences Discuss., 10, 3431–3453, doi:10.5194/bgd-10-3431-2013, 2013.

Booth, J. A. T., McPhee-Shaw, E. E., Chua, P., Kingsley, E., Denny, M., Phillips, R., Bograd, S. J., Zeidberg, L. D., and Gilly, W. F.: Natural intrusions of hypoxic, low pH water into nearshore marine environments on the California coast, Cont. Shelf. Res., 45, 108–115, doi:10.1016/j.csr.2012.06.009, 2012.

Boyd, P. W.: Beyond ocean acidification, Nat. Geosci., 4, 273–274, 2011.

Buckley, L. B. and Kingsolver, J. G.: Functional and phylogenetic approaches to forecasting species' responses to climate change, Annu. Rev. Ecol. Evol. S., 43, 205–226, doi:10.1146/annurev-ecolsys-110411-160516, 2012.

Chan, F., Barth, J., Blanchette, C., Chavez, F., Cheriton, O., Friederich, G., Gaylord, B., Gouhier, T., Hill, T. M., Hofmann, G. E., McManus, M., Menge, B. A., Russell, A. D., Sanford, E., and Washburn, L.: Widespread detection of coastal ocean acidification across the California Current System, in preparation, 2013.

Checkley Jr., D. M. and Barth, J. A.: Patterns and processes in the California Current System, Prog. Oceanogr., 83, 49–64, doi:10.1016/j.pocean.2009.07.028, 2009.

Chown, S. L.: Trait-based approaches to conservation physiology: forecasting environmental change risks from the bottom up, Philos. T. R. Soc. B, 367, 1615–1627, doi:10.1098/rstb.2011.0422, 2012.

Chown, S. L. and Gaston, K. J.: Macrophysiology for a changing world, P. Roy. Soc. B-Biol. Sci., 275, 1469–1478, 2008.

Costanza, R., d'Arge, R., de Groot, R., Farber, S., Grasso, M., Hannon, B., Limburg, K., Naeem, S., O'Neil, R. V., Paruelo, J., Raskin, R. G., Sutton, P., and van den Belt, M.: The value of the world's ecosystem services and natural capital, Nature, 387, 253–260, 1997.

Crim, R. N., Sunday, J. M., and Harley, C. D. G.: Elevated seawater CO₂ concentrations impair larval development and reduce larval survival in endangered northern abalone (*Haliotis kamtschatkana*), J. Exp. Mar. Biol. Ecol., 400, 272–277, doi:10.1016/j.jembe.2011.02.002, 2011.

Exploring local adaptation and the ocean acidification seascape

G. E. Hofmann et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Dawson, T. P., Jackson, S. T., House, J. I., Prentice, I. C., and Mace, G. M.: Beyond predictions: biodiversity conservation in a changing climate, *Science*, 332, 53–58, doi:10.1126/science.1200303, 2011.

De Wit, P. and Palumbi, S. R.: Transcriptome-wide polymorphisms of red abalone (*Haliotis rufescens*) reveal patterns of gene flow and local adaptation, *Mol. Ecol.*, 22, 2884–2897, doi:10.1111/mec.12081, 2012.

Doney, S. C., Fabry, V. J., Feely, R. A., and Kleypas, J. A.: Ocean acidification: the other CO₂ problem, *Annual Review of Marine Science*, 1, 169–192, 2009.

Dore, J. E., Lukas, R., Sadler, D. W., Church, M. J., and Karl, D. M.: Physical and biogeochemical modulation of ocean acidification in the central North Pacific, *P. Natl. Acad. Sci. USA*, 106, 12235–12240, 2009.

Edmands, S., Moberg, P. E., and Burton, R. S.: Allozyme and mitochondrial DNA evidence of population subdivision in the purple sea urchin *Strongylocentrotus purpuratus*, *Mar. Biol.*, 126, 443–450, 1996.

Evans, T. G. and Hofmann, G. E.: Defining the limits of physiological plasticity: how gene expression can assess and predict the consequences of ocean change, *Philos. T. R. Soc. B*, 367, 1733–1745, doi:10.1098/rstb.2012.0019, 2012.

Evans, T. G., Chan, F., Menge, B. A., and Hofmann, G. E.: Transcriptomic responses to ocean acidification in larval sea urchins from a naturally variable pH environment, *Mol. Ecol.*, 22, 1609–1625, doi:10.1111/mec.12188, 2013.

Fassbender, A. J., Sabine, C. L., Feely, R. A., Langdon, C., and Mordy, C. W.: Inorganic carbon dynamics during northern California coastal upwelling, *Cont. Shelf. Res.*, 31, 1180–1192, doi:10.1016/j.csr.2011.04.006, 2011.

Feely, R. A., Sabine, C. L., Hernandez-Ayon, J. M., Ianson, D., and Hales, B.: Evidence for upwelling of corrosive “acidified” water onto the continental shelf, *Science*, 20, 1490–1492, doi:10.1126/science.1155676, 2008.

Feely, R. A., Doney, S. C., and Cooley, S. R.: Ocean acidification: present conditions and future changes in a high-CO₂ world, *Oceanography*, 22, 37–47, 2009.

Frieder, C. A., Nam, S. H., Martz, T. R., and Levin, L. A.: High temporal and spatial variability of dissolved oxygen and pH in a nearshore California kelp forest, *Biogeosciences*, 9, 3917–3930, doi:10.5194/bg-9-3917-2012, 2012.

Exploring local adaptation and the ocean acidification seascape

G. E. Hofmann et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Gaylord, B., Hill, T. M., Sanford, E., Lenz, E. A., Jacobs, L. A., Sato, K. N., Russell, A. D., and Hettinger, A.: Functional impacts of ocean acidification in an ecologically critical foundation species, *J. Exp. Biol.*, 214, 2586–2594, 2011.

Gillson, L., Dawson, T. P., Jack, S., and McGeoch, M. A.: Accommodating climate change contingencies in conservation strategy, *Trends Ecol. Evol.*, 28, 135–142, doi:10.1016/j.tree.2012.10.008, 2013.

Gouhier, T. C., Guichard, F. D. R., and Menge, B. A.: Ecological processes can synchronize marine population dynamics over continental scales, *P. Natl. Acad. Sci. USA*, 107, 8281–8286, doi:10.1073/pnas.0914588107, 2010.

Gruber, N., Hauri, C., Lachkar, Z., Loher, D., Froelicher, T. L., and Plattner, G.-K.: Rapid progression of ocean acidification in the California Current System, *Science*, 337, 220–223, doi:10.1126/science.1216773, 2012.

Hall-Spencer, J. M., Rodolfo-Metalpa, R., Martin, S., Ransome, E., Fine, M., Turner, S. M., Rowley, S., Tedesco, D., and Buia, M.-C.: Volcanic carbon dioxide vents show ecosystem effects of ocean acidification, *Nature*, 454, 96–99, doi:10.1038/nature07050, 2008.

Harris, K. E., DeGrandpre, M. D., and Hales, B.: Aragonite saturation state dynamics in a coastal upwelling zone, *Geophys. Res. Lett.*, 40, 1–6, 2013.

Harvey, B. P., Gwynn-Jones, D., and Moore, P. J.: Meta-analysis reveals complex marine biological responses to the interactive effects of ocean acidification and warming, *Ecol. Evol.*, 3, 1016–1030, doi:10.1002/ece3.516, 2013.

Hauri, C., Gruber, N., Plattner, G.-K., Alin, S., Feely, R. A., Hales, B., and Wheeler, P. A.: Ocean acidification in the California Current System, *Oceanography*, 22, 60–71, 2009.

Hauri, C., Gruber, N., Vogt, M., Doney, S. C., Feely, R. A., Lachkar, Z., Leinweber, A., McDonnell, A. M. P., Munnich, M., and Plattner, G.-K.: Spatiotemporal variability and long-term trends of ocean acidification in the California Current System, *Biogeosciences*, 10, 193–216, doi:10.5194/bg-10-193-2013, 2013.

Helmuth, B.: From cells to coastlines: how can we use physiology to forecast the impacts of climate change?, *J. Exp. Biol.*, 212, 753–760, 2009.

Hettinger, A., Sanford, E., Hill, T. M., Russell, A. D., Sato, K. N. S., Hoey, J., Forsch, M., Page, H. N., and Gaylord, B.: Persistent carry-over effects of planktonic exposure to ocean acidification in the Olympia oyster, *Ecology*, 93, 2758–2768, 2012.

Hoffmann, A. A. and Sgrò, C. M.: Climate change and evolutionary adaptation, *Nature*, 470, 479–485, 2011.

Exploring local adaptation and the ocean acidification seascape

G. E. Hofmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Hofmann, G. E., Smith, J. E., Johnson, K. S., Send, U., Levin, L. A., Micheli, F., Paytan, A., Price, N. N., Peterson, B., Takeshita, Y., Matson, P. G., Crook, E. D., Kroeker, K. J., Gambi, M. C., Rivest, E. B., Frieder, C. A., Yu, P. C., and Martz, T. R.: High-frequency dynamics of ocean pH: a multi-ecosystem comparison, *PLoS ONE*, 6, e28983, doi:10.1371/journal.pone.0028983, 2011.

Hofmann, G. E., Blanchette, C. A., Rivest, E. B., and Kapsenberg, L.: Taking the pulse of marine ecosystems: the importance of coupling long-term physical and biological observations in the context of global change biology, *Oceanography*, in press, 2013.

Huey, R. B., Kearney, M. R., Krockenberger, A., Holtum, J. A. M., Jess, M., and Williams, S. E.: Predicting organismal vulnerability to climate warming: roles of behaviour, physiology and adaptation, *Philos. T. R. Soc. B*, 367, 1665–1679, 2012.

Ibáñez, I., Gornish, E. S., Buckley, L., Debinski, D. M., Hellmann, J., Helmuth, B., HilleRis-Lambers, J., Latimer, A. M., Miller-Rushing, A. J., and Uriarte, M.: Moving forward in global-change ecology: capitalizing on natural variability, *Ecology and Evolution*, 3, 170–181, doi:10.1002/ece3.433, 2013.

Jacobs, D. K., Haney, T. A., and Louie, K. D.: Genes, diversity, and geologic process on the pacific coast, *Annu. Rev. Earth Pl. Sc.*, 32, 601–652, 2004.

Johansen, H. W.: *Coralline Algae, a First Synthesis*, CRC Press, Inc., Boca Raton, Florida, 239 pp., 1981.

Kawecki, T. J. and Ebert, D.: Conceptual issues in local adaptation, *Ecol. Lett.*, 7, 1225–1241, 2004.

Kelly, M. W. and Hofmann, G. E.: Adaptation and the physiology of ocean acidification, *Funct. Ecol.*, doi:10.1111/j.1365-2435.2012.02061.x, 2012.

Kelly, M. W., Padilla-Gamiño, J. L., and Hofmann, G. E.: Natural variation, and the capacity to adapt to high $p\text{CO}_2$ in a keystone species, *Glob. Change Biol.*, 19, 2536–2546, doi:10.1111/gcb.12251, 2013.

Kelly, R. P., Foley, M. M., Fisher, W. S., Feely, R. A., Halpern, B. S., Waldbusser, G. G., and Caldwell, M. R.: Mitigating local causes of ocean acidification with existing laws, *Science*, 332, 1036–1037, doi:10.1126/science.1203815, 2011.

Koch, M., Bowes, G., Ross, C., and Zhang, X. H.: Climate change and ocean acidification effects on seagrasses and marine macroalgae, *Glob. Change Biol.*, 19, 103–132, doi:10.1111/j.1365-2486.2012.02791.x, 2013.

Exploring local adaptation and the ocean acidification seascape

G. E. Hofmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Kroeker, K. J., Micheli, F., Gambi, M. C., and Martz, T. R.: Divergent ecosystem responses within a benthic marine community to ocean acidification, *Proc. Natl. Acad. Sci. USA*, 108, 14515–14520, doi:10.1073/pnas.1107789108, 2011.
- 5 Kroeker, K. J., Kordas, R. L., Crim, R., Hendriks, I. E., Ramajo, L., Singh, G. S., Duarte, C. M., and Gattuso, J.-P.: Impacts of ocean acidification on marine organisms: quantifying sensitivities and interaction with warming, *Glob. Change Biol.*, 19, 1884–1896, doi:10.1111/gcb.12179, 2013.
- LaVigne, M., Hill, T. M., Sanford, E., Gaylord, B., Russell, A. D., Lenz, E. A., Hosfelt, J. D., and Young, M. K.: The elemental composition of purple sea urchin (*Strongylocentrotus purpuratus*) calcite and potential effects of $p\text{CO}_2$ during early life stages, *Biogeosciences*, 10, 3465–3477, doi:10.5194/bg-10-3465-2013, 2013.
- 10 Martz, T. R., Connery, J. G., and Johnson, K. S.: Testing the Honeywell Durafet® for seawater pH applications, *Limnol. Oceanogr.-Meth.*, 8, 172–184, 2010.
- Matson, P. G., Yu, P. C., Sewell, M. A., and Hofmann, G. E.: Development under elevated $p\text{CO}_2$ conditions does not affect lipid utilization and protein content in early life-history stages of the purple sea urchin, *Strongylocentrotus purpuratus*, *Biol. Bull.*, 223, 312–327, 2012.
- 15 Menge, B. A. and Menge, D. N. L.: Dynamics of coastal meta-ecosystems: the intermittent upwelling hypothesis and a test in rocky intertidal regions, *Ecol. Monogr.*, doi:10.1890/12-1706.1, 2013.
- 20 Menge, B. A., Chan, F., and Lubchenco, J.: Response of a rocky intertidal ecosystem engineer and community dominant to climate change, *Ecol. Lett.*, 11, 151–162, 2008.
- Miklasz, K. A.: Physical Constraints on the Size and Shape of Microalgae, Stanford University, 268 pp., 2012.
- Navarrete, S. A., Broitman, B. R., and Menge, B. A.: Interhemispheric comparison of recruitment to intertidal communities: pattern persistence and scales of variation, *Ecology*, 89, 1308–1322, doi:10.1890/07-0728.1, 2008.
- 25 O'Donnell, M. J., Hammond, L. M., and Hofmann, G. E.: Predicted impact of ocean acidification on a marine invertebrate: elevated CO_2 alters response to thermal stress in sea urchin larvae, *Mar. Biol.*, 156, 439–446, 2009.
- 30 O'Donnell, M. J., Todgham, A. E., Sewell, M. A., Hammond, L. M., Ruggiero, K., Fanguie, N. A., Zippay, M. L., and Hofmann, G. E.: Ocean acidification alters skeletogenesis and gene expression in larval sea urchins, *Mar. Ecol.-Prog. Ser.*, 398, 157–171, doi:10.3354/meps08346, 2010.

Exploring local adaptation and the ocean acidification seascape

G. E. Hofmann et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

- O'Donnell, M. J., George, M. N., and Carrington, E.: Mussel byssus attachment weakened by ocean acidification, *Nature Climate Change*, 3, 587–590, 2013.
- Orr, J. C., Fabry, V. J., Aumont, O., Bopp, L., Doney, S. C., Feely, R. A., Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F., Key, R. M., Lindsay, K., Maier-Reimer, E., Matear, R., Monfray, P., Mouchet, A., Najjar, R. G., Plattner, G.-K., Rodgers, K. B., Sabine, C. L., Sarmiento, J. L., Schlitzer, R., Slater, R. D., Totterdell, I. J., Weirig, M.-F., Yamanaka, Y., and Yool, A.: Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms, *Nature*, 437, 681–686, 2005.
- Padilla-Gamiño, J. L., Kelly, M. W., Evans, T. G., and Hofmann, G. E.: Temperature and CO₂ additively regulate physiology, morphology and genomic responses of larval sea urchins, *Strongylocentrotus purpuratus*, *P. Roy. Soc. B-Biol. Sci.*, 280, doi:10.1098/rspb.2013.0155, 2013.
- Pespeni, M. H., Garfield, D. A., Manier, M. K., and Palumbi, S. R.: Genome-wide polymorphisms show unexpected targets of natural selection, *P. Roy. Soc. B-Biol. Sci.*, 279, 1412–1420, doi:10.1098/rspb.2011.1823, 2012.
- Pespeni, M. H., Sanford, E., Gaylord, B., Hill, T. M., Hosfelt, J. D., Jaris, H. K., LaVigne, M., Lenz, E. A., Russell, A. D., Young, M. K., and Palumbi, S. R.: Evolutionary change during experimental ocean acidification, *Proc. Natl. Acad. Sci. USA*, 10, 6937–942, doi:10.1073/pnas.1220673110, 2013a.
- Pespeni, M. H., Barney, B. T., and Palumbi, S. R.: Differences in the regulation of growth and biomineralization genes revealed through long-term common-garden acclimation and experimental genomics in the purple sea urchin, *Evolution*, 67, 1901–1914, doi:10.1111/evo.12036, 2013b.
- Pettorelli, N.: Climate change as a main driver of ecological research, *J. Appl. Ecol.*, 49, 542–545, doi:10.1111/j.1365-2664.2012.02146.x, 2012.
- Pickett, S. T. A.: Space-for-time substitution as an alternative to long-term studies, in: *Long-Term Studies in Ecology: Approaches and Alternatives*, edited by: Likens, G. E., Springer-Verlag, New York, 110–135, 1989.
- Price, N. N., Martz, T. R., Brainard, R. E., and Smith, J. E.: Diel variability in seawater pH relates to calcification and benthic community structure on coral reefs, *PLoS ONE*, 7, e43843, doi:10.1371/journal.pone.0043843, 2012.

Exploring local adaptation and the ocean acidification seascape

G. E. Hofmann et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

- Provoost, P., van Heuven, S., Soetaert, K., Laane, R. W. P. M., and Middelburg, J. J.: Seasonal and long-term changes in pH in the Dutch coastal zone, *Biogeosciences*, 7, 3869–3878, doi:10.5194/bg-7-3869-2010, 2010.
- Sanford, E. and Kelly, M. W.: Local adaptation in marine invertebrates, *Annual Review of Marine Science*, 3, 509–535, doi:10.1146/annurev-marine-120709-142756, 2011.
- Steinberg, D. K., Martinson, D. G., and Costa, D. P.: Two decades of pelagic ecology in the Western Antarctic Peninsula, *Oceanography*, 25, 56–67, 2012.
- Sunday, J. M., Crim, R. N., Harley, C. D. G., and Hart, M. W.: Quantifying rates of evolutionary adaptation in response to ocean acidification, *PLoS ONE*, 6, e22881, doi:10.1371/journal.pone.0022881, 2011.
- Tatters, A. O., Fu, F. X., and Hutchins, D. A.: High CO₂ and silicate limitation synergistically increase the toxicity of *Pseudonitzschia fraudulenta*, *PLoS ONE*, 7, e32116, doi:10.1371/journal.pone.0032116, 2012.
- Thomsen, J., Casties, I., Pansch, C., Körtzinger, A., and Melzner, F.: Food availability outweighs ocean acidification effects in juvenile *Mytilus edulis*: laboratory and field experiments, *Glob. Change Biol.*, 19, 1017–1027, doi:10.1111/gcb.12109, 2013.
- Visser, M. E.: Keeping up with a warming world, assessing the rate of adaptation to climate change, *P. R. Soc. B-Biol. Sci.*, 275, 649–659, 2008.
- Wernberg, T., Smale, D. A., Tuya, F., Thomsen, M. S., Langlois, T. J., de Bettignies, T., Bennett, S., and Rousseaux, C. S.: An extreme climatic event alters marine ecosystem structure in a global biodiversity hotspot, *Nature Climate Change*, 3, 78–82, 2013.
- Williams, S. E., Shoo, L. P., Isaac, J. L., Hoffmann, A. A., and Langham, G.: Towards an integrated framework for assessing the vulnerability of species to climate change, *PLoS Biology*, 6, e325, doi:10.1371/journal.pbio.0060325, 2008.
- Wootton, J. T. and Pfister, C. A.: Carbon system measurements and potential climatic drivers at a site of rapidly declining ocean pH, *PLoS ONE*, 7, e53396, doi:10.1371/journal.pone.0053396, 2012.
- Wootton, J. T., Pfister, C. A., and Forester, J. D.: Dynamic patterns and ecological impacts of declining ocean pH in a high-resolution multi-year dataset, *P. Natl. Acad. Sci. USA*, 105, 18848–18853, 2008.

Yu, P. C., Matson, P. G., Martz, T. R., and Hofmann, G. E.: The ocean acidification seascape and its relationship to the performance of calcifying marine invertebrates: laboratory experiments on the development of urchin larvae framed by environmentally-relevant $p\text{CO}_2/\text{pH}$, J. Exp. Mar. Biol. Ecol., 400, 288–295, doi:10.1016/j.jembe.2011.02.016, 2011.

BGD

10, 11825–11856, 2013

Exploring local adaptation and the ocean acidification seascape

G. E. Hofmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Exploring local adaptation and the ocean acidification seascape

G. E. Hofmann et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Table 1. Ocean acidification studies on functional traits of organisms in the California Current Large Marine Ecosystem.

Taxonomic Group	Species name	Life Stage	Variable Measured	Response	Source
Invertebrate	<i>Strongylocentrotus franciscanus</i>	Larvae	Transcription of <i>hsp70</i>	Larvae that were raised under enhanced CO ₂ displayed compromised expression of the thermally induced molecular chaperone <i>hsp70</i> .	O'Donnell et al. (2009)
Invertebrate	<i>Lytechinus pictus</i>	Larvae	Morphometrics Transcriptomics	Larvae cultured under high CO ₂ were smaller and had a more triangular body. Downregulation of genes related to metabolism and biomineralization. Upregulation of a few genes related to ion regulation and acid-base pathways.	O'Donnell et al. (2010)
Invertebrate	<i>Mytilus californianus</i>	Larvae	Shell strength and size	Larvae raised under high CO ₂ developed thinner, weaker and smaller shells as compared to controls.	Gaylord et al. (2011)
Invertebrate	<i>Mytilus trossulus</i> and <i>Strongylocentrotus franciscanus</i>	Larvae	Developmental rate	Decrease in larval body size in both the sea urchin and the mussel	Sunday et al. (2011)
Invertebrate	<i>Haliotis kamtschatkana</i>	Larvae	Development, size and survivorship	Exposure to high CO ₂ during development resulted in lower survival, abnormal development and smaller size in the northern abalone	Crim et al. (2011)
Invertebrate	<i>Strongylocentrotus purpuratus</i>	Larvae	Development Size	No differences in development between low and high CO ₂ treatments. Smaller larvae in the high CO ₂ treatment.	Yu et al. (2011)
Microbes	Nitrifiers	n/a	Ammonia oxidation rates	Microbial nitrification rates decreased when pH was experimentally reduced.	Beman et al. (2011)
Invertebrate	<i>Ostreola conchaphila</i>	Larvae Juvenile	Shell growth rate, shell area	Carryover effects of water chemistry experienced at early stage (larvae) to the later post-larval juvenile stage. Juveniles exposed to acidified conditions in their larval stages failed to attain the same size as juveniles derived from larvae exposed to less acidic conditions.	Hettinger et al. (2012)

Exploring local adaptation and the ocean acidification seascape

G. E. Hofmann et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 1. Continued.

Taxonomic Group	Species name	Life Stage	Variable Measured	Response	Source
Algae	<i>Pseudo-nitzshia fraudulenta</i>	n/a	Domoic acid production, cellular Si:C ratios	Strong synergism between high CO ₂ levels and silicate-limited growth, which greatly increases cellular toxicity relative to growth under lower CO ₂ conditions.	Tatters et al. (2012)
Invertebrate	<i>Strongylocentrotus purpuratus</i>	Juvenile Adults	Mg/Ca, Sr/Ca	Composition of skeleton precipitated during early and adult life history stages appears relative robust to spatial gradients and predicted changes in seawater chemistry.	LaVigne et al. (2012)
Invertebrate	<i>Strongylocentrotus purpuratus</i>	Larvae	Morphometrics and biochemical responses	Lipid utilization rates and protein content did not vary with pCO ₂ , larval growth was reduced at elevated pCO ₂ despite similar rates of energy utilization, and relationships between egg phospholipid content and larval length were found under control but not high pCO ₂ conditions.	Matson et al. (2012)
Invertebrate	<i>Strongylocentrotus purpuratus</i>	Larvae	Transcriptomics	Upregulation of several calcium transporters and binding proteins in a population of urchins from an intertidal site characterized by low and fluctuating pH.	Evans et al. (2013a)
Invertebrate	<i>Strongylocentrotus purpuratus</i>	Larvae	Development, Size, Respiration, Transcriptomics	Additive effects of high temperature and high CO ₂ conditions induce metabolic depression and a major downregulation of histone encoding genes. No differences in development between CO ₂ treatments were found.	Padilla-Gamiño et al. (2013)
Invertebrate	<i>Strongylocentrotus purpuratus</i>	Larvae	Heritability, Genetic Variation, Size	Genetic variation in the response to ocean acidification, indicating potential for adaptation. Smaller larvae under high CO ₂ conditions.	Kelly et al. (2013)
Invertebrate	<i>Mytilus trossulus</i>	Adults	Strength of byssal threads	Threads grown under high pCO ₂ conditions exhibited compromised mechanical characteristics.	O'Donnell et al. (2013)




Fig. 1. Map of the OMEGAS study sites in the California Current Large Marine Ecosystem. The OMEGAS primary study sites are from north to south: Fogarty Creek OR (FC), Strawberry Hill (SH), Van Damme (VD), Bodega Marine Reserve (BMR), Terrace Point (TPT) and Lompoc Landing (LOL). Note: Colors in the image sea surface temperature (degree C) from the NOAA CoastWatch database.

Exploring local adaptation and the ocean acidification seascape

G. E. Hofmann et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Exploring local adaptation and the ocean acidification seascape

G. E. Hofmann et al.




Fig. 2a. Cumulative frequencies of exposure to pH (total-scale) less than 7.8 (**A**) and 7.7 (**B**) as recorded at 10 min intervals by in-situ Durafet[®]-based pH sensors deployed in intertidal environment between April and September (maximum record length). Exposure frequencies at a given pH threshold are normalized to expanding windows of maximum to minimum in-situ water temperature to illustrate the response of pH to the progressive strengthening of seasonal upwelling at each site.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 2b. Continued.

Exploring local adaptation and the ocean acidification seascape

G. E. Hofmann et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 3. Fertilization success of *Strongylocentrotus purpuratus* gametes across populations and $p\text{CO}_2$ treatments. Adult urchins were collected from two locations that spanned the CCLME and the OMEGAS study sites: Fogarty Creek, OR (FC) and Bodega Marine Reserve, CA (BMR). Adult pairs from the same site were spawned in the laboratory (FC $N = 12$ pairs; BMR $N = 8$ pairs). Eggs were fertilized at 14°C under control ($400\ \mu\text{atm}$) and elevated ($800\ \mu\text{atm}$, $1200\ \mu\text{atm}$) $p\text{CO}_2$ conditions at an ecologically relevant sperm concentration of 7×10^3 sperm mL^{-1} . Gametes were allowed a 30 s contact time and at least 200 eggs were scored per pair after the second cleavage. There was no significant effect of $p\text{CO}_2$ by population (two-way ANOVA; $p\text{CO}_2 \times \text{site}$ effect $p = 0.8577$; $p\text{CO}_2$ effect $p = 0.3168$; site effect $p = 0.7147$). A similar pattern was observed at higher and lower sperm concentrations (data not shown).

Exploring local adaptation and the ocean acidification seascape

G. E. Hofmann et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A. Respiration in larval urchins




Photo Credit – J. Padilla-Gamiño

B. Mg/Ca and Sr/Ca in urchins




Photo Credit – G. Hofmann

C. Growth in intertidal mussels




Photo Credit – C. Blanchette

Fig. 4. Responses of marine calcifiers at the OMEGAS sites. **(A)** respiration rates in larval sea urchins of *Strongylocentrotus purpuratus*; Sea urchin larvae from the Santa Barbara and Van Damme were raised at 13 °C under high (1100 μ atm) and low (400 μ atm) p CO₂ using a flow-through CO₂ mixing system. Rates of oxygen consumption were obtained 92 h after fertilization in early pluteus larvae. Larval respiratory rates did not differ among sites or p CO₂ treatments. **(B)** Biomineral composition of the spines of adult *S. purpuratus*; data shown are mean Mg/Ca (upper panel) and Sr/Ca (lower panel) ratios for spines collected from adult urchins at OMEGAS study sites: Purisma Point, CA (PP) near Lompoc Landing, Terrance Point CA (TP), Van Damme State Park (VD) and Fogarty Creek, OR (FC); **(C)** Growth of intertidal mussels, *Mytilus californianus*, during the 2011 summer upwelling season. Mussels ~ 40 mm in length were collected in April 2011 from the OMEGAS sites, notched at the posterior lip of the shell, and outplanted back to each site. Mussels were retrieved in October 2011, and growth was measured as the increase in shell length. Growth rate is expressed as mm day^{-1} , standardized by initial length; results shown here are scaled by the number of days for the study duration at each site and the individual mussel length at study initiation. (FC = Fogarty Creek, SH = Strawberry Hill, VD = Van Damme, BO = Bodega, TP = Terrace Point, LL = Lompoc).

Exploring local adaptation and the ocean acidification seascape

G. E. Hofmann et al.




Fig. 5. Breeding experiment conducted using purple urchin adults collected from two OMEGAS study sites. **(A)** Cross design: we crossed adult urchins from two sites in California, USA, known to differ in their carbonate chemistry regimes (OMEGAS sites – northern CA = Van Damme; southern CA = Santa Barbara). In the lab, we crossed each male with two females from his own site, and two females from the opposite site, and split embryos from each cross into low and high $p\text{CO}_2$ rearing conditions. **(B)** Total larval length (\pm SD) of *S. purpuratus* raised for five days under low $p\text{CO}_2$ and high $p\text{CO}_2$ conditions. Offspring of northern CA sires are larger than the offspring of southern CA sires under high $p\text{CO}_2$ (pMCMC = 0.012) but not under low $p\text{CO}_2$ (pMCMC = 0.27). Modified from Kelly et al., 2013.