

**Carbon variability in
Southern Ocean**

L. Resplandy et al.

Observed small spatial scale and seasonal variability of the CO₂-system in the Southern Ocean

L. Resplandy^{1,*}, J. Boutin¹, and L. Merlivat¹

¹LOCEAN (CNRS, IRD, MNHN, UPMC), France

* now at: Laboratoire des sciences du climat et de l'environnement (LSCE) – UMR 8212, France

Received: 19 June 2013 – Accepted: 1 July 2013 – Published: 22 August 2013

Correspondence to: L. Resplandy (laure.resplandy@lsce.ipsl.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

The considerable uncertainties in the carbon budget of the Southern Ocean are largely attributed to unresolved variability, in particular at seasonal time scale and small spatial scale (~ 100 km). In this study, the variability of surface $p\text{CO}_2$ and DIC at seasonal and small-spatial scales is examined using a dataset of surface drifters including $\sim 80\,000$ measurements at high spatio-temporal resolution. On spatial scales of 100 km, we find gradients ranging from 5 to 50 μatm for $p\text{CO}_2$ and 2 to 30 $\mu\text{mol kg}^{-1}$ for DIC, with highest values in energetic and frontal regions. This result is supported by a second estimate obtained with SST satellite images and local DIC/SST relationships derived from drifters observations. We find that dynamical processes drives the variability of DIC at small spatial scale in most regions of the Southern Ocean, the cascade of large-scale gradients down to small spatial scales leading to gradients up to 15 $\mu\text{mol kg}^{-1}$ over 100 km. Although the role of biological activity is more localized, it enhances the variability up to 30 $\mu\text{mol kg}^{-1}$ over 100 km. The seasonal cycle of surface DIC is reconstructed following Mahadevan et al. (2011), using an annual climatology of DIC and a monthly climatology of mixed layer depth. This method is evaluated using drifters observations and proves to be a reasonable first-order estimate of the seasonality in the Southern Ocean, which could be used to validate models simulations. We find that small spatial scales structures are a non negligible source of variability for DIC, with amplitudes of about a third of the variations associated with the seasonality and up to 10 times the magnitude of large-scale gradients. The amplitude of small-scale variability reported here should be kept in mind when inferring temporal changes (seasonality, inter-annual variability, decadal trends) of the carbon budget from low resolution observations and models.

Carbon variability in Southern Ocean

L. Resplandy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

The Southern Ocean is a key region for the global carbon cycle and the climate system, as it accounts for about 25–30 % of the total anthropogenic carbon uptake (Orr et al., 2001; Mikaloff Fletcher et al., 2006; Gruber et al., 2009; Takahashi et al., 2009). The carbon flux in this region is governed by the balance between a large uptake of anthropogenic CO₂ and a substantial outgassing of natural CO₂ (Lovenduski et al., 2008; Gruber et al., 2009). Despite its importance, the Southern Ocean remains the region where uncertainties on the air–sea CO₂ flux and the carbon budget are the highest (e.g. Gruber et al., 2009). Quantifying this subtle balance and its sensitivity to dynamical and biological changes is therefore one of the most challenging issues in assessing today's global carbon budget.

The recent syntheses of Gruber et al. (2009) and Lenton et al. (2013) compared the CO₂ uptake by the Southern Ocean estimated from various models (ocean biogeochemical models, inverse atmospheric and oceanic models) with those derived from observations. Although the mean of all models agree surprisingly well with observations to an uptake of $\sim 0.34 \pm 0.2 \text{ PgC yr}^{-1}$ in both Gruber et al. (2009) and Lenton et al. (2013), individual model results spread over a wide range, between 0.15 and 0.72 PgC yr⁻¹ in Gruber et al. (2009) and between -0.3 (i.e. outgazing) and 0.3 PgC yr⁻¹ in Lenton et al. (2013).

These large discrepancies in the flux are also found between the various estimates derived from observations at the regional level and on seasonal timescales. For example, Barbero et al. (2011) recomputed the CO₂ ocean uptake in the Southern Ocean Pacific sector using $p\text{CO}_2$ maps extrapolated from observations by McNeil et al. (2007); Boutin et al. (2008); Takahashi et al. (2009); Barbero et al. (2011). From these four methods, the oceanic CO₂ uptake varies between 0.05 and 0.50 PgC yr⁻¹, with highest discrepancies (up to 0.8 PgC yr⁻¹) during winter.

These large uncertainties are primarily attributed to unresolved spatio-temporal variability and to high wind speeds that translates moderate errors on $p\text{CO}_2$ into large er-

BGD

10, 13855–13895, 2013

Carbon variability in Southern Ocean

L. Resplandy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Carbon variability in Southern Ocean

L. Resplandy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



rors in air–sea CO₂ fluxes. In this paper, we focus on the issue of unresolved variability. The Southern Ocean is remote and hardly accessible in winter leading to a very sparse spatio-temporal coverage of hydrographic observations, including measurements of surface pCO₂ (Takahashi et al., 2009; Pfeil et al., 2012). Undersampling biases are aggravated by the high variability this oceanic region displays over a wide range of temporal and spatial scales. The combination of various observation- and model-based methods have recently enabled a better quantification of the air–sea CO₂ flux in the Southern Ocean and its variability at large spatial scale (e.g. Gruber et al., 2009) and inter-annual time scale (e.g. Séférian et al., 2013). Substantial variability was captured at decadal time-scales, with estimates derived from observations and models of the order of 0.05 to 0.15 PgC yr⁻¹ (Metzl, 2009; Séférian et al., 2013; Brix et al., 2013) and attributed to sea–ice interactions leading to deep mixing events (Séférian et al., 2013) and the Southern Annular Mode (Brix et al., 2013; Séférian et al., 2013).

The variability introduced at smaller spatial and shorter temporal scales remains largely unknown. Periodic surveys in the Indian sector and a 10 yr time-series located south of New Zealand have revealed relatively large variations of oceanic surface pCO₂ at seasonal time scale (10–40 μatm, Metzl et al., 2006; Brix et al., 2013). Although these surveys provide very useful information, the seasonality of the CO₂ system is still poorly constrained at the basin scale, complicating the interpretation of observations and the evaluation of ocean models in this region. Furthermore, the Southern Ocean is one of the most energetic regions of the world ocean, with numerous mesoscale eddies modulating the dynamical transport of biogeochemical tracers including carbon (Rintoul et al., 2001; Ito et al., 2004; Tortell et al., 2011) and the biological activity (Read et al., 2007; Tortell et al., 2011). The few observations of these small spatial scales are however too sparse to assess the role of small-scale structures on the carbon budget.

In this study, we estimate the variability of the CO₂ system introduced by small spatial scale structures (~ 100 km) and the seasonality in the Southern Ocean. We combine an extensive data set of surface drifters including ~ 80 000 yr-round measurements at high spatio-temporal resolution with satellite and in-situ gridded products to quantify

Carbon variability in
Southern Ocean

L. Resplandy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the small scale variability of $p\text{CO}_2$ and dissolved inorganic carbon (DIC), identify the source of this variability (dynamical and/or biological) and compare it to the variability introduced by regional contrasts and the seasonality. Section 2 describes the observations and the various methods used. Section 3 examines the variability of surface $p\text{CO}_2$ and DIC in a case study around Crozet Island before extending the analysis to the Southern Ocean. Finally, Sect. 4 synthesizes and discusses the main results.

2 Methods

2.1 Observations

2.1.1 Carioca drifters

Between 2003 and 2009, 9 autonomous lagrangian CARbon Interface Ocean Atmosphere (Carioca) drifters were deployed in the Southern Ocean, acquiring an extensive dataset of 80 537 individual observations (Fig. 1a). Carioca drifters measure hourly, at a depth of 2 m, CO_2 fugacity ($f\text{CO}_2$), sea surface temperature (SST), sea surface salinity (SSS) and fluorescence (Copin-Montégut et al., 2004; Boutin et al., 2008). Extensive validation of Carioca measurements in the Southern Ocean indicate an absolute (relative) precision close to 3 (1) microatm (see Appendix of Boutin et al., 2008). Alkalinity is derived from SSS and SST using Lee et al. (2006) formulation; DIC is derived from $f\text{CO}_2$, alkalinity, SST and SSS using CO_2 dissociation constants and solubility as described in Boutin et al. (2008). The unit for DIC is $\mu\text{mol kg}^{-1}$. The relative precision of successive DIC values is expected to be $0.5 \mu\text{mol kg}^{-1}$. In order to avoid artificial diurnal cycle of fluorescence created by the quenching effect, we consider daily average fluorescence. Finally, to allow comparison, floats observations and the Glodap climatology were normalised to the year 2005 assuming linear drifts of $0.7 \mu\text{mol kg}^{-1} \text{yr}^{-1}$ for DIC and $1.4 \mu\text{atm yr}^{-1}$ for $p\text{CO}_2$ (Takahashi et al., 2009; Boutin and Merlivat, 2009).

2.1.2 Satellite data and global climatologies

We use 3-D-gridded DIC fields from Glodap (Key et al., 2004, data available online at: cdiac.ornl.gov/ftp/oceans/GLODAP) and Satellite SSTs from Advanced Microwave Scanning Radiometer Earth Observing System (AMSR-E, www.gbcc.msfc.nasa.gov/AMSR/), both collocated in time and space with drifters measurements using nearest neighbor interpolation. Fronts and transport barriers delimiting small-scale structures are computed in the subantarctic frontal zone around the Crozet Plateau. This is performed using Lagrangian coherent structures embedded in the velocity field with finite-size Lyapunov exponent technique (d'Ovidio et al., 2004; Lehahn et al., 2007). Velocity fields are from Ssalto/Duacs Aviso altimetry product (available online at www.aviso.oceanobs.com/en/data/product-information/duacs/).

2.2 Assessing small-scale variability: SV100km diagnostic

We quantify the variability of $p\text{CO}_2$ and DIC at small-scales using the 9 drifters extensive dataset. Carioca drifters sample with a horizontal resolution of 1 to 3 km (drifting speed of $\sim 0.3\text{--}0.9\text{ ms}^{-1}$ and hourly sampling), hence distinctly capturing the variability at the mesoscale ($\sim 100\text{ km}$). The contribution of mesoscale was disentangled from the large-scale and seasonal signal also captured by the drifters using a fast Fourier transform 20-day high-pass filter. This allows to filter out variations with time scales larger than 20 days (seasonal signal) and with spatial scales larger than $\sim 500\text{--}1000\text{ km}$ (distance covered with a drifting speed of $\sim 0.3\text{--}0.9\text{ ms}^{-1}$ in 20 days). In the following, \hat{X} denotes the anomaly of the field X associated with small scales obtained with the high-pass filter. The spatial variability of $p\text{CO}_2$ and DIC at small spatial scale is then evaluated using the diagnostic SV100km that quantifies the spatial variability on scales of 100 km (see Sect. 3.1.2). At each point t of the trajectory, SV100km is defined as the range of variation in a $100\text{ km} \times 100\text{ km}$ box centered on the point:

$$\text{SV100km}(t) = \max(\hat{X}(t, 100\text{ km})) - \min(\hat{X}(t, 100\text{ km})) \quad (1)$$

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



where $\widehat{X}(t, 100 \text{ km})$ is the time-series of the high-pass filtered field X included in a box of $100 \text{ km} \times 100 \text{ km}$ centered on t .

SV100km was also evaluated using a second method combining drifters measurements and satellite observations (see Sect. 3.3). This method relies on the frequent linear correlation found locally between $\widehat{\text{DIC}}$ and $\widehat{\text{SST}}$ on small spatial scales. The existence of “local” relationship between a carbon variable (namely ρCO_2) and SST was identified by Chen et al. (2007) across one eddy in the tropical Pacific. Here, we derive such “local” linear correlations between $\widehat{\text{DIC}}$ and $\widehat{\text{SST}}$ using 20-day sliding windows covering the complete dataset (Eq. 2). 2-D maps of DIC anomaly (noted $\widehat{\text{DIC}}_{\text{sat}}$) were then reconstructed by applying the local linear relationship obtained from in-situ observations to $\widehat{\text{SST}}_{\text{sat}}$ derived from AMSR-E satellite observations in $100 \text{ km} \times 100 \text{ km}$ boxes around the float location (Eq. 3). Here, $\widehat{\text{SST}}_{\text{sat}}$ is the SST anomaly obtained by removing the mean SST in each $100 \text{ km} \times 100 \text{ km}$ box. The reconstruction was performed only for correlation coefficients $r^2 \geq 0.7$ (see on Fig. 6). Here, $\text{SV100km}_{\text{sat}}$ is computed from a 2-D spatial map of DIC anomaly at a given time in a $100 \text{ km} \times 100 \text{ km}$ box around the point i (Eq. 4), whereas in the first method SV100km is computed from the time-series of DIC included in the same $100 \text{ km} \times 100 \text{ km}$ box (Eq. 1).

$$\widehat{\text{DIC}} = a \cdot \widehat{\text{SST}} + b, \quad (2)$$

$$\widehat{\text{DIC}}_{\text{sat}} = a \cdot \widehat{\text{SST}}_{\text{sat}} + b \quad (3)$$

$$\text{SV100km}_{\text{sat}}(i) = \max(\widehat{X}(i, 100 \text{ km}_{\text{sat}})) - \min(\widehat{X}(i, 100 \text{ km}_{\text{sat}})) \quad (4)$$

where , $\widehat{X}(i, 100 \text{ km})$ is the map of \widehat{X} included in a box of $100 \text{ km} \times 100 \text{ km}$ centered on i .

2.3 Drivers of small-scale variability: Principal Component Analysis

The mechanisms driving the variability at small-scale quantified by SV100km are examined using a Principal Component Analysis (PCA). PCA, also known as empirical

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Carbon variability in Southern Ocean

L. Resplandy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



orthogonal function analysis, has been widely used to explore the major factors that give rise to variability patterns in geosciences and in particular to surface $p\text{CO}_2$ variability (Dandonneau, 1995; Murata, 2006; Lohrenz et al., 2009). Here, we use this technique to examine how the variability at small spatial scales in the CO_2 system and more specifically in DIC concentrations relates to the variability of SST and fluorescence, a signature of biological processes. The choice of DIC over $p\text{CO}_2$ to make this analysis aims at disentangling the contribution of dynamical processes that influences $p\text{CO}_2$ through DIC and SST from the thermodynamical effect that only affects $p\text{CO}_2$. Furthermore, model- and observation-based studies concur with the fact that DIC is the main driver of surface $p\text{CO}_2$ variability in the Southern Ocean (Metzl et al., 2006; Lovenduski et al., 2008; Metzl, 2009; Brix et al., 2013; S  f  rian et al., 2013). The objective here is not to directly explain variations of DIC by variations in SST and fluorescence (Fluo) but to relate the small-scale spatial variability in DIC to either dynamical or biological factors and identify where and when those variations are dominant. Indeed, surface DIC concentrations generally vary in association with physical and/or biological factors, such as vertical and lateral dynamical transport or biological uptake by photosynthesis. The factors driving the variability of DIC at small-scales are likely to be different depending on the region and time of year. We therefore chose not to compute one PCA with the complete Carioca data set. Instead, we compute “local PCAs” over segments of the drifters trajectory to identify the factors driving the variability at small spatial scales. This approach differs from those of Dandonneau (1995); Murata (2006) and Lohrenz et al. (2009) that computed one PCA with their complete data set on relatively smaller regions.

Similarly to the method used to assess small-scale variability, PCAs are performed on $\widehat{\text{Fluo}}$, $\widehat{\text{SST}}$ and $\widehat{\text{DIC}}$ (i.e. filtered from variations with scales > 20 days and > 500 – 1000 km, see Sect. 2.2) and normalized to an average of 0 and a standard deviation of 1 (noted $\widehat{\text{Fluo}}_n$, $\widehat{\text{SST}}_n$ and $\widehat{\text{DIC}}_n$). At each point of the drifters trajectory, a PCA is computed over a 20-day window centered on the point with a data set of 480 rows (20-day hourly measurements) and 3 columns ($\widehat{\text{Fluo}}_n$, $\widehat{\text{SST}}_n$ and $\widehat{\text{DIC}}_n$). Each of the

~ 80 000 PCAs gives 3 “local principal components” (noted PC1, PC2 and PC3) and the variance explained by each of the PCs. For example, at 11.3° W/−45.3° N the PCA gives the following result:

$$PC1 = 0.63083 \cdot \widehat{SST}_n + 0.46049 \cdot \widehat{Fluo}_n - 0.62450 \cdot \widehat{DIC}_n,$$

$$PC2 = 0.30271 \cdot \widehat{SST}_n - 0.88713 \cdot \widehat{Fluo}_n - 0.34838 \cdot \widehat{DIC}_n,$$

$$PC3 = 0.460492 \cdot \widehat{SST}_n - 0.887132 \cdot \widehat{Fluo}_n + 0.030727 \cdot \widehat{DIC}_n,$$

with PC1, PC2 and PC3 respectively explains 73 %, 22 % and 5 % of the variance.

The contribution of the normalized variables to the PC is given by their loading (for example −0.6245 for DIC_n on PC1), its sign having the same meaning as the sign of a correlation coefficient. In this example, most of the variability is explained by PC1 (73 %) and is characterized by opposed variations of SST and DIC. In the following, only PC1 and PC2 that are significantly contributing to DIC small-scale variability, i.e. with explained variance greater than 20 % and cumulative contribution exceeding 80 %, are considered (see Sects. 3.1.1 and 3.2.2). Note that PCAs were not performed along the drifter 01110, for which fluorescence was not available (Fig. 1a).

3 Results

The variability of the CO₂ system at small spatial scale (i.e. ~ 100 km) and the processes driving this variability are first presented in a case study region around the Crozet Plateau. Then, the results are extended to the Southern Ocean using the observations of the nine drifters.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.1 Observed variability of the CO₂ system at small-scale

3.1.1 Case study – subantarctic frontal zone and Crozet Plateau

The case study is focused on a drifter drifting in the southwest Indian Ocean, more specifically in the subantarctic frontal zone and across the Crozet Plateau (20–60° E, 50–42° S), between December 2006 and April 2007 (Fig. 1a). The drifter followed a path characteristic of surface drifters in this area (Pollard et al., 2007). After drifting along the Subantarctic Front (SAF), it crossed the Southwest Indian Ridge in January 2007, where the Polar Front (PF) and SAF combine in what is the major pathway of the ACC across the ridge (35° E–48° S, Fig. 2). After following the SAF, the drifter headed north along the Subantarctic Front between the Del Caño Rise and Crozet in March (~ 48° E) and then drifted east through the Crozet Plateau in late April (Fig. 2). The evolution of $p\text{CO}_2$, DIC, SST and Fluo along this trajectory are modulated by the presence of the SAF, separating cold/DIC-rich southern waters from warmer/DIC-poor northern waters, and the presence of the Crozet Plateau, known to be biologically active with a bloom occurring annually between September and January (Venables et al., 2007). For example, high surface $p\text{CO}_2$ (up to 388 μatm) are maintained west of the ridge by either high DIC/low SST southern waters or relatively high SST/low DIC northern waters (Fig. 2a, c and g). In contrast, low $p\text{CO}_2$ sampled north of Crozet are associated with low DIC/high Fluo waters (Fig. 2a, c, e and g).

The variability at small spatial scales is highlighted by filtering out scales > 20 days and > 500–1000 km (see Sect. 3.1.2). The resulting anomalies in $p\text{CO}_2$, DIC, SST and fluorescence fields highlight the hotspots of variability at small spatial scales (Fig. 2 right panels). Although the position of the fronts indicated here is climatological and not co-located in time with the position of the drifter, most hotspots of variability are located in the vicinity of the front. This is not surprising as density fronts are a major generator of mesoscale eddies and potential dynamical and biological boundaries with contrasted properties on either side. The strength of the CO₂-system variability at small-spatial scales is estimated using the diagnostic SV100km that quantifies the

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



variability in 100 km × 100 km boxes (see Sect. 3.1.2 for details). Highest SV100km are found north of Crozet with small-scale variations of $p\text{CO}_2 \geq 15 \mu\text{atm}$ and small-scale variations of $\text{DIC} \geq 12 \mu\text{mol kg}^{-1}$ (Fig. 2i, j). In contrast, areas of lowest variability are found south of Del Caño Rise, where SV100km is of the order of 5 μatm and 3 $\mu\text{mol kg}^{-1}$ for $p\text{CO}_2$ and DIC respectively (Fig. 2i, j).

3.1.2 A first estimate for the Southern Ocean

The variability of $p\text{CO}_2$ and DIC at small spatial scale is estimated using the diagnostic SV100km computed along the nine drifters trajectories (Fig. 3a, b). Although the drifter in the case study crossed the SAF and the Crozet Plateau, the variability identified in this area is relatively low compared to other regions of the Southern Ocean. Highest variability of the CO_2 system is found in the energetic region of the Aghulas, where $p\text{CO}_2$ and DIC horizontal gradients respectively reach values ≥ 20 (up to 50) μatm and ≥ 10 (up to 30) $\mu\text{mol kg}^{-1}$ on spatial scales of 100 km (Fig. 3a, b). drifters also captured relatively high variability at small spatial scale close to frontal zones (PF, STF and SAF), where SV100km is of the order of 15–20 μatm and 10–15 $\mu\text{mol kg}^{-1}$ for $p\text{CO}_2$ and DIC respectively. High variability is found in regions of lower eddy kinetic energy (EKE) in the southern Pacific (150° W, 40–45° S and 120° W, 45–50° S). However when the two drifters 03740 and 01110 sampled this area in 2004, the SAF was located north of its climatological position, which probably explain the high variability captured in this region by the drifters (Barbero et al., 2011). Elsewhere, the variability typically ranges between 5 and 15 μatm and 2 and 10 $\mu\text{mol kg}^{-1}$.

3.2 Drivers of DIC variability at small-scale

Here we examine how the variability of surface DIC at small spatial scales relates to dynamical processes and biological activity. The respective role of dynamics and biology is identified by correlating observed variations of SST and fluorescence to variations of DIC surface concentrations, using locally derived principal component analyses (PCAs,

BGD

10, 13855–13895, 2013

Carbon variability in Southern Ocean

L. Resplandy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



see Sect. 2.3). The analysis of the PCAs along eight of the drifters reveals three major cases: the variability of DIC at small-scale is either dominated by (1) ocean dynamical processes, (2) biological uptake of DIC or (3) a combination of dynamical and biological processes. Note that we neglect the influence of air–sea fluxes by less than 15%. We discriminate between the three cases using the loadings of the PCs that are strongly correlated to DIC (i.e. with $|\text{DIC}_n \text{ loading}| > 0.3$). In the *dynamics-dominated case* (noted DYN), the variability at small-scale is dominated by the signature of anticorrelated large-scale gradients (vertical and/or lateral) of DIC and SST. Indeed, increasing depth and increasing latitude are characterized by decreasing temperature and increasing DIC (Fig. 1e). This case is hence identified by anticorrelated loadings on DIC and SST and a low loading of fluorescence compared to SST ($|\text{SST}_n \text{ loading}| > |\text{Fluo}_n \text{ loading}| + 0.3$). The *biology-dominated case* (noted BIO) is identified with anticorrelated loadings on DIC and fluorescence (DIC uptake by biological activity) and low loadings on SST compared to fluorescence (i.e. $|\text{SST}_n \text{ loading}| < |\text{Fluo}_n \text{ loading}| + 0.3$). Finally, the *dynamics-biology combined case* (noted DYN + BIO) correspond to anticorrelated SST and DIC, anticorrelated fluorescence and DIC and similar loadings on SST and fluorescence (i.e. $|\text{SST}_n \text{ loading}| - |\text{Fluo}_n \text{ loading}| \leq 0.3$).

3.2.1 Case study – subantarctic frontal zone and Crozet Plateau

The PCA reveals that the majority of the variance at small spatial scales in the case study area is explained by the first and second principal components (PC1 and PC2). Of the total variance, ~ 50 to 70 % is accounted for by PC1, 20 to 30 % by PC2 leading to a cumulated variance ≥ 80 % (Fig. 4a and b). The contribution of PC3 is significantly lower with a maximum explained variance of 20 % at ~ 33° E (not shown). The fact that PC3 explains a non-negligible part of the variance highlights that SST and fluorescence are not sufficient to properly explain all variations of the DIC concentration at small-spatial scales. However, the objective here is not to directly explain variations of DIC by variations in SST and fluorescence but to identify if the variability at small-scale can be correlated to either ocean dynamics or biological uptake of DIC. In this region,

BGD

10, 13855–13895, 2013

Carbon variability in Southern Ocean

L. Resplandy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Carbon variability in Southern Ocean

L. Resplandy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



only PC1 is strongly related to the variations of DIC ($|\text{loadings DIC}_n| \leq 0.7$, Fig. 4c). In contrast, DIC contribution to PC2 is weak ($|\text{loadings DIC}_n| \leq 0.3$), indicating that PC2 primarily reflects the joint variations of SST and fluorescence (Fig. 4d, f and h). In the following we therefore focus on PC1 that explains most of the DIC variability at small spatial scale.

In this region, PC1 opposes SST and DIC i.e. a decrease in DIC is correlated to an increase in SST (Fig. 4c and e). This negative correlation arises from the opposed large-scale horizontal and vertical gradients of SST and DIC (Fig. 1). What is identified here is the strong signature of large-scale patterns on the variability at smaller spatial scales. In the frontal zone, large-scale gradient of SST and DIC are sharpened while mesoscale eddies generated by the instability of the density front transport these properties across the front. In addition, fluorescence is strongly anti-correlated to DIC in particular on the Crozet Plateau, where the SV100km of ρCO_2 and DIC are high (Fig. 4c and g and Fig. 2j). Positive anomaly of fluorescence are associated with positive anomalies of SST indicating that biological activity is higher north of the SAF (Fig. 2f and h). The dominant drivers of DIC variability are identified using criteria on DIC, SST and fluorescence contribution to PC1 (see details in Sect. 2.3). We find that the ocean dynamics is the major factor explaining the observed variability of DIC at small spatial scale (DYN on Fig. 4i and combines with biological activity north of Crozet (DYN+BIO on Fig. 4i and Fig. 2j). This suggests that the impact of biological activity is more localized than the impact of ocean circulation but plays a crucial role by enhancing the amplitude of DIC variability at small-spatial scales.

3.2.2 Southern Ocean

Similarly to the case study, the PCA of DIC, SST and fluorescence reveals that most of the variability is explained by PC1 and PC2: PC1 accounts for 50 to 80 % of the variance (Fig. 5a) and PC2 for 10 to 50 % (Fig. 5b), leading to a cumulated variance explained by PC1 and PC2 ≥ 80 % in more than 85 % of the points (Fig. 5c). The contribution of PC3 is significantly lower, with an explained variance ≤ 20 % in more than

75% of the points. For clarity, we show here the processes driving the DIC distribution at small spatial scales identified from PCAs loadings, while the details of PC1, PC2 and PC3 loadings are shown in Appendix A (Figs. A1–A3).

From PC1, ocean dynamics is identified as the major mechanism setting DIC patterns at small scale, in particular between the Polar Front and the Subtropical Front (DYN on Fig. 5d). In these regions, PC1 primarily reflects the large-scale gradients opposing variations of DIC and SST (Fig. A1). In contrast, biological activity emerges as the dominant driver in the region of the Aghulas retroflection (BIO on Fig. 5d). Areas where biological activity dominates are characterized by positive contributions of fluorescence and negative contributions of SST and DIC (Fig. A1). The correlated variations of SST and DIC in these regions contrast with the anti-correlation observed everywhere else (Figs. A1a, b and 5). This suggests that, in these regions, DIC of cold and carbon-rich southern waters has been consumed by biological activity, therefore inverting the signature of the large-scale opposed gradients of DIC and SST. It is interesting to note that the dominance of biological activity depends more on the region than on the season of sampling, i.e. BIO is identified during winter as well as during summer (Figs. 1b and 5d). Indeed, seasonal variations of DIC, and hence the variations associated with the seasonal bloom, have been filtered out from this analysis. What is highlighted here is the role of biological activity in introducing variability on scales of 100 km.

Finally, in many regions, a combination of both biological activity and ocean circulation influences the DIC distribution at small spatial scales (DYN + BIO on Fig. 5d). In these regions, fluorescence and SST contributions to PC1 are large and negatively correlated to DIC. This suggests that ocean dynamics maintains opposed large-scale gradients of DIC and SST down to small spatial scales, while biological activity could play a significant role in the observed variability at small-scale (Fig. A1). The co-dominance of both factors is identified mainly along frontal areas or in regions of high EKE (Fig. 3). In one of these regions located between 142° W and 134° W in the Pacific sector, the joint contributions of biological activity and ocean circulation identified with the PCA is

BGD

10, 13855–13895, 2013

Carbon variability in Southern Ocean

L. Resplandy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Carbon variability in Southern Ocean

L. Resplandy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



strongly supported by the work of Barbero et al. (2011). In their work, Barbero et al. (2011) combined satellite chlorophyll *a* concentrations and net community primary production inferred from diel cycles of in-situ DIC concentrations in the southern Pacific (Boutin and Merlivat, 2009). They found that the drifter 03740 crossed patches of higher primary production between 142° W and 134° W, hence confirming the contribution of biological production in this region (see Fig. 3 Barbero et al., 2011). They also pointed out that this patches of biological activity were located along the SAF, which was at that time located north of its climatological position. This supports that the ocean dynamics enhanced by the presence of the front was a key player in setting DIC variability.

The dominant drivers identified from PC2 mostly correspond to the contribution of ocean dynamics (resp. biological activity) in regions where biological activity (resp. ocean dynamics) dominates PC1 (Fig. 5d, e). Note that segments where neither ocean circulation nor biological activity can explain variations of DIC, SST and fluorescence on PC1 and PC2 correspond to either areas where salinity plays a significant role or areas where no strong signature could be identified (indicated in brown on Fig. 5d, e).

3.3 How can satellite SST constrain the estimate of DIC variability at small-scale?

The predominance of ocean dynamics and the maintenance of opposed gradients of DIC and SST down to small spatial scales suggests that SST can be used as a proxy of DIC variability on scales of 100 km. In-situ DIC and SST are strongly linearly correlated in particular in regions where ocean circulation was identified as the dominant mechanism (see Sect. 3.1.2, Figs. 6a and 5d, e). Note that in contrast, surface $p\text{CO}_2$ and SST are only poorly correlated (Fig. 6b). Maps of DIC were reconstructed using the linear relationship derived locally between in-situ SST and DIC and AMSR-E SST maps.

The method is illustrated in the case-study area (Fig. 7). In Autumn 2007, the drifter 13060 followed the SAF highlighted by the presence of a sharp satellite SST gradient and a dynamical boundary indicated by altimetry (black contours on Fig. 7). Between

Carbon variability in
Southern Ocean

L. Resplandy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



18 March and 7 April the drifter trajectory was wrapped around two mesoscale eddies (noted E1 and E2), allowing the drifter to sample northern warm DIC-poor waters trapped in E1 and southern cold DIC-rich waters trapped in E2 (Fig. 7a, c and d). In this 20-day window centered on 28 March, in-situ \widehat{DIC} and \widehat{SST} are strongly anticorrelated ($\widehat{DIC} = -9.69 \cdot \widehat{SST} + 0.09$, with $r^2 > 0.86$). In addition, the SST sampled by the drifter and the SST observed by the satellite AMSR interpolated in space and time at the drifter location are in good agreement (Fig. 7a and b). This gives us confidence that the map of \widehat{DIC}_{sat} reconstructed from satellite SST and the local linear correlation obtained between \widehat{DIC} and \widehat{SST} provides a good estimate of the DIC variability at small spatial scale (Fig. 7e). In the box covered by the drifter (45 to 49° E; 48 to 45° S), the reconstructed \widehat{DIC}_{sat} varies roughly between -15 and $20 \mu\text{mol kg}^{-1}$. $SV100\text{km}_{sat}$, estimated in $100\text{km} \times 100\text{km}$ boxes, along the drifter trajectory has an average value of $10 \mu\text{mol kg}^{-1}$, with higher values (up to $15 \mu\text{mol kg}^{-1}$) in the vicinity of SAF and lower values (down to $5 \mu\text{mol kg}^{-1}$) further from the SAF. This estimate of the variability of DIC, which relies on the instantaneous spatial distribution of SST and a linear correlation between DIC and SST, is relatively independent from the variability estimated using the high-pass filtered time-series sampled by the drifters, the only link between the two being the linear correlation based on the time-series. This second estimate given by $SV100\text{km}_{sat}$ (5 to $15 \mu\text{mol kg}^{-1}$) is very similar to the $SV100\text{km}$ estimated from the time-series in this area (5 to $12 \mu\text{mol kg}^{-1}$ on Fig. 2j).

To strengthen the results obtained with the $SV100\text{km}$ derived from time-series, this second estimate is computed using all SST satellite images co-located with drifters trajectories available when the correlation coefficient r^2 exceeds 0.7 (Fig. 3c). This estimate confirms the order of magnitude of DIC variability at small spatial scales obtained from the time-series (Fig. 3b), with typical gradients of 2 – $10 \mu\text{mol kg}^{-1}$, increasing to 10 – $15 \mu\text{mol kg}^{-1}$ in frontal areas and up to 15 – $30 \mu\text{mol kg}^{-1}$ in regions of high eddy kinetic energy or where biological activity was identified as a driving mechanism of DIC variability (Figs. 3b, c and 5d).

3.4 Does small-scale matters compared to large-scale and seasonal variability?

This variability of DIC identified at small spatial scales can be compared to the variability at larger spatial and temporal scales. Large-scale regional and latitudinal contrasts highlighted by the Glodap database primarily arise from the thermal effect on CO₂ solubility, the Revelle factor and the large scale circulation (Fig. 1e, Williams and Follows, 2011). DIC temporal variability is dominated by the seasonality arising from the deepening of the mixed layer (ML) in winter and the biological uptake of DIC in summer. The extensive data set presented here captured the variability introduced by these various spatial and temporal scales (colored lines on Fig. 8). In the previous sections, the variability arising from small-scale structures was identified by filtering out temporal scales longer than 20 days and spatial scales larger than 500–1000 km (see Sect. 3.1.2). It is however relatively difficult to disentangle seasonal variations from large-scale regional variations along the drifters trajectories.

To estimate the variability introduced by large-scale regional contrasts in drifters measurements, we used the Glodap annual value interpolated along the drifter trajectory (light grey on Fig. 8). Interestingly, these time-series reconstructed from the interpolation of Glodap DIC at the drifter location reproduce a large part of the variations captured by the drifters, suggesting that large scale regional contrasts accounts for a large part of the low-frequency variability sampled by the drifters (Fig. 8). In particular, we note that the large-scale gradients are well resolved by Glodap along drifters trajectories in the Atlantic and Indian sectors, where more observations are available (Fig. 8a–d and i). From this reconstructed time-series we estimate large-scale regional gradients to be of the order of ~ 50 to $100 \mu\text{mol kg}^{-1}$ over spatial scales of ~ 5000 – $10\,000$ km (i.e. temporal scales of ~ 6 months along the drifter trajectories), which is much smaller than the variability estimated at small scale in this study (2 – $30 \mu\text{mol kg}^{-1}$ over 100 km).

To date, no climatology describe the seasonal cycle of DIC. To reconstruct the seasonal cycle of surface DIC, we used the 3-D Glodap annual values and a monthly clima-

BGD

10, 13855–13895, 2013

Carbon variability in Southern Ocean

L. Resplandy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 tology of mixed layer depth derived from Argo drifters (Sallée et al., 2010) (Fig. 9a and b). As in Mahadevan et al. (2011), the seasonality is estimated by averaging the Glodap values within the mixed layer. In winter, the ML deepens and surface DIC concentrations increase when averaged with high-DIC subsurface concentrations. In summer, 10 restratification leads to lower estimates of surface DIC concentrations due to lower mixing with subsurface waters, hence mimicking the biological uptake of DIC. Two estimates of the seasonal cycles are obtained: the first one uses the mean value of the mixed layer depth in the $1^\circ \times 1^\circ$ box around the drifter sample (black lines on Fig. 8), whereas the second uses the maximum value of the mixed layer in that same box 15 (dark grey lines on Fig. 8). The comparison between the two gives an estimation of the method's sensitivity to interannual and small-scale variability of the ML. This reconstruction assumes that: (1) on annual scale, summertime biological uptake of DIC is counterbalanced by wintertime physical input of DIC associated with the deepening of the ML; (2) the biological uptake of DIC is synchronous with the ML restratification and (3) Glodap values correspond to well-stratified low-DIC summer conditions. We expect hypotheses 1 and 3 to be relatively valid in the Southern Ocean because of the predominant effect of winter mixing at high latitudes and because most of the profiles used to derive Glodap were sampled between October and April in this region (see Appendix Fig. A4). In contrast, hypothesis 2 may introduce lags in DIC seasonality.

20 The reconstructed seasonal DIC interpolated along the drifters trajectories captures most of the low frequency signal observed in-situ by the drifters (dark grey lines vs. colored lines on Fig. 8), which suggests that Glodap spatial variability and seasonal variability reconstructed from the ML climatology give a useful estimate of the regional patterns and seasonality of the DIC in the Southern Ocean. Note that part of the signal 25 captured by the drifters however remains unexplained, most probably because of the interannual and small-scale variability, the assumptions made to compute the seasonal cycle (see 3 hypotheses above) and the biases introduced by undersampling in the ML climatology and the Glodap database. For example, largest DIC differences (up to $25\text{--}50\ \mu\text{mol kg}^{-1}$) between drifters measurements and the reconstructed time-series are

**Carbon variability in
Southern Ocean**

L. Resplandy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



observed in the Pacific sector and South of Tasmania (drifters 1110 and 3739) that are poorly sampled by Glodap (Fig. 8). When applied to the whole Southern Ocean, this estimate gives an amplitude of DIC seasonal variations (winter max – summer min) of the order of 10–50 $\mu\text{mol kg}^{-1}$ (Fig. 9c). Our estimate is in agreement with the seasonality observed in the Indian sector during the OISO cruises (60° E, 50–55° S), where it amounts to $\sim 10\text{--}30 \mu\text{mol kg}^{-1}$ (Metzl et al., 2006) and in the Pacific sector at the Munida time-series south of New Zealand (171° E, 46° S), where it reaches 40 $\mu\text{mol kg}^{-1}$ (Brix et al., 2013).

4 Discussion and conclusions

In this study, the $p\text{CO}_2$ and DIC variability at small spatial scale in the Southern Ocean is estimated using an extensive dataset of surface drifters measurements. Typical gradients on spatial scales of 100 km range from 5 to 50 μatm for $p\text{CO}_2$ and 2 to 30 $\mu\text{mol kg}^{-1}$ for DIC. The result of DIC variability is supported by a second estimate based on SST satellite images and DIC/SST relationships at spatial scales of ~ 100 km derived from drifters observations. Regions with highest variability of the CO_2 system mostly correspond to regions of high eddy kinetic energy or frontal zones. These estimates of the CO_2 system variability at small-scale are comparable to estimates obtained in the tropical Pacific (up to 30 μatm on scales of 50 km, Chen et al., 2007) northeast Atlantic (up to 25 μatm on scales of 25 km, Resplandy et al., 2009).

Using a principal component analysis we find that the DIC variability at small-scale mostly arises from ocean dynamics. Although the circulation is identified as the first mechanism generating variability at small spatial scales, it results in relatively low variability with typical gradients of $\sim 2\text{--}10 \mu\text{mol kg}^{-1}$ over scales of 100 km, reaching values up to 10–15 $\mu\text{mol kg}^{-1}$ in regions of fronts and high eddy kinetic energy. In contrast, when biological activity is identified as a key process in DIC distribution at small spatial scale, gradients are enhanced with values up to 10–30 $\mu\text{mol kg}^{-1}$ over scales of 100 km. This is in agreement with previous in-situ observations that highlighted higher

BGD

10, 13855–13895, 2013

Carbon variability in Southern Ocean

L. Resplandy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



spatial variability of biogeochemical parameters when primary productivity is enhanced (Metzl et al., 2006; Rangama et al., 2005).

Modeling studies showed that both horizontal and vertical advection associated with small-scale structures can affect the distribution of DIC at small spatial scales and hence the surface $p\text{CO}_2$ (Mahadevan et al., 2004; Resplandy et al., 2009). Horizontally, the ocean circulation advects large-scale gradients down to small spatial scales. It is now widely accepted that mesoscale eddies, which dominate the variability at scales of the order of 100 km, horizontally transport and stir tracers such as DIC and temperature, resulting in a cascade of tracer variance to smaller scales (Abraham et al., 2000; Lévy and Klein, 2004). In addition, density gradients associated with small-scale features are associated with large vertical velocities (Capet et al., 2008; Lévy et al., 2012), that could transport deeper waters of high DIC concentration to the surface. It is generally very difficult to assess the contribution of vertical advection purely from observations. More particularly, the data set used here does not allow to discriminate between the contribution of horizontal and vertical advection. First, large-scale horizontal and vertical gradients in the Southern Ocean are associated with a similar signature of increasing DIC concentration with decreasing temperature that can not be separated with the principal component analysis. In addition, our observations only describe surface waters and provide no information on vertical gradients. Nevertheless, the fact that dynamical advection alone results in relatively low variability in DIC except in zones of strong horizontal gradients (frontal zones) strongly suggests that the horizontal component of advection plays a major role in setting the variability at small spatial scales. The impact of episodic vertical mixing events, such as eddies or storms, on surface $p\text{CO}_2$ has been estimated in Mahadevan et al. (2011) using global monthly climatological datasets. They found that vertical mixing events could induce changes in DIC through the vertical input of DIC-rich subsurface waters and the enhancement of biological uptake. In agreement with previous studies based on observations (Chen et al., 2007) and models (Mahadevan et al., 2004; Resplandy et al., 2009), they found a rel-

BGD

10, 13855–13895, 2013

Carbon variability in Southern Ocean

L. Resplandy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



atively low impact on $p\text{CO}_2$ (< 5%) due to compensations between biological uptake, DIC input and thermal effects.

This study confirms that seasonal variations are the dominant mode of variability of the CO_2 system in the Southern Ocean (Lenton et al., 2006). However, we find that small spatial scales structures are a non negligible source of variability for DIC, with amplitudes of about a third of the variations associated with the seasonality (2–30 $\mu\text{mol kg}^{-1}$ vs. 10–50 $\mu\text{mol kg}^{-1}$). In addition, the amplitude of this variability is at least of the same order of magnitude as large-scale gradients and can be up to ~ 10 times larger in regions of high eddy kinetic energy and biological activity. These small scale processes are not captured by on-board observations and climate models used to derive carbon budgets. This issue combines with the fact that seasonal cycles of $p\text{CO}_2$ and DIC in the Southern Ocean are still poorly described due to the lack of shipboard observations mainly during winter (Metzl et al., 2006). This has strong implication on our understanding of the carbon budget derived from observations (Barbero et al., 2011) but also from model simulations (Lenton et al., 2013), whose evaluation rely on these observations. Direct measurements being scarce, an alternative is to derive DIC from observed $p\text{CO}_2$, temperature and salinity fields. Although numerous observations of surface $p\text{CO}_2$ are available (Takahashi et al., 2009; Pfeil et al., 2012), the amount of DIC observations that can be derived is drastically lower. In Lenton et al. (2012), 75 000 surface DIC estimates could be computed from the 4.4 million individual measurements of the global database of Takahashi et al. (2009) after removing non collocated measurements of $p\text{CO}_2$, SST and SSS and coastal data. Among these data, only few are located in the Southern Ocean, thus limiting the assessment of the seasonal cycle and small-scale variability of DIC.

In this work, the seasonal cycle of surface DIC is reconstructed from the annual Glodap data set and a mixed layer climatology. This methodology has been previously used in Mahadevan et al. (2011) at global scale but was not validated. Here, we provide a validation of this method in the Southern Ocean using the months- to year-long high frequency time-series of the nine drifters deployed in the area. The fair agreement

BGD

10, 13855–13895, 2013

Carbon variability in Southern Ocean

L. Resplandy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Carbon variability in Southern Ocean

L. Resplandy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



between in-situ observations and the reconstructed seasonal DIC suggests that this method gives a first-order estimates of the seasonal amplitude of surface DIC in the Southern Ocean that could be used for model evaluation. However, it should be kept in mind that the ability of this method is lower in poorly sampled regions, such as the Pacific sector and that it provides stronger constraint on the timing of wintertime DIC enrichment than on the timing of summertime DIC consumption. Despite the limitations of the method, our finding provide a first step to better understand the contribution and amplitude of variability introduced by small-scale structures versus the variability introduced by large-scale contrasts, seasonality and inter-annual to decadal variability. This study suggests that care should be taken when inferring temporal changes (seasonality, inter-annual variability, decadal trends) from sparse observations. Indeed, these observations and the methods used don't take into account the variability at small-spatial scales, while our results highlight that the various spatial and temporal sources of variability can be of the same order of magnitude in some regions of the Southern Ocean.

Appendix A

Principal Component Analysis loadings of PC1, PC2 and PC3 obtained with DIC, SST and fluorescence (Figs. A1–A3). The loadings are shown only when the PC relates strongly to the variations of DIC and when it explains a significant part of the variance at small-scale (i.e. when the carbon variable loading on this PC is ≥ 0.3 and the explained variance of this PC is $\geq 20\%$, see Sect. 2.3).

Acknowledgements. Support was provided by the European Integrated project CARBOOCEAN and EU FP7 project CARBOCHANGE “Changes in carbon uptake and emissions by oceans in a changing climate”, which received funding from the European Commissionâ Seventh Framework Programme under grant agreement no. 264879. We thank L. Beaumont and T. Danguy from technical division of Institut National des Sciences de l'Univers for supervising CARIOCA assembly and performing thorough CARIOCA calibrations. We

sincerely thank Francesco d'Ovidio for providing and maintaining the LAGRANGIAN Manifolds and Trajectories Analyser (Lamta) toolbox. Altimeter products were produced by Ssalto/Duacs and distributed by AVISO with support from the Centre National d'Études Spatiales.



The publication of this article
is financed by CNRS-INSU.

References

- Abraham, E. R., Law, C. S., Boyd, P. W., Lavender, S. J., Maldonado, M. T., and Bowie, A. R.: Importance of stirring in the development of an iron-fertilized phytoplankton bloom, *Nature*, 407, 727–730, doi:10.1038/35037555, 2000. 13874
- Barbero, L., Boutin, J., Merlivat, L., Martin, N., Takahashi, T., Sutherland, S. C., and Wanninkhof, R.: Importance of water mass formation regions for the air-sea CO₂ flux estimate in the Southern Ocean, *Global Biogeochem. Cy.*, 25, GB1005, doi:10.1029/2010GB003818, 2011. 13857, 13865, 13869, 13875
- Boutin, J. and Merlivat, L.: New in situ estimates of carbon biological production rates in the Southern Ocean from CARIOCA drifter measurements, *Geophys. Res. Lett.*, 36, L13608, doi:10.1029/2009GL038307, 2009. 13859, 13869
- Boutin, J., Merlivat, L., Hénocq, C., Martin, N., and Sallée, J. B.: Air-sea CO₂ flux variability in frontal regions of the Southern Ocean from CARIOCA drifters, *Limnol. Oceanogr.*, 53, 2062–2079, 2008. 13857, 13859
- Brix, H., Currie, K. I., and Mikaloff Fletcher, S. E.: Seasonal variability of the carbon cycle in subantarctic surface water in the South West Pacific, *Global Biogeochem. Cy.*, 27, 200–211, doi:10.1002/gbc.20023., 2013. 13858, 13862, 13873
- Capet, X., Colas, F., Penven, P., Marchesiello, P., and McWilliams, J. C.: Eastern boundary subtropical upwelling systems, in: *Eddy Resolving Ocean Modeling*, vol. 117, edited by: Hecht, M. W. and Hasumi, H., Geophysical Monograph Series, Washington DC, 131–147, 2008. 13874

Carbon variability in Southern Ocean

L. Resplandy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Carbon variability in Southern Ocean

L. Resplandy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Chen, F., Cai, W.-J., Benitez-Nelson, C., and Wang, Y.: Sea surface “pco₂-sst” relationships across a cold-core cyclonic eddy: implications for understanding regional variability and air–sea gas exchange, *Geophys. Res. Lett.*, 34, L10603, doi:10.1029/2006GL028058, 2007. 13861, 13873, 13874

5 Copin-Montégut, C., Begovic, M., and Merlivat, L.: Variability of the partial pressure of CO₂ on diel to annual time scales in the northwestern Mediterranean Sea, *Sci. Mar.*, 85, 169–189, doi:10.1016/j.marchem.2003.10.005, 2004. 13859

Dandonneau, Y.: Sea-surface partial pressure of carbon dioxide in the eastern equatorial Pacific (August 1991 to October 1992): a multivariate analysis of physical and biological factors, *Deep-Sea Res. Pt. II*, 42, 349–364, doi:10.1016/0967-0645(95)00017-K, 1995. 13862

10 d’Ovidio, F., Fernández, V., Hernández-García, E., and López, C.: Mixing structures in the Mediterranean Sea from finite-size Lyapunov exponents, *Geophys. Res. Lett.*, 31, L17203, doi:10.1029/2004GL020328, 2004. 13860, 13889

15 Gruber, N., Gloor, M., Fletcher, S. E. M., Doney, S. C., Dutkiewicz, S., Follows, M. J., Gerber, M., Jacobson, A. R., Joos, F., Lindsay, K., Menemenlis, D., Mouchet, A., Müller, S. A., Sarmiento, J. L., and Takahashi, T.: Oceanic sources, sinks, and transport of atmospheric CO₂, *Global Biogeochem. Cy.*, 23, GB1005, doi:10.1029/2008GB003349, 2009. 13857, 13858

Ito, T., Marshall, J., and Follows, M.: What controls the uptake of transient tracers in the Southern Ocean?, *Global Biogeochem. Cy.*, 18, GB2021, doi:10.1029/2003GB002103, 2004. 13858

20 Key, R. M., Kozyr, A., Sabine, C. L., Lee, K., Wanninkhof, R., Bullister, J. L., Feely, R. A., Millero, F. J., Mordy, C., and Peng, T.-H.: A global ocean carbon climatology: results from Global Data Analysis Project (GLODAP), *Global Biogeochem. Cy.*, 18, GB4031, doi:10.1029/2004GB002247, 2004. 13860

25 Lee, K., Tong, L. T., Millero, F. J., Sabine, C. L., Dickson, A. G., Goyet, C., Park, G.-H., Wanninkhof, R., Feely, R. A., and Key, R. M.: Global relationships of total alkalinity with salinity and temperature in surface waters of the world’s oceans, *Geophys. Res. Lett.*, 33, L19605, doi:10.1029/2006GL027207, 2006. 13859

30 Lehahn, Y., d’Ovidio, F., Lévy, M., and Heitzel, E.: Stirring of the Northeast Atlantic spring bloom: a lagrangian analysis based on multi-satellite data, *J. Geophys. Res.*, 112, C08005, doi:10.1029/2006JC003927, 2007. 13860

Carbon variability in
Southern Ocean

L. Resplandy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Lenton, A., Matear, R. J., and Tilbrook, B.: Design of an observational strategy for quantifying the Southern Ocean uptake of CO₂, *Global Biogeochem. Cy.*, 20, GB4010, doi:10.1029/2005GB002620, 2006. 13875
- 5 Lenton, A., Metzl, N., Takahashi, T., Kuchinke, M., Matear, R. J., Roy, T., Sutherland, S. C., Sweeney, C., and Tilbrook, B.: The observed evolution of oceanic pCO₂ and its drivers over the last two decades, *Global Biogeochem. Cy.*, 26, GB2021, doi:10.1029/2011GB004095, 2012. 13875
- 10 Lenton, A., Tilbrook, B., Law, R. M., Bakker, D., Doney, S. C., Gruber, N., Ishii, M., Hoppema, M., Lovenduski, N. S., Matear, R. J., McNeil, B. I., Metzl, N., Mikaloff Fletcher, S. E., Monteiro, P. M. S., Rödenbeck, C., Sweeney, C., and Takahashi, T.: Sea–air CO₂ fluxes in the Southern Ocean for the period 1990–2009, *Biogeosciences*, 10, 4037–4054, doi:10.5194/bg-10-4037-2013, 2013. 13857, 13875
- 15 Lévy, M. and Klein, P.: Does the low frequency variability of mesoscale dynamics explain a part of the phytoplankton and zooplankton spectral variability?, *P. R. Soc. London*, 460, 1673–1683, doi:10.1098/rspa.2003.1219, 2004. 13874
- Lévy, M., Ferrari, R., Franks, P., Martin, A., and Riviere, P.: Bringing physics to life at the submesoscale, *Geophys. Res. Lett.*, 39, L14602, doi:10.1029/2012GL052756, 2012. 13874
- 20 Lohrenz, S. E., Cai, W., Chen, F., Chen, X., and Tuel, M.: Seasonal variability in air–sea fluxes of CO₂ in a river influenced coastal margin, *J. Geophys. Res.*, 115, C10034, doi:10.1029/2009JC005608, 2009. 13862
- Lovenduski, N. S., Gruber, N., and Doney, S. C.: Toward a mechanistic understanding of the decadal trends in the Southern Ocean carbon sink, *Global Biogeochem. Cy.*, 22, GB3016, doi:10.1029/2007GB003139, 2008. 13857, 13862
- 25 Mahadevan, A., Lévy, M., and Mémerly, L.: Mesoscale variability of sea surface pCO₂: What does it respond to?, *Global Biogeochem. Cy.*, 18, GB1017, doi:10.1029/2003GB002102, 2004. 13874
- Mahadevan, A., Tagliabue, A., Bopp, L., Lenton, A., Mémerly, L., and Lévy, M.: Impact of episodic vertical fluxes on sea surface pCO₂, *Philos. T. R. Soc. A*, 369, 2009–2025, doi:10.1098/rsta.2010.0340, 2011. 13856, 13872, 13874, 13875
- 30 McNeil, B. I., Metzl, N., Key, R. M., Matear, R. J., and Corbiere, A.: An empirical estimate of the southern ocean air–sea CO₂ flux, *Global Biogeochem. Cy.*, 21, GB3011, doi:10.1029/2007GB002991, 2007. 13857

Carbon variability in Southern Ocean

L. Resplandy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Metzl, N.: Decadal increase of oceanic carbon dioxide in Southern Indian Ocean surface waters (1991–2007), *Deep-Sea Res. Pt. II*, 56, 607–619, doi:10.1016/j.dsr2.2008.12.007, 2009. 13858, 13862

Metzl, N., Brunet, C., Jabaud-Jan, A., Poisson, A., and Schauer, B.: Summer and winter air–sea $\{CO_2\}$ fluxes in the Southern Ocean, *Deep-Sea Res. Pt. I*, 53, 1548–1563, doi:10.1016/j.dsr.2006.07.006, 2006. 13858, 13862, 13873, 13874, 13875

Mikaloff Fletcher, S. E., Gruber, N., Jacobson, A. R., Gloor, M., Doney, S. C., Dutkiewicz, S., Gerber, M., Follows, M., Joos, F., Lindsay, K., Menemenlis, D., Mouchet, A., Müller, S. A., and Sarmiento, J. L.: Inverse estimates of anthropogenic CO_2 uptake, transport, and storage by the ocean, *Global Biogeochem. Cy.*, 20, GB2002, doi:10.1029/2005GB002530, 2006. 13857

Murata, A.: Increased surface seawater pCO_2 in the eastern Bering Sea shelf: an effect of blooms of *coccolithophorid Emiliania huxleyi*?, *Global Biogeochem. Cy.*, 20, GB4006, doi:10.1029/2005GB002615, 2006. 13862

Orr, J. C., Maier-Reimer, E., Mikolajewicz, U., Monfray, P., Sarmiento, J. L., Toggweiler, J. R., Taylor, N. K., Palmer, J., Gruber, N., Sabine, C. L., Le Quéré, C., Key, R. M., and Boutin, J.: Estimates of anthropogenic carbon uptake from four three-dimensional global ocean models, *Global Biogeochem. Cy.*, 15, 43–60, doi:10.1029/2000GB001273, 2001. 13857

Pfeil, B., Olsen, A., Bakker, D. C. E., Hankin, S., Koyuk, H., Kozyr, A., Malczyk, J., Manke, A., Metzl, N., Sabine, C. L., Akl, J., Alin, S. R., Bates, N., Bellerby, R. G. J., Borges, A., Boutin, J., Brown, P. J., Cai, W.-J., Chavez, F. P., Chen, A., Cosca, C., Fassbender, A. J., Feely, R. A., González-Dávila, M., Goyet, C., Hales, B., Hardman-Mountford, N., Heinze, C., Hood, M., Hoppema, M., Hunt, C. W., Hydes, D., Ishii, M., Johannessen, T., Jones, S. D., Key, R. M., Körtzinger, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lenton, A., Lourantou, A., Merlivat, L., Midorikawa, T., Mintrop, L., Miyazaki, C., Murata, A., Nakadate, A., Nakano, Y., Nakaoka, S., Nojiri, Y., Omar, A. M., Padin, X. A., Park, G.-H., Paterson, K., Perez, F. F., Pierrot, D., Poisson, A., Ríos, A. F., Santana-Casiano, J. M., Salisbury, J., Sarma, V. V. S. S., Schlitzer, R., Schneider, B., Schuster, U., Sieger, R., Skjelvan, I., Steinhoff, T., Suzuki, T., Takahashi, T., Tedesco, K., Telszewski, M., Thomas, H., Tilbrook, B., Tjiputra, J., Vandemark, D., Veness, T., Wanninkhof, R., Watson, A. J., Weiss, R., Wong, C. S., and Yoshikawa-Inoue, H.: A uniform, quality controlled Surface Ocean CO_2 Atlas (SOCAT), *Earth Syst. Sci. Data*, 5, 125–143, doi:10.5194/essd-5-125-2013, 2013. 13858, 13875

Carbon variability in
Southern Ocean

L. Resplandy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Pollard, T., R., Venables, H. J., Read, J. F., and Allen, J. T.: Large-scale circulation around the Crozet Plateau controls an annual phytoplankton bloom in the Crozet Basin, *Deep-Sea Res. Pt. II*, 54, 1915–1929, doi:10.1016/j.dsr2.2007.06.012, 2007. 13864
- Rangama, Y., Boutin, J., Etcheto, J., Merlivat, L., Takahashi, T., Delille, B., Frankignoulle, M., and Bakker, D. C. E.: Variability of the net air–sea CO₂ flux inferred from shipboard and satellite measurements in the Southern Ocean south of Tasmania and New Zealand, *J. Geophys. Res.*, 110, C09005, doi:10.1029/2004JC002619, 2005. 13874
- Read, J., Pollard, R., and Allen, J.: Sub-mesoscale structure and the development of an eddy in the subantarctic front north of the Crozet Islands, *Deep-Sea Res. Pt. II*, 54, 1930–1948, doi:10.1016/j.dsr2.2007.06.013, 2007. 13858
- Resplandy, L., Lévy, M., d'Ovidio, F., and Merlivat, L.: Impact of submesoscale variability in estimating the air–sea CO₂ exchange: results from a model study of the POMME experiment, *Global Biogeochem. Cy.*, 23, GB1017, doi:10.1029/2008GB003239, 2009. 13873, 13874
- Rintoul, S. R., Hughes, C., and Olbers, D.: The Antarctic circumpolar current system, in: *Ocean Circulation and Climate*, edited by: Siedler, G., Church, J., and Gould, J., Academic Press, London, 271–302, 2001. 13858
- Sallée, B. J., Speer, K., and Rintoul, S. R.: Zonally asymmetric response of the Southern Ocean mixed-layer depth to the Southern Annular Mode, *Nat. Geosci.*, 3, 273–279, 2010. 13872, 13891
- Séférian, R., Bopp, L., Swingedouw, D., and Servonnat, J.: Dynamical and biogeochemical control on the decadal variability of ocean carbon fluxes, *Earth Syst. Dynam.*, 4, 109–127, doi:10.5194/esd-4-109-2013, 2013. 13858, 13862
- Takahashia, T., Sutherland, S. C., Wanninkhof, R., Sweeney, C., Feely, R. A., Chipman, D. W., Hales, B., Friederich, G., Chavez, F., Sabine, C., Watson, A., Bakker, D. C. E., Schuster, U., Metzl, N., Yoshikawa-Inoue, H., Ishii, M., Midorikawa, T., Nojiri, Y., Körtzinger, A., Steinhoff, T., Hoppema, M., Olafsson, J., Arnarson, T. S., Tilbrook, B., Johannessen, T., Olsen, A., Bellerby, R., Wong, C. S., Delille, B., Bates, N. R., and de Baar, H. J. W.: Climatological mean and decadal change in surface ocean *p*CO₂, and net sea-air CO₂ flux over the global oceans, *Deep-Sea Res- Pt. II*, 56, 554–577, doi:10.1016/j.dsr2.2008.12.009, 2009. 13857, 13858, 13859, 13875
- Tortell, P. D., Guéguen, C., Long, M. C., Payne, C. D., Lee, P., and DiTullio, G. R.: Spatial variability and temporal dynamics of surface water *p*CO₂, O₂/Ar and dimethylsulfide in the Ross

Sea, Antarctica, Deep-Sea Res. Pt. I, 58, 241–259, doi:10.1016/j.dsr.2010.12.006, 2011. 13858

Venables, H. J., Pollard, T. R., and Popova, E. E.: Physical conditions controlling the development of a regular phytoplankton bloom north of the Crozet Plateau, Southern Ocean, Deep-Sea Res. Pt. II, 54, 1949–1965, doi:10.1016/j.dsr2.2007.06.014, 2007. 13864

Williams, R. G. and Follows, M. J.: Ocean Dynamics and the Carbon Cycle, Cambridge University Press, 2011. 13871

BGD

10, 13855–13895, 2013

Carbon variability in Southern Ocean

L. Resplandy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 1. (a) drifters trajectories ; (b) drifters sampling month; (c) drifters $p\text{CO}_2$ (in μatm); (d) drifters DIC (in μmolkg^{-1}) and (e) DIC in Glodap database (in μmolkg^{-1}). A correction of $0.7 \mu\text{molkg}^{-1}\text{yr}^{-1}$ has been applied to reference all DIC observations to year 2005.




Fig. 2. Case study: drifter 13060 in the Crozet Plateau area. **(a, c, e, g)** surface $p\text{CO}_2$, SST, fluorescence and DIC; **(b, d, f, h)** 20-day high-pass filtered surface $p\text{CO}_2$, SST, fluorescence and DIC; **(i–j)** SV100km quantifying small scale spatial variability of $p\text{CO}_2$ and DIC.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)





Fig. 3. Eddy kinetic energy (EKE, grey scale) estimated from satellite altimetry and mesoscale variability **(a)** SV100km of in-situ $p\text{CO}_2$ (in μatm) and **(b)** SV100km of in-situ DIC (in $\mu\text{mol kg}^{-1}$) estimated from the standard deviation of high-pass filtered fields in $100\text{ km} \times 100\text{ km}$ boxes along drifters trajectories (see Sect. 3.1.2 for further details), **(c)** SV100km_{sat} of DIC (in $\mu\text{mol kg}^{-1}$) reconstructed from satellite SST and the local linear correlations between SST and DIC deduced from drifters data. DIC was reconstructed only for correlation coefficient $r^2 > 0$. (Fig. 6), hence explaining the lower spatial coverage of panel **(c)** compared to panel **(b)**. EKE was estimated in 2006 from satellite altimetry as $(u - \bar{u})^2 + (v - \bar{v})^2$, with u and v the zonal and meridional surface current velocities and where the overbar denotes the annual mean.




Fig. 4. Principal component analysis in the case study area: explained variance by (a) PC1 and (b) PC2; DIC loadings for (c) PC1 and (d) PC2; SST loadings for (e) PC1 and (f) PC2; fluorescence loadings for (g) PC1 and (h) PC2 and (i) dominant processes influencing DIC identified using PC1 loadings: ocean dynamics (DYN), biological uptake of DIC (BIO) and others (NO).

Carbon variability in
Southern Ocean

L. Resplandy et al.




Fig. 5. “Local” PCA for DIC, SST and fluorescence. **(a–c)** Variance explained by **(a)** PC1, **(b)** PC2 and **(c)** the cumulated variance explained by PC1 and PC2. **(d–e)** Dominant processes influencing the variability at small spatial scales: ocean dynamics (DYN), biological uptake of DIC (BIO) and others (NO) were identified using **(d)** PC1 and **(e)** PC2 loadings.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Carbon variability in
Southern Ocean

L. Resplandy et al.




Fig. 6. Correlation coefficient (r^2) for the linear correlation between **(a)** SST and DIC; **(b)** SST and $p\text{CO}_2$.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Carbon variability in Southern Ocean

L. Resplandy et al.




Fig. 7. Case study: **(a)** SST measured by drifter 13 060 between 18 March and 7 April 2007; **(b)** satellite AMSR SST collocated in time and space with the drifter; **(c)** DIC measured by drifter 13 060; **(d)** satellite AMSR SST on 28 March 2007; **(e)** AMSR \widehat{SST}_{sat} on 28 March 2007 and **(f)** \widehat{DIC}_{sat} on 28 March 2007 reconstructed from \widehat{SST}_{sat} and a DIC/SST linear relationship derived from drifter measurements ($\widehat{DIC} = -9.69 \cdot \widehat{SST} + 0.09$, with $r^2 > 0.86$). Dynamical boundaries derived from altimetry, i.e. unstable manifolds are indicated in black (d’Ovidio et al., 2004). The drifter trajectory (magenta) is wrapped around two eddies, noted E1 and E2. Grey contours indicate the bathymetry.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 8. DIC along Carioca drifters trajectories: DIC deduced from the $p\text{CO}_2$ sampled by the drifters (colored lines) is compared to the “annual Glodap DIC” and two estimates of the “seasonal Glodap DIC” along the drifters location. The “annual Glodap DIC” (light grey) shows the large scale gradients of surface DIC in Glodap in a 5° by 5° box centered on the drifter location. The two “seasonal Glodap DIC” estimates were computed by averaging the vertical profile of Glodap DIC over monthly mixed layer depths. The “seasonal Glodap DIC mean” and “seasonal Glodap DIC max” respectively correspond to the mean and maximum values obtained in each 5° by 5° boxes centered on the drifters trajectories (see Sect. 3.4), which gives an estimation of the seasonal signal on surface DIC concentrations.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)





Fig. 9. (a) Vertical gradient of DIC between the surface and 200 m in Glodap database; (b) seasonal amplitude (annual max–annual min) of ML depth in the climatology derived from Argo drifters of Sallée et al. (2010); (c) DIC seasonal amplitude (annual max–annual min) reconstructed by averaging Glodap annual values over the ML monthly climatology. See text Sect. 3.4 for further details.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Carbon variability in
Southern Ocean

L. Resplandy et al.

**Fig. A1.** PC1 loadings for DIC, SST and fluorescence.[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Carbon variability in Southern Ocean

L. Resplandy et al.




Fig. A2. PC2 loadings for DIC, SST and fluorescence.

[Title Page](#)

[Abstract](#) [Introduction](#)

[Conclusions](#) [References](#)

[Tables](#) [Figures](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

[Back](#) [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)






Fig. A3. PC3 loadings for DIC, SST and fluorescence.

[Title Page](#)

Abstract	Introduction
Conclusions	References
Tables	Figures

⏪	⏩
◀	▶
Back	Close

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Carbon variability in Southern Ocean

L. Resplandy et al.




Fig. A4. Distribution and sampling month of DIC used to derive the 3-D-gridded Glodap product in the Southern Ocean.

[Title Page](#)

[Abstract](#) [Introduction](#)

[Conclusions](#) [References](#)

[Tables](#) [Figures](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

[Back](#) [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

