

Table S1: Full summary of Earth system models including model names, history of development, number of soil/litter pools, temperature/moisture functions, link to nitrogen cycling, the CMIP5 variables used to construct total soil carbon, and the years analyzed for each CMIP5 experiment. (adapted from Todd-Brown et al., 2013)

Model Center	Model name	Soil model history	Litter	Soil	Temperature	Moisture	Nitrogen	Reported CMIP5 variables			Years considered for analysis	
								cSoil	cLitter	cCwd	Historical	RCP 8.5
Beijing Climate Center, China Meteorological Administration (BCC)	BCC-CSM1.1 BCC-CSM1.1(m) (Wu et al., submitted)	AVIM2 (Huang <i>et al.</i> , 2007; Ji <i>et al.</i> , 2008) CEVSA (Cao & Woodward, 1998) CENTURY (Parton <i>et al.</i> , 1987, 1993)	2	6	hill <sup>a</sup>	hill	yes	X	X	--	1850-2005	2006-2099
College of Global Change and Earth System Science, Beijing Normal University (GCESS)	BNU-ESM (Personal communication Qian Zhang, 2013)	CoLM (Dai <i>et al.</i> , 2003, 2004) LPJ-DyN (Xu-Ri & Prentice, 2008) LPJ-DGVM (Sitch <i>et al.</i> , 2003) DEMETER-1 (Foley, 1995) CENTURY (Parton <i>et al.</i> , 1987, 1992)	2	2	Arrhenius	increasing	no <sup>d</sup>	X	X	--	1850-2005	2006-2100
Canadian Centre for Climate Modeling and Analysis (CCCMA)	CanESM2 (CMIP5 output)	CTEM1 (Arora, 2003; Arora & Boer, 2005; Arora <i>et al.</i> , 2011)	1 <sup>b</sup>	1 <sup>b</sup>	Q10 <sup>c</sup>	hill	no	X	X	--	1850-2005	2006-2100
National Center for Atmospheric Research (NCAR)	CCSM4 (Gent <i>et al.</i> , 2011)	CLM4 (Oleson <i>et al.</i> , 2008) CN (Thornton <i>et al.</i> , 2007) Biome-BCG 4.1.2 (Kimball <i>et al.</i> , 1997; Thornton, 1998; Thornton <i>et al.</i> , 2002; Thornton & Rosenbloom, 2005) (Olson, 1963; Veen &	4	3	Arrhenius	increasing	yes	X	X	X	1850-2005	2006-2100

Model Center	Model name	Soil model history	Litter	Soil	Temperature	Moisture	Nitrogen	Reported CMIP5 variables			Years considered for analysis	
								cSoil	cLitter	cCwd	Historical	RCP 8.5
		Paul, 1981; Veen <i>et al.</i> , 1984)										
Community Earth System Model Contributors (NSF-DOE-NCAR)	CESM1(BGC) CESM1(CAM5) CESM1(FASTCHEM) CESM1(WACCM) (Keppel-Aleks <i>et al.</i> , 2013; Lindsay <i>et al.</i> , 2013)	CLM4(Oleson <i>et al.</i> , 2008) CN (Thornton <i>et al.</i> , 2007) Biome-BCG 4.1.2 (Kimbball <i>et al.</i> , 1997; Thornton, 1998; Thornton <i>et al.</i> , 2002; Thornton & Rosenbloom, 2005) (Olson, 1963; Veen & Paul, 1981; Veen <i>et al.</i> , 1984)	4	3	Arrhenius	increasing	yes	X	X	X	1850-2005	2006-2100
NOAA Geophysical Fluid Dynamics Laboratory (NOAA GFDL)	GFDL-ESM2G GFDL-ESM2M (CMIP5 output)	LM3 (LM3p7_cESM, M45) (Shevliakova <i>et al.</i> , 2009) ED (Moorcroft <i>et al.</i> , 2001) (Bolker <i>et al.</i> , 1998) CENTURY (Parton <i>et al.</i> , 1987)	--	2	hill	increasing	no	X	--	--	1861-2005	2006-2100
Met Office Hadley Centre (additional HadGEM2-ES realizations contributed by Instituto Nacional de Pesquisas Espaciais) (MOHC additional realizations by INPE)	HadGEM2-ES HadGEM2-CC (Jones <i>et al.</i> , 2011)	(Collins <i>et al.</i> , 2011; Martin <i>et al.</i> , 2011) TRIFFID (Cox, 2001)	--	4	Q10	hill	no	X	--	--	1860-2005	2006-2099
Institute for Numerical Mathematics (INM)	INM-CM4 (Volodin <i>et al.</i> , 2010)	(Volodin, 2007) LSM (Bunnell <i>et al.</i> , 1977; Bonan, 1995, 1996)	--	1 <sup>b</sup>	Q10 <sup>c</sup>	hill	no	X	--	--	1850-2005	2006-2100
Institut Pierre-Simon Laplace	IPSL-CM5A-LR IPSL-CM5A-MR	ORCHIDEE ( <a href="http://orchid.ipsl.jussieu.fr">http://orchid.ipsl.jussieu.fr</a> )	3	4	Q10	increasing	no	X	X	--	1850-2005	2006-2100

Model Center	Model name	Soil model history	Litter	Soil	Temperature	Moisture	Nitrogen	Reported CMIP5 variables			Years considered for analysis	
								cSoil	cLitter	cCwd	Historical	RCP 8.5
	IPSL-CM5B-LR ( <a href="http://icmc.ipsl.fr">http://icmc.ipsl.fr</a> )	STOMATE (Krinner <i>et al.</i> , 2005) CENTURY (Parton <i>et al.</i> , 1988)										
Japan Agency for Marine-Earth Science and Technology, Atmosphere and Ocean Research Institute (The University of Tokyo), and National Institute for Environmental Studies (MIROC)	MIROC-ESM MIROC-ESM-CHEM (Watanabe <i>et al.</i> , 2011)	SEIB-DGVM (Sato <i>et al.</i> , 2007) Roth-C (Coleman & Jenkinson, 1999) DEMETER-1 (Foley, 1995) CENTURY (Parton <i>et al.</i> , 1987, 1992)	--	2	Arrhenius	increasing	no	X	X	--	1850-2005	2006-2100
Max-Planck-Institut für Meteorologie (Max Planck Institute for Meteorology) (MPI-M)	MPI-ESM-LR MPI-ESM-MR (CMIP5 output)	JSBACH (Raddatz <i>et al.</i> , 2007) BETHY (Knorr, 2000) CENTURY (Parton <i>et al.</i> , 1993)	--	3	Q10	increasing	no	X	X	--	1850-2005	2006-2100
Norwegian Climate Centre (NCC)	NorESM1-ME NorESM1-M (Tjiputra <i>et al.</i> , in prep)	CLM4 (Oleson <i>et al.</i> , 2008) CN (Thornton <i>et al.</i> , 2007) Biome-BCG 4.1.2 (Kimball <i>et al.</i> , 1997; Thornton, 1998; Thornton <i>et al.</i> , 2002; Thornton & Rosenbloom, 2005) (Olson, 1963; Veen & Paul, 1981; Veen <i>et al.</i> , 1984)	4	3	Arrhenius	increasing	yes	X	X	X	1850-2005	2006-2100

<sup>a</sup> We define a hill function as a function that increases to a maximum and then decreases.

<sup>b</sup> Turnover parameterization dependent on biome or vegetation type.

<sup>c</sup> Q<sub>10</sub> value dependent on temperature.

<sup>d</sup> Nitrogen option available but not turned on for CMIP5 run

	CESM1-BGC	NorESM1-M	BNU-ESM	BCC-CSM1-1-M	HadGEM2-ES	IPSL-CM5A-MR	GFDL-ESM2G	CanESM2	InM-CM4	MIROC-ESM	MPI-ESM-MR	Multi-model mean
Tundra and boreal	3.9	4.8	4.6	3.1	7.8	6.5	4.6	7.3	3.1	8.0	5.6	5.4
Other biomes	3.7	3.9	4.2	4.0	5.9	6.0	3.5	5.6	3.1	5.5	4.9	4.6
Global	3.8	4.1	4.3	3.8	6.4	6.1	3.8	6.1	3.1	6.2	5.1	4.8

Table S2: Mean soil temperature change for tundra and boreal forest biomes, all other biomes, and the globe for each ESM. Soil temperatures are area weighted means for the top 10 cm of the reported soil column.

	$\Delta C = C_{start}$	$\Delta C = \left( \frac{I_{start}}{k_{start}} \frac{k_{end}}{I_{end}} - 1 \right) C_{start}$ (Eq 2)	$\Delta C = \frac{I_{start}}{I_{end}}$	$\Delta C = \frac{k_{end}}{k_{start}}$
CESM1-BGC	0.100	0.735	0.153	NS
NorESM1-M	0.357	0.768	0.095	0.002
BNU-ESM	0.070	0.426	0.314	0.026
BCC-CSM1.1-M	0.388	0.580	0.221	0.118
HadGEM2-ES	0.032	0.762	0.215	0.108
IPSL-CM5A-MR	0.030	0.048	0.171	0.097
GFDL-ESM2G	0.346	0.664	0.105	0.123
CanESM2	0.004	0.870	0.333	0.079
INM-CM4	0.013	0.617	0.296	0.028
MIROC-ESM	0.060	NS	0.022	0.098
MPI-ESM-MR	0.008	0.038	0.029	0.001

Table S3: Within-model explanation of variation ( $R^2$ ) in the change in soil carbon over the 21<sup>th</sup> century by modern soil carbon, Eq 2) relative change in soil inputs, and relative change in decomposition. Decomposition rate ( $k$ ) was calculated from global heterotrophic respiration divided by soil carbon stock. Regressions used gridded mean differences (2090-2099 versus 1997-2006) and initial gridded means (1997-2006). All models have  $p < 0.05$ , unless marked as not significant (NS).

	CESM1-BGC	NorESM1-M	BNU-ESM	BCC-CSM1.1-M	HadGEM2-ES	IPSL-CM5A-MR	GFDL-ESM2G	CanESM2	INM-CM4	MIROC-ESM	MPI-ESM-MR
$R = kC$ (Eq. 3)	1/k [yr]	14.5	14.8	22.9	28.2	20.2	31.0	72.5	42.6	36.3	87.6
	$R^2$	0.658	0.665	0.029	0.065	0.364	0.119	0.033	0.084	0.131	0.004 <sup>a</sup>
$R = kQ_{10}^{(T-15)/10}C$	1/k [yr]	15.9	15.6	16.7	19.1	16.9	18.5	15.0	26.2	24.3	42.9
	$Q_{10}$	1.55	1.52	2.18	2.08	1.67	1.56	2.29	1.79	2.09	2.13
	$R^2$	0.909	0.915	0.931	0.928	0.796	0.754	0.622	0.834	0.727	0.677
$R = kQ_{10}^{(T-15)/10}W^bC$ (Eq. 5)	1/k [yr]	12.9	11.7	7.8	13.7	9.4	15.5	11.7	16.7	23.7	39.6
	$Q_{10}$	1.54	1.52	2.12	2.06	1.78	1.71	2.27	1.63	2.10	2.16
	b	0.15	0.08	0.61	0.20	1.06	0.60	0.53	0.20	0.03	0.27
	$R^2$	0.916	0.918	0.961	0.936	0.910	0.932	0.717	0.908	0.729	0.700

Table S4: Reduced complexity model parameters and fit to gridded mid-19<sup>th</sup> century 10-year mean. The first RCM depends solely on the soil carbon stocks and global decomposition rate (Eq. 3). The second RCM depends on soil carbon stocks, intrinsic decomposition rate, and temperature sensitivity. The final RCM depends on soil carbon stocks, intrinsic decomposition rate, temperature sensitivity, and moisture sensitivity (Eq. 5). All RCMs are optimized to their gridded initial 10-year *historic* mean and their associated  $R^2$ -fits are reported. All associated  $p < 10^{-11}$  except <sup>(a)</sup> where  $p = 0.007$ . HadGEM2-ES and GFDL-ESM2G are measured at 1860-1869 instead of 1850-1859 where their *historic* scenarios begin.




Figure S1: Cluster analysis of model outputs based on the spatial pattern of change in soil carbon between the 10-year means from 1997-2006 and 2090-2099.

Dissimilarities within model centers are as follows: BCC <3% dissimilar, CESM1 <1%, GFDL <3%, HadGEM2 <2%, IPSL <3%, MIROC <6%, MPI <2%, and NorESM1 <1%. The entire NorESM-CCSM-CESM1 cluster is <2% dissimilar.




Figure S2: Modern soil carbon density [ $\text{kg m}^{-2}$ ] for all models (10-year average; 1997-2006).




Figure S3: Modern NPP [ $\text{kg m}^{-2} \text{yr}^{-1}$ ] for all models (10-year average; 1997-2006).




Figure S4: Absolute change in NPP [ $\text{kg m}^{-2} \text{yr}^{-1}$ ] over the 21<sup>st</sup> century for all models (difference in 10-year means; 2090-2099 minus 1997-2006).




Figure S5: Relative change in NPP [%] over the 21<sup>st</sup> century for all models (difference in 10-year means; 2090-2099 minus 1997-2006).




Figure S6: Absolute change in soil carbon turnover time [yr] over the 21<sup>st</sup> century for all models (difference in 10-year means; 2090-2099 minus 1997-2006).




Figure S7: Relative change in soil carbon turnover time [%] over the 21<sup>st</sup> century for all models (difference in 10-year means; 2090-2099 minus 1997-2006).




Figure S8: Modern soil carbon turnover time [yr] for all models (10-year average; 1997-2006).




Figure S9: Absolute change in soil temperature [ $^{\circ}\text{C}$ ] over the 21<sup>st</sup> century for all models (difference in 10-year means; 2090-2099 minus 1997-2006).




Figure S10: Absolute change in soil water [ $\text{kg water m}^{-2} \text{ kg}^{-1} \text{ max. water m}^{-2}$ ] over the 21<sup>st</sup> century for all models (difference in 10-year means; 2090-2099 minus 1997-2006).




Figure S11: Modern normalized soil water [ $\text{kg water m}^{-2}$   $\text{kg}^{-1}$  max. water  $\text{m}^2$ ] for all models (10-year average; 1997-2006).




Figure S12: The change in soil carbon over the 21<sup>st</sup> century as a function of the a)

change in steady state soil carbon  $\left(\frac{I_{end}}{k_{end}} - \frac{I_{start}}{k_{start}}\right)$  and b) change in steady state relative to the initial steady state  $\left(\frac{I_{end} \times k_{start}}{k_{end} \times I_{start}} - 1\right)$ . The models are represented as follows; a: CESM1-BGC, b: NorESM1-M, c: BNU-ESM, d: BCC-CSM1.1-M, e: HadGEM2-ES, f: IPSL-CM5A-MR, g: GFDL-ESM2G, h: CanESM2, i: INM-CM4, j: MIROC-ESM, and k: MPI-ESM-MR.

- Arora VK (2003) Simulating energy and carbon fluxes over winter wheat using coupled land surface and terrestrial ecosystem models. *Agricultural and Forest Meteorology*, **118**, 21–47.
- Arora VK, Boer GJ (2005) A parameterization of leaf phenology for the terrestrial ecosystem component of climate models. *Global Change Biology*, **11**, 39–59.
- Arora VK, Scinocca JF, Boer GJ *et al.* (2011) Carbon emission limits required to satisfy future representative concentration pathways of greenhouse gases. *Geophysical Research Letters*, **38**, L05805.
- Bolker BM, Pacala SW, Parton Jr. WJ (1998) Linear analysis of soil decomposition: Insights from the Century model. *Ecological Applications*, **8**, 425–439.
- Bonan GB (1995) Land-atmosphere CO<sub>2</sub> exchange simulated by a land surface process model coupled to an atmospheric general circulation model. *Journal of Geophysical Research*, **100**, 2817–2831.
- Bonan GB (1996) A land surface model (LSM version 1.0) for ecological, hydrological and atmospheric studies: technical description and user's guide. *NCAR technical note NCAR/TN-417+STR*.
- Bunnell FL, Tait DEN, Flanagan PW, Van Clever K (1977) Microbial respiration and substrate weight loss—I. *Soil Biology and Biochemistry*, **9**, 33–40.
- Cao M, Woodward FI (1998) Net primary and ecosystem production and carbon stocks of terrestrial ecosystems and their responses to climate change. *Global Change Biology*, **4**, 185–198.
- Coleman K, Jenkinson DS (1999) ROTH-C-26.3, A model for the turnover of carbon in soils. *Model description and windows user guide*.
- Collins WJ, Bellouin N, Doutriaux-Boucher M *et al.* (2011) Development and evaluation of an Earth-System model – HadGEM2. *Geoscientific Model Development*, **4**, 1051–1075.
- Cox PM (2001) Description of the “TRIFFID” dynamic global vegetation model. *Hadley Centre technical note 24*.
- Dai Y, Zeng X, Dickinson RE *et al.* (2003) The Common Land Model. *Bulletin of the American Meteorological Society*, **84**, 1013–1023.

- Dai Y, Dickinson RE, Wang Y-P (2004) A Two-Big-Leaf Model for Canopy Temperature, Photosynthesis, and Stomatal Conductance. *Journal of Climate*, **17**, 2281–2299.
- Foley JA (1995) An equilibrium model of the terrestrial carbon budget. *Tellus B*, **47**, 310–319.
- Gent PR, Danabasoglu G, Donner LJ *et al.* (2011) The Community Climate System Model version 4. *Journal of Climate*, **24**, 4973–4991.
- Huang M, Ji J, Li K, Liu Y, Yang F, Tao B (2007) The ecosystem carbon accumulation after conversion of grasslands to pine plantations in subtropical red soil of South China. *Tellus B*, **59**, 439–448.
- Ji J, Huang M, Li K (2008) Prediction of carbon exchanges between China terrestrial ecosystem and atmosphere in 21<sup>st</sup> century. *Science in China Series D: Earth Sciences*, **51**, 885–898.
- Jones CD, Hughes JK, Bellouin N *et al.* (2011) The HadGEM2-ES implementation of CMIP5 centennial simulations. *Geoscientific Model Development*, **4**, 543–570.
- Keppel-Aleks G, Randerson JT, Lindsay K *et al.* (2013) Atmospheric carbon dioxide variability in the Community Earth System Model: Evaluation and transient dynamics during the 20th and 21st centuries. *Journal of Climate*.
- Kimball JS, Thornton PE, White MA, Running SW (1997) Simulating forest productivity and surface-atmosphere carbon exchange in the BOREAS study region. *Tree Physiology*, **17**, 589–599.
- Knorr W (2000) Annual and interannual CO<sub>2</sub> exchanges of the terrestrial biosphere: process-based simulations and uncertainties. *Global Ecology and Biogeography*, **9**, 225–252.
- Krinner G, Viovy N, de Noblet-Ducoudré N *et al.* (2005) A dynamic global vegetation model for studies of the coupled atmosphere-biosphere system. *Global Biogeochemical Cycles*, **19**, GB1015.
- Lindsay K, Bonan GB, Doney SC *et al.* (2013) Preindustrial control and 20th century carbon cycle experiments with the Earth system model CESM1-(BGC). *Journal of Climate*, **in review**.
- Martin GM, Bellouin N, Collins WJ *et al.* (2011) The HadGEM2 family of Met Office Unified Model climate configurations. *Geoscientific Model Development*, **4**, 723–757.

- Moorcroft PR, Hurt G C, Pacala SW (2001) A method for scaling vegetation dynamics: The Ecosystem Demography model (ED). *Ecological Monographs*, **71**, 557–586.
- Oleson KW, Niu G-Y, Yang Z-L *et al.* (2008) Improvements to the Community Land Model and their impact on the hydrological cycle. *Journal of Geophysical Research*, **113**, G01021.
- Olson JS (1963) Energy storage and the balance of producers and decomposers in ecological systems. *Ecology*, **44**, 322–331.
- Parton WJ, Schimel DS, Cole C V., Ojima DS (1987) Analysis of factors controlling soil organic matter levels in great plains grasslands. *Soil Science Society of America Journal*, **51**, 1173–1179.
- Parton WJ, Stewart JWB, Cole C V. (1988) Dynamics of C, N, P and S in grassland soils: a model. *Biogeochemistry*, **5**, 109–131.
- Parton WJ, Running SW, Walker B (1992) A toy terrestrial carbon flow model. *University Corp. for Atmospheric Research, Modeling the Earth System*, SEE N94-30616 08-45, 303–314.
- Parton WJ, Scurlock JMO, Ojima DS *et al.* (1993) Observations and modeling of biomass and soil organic matter dynamics for the grassland biome worldwide. *Global Biogeochemical Cycles*, **7**, 785–809.
- Raddatz TJ, Reick CH, Knorr W *et al.* (2007) Will the tropical land biosphere dominate the climate–carbon cycle feedback during the twenty-first century? *Climate Dynamics*, **29**, 565–574.
- Sato H, Itoh A, Kohyama T (2007) SEIB–DGVM: A new dynamic global vegetation model using a spatially explicit individual-based approach. *Ecological Modelling*, **200**, 279–307.
- Shevliakova E, Pacala SW, Malyshev S *et al.* (2009) Carbon cycling under 300 years of land use change: Importance of the secondary vegetation sink. *Global Biogeochemical Cycles*, **23**, GB2022.
- Sitch S, Smith B, Prentice IC *et al.* (2003) Evaluation of ecosystem dynamics, plant geography and terrestrial carbon cycling in the LPJ dynamic global vegetation model. *Global Change Biology*, **9**, 161–185.
- Thornton PE (1998) *Regional ecosystem simulation: combining surface- and satellite-based observations to study linkages between terrestrial energy and mass budgets*. Univ. of Montana, Missoula.

Thornton PE, Rosenbloom NA (2005) Ecosystem model spin-up: Estimating steady state conditions in a coupled terrestrial carbon and nitrogen cycle model. *Ecological Modelling*, **189**, 25–48.

Thornton PE, Law BE, Gholz HL *et al.* (2002) Modeling and measuring the effects of disturbance history and climate on carbon and water budgets in evergreen needleleaf forests. *Agricultural and Forest Meteorology*, **113**, 185–222.

Thornton PE, Lamarque J-F, Rosenbloom NA, Mahowald NM (2007) Influence of carbon-nitrogen cycle coupling on land model response to CO<sub>2</sub> fertilization and climate variability. *Global Biogeochemical Cycles*, **21**, GB4018.

Tjiputra JF, Roelandt C, Bentsen M *et al.* (2013) Evaluation of the carbon cycle components in the Norwegian Earth System Model (NorESM). *Geoscientific Model Development*, **6**, 301–325.

Todd-Brown KEO, Randerson JT, Post WM, Hoffman FM, Tarnocai C, Schuur EAG, Allison SD (2013) Causes of variation in soil carbon simulations from CMIP5 Earth system models and comparison with observations. *Biogeosciences*, **10**, 1717–1736.

Veen JA Van, Paul EA (1981) Organic carbon dynamics in grassland soils. 1. Background information and computer simulation. *Canadian Journal of Soil Science*, **61**, 185–201.

Veen JA, Ladd JN, Frissel MJ (1984) Modelling C and N turnover through the microbial biomass in soil. *Plant and Soil*, **76**, 257–274.

Volodin EM (2007) Atmosphere-ocean general circulation model with the carbon cycle. *Izvestiya, Atmospheric and Oceanic Physics*, **43**, 266–280.

Volodin EM, Dianskii NA, Gusev A V. (2010) Simulating present-day climate with the INMCM4.0 coupled model of the atmospheric and oceanic general circulations. *Izvestiya, Atmospheric and Oceanic Physics*, **46**, 414–431.

Watanabe S, Hajima T, Sudo K *et al.* (2011) MIROC-ESM 2010: model description and basic results of CMIP5-20c3m experiments. *Geoscientific Model Development*, **4**, 845–872.

Wu T, Li W, Ji J *et al.* (2013) Global carbon budgets simulated by the Beijing Climate Center Climate System Model for the last century. *Journal of Geophysical Research: Atmospheres*, **118**, 1–22.

Xu-Ri, Prentice IC (2008) Terrestrial nitrogen cycle simulation with a dynamic global vegetation model. *Global Change Biology*, **14**, 1745–1764.