

**A model for variable
phytoplankton
stoichiometry**

J. A. Bonachela et al.

A model for variable phytoplankton stoichiometry based on cell protein regulation

J. A. Bonachela¹, S. D. Allison², A. C. Martiny², and S. A. Levin¹

¹Department of Ecology and Evolutionary Biology, Princeton University, Princeton, New Jersey, 08544, USA

²Department of Ecology and Evolutionary Biology & Department of Earth System Science, University of California, Irvine, California, 92697, USA

Received: 25 January 2013 – Accepted: 6 February 2013 – Published: 21 February 2013

Correspondence to: J. A. Bonachela (jabo@princeton.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

The elemental ratios of marine phytoplankton emerge from complex interactions between the biotic and abiotic components of the ocean, and reflect the plastic response of individuals to changes in their environment. The stoichiometry of phytoplankton is, thus, dynamic and dependent on the physiological state of the cell. We present a theoretical model for the dynamics of the carbon, nitrogen and phosphorus contents of a phytoplankton population. By representing the regulatory processes controlling nutrient uptake, and focusing on the relation between nutrient content and protein synthesis, our model qualitatively replicates existing experimental observations for nutrient content and ratios. The population described by our model takes up nutrients in proportions that match the input ratios for a broad range of growth conditions. In addition, there are two zones of single-nutrient limitation separated by a wide zone of co-limitation. Within the co-limitation zone, a single point can be identified where nutrients are supplied in an optimal ratio. The existence of a wide co-limitation zone affects the standard picture for species competing for nitrogen and phosphorus, which shows here a much richer pattern. However, additional comprehensive laboratory experiments are needed to test our predictions. Our model contributes to the understanding of the global cycles of oceanic nitrogen and phosphorus, as well as the elemental ratios of these nutrients in phytoplankton populations.

1 Introduction

Marine phytoplankton take up and assimilate inorganic nutrients, thereby altering nutrient ratios in the ocean. Phytoplankton stoichiometry is, in turn, influenced by environmental factors, as individual cells can regulate their element ratios in response to changes in growth conditions (Rhee, 1978; Goldman et al., 1979). This regulatory capability is controlled by the cell's physiological response traits and, therefore, by evolution. Consequently, species are characterized by different stoichiometries (Geider and

BGD

10, 3241–3279, 2013

A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



La Roche, 2002; Klausmeier et al., 2004a), which may contribute to the maintenance of oceanic biodiversity (Göthlich and Oschlies, 2012). Thus, the mechanisms underlying phytoplankton ratios of important elements such as carbon, nitrogen and phosphorus are essential for understanding the biogeochemical cycles of these nutrients, and their role in shaping phytoplankton community composition.

Previous laboratory work has focused mainly on the response of cellular contents and stoichiometry to changes in (i) nitrogen and phosphorus inputs and (ii) dilution rates, in continuous cultures (see Fig. 1 and Tables 1 and 2). All this experimental work agrees on some important phenomenology; however, the multitude of different species and environmental conditions used in experiments has led to a lack of consensus with respect to some trends (e.g. phosphorus quota vs dilution rate, Fig. 1c) or, especially, the specific shape of the functional dependence. This has, in consequence, limited the ability of those experiments to give generalized answers to fundamental questions. For instance, how do the cell contents of the different nutrients (or quotas) interact with each other, and how does that feedback influence growth and stoichiometry regulation? How do the answers to the previous questions affect the condition(s) under which growth is limited by multiple nutrients, and the competitive abilities of the cell?

Theorists have developed models intended to reproduce experimental results and identify controls on phytoplankton stoichiometry. In accordance with early experiments (Droop, 1968; Rhee, 1978; Terry et al., 1985), one group of models assumes that growth rate, μ , depends *exclusively* on the most limiting nutrient (Liebig's law) following a hyperbolic function f (Droop's law): $\mu = \min(f(N), f(P))$ (Fig. 2a). These "threshold models" impose, thus, a single functional form f for the growth rate that depends exclusively on whichever nutrient limits growth (Legović and Cruzado, 1997; Klausmeier et al., 2004a,b, 2007). In consequence, models following this approach predict a symmetric response of the nitrogen and phosphorus quotas to changes in the dilution rate of simulated chemostats (see Fig. 2b). However, a different trend (i.e. asymmetry) for the quotas has been observed in some experiments (Fig. 1b, c) (Elrifi and Turpin, 1985). To reproduce this trend, some threshold models impose interdependency of the

BGD

10, 3241–3279, 2013

A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



nutrient contents by using phenomenological expressions that assume that the acquisition of one nutrient is limited by the other nutrient (Roelke et al., 1999; Bougaran et al., 2010). A second group of models also assumes biochemical independence of the nutrients and a common functional form for the response of growth to each nutrient quota; however, growth depends on the product of those functions (e.g. $\mu = f(N)f(P)$), allowing for limitation by multiple nutrients (Terry, 1982; Saito et al., 2008). A last group of models avoids assumptions about the functional dependence of growth rate on quotas by following a more mechanistic approach. In these models, equations account for essential physiological processes through growth, uptake or respiration terms. This group includes simple models devised to reproduce both marine and terrestrial plant behavior (Ågren, 2004), as well as more sophisticated ones able to replicate experimental results from different phytoplankton strains (Pahlow and Oschlies, 2009). The latter example, however, assumes optimal nutrient uptake conditions (Smith and Yamanaka, 2007), which implicitly imposes an evolutionary strategy for the phytoplankton species that yields a fitness maximum, regardless of changes in environmental or predatory pressures. Furthermore, this model also makes use of expressions that impose quota interactions at the level of nutrient uptake and instantaneous cell acclimation.

These modeling efforts to introduce a dependence of nutrient uptake on cell quotas acknowledge the well-documented importance of protein regulation in determining cell nutrient content. For example, phytoplankton show changes in the number of nutrient-uptake proteins according to the environmental nutrient concentration (McCarthy and Goldman, 1979; Gotham and Rhee, 1981a,b), which ultimately can be translated into a cell quota dependence for uptake proteins (Morel, 1987). Thus, the different resource allocation strategies followed by the cell to perform this regulation influence crucially its stoichiometry. Conversely, the response capacity of the cell is determined by the phosphorus cell content, which is required for ribosomes (P-rich molecules) in order to synthesize uptake proteins. Thus, accounting for this feedback between resource allocation/protein regulation and nutrient content is essential for predicting phytoplankton stoichiometry.

BGD

10, 3241–3279, 2013

A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The models above have included resource allocation in different ways. Klausmeier et al., for instance, discern between uptake and assembly machineries; that is, protein/chloroplasts and ribosomes, respectively (Klausmeier et al., 2004a). This model imposes a fixed stoichiometry for each of these types, and allows the cell to allocate resources as constrained by a trade-off between uptake and assembly. A similar functional distinction is made in (Pahlow and Oschlies, 2009), although the stoichiometry and trade-off between the different functional machinery is less constrained. Although existing models are able to replicate most of the patterns summarized in Fig. 1 and Tables 1 and 2, the use of prescribed phenomenological forms for growth or quota interactions limits their potential to study the feedback between quotas, cell growth, and stoichiometry.

In this paper, we aim to improve our understanding of how species stoichiometry emerges from the interplay between environment and cell physiology (and, ultimately, evolution). To this end, we use recent theoretical advances in the relation between nutrient uptake and cellular allocation patterns to build a theoretical model that, in contrast to previous models, incorporates the key dependence between dynamic uptake-protein regulation and nutrient availability. We avoid the use of optimality assumptions or imposing explicit dependencies between quotas, uptake and growth, by using simple expressions representing plausible physiological mechanisms such as protein regulation. Thus, our model describes phytoplankton cells for which protein regulation (and hence cell stoichiometry and growth) is determined by the nutritional history of the cell. This regulation allows for a dynamic acclimation to nutrient variability by altering the number of uptake proteins. We first examine the ability of the model to reproduce qualitatively the existing experimental data. We then explore the conditions under which nitrogen or phosphorus limit (or co-limit) phytoplankton growth, and speculate on how the classic predictions of resource competition theory are affected by the resulting interactions between nitrogen and phosphorus in our model. Finally, we propose an experimental setup able to verify the theoretical hypotheses presented here.

2 Methods

Our model is composed of dynamic equations for the population content of organic carbon, C, nitrogen, N, and phosphorus, P. Positive terms describe the basic processes contributing to increases in population levels of the considered elements, and negative terms account for the decreases. Regulation of protein production is the key mechanism underlying the dynamics of the population, including explicit equations for the regulation of nutrient uptake proteins. This regulation is controlled by two master functions: a function F that represents the protein synthesis and the cues that influence it; and a function G that accounts for the availability of ribosomes to carry out that synthesis.

We consider chemostat conditions, which allows the system to reach steady state as cells are washed out during the dilution process. We use these continuous culture conditions for the sake of simplicity, as this facilitates a better comparison with existing experimental and theoretical work; nonetheless, the model and results can be easily extended to other environmental conditions.

(i) *Equations for organic nitrogen and phosphorus*: Population N and P increase at a rate V_N and V_P , respectively, due to nutrient uptake, and decrease due to the washout or dilution process of the chemostat (at a rate w) or through other losses, such as leakage, at rates R_N and R_P , respectively:

$$\frac{dN}{dt} = V_N(t) - (R_N + w)N(t), \quad (1)$$

$$\frac{dP}{dt} = V_P(t) - (R_P + w)P(t). \quad (2)$$

BGD

10, 3241–3279, 2013

A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The population uptake rate for the different nutrients is given by:

$$V_N(t) = \frac{V_{\max_N}(t) [N]}{[N] + \tilde{K}_N} B(t), \quad (3)$$

$$V_P(t) = \frac{V_{\max_P}(t) [P]}{[P] + \tilde{K}_P} B(t), \quad (4)$$

- 5 (see symbols and units in Table 3). B is the number of cells in the population, V_{\max} is the maximum uptake rate of a cell, and \tilde{K} is an effective form for the half-saturation constant accounting for a boundary layer in which the local nutrient concentration is smaller than the bulk nutrient concentration (Pasciak and Gavis, 1974; Mierle, 1985) (see derivation in, e.g. Armstrong, 2008; Bonachela et al., 2011):

$$10 \quad \tilde{K}_N(t) = K_N \left(1 + \frac{V_{\max_N}(t)}{4\pi D_N r_c K_N} \right), \quad (5)$$

$$\tilde{K}_P(t) = K_P \left(1 + \frac{V_{\max_P}(t)}{4\pi D_P r_c K_P} \right), \quad (6)$$

- where K_N and K_P are the standard half-saturation constants associated with the uptake of N and P, respectively, D_N and D_P are the diffusivity of nitrogen and phosphorus in the medium, respectively, and r_c is the cell radius. Note that the maximum uptake rate is proportional to the total number of uptake proteins that the cell accumulates at its surface (see Appendix). As explained below, that number changes according to environmental conditions. These variations entail changes in \tilde{K} , which accounts for the observed differences in the half-saturation constant of single species subject to variations in environmental nutrient concentrations (Pasciak and Gavis, 1974)¹.

¹Also note that this expression predicts improved uptake abilities for small cells when it is formulated in terms of a *normalized* maximum uptake rate, V_{\max}/r_c^2 (Armstrong, 2008).

A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(ii) *Equations for the number of uptake proteins*: in our model, phytoplankton are able to regulate the number of proteins, n , they allocate for the uptake of the different nutrients. The cell's content of the nutrient (or quota, Q) is the key factor controlling this regulatory process (Morel, 1987; Song and Ward, 2007; Flynn, 2008). As documented in experimental work for the uptake of a single nutrient (either nitrogen or phosphorus), a population shows an increased number of uptake proteins (or population V_{\max}) when the quota is low, compensating for low uptake rates, and a lower V_{\max} when the nutrient is abundant (McCarthy and Goldman, 1979; Gotham and Rhee, 1981a,b; Riegman and Mur, 1984; Dyhrman and Palenik, 2001). Thus, in oligotrophic conditions the cell allocates more uptake proteins in order to increase its absorbing area and, therefore, the probability of a successful encounter between the scarce nutrient ions and the protein at the cell membrane; on the other hand, the cell may down-regulate the synthesis of these proteins when the internal concentration of the nutrient reaches the storage maximum Q_{\max} in order to decrease biosynthesis and maintenance costs. This is eventually translated into an effective increase in affinity under oligotrophic conditions, and a decrease in eutrophic environments (Bonachela et al., 2011). This mechanism can be encoded in a simple way by using the “protein expression” function, F , given by (Bonachela et al., 2011):

$$F \left(\frac{Q_{\max} - Q}{Q_{\max} - Q_{\min}} \right) = \frac{2}{1 + e^{-k_F \frac{Q_{\max} - Q}{Q_{\max} - Q_{\min}}}} - 1, \quad (7)$$

where k_F is a free parameter controlling the shape of the function and Q_{\min} is the minimum amount of nutrient required for the cell to grow. We use here a generic sigmoid function, which is motivated by the identification of F with processes involving gene expression – traditionally represented by Hill (sigmoid) functions (Alon, 2007); however, any other normalized function of similar trend does not alter the qualitative behavior of the model (Bonachela et al., 2011). In addition, it is possible to add other (competing) strategies to the regulation, such as up-regulation of protein synthesis when uptake activity is high and down-regulation when it is low. As justified in (Bonachela et al.,

2011), these modifications do not alter the qualitative behavior of the number of uptake proteins. In order to constrain uptake protein synthesis when nutrient levels drop below growth minimum requirements, we also impose the condition $F = 0$ when Q falls below Q_{\min} ².

Likewise, our model represents a dependence of protein expression on phosphorus: as explained above, phosphorus is a major component of ribosomes, essential for the synthesis of any kind of protein, and also required in ATP, ADP and other forms of energy storage for the cell (see Sterner and Elser, 2002, or discussion in Geider and La Roche, 2002; Bougaran et al., 2010 and references therein). Thus, we include the “protein repression” function, G , which is strictly a function of the internal phosphorus content of the cell, Q_P :

$$G \left(\frac{Q_P - Q_{P_{\min}}}{Q_{P_{\max}} - Q_{P_{\min}}} \right) = \frac{1}{1 + e^{-k_{G,1} \left(\frac{Q_P - Q_{P_{\min}}}{Q_{P_{\max}} - Q_{P_{\min}}} - k_{G,2} \right)}}, \quad (8)$$

where $k_{G,1}$ and $k_{G,2}$ are shape factors that help establish the range and boundaries of the function. The lower the content of phosphorus, the stronger the influence of the function on the regulation of the synthesis of proteins. Note that the repression function depends only on Q_P because we assume that phosphorus (through ribosomal RNA) is the ultimate limiting nutrient for the synthesis of proteins in our model. Any cellular nitrogen will be allocated to proteins as long as Q_N is above the minimum required to grow, $Q_{N_{\min}}$.

As a last constraint, we account for the physical impossibility for the cell to accumulate on its surface more uptake proteins than allowed by the available space. Thus, we introduce in the equations for both n a Heaviside function, H , that depends on the ratio of absorbing area to total area, A_{rel} (see Appendix for mathematical definitions). This

²As commented in Results, however, this limit is never reached in simulations, and quotas reach larger *effective* extreme values instead.

BGD

10, 3241–3279, 2013

A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



constraint also introduces an effective competition between the N-uptake and P-uptake machinery for space at the membrane.

If we take into account the contribution of all the factors above, the equations for the number of uptake sites are:

$$5 \quad \frac{dn_N}{dt} = v_N H(1 - A_{\text{rel}}(t)) F \left(\frac{Q_{N_{\text{max}}} - Q_N(t)}{Q_{N_{\text{max}}} - Q_{N_{\text{min}}}} \right) G \left(\frac{Q_P(t) - Q_{P_{\text{min}}}}{Q_{P_{\text{max}}} - Q_{P_{\text{min}}}} \right) B(t) - wn_N(t), \quad (9)$$

$$\frac{dn_P}{dt} = v_P H(1 - A_{\text{rel}}(t)) F \left(\frac{Q_{P_{\text{max}}} - Q_P(t)}{Q_{P_{\text{max}}} - Q_{P_{\text{min}}}} \right) G \left(\frac{Q_P(t) - Q_{P_{\text{min}}}}{Q_{P_{\text{max}}} - Q_{P_{\text{min}}}} \right) B(t) - wn_P(t), \quad (10)$$

10 where v_X is the maximum number of proteins for the uptake of nutrient X that a cell can synthesize per unit time (see Appendix). Because the change in n with time depends on the quotas, the maximum uptake rate at any time depends on the nutritional history of the cell.

Figure 3 shows the dependence of these equations on the quotas of the different nutrients. The use of the expression and repression functions imposes the allocation strategy of the cell under nutrient limitation. When nitrogen is scarce and the cell is N-limited, the cell prioritizes the synthesis of N-uptake proteins in order to increase Q_N and, ultimately, increase growth (see below). Only when the N-quota falls below the survival threshold of the minimum quota, $Q_{N_{\text{min}}}$, does the cell down-regulate N-protein synthesis. Quotas below $Q_{N_{\text{min}}}$ result in decreasing uptake and growth rates and, ultimately, starvation. On the other hand, phosphorus is the key regulator of biosynthesis due to its role in ribosomes and energy reserves; therefore, P scarcity entails a stronger limitation on the synthesis of proteins, even for quotas above the minimum. Conversely, luxury consumption of a nutrient leads to the down-regulation of the synthesis of its uptake proteins, with cell growth being limited only by the availability of the reciprocal nutrient. Note that the different dependencies of n_N and n_P on the quotas break the symmetry expected a priori from Eqs. (1)–(6) for N and P.

BGD

10, 3241–3279, 2013

A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(iii) *Equation for Organic Carbon*: the dynamics of the population organic carbon takes into account a photosynthetic term (only source of organic C), maintenance costs and the loss due to dilution:

$$\frac{dC}{dt} = \left[P_{\max} F \left(\frac{Q_N(t) - Q_{N_{\min}}}{Q_{N_{\max}} - Q_{N_{\min}}} \right) G \left(\frac{Q_P(t) - Q_{P_{\min}}}{Q_{P_{\max}} - Q_{P_{\min}}} \right) - M_C \left(\frac{V_N(t)}{N(t)} + \frac{V_P(t)}{P(t)} \right) - R_C - w \right] C(t), \quad (11)$$

where P_{\max} represents the maximum photosynthetic rate, M_C is a maintenance rate and R_C a respiration rate. The first term accounts for the synthesis of photosynthetic proteins. Following e.g. (Geider et al., 1998), nitrogen quota plays a main role in the photosynthetic rate because chloroplasts are protein- (i.e. nitrogen-) rich. In our model, the protein expression and repression functions regulate photosynthesis; in the case of photosynthetic proteins (e.g. RuBisCO), both F and G increase with the internal content of the nutrient in the cell, as large internal levels of N and P are a proxy for favorable growth conditions (see Fig. 4a). The second term accounts for the maintenance cost associated with uptake proteins, cost represented by the nutrient-specific uptake rate³. The third term accounts for the respiration rate related to the rest of the cell metabolism, and the last term is the mortality or dilution term. The use of a constant P_{\max} or a linear respiration term is the consequence of simplification, assuming constant and ideal irradiance and temperature conditions for photosynthesis. This representation of C acquisition contrasts with the explicit terms for nutrient uptake considered in all the equations above. Thus, our model focuses on nutrient acquisition as the main regulator of the population stoichiometry.

The per-capita rate of change for carbon in the population is used to calculate the growth rate, $\mu = \frac{1}{C} \frac{dC}{dt} + w$. Consequently, the organic carbon per cell in the population

³Note that both V_X and Q_X – and, therefore, X itself – are increasing functions of the growth rate, with $Q_X \in [Q_{X_{\min}}^{\text{eff}}, Q_{X_{\max}}^{\text{eff}}]$ and $V_X \in [0, V_{\max}^{\text{lo}}]$ (see Sect. 3). Thus, the maintenance term is well defined into $[0, V_{\max}^{\text{lo}}/Q_{X_{\max}}^{\text{eff}}]$.

A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



remains fixed. This allows us to track the number of cells in the population:

$$\frac{dB}{dt} = (\mu(t) - w) B(t) \quad (12)$$

and, therefore, to switch from a population-dynamics to a biomass-dynamics approach when required. In this paper, we use Eq. (11) only to calculate the growth rate of the population, and focus on $B(t)$ to characterize its abundance. Thus, the behaviors of $Q_N = N/B$ and $Q_P = P/B$ are qualitatively similar to those of N/C and P/C , respectively.

(iv) *Equations for Inorganic Nitrogen and Phosphorus*: to monitor changes in the inorganic nutrients present in the chemostat, we take into account the inflow of fresh nutrient (the only positive term in the equations), the outflow due to dilution, and the consumption of the nutrient by the population through the uptake term:

$$\frac{d[N]}{dt} = w ([N_0] - [N]) - V_N(t), \quad (13)$$

$$\frac{d[P]}{dt} = w ([P_0] - [P]) - V_P(t), \quad (14)$$

where $[N_0]$ and $[P_0]$ are the concentrations of nutrient in the fresh medium that enter the vessel at the dilution rate.

A graphical summary of the physiological processes and interactions considered in the model can be found in Fig. 4b.

(v) *Simulation Set-Up*: we varied the input ratio, $[N_0] : [P_0]$, and the dilution rate, w , of the chemostat in simulations to study the reaction of phytoplankton to changes in environmental conditions that may alter nutrient availability. We explored the consequences for phytoplankton stoichiometry by integrating numerically the dynamic equations of the population-level model, Eqs. (1)–(14). Although the model is applicable to any phytoplankton strain, for the sake of concreteness we used data available in the literature to parametrize our population in accordance with a generic *Synechococcus* species (see

BGD

10, 3241–3279, 2013

A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



actual values in Table 3). For each pair of ($w, [N_0] : [P_0]$), we let the system reach stationary state and calculated different observables such as the number of cells, uptake and growth rates, or nutrient content of the population. We modified the input ratio by altering one nutrient input concentration while fixing the other. This procedure allowed us to compare the qualitative behavior of the model with most of the available experimental and theoretical work. Following (Rhee, 1978), we fixed $[P_0]$ and varied $[N_0]$ in our analysis.

3 Results

We first explore the behavior of the per-capita nutrient content (or quota, Q) of nitrogen and phosphorus when the relative input concentration of the nutrients is varied. For low relative input nitrogen (i.e. low $[N_0] : [P_0]$) Q_N remains fixed at low value that depends on the dilution rate of the chemostat, while phosphorus reaches saturating levels in the cell (Fig. 5a). Cell growth is, then, limited solely by the availability of nitrogen, because changes in the input ratio do not affect the stationary value of the quota of a nutrient when it is the only limiting factor in the chemostat. As nitrogen input increases, population growth drives inorganic phosphorus to lower values. In consequence, phosphorus becomes limiting *as well*, and the stationary value of Q_N grows with $[N_0] : [P_0]$ until the realized maximum storage capacity, $Q_{N_{\max}}^{\text{eff}}$, is reached. This effective maximum value differs from $Q_{N_{\max}}$ due to the presence of the loss rate R_N and the down-regulation of η_N , which affect growth even for large concentrations of the nutrient (see Eq. 3 under stationary conditions). Q_P shows an analogous pattern as the relative phosphorus input increases (i.e. $[N_0] : [P_0]$ decreases).

Consequently, in the representation of Q_N against Q_P (Fig. 5b), zones of exclusive limitation by nitrogen are characterized by the vertical sections of the curve, while zones of exclusive limitation by phosphorus constitute the horizontal parts. For the rest of the points on the curve, cells are co-limited by both nutrients. The range of co-limitation decreases as the dilution rate increases.

BGD

10, 3241–3279, 2013

A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



In this broad zone of co-limitation (ranging from $[N_0 : P_0] \sim 20$ to $[N_0 : P_0] \sim 70$ for the smallest w in Fig. 5), the population $N : P$ matches the input ratio (Fig. 6a). Deviations from this relationship occur at low input ratios because only nitrogen is limiting and Q_N remains at its effective minimum for a given dilution rate whereas Q_P remains fixed at its effective maximum value (Fig. 5a). The converse is true for low relative phosphorus ratios (high $[N_0] : [P_0]$). These deviation patterns are more pronounced as the dilution rate increases. However, there is a single input ratio at which cells are able to incorporate both nutrients in a proportion that exactly matches the input ratio regardless of the value of the dilution rate, resulting in the reduction of both inorganic nutrients to very low concentrations. This is an *optimal nutrient ratio*, as the cell is able to draw down nutrients to low levels even at high dilution rates. The ratio at which this happens, $[N_0] : [P_0]_{\text{opt}}$, can be determined by representing $(Q_N : Q_P) / ([N_0] : [P_0])$ against the input ratio for different values of the dilution rate; all these curves intersect at the ordinate 1, with the abscissa corresponding to $[N_0] : [P_0]_{\text{opt}} \sim 26$ in this example (Fig. 6b). This optimal nutrient ratio coincides with the cell quota ratio during exponential growth, and can be calculated from its theoretical definition (see Appendix).

Variations in the dilution rate affect nitrogen and phosphorus quotas differently (Fig. 7). Q_N is an increasing linear or convex function of the dilution rate, depending on the limiting nutrient; on the other hand, Q_P is a more complicated function that shows convexity when phosphorus is limiting and non-monotonicity otherwise, with a range of change wider than that of Q_N (Fig. 7a, b). This pattern translates into a growth rate that, for any input ratio, increases as the quota of the limiting nutrient(s) increases, saturating to a maximum value around $\mu_{\text{max}} = \mu_{N_{\text{max}}} = \mu_{P_{\text{max}}} = 0.825 \pm 0.005$ (Fig. 7c, d). The projected quota at zero growth decreases with declining relative supply of each nutrient, until reaching $Q_{\text{min}}^{\text{eff}}$. On the other hand, the quota associated with maximum growth increases as the relative input of the corresponding nutrient increases, until it reaches $Q_{\text{max}}^{\text{eff}}$. The latter behavior is more marked in the case of phosphorus than for nitrogen.

We also determined the uptake rate of the limiting nutrient, which follows a hyperbolic functional form depending on the external nutrient concentration (see Fig. 8). The maximum uptake rate equations, Eqs. (9) and (10), introduce a dependence of V_{\max} on environmental conditions (Bonachela et al., 2011). More specifically, the stationary value of V_{\max} for the limiting nutrient decreases as the dilution rate increases, reaching a lower plateau at dilution rates close to the maximum growth rate. At their minimum values, $V_{\max_N}^{\text{lo}} \sim 2.2 \times 10^{-14} \text{ mol cell}^{-1} \text{ d}^{-1}$ and $V_{\max_P}^{\text{lo}} \sim 8.5 \times 10^{-16} \text{ mol cell}^{-1} \text{ d}^{-1}$. Thus, as environmental conditions improve, the cell allocates more resources to growth and less to uptake (see Eqs. 9–11, or Figs. 3 and 4a).

4 Discussion

4.1 Model validation

If we aspire to understand the interactions between quotas and their effect on cell growth and stoichiometry, our model must be able to predict realistic behavior with as few assumptions as possible. Indeed, our model makes predictions that match qualitatively most of the phenomenology observed experimentally in phytoplankton populations subject to changing input nutrient ratios or dilution rates. The behavior we observe for the quotas, with Q_N increasing and Q_P decreasing with the input ratio (Fig. 5a) is described in the early work by Rhee (Rhee, 1978) and Terry et al. (Terry et al., 1985), although in those cases maximum storage limits (corresponding to upper parts of the curves in Fig. 5a) were seemingly never reached. In our model, these effective maximum (i.e. *saturation*) values for Q are reached as a result of the dynamic equation for each nutrient's uptake protein synthesis. This physiological range for the quotas imposes limits on protein regulation (see Eqs. 9–11) and, ultimately, on the element ratios. That is the case with the N : P ratio (Fig. 6a), for which plateaus at both the upper and lower part of the curve are the result of the cell reaching its extreme quota values ($Q_{N_{\max}}^{\text{eff}}$ and $Q_{P_{\min}}^{\text{eff}}$ in the case of the upper plateau, and $Q_{N_{\min}}^{\text{eff}}$ and $Q_{P_{\max}}^{\text{eff}}$ for the

BGD

10, 3241–3279, 2013

A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



lower one). This is analogous, for instance, to the situation described in Fig. 3 – compilation of algal cultures – and Fig. 4 in (Hall et al., 2005) (see Fig. 1d). On the other hand, our protein regulation mechanism causes an asymmetric reaction of phytoplankton N and P to changes in dilution rate (Fig. 7a, b) through different dependencies of Eqs. (9) and (10) (or dV_{\max_N}/dt and dV_{\max_P}/dt , respectively) on Q_N and Q_P (see Fig. 3). Asymmetry, observed repeatedly in experiments (Terry, 1982; Elrifi and Turpin, 1985), is introduced in other models by using terms devised specifically to that end (see e.g. Bougaran et al., 2010).

A last important experimental observation is the hyperbolic dependence of growth on the limiting quotas (see Fig. 1 or corresponding references in Table 1). Imposed in threshold (Droop-like) models (Fig. 2a) (Klausmeier et al., 2004a,b, 2007; Bougaran et al., 2010), this behavior is an emergent property of mechanistic approaches (see (Pahlow and Oschlies, 2009) or Fig. 7c, d). Thus, in each zone of single limitation, threshold models show growth curves that depend only on the nutrient quota and not on the input ratio. Moreover, they impose $\mu_{N_{\max}} = \mu_{P_{\max}}$, in accordance with experiments (Rhee, 1978; Elrifi and Turpin, 1985). In our case, (also expected from Pahlow and Oschlies, 2009), there are marked differences between the single-limitation and co-limitation zones: in the single-limitation zones we observe a threshold-like behavior, in which the growth rate curve depends only on the (non-saturated) quota, and different input ratio curves collapse to a single curve ($[N_0] : [P_0] < 16$ in Fig. 7c and $[N_0] : [P_0] > 60$ in Fig. 7d); in the zone of co-limitation, however, the growth rate depends on both quotas, and the curves change with the value of the input ratio (rest of $[N_0] : [P_0]$ in Fig. 7c, d). The equality of $\mu_{N_{\max}}$ and $\mu_{P_{\max}}$ results from the dynamics of our model, as do the specific values of $\mu_{N_{\max}}$, $\mu_{P_{\max}}$, $V_{\max_N}^{\text{lo}}$, and $V_{\max_P}^{\text{lo}}$. These values, contrarily to other models, are not imposed *a priori* in our case; furthermore, they are in reasonable agreement with static maximum uptake rates measured for *Synechococcus* (Healey, 1985).

A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

4.2 Quota interactions

In addition to replicating empirical findings, our model approaches cell quota interactions differently from other models. The interplay between quotas is a key determinant of phytoplankton stoichiometry, as it decides the response of cells to changes in the environment and, ultimately, cellular elemental ratios. Most other models introduce quota interactions through phenomenological expressions relating V_{\max_N} and V_{\max_P} to cell quotas. They impose either an independent limitation (see “dynamic uptake version” in Klausmeier et al., 2004b), or different kinds of cross-limitations (Bougaran et al., 2010; Pahlow and Oschlies, 2009). These expressions assume a direct and instantaneous dependence of maximum uptake rates on nutrient quotas. However, there is no known physiological mechanism by which a cell can instantaneously change its uptake potential. Instead, such adjustments occur through allocation processes that alter the synthesis and degradation rates for uptake proteins (Caperon, 1969; Song and Ward, 2007; Klausmeier et al., 2007). We capture these allocation strategies in our model by allowing nutrient quotas to determine the production rate of uptake sites (Eqs. 9–10) rather than the standing pool of uptake sites (which is proportional to V_{\max}). Thus, our model uses the quotas to specify how cellular resources are allocated to N versus P acquisition.

Therefore, nutrient interactions result in our model from the role of each quota in the expression and repression functions. We avoid imposing phenomenological expressions by constructing these regulatory functions from simple biological arguments. Both F and G encode the physiological responses triggered by changes in the quota of the different nutrients, which determine the rates of resource acquisition for the cell through photosynthesis and nutrient uptake⁴. Equations (9) and (10) adjust the uptake strategy by altering the production of uptake proteins in the cell, which allows for representation

⁴Moreover, the opposite response of the regulating functions to quota changes introduce a trade-off between these two processes (see Figs. 3 and 4a), as well as a trade-off between the uptake of the two nutrients evidenced by Fig. 5b.

BGD

10, 3241–3279, 2013

A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of lags in cellular responses. These simple but mechanistic dynamic equations confer plasticity to the population, allowing it to adjust its stoichiometry in a changing environment. Moreover, we avoid using assumptions about the optimal character of uptake or growth (Klausmeier et al., 2004b, 2007; Smith and Yamanaka, 2007; Pahlow and Oschlies, 2009), thus setting phytoplankton cells in a less-constrained evolutionary context at the expense of a larger parameter space.

Models that, on the contrary, assume no interaction between nutrients show a sharp transition between nitrogen and phosphorus limitation, with a single co-limitation point given by $[N_0] : [P_0]_{\text{opt}}$ (Fig. 2d) (Tilman, 1982). Based on Droop's law, that constant value equals $Q_{N_{\text{min}}} / Q_{P_{\text{min}}}$ (Rhee and Gotham, 1980; Klausmeier et al., 2007). For cells described by our model, nitrogen and phosphorus are "interacting essential resources" (Tilman, 1982), showing single-limitation zones but also a wide range for co-limitation (Fig. 5). Consequently, scarcity in one nutrient can be partially compensated for by an increased concentration of the other nutrient (Tilman, 1982). Moreover, this co-limitation range depends in our model on the environmental conditions (see Fig. 6a). Experimental work has reported a dependence of co-limitation on the environment, particularly the dilution rate (Kunikane et al., 1984; Elrifi and Turpin, 1985; Terry et al., 1985) and irradiance (Healey, 1985). In addition, our model predicts that the optimal ratio, a specific point in the co-limitation region, is $[N_0] : [P_0]_{\text{opt}} = Q_{N_{\text{max}}}^{\text{eff}} / Q_{P_{\text{max}}}^{\text{eff}}$ (see Appendix). The latter expression, shared by the model in (Pahlow and Oschlies, 2009), is not based on any imposed form for the growth rate, and matches the theoretical definition of the optimal ratio introduced in the Results section (ratio shown by the cell during exponential growth). Furthermore, note that the optimal ratio is far from being related to the Redfield ratio (N : P=16) (Klausmeier et al., 2004a), which in our model is just one more point in the co-limitation zone.

BGD

10, 3241–3279, 2013

A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4.3 Implications of a broad co-limitation region

As explained in Tilman (1982), Rhee (1978), the outcome of competition between species growing on two non-interacting essential resources depends on the relative distance between their respective co-limitation points. Let us consider the case of nitrogen and phosphorus. When two species, A and B (with single transition points $(N : P)_{opt_A}$ and $(N : P)_{opt_B}$, respectively), are placed in the same chemostat and these nutrients are considered perfectly essential (like in threshold models), the theoretical outcome can be determined using the distance of each species' stoichiometric ratio from its respective optimum, $\Delta = [(N : P)_{opt} - (N : P)] / (N : P)_{opt}$. If Δ_A and Δ_B have different signs, each species will be limited by different nutrients and co-existence is possible (see Fig. 9a). If both have the same sign, the species that is closer to its optimum ratio (i.e. with smaller $|\Delta|$) will, in principle, out-compete the other.

As an important consequence of Fig. 5b, we speculate that a wide region of co-limitation will affect the predicted outcome of phytoplankton species competition. Indeed, the potential outcome in our model is more complicated (Fig. 9b). Opportunities for co-existence do still depend on the competing species being limited by reciprocal nutrients (i.e. different sign for their Δ). Δ is still important because it indicates which nutrient is more influential to the growth of each species. However, the input ratio at which the quota of a nutrient reaches its maximum (i.e. saturation) is also important, because it determines the zone where the cell is limited exclusively by the reciprocal nutrient. Co-limited phytoplankton have competitive advantage over those limited by a single nutrient (Tilman, 1982). Thus, saturation points influence the outcome of competition experiments in our model, introducing competitive exclusion in parts of the diagram where threshold models predict co-existence (see shaded parts in Fig. 9). Furthermore, in our model both consumption ability (or, ultimately, V_{max}) and saturation points depend on environmental conditions and, thus, must be accounted for. This is in contrast to descriptions in which nitrogen and phosphorus do not interact, for which only the (fixed) optimal ratios of consumption rates are needed (Tilman, 1982).

BGD

10, 3241–3279, 2013

A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Thus, if real phytoplankton are co-limited by nutrients across a range of input ratios, models featuring wide co-limitation zones (like ours or the one in Pahlow and Oschlies, 2009) will best describe competitive outcomes. Moreover, the cell plasticity included in our description allows for predictions about species dominance and co-existence in realistic, dynamic environments.

4.4 Experimental test

New experiments are needed to confirm the relationship proposed here between quota interactions, growth, uptake and, eventually, stoichiometry, as no clearly unequivocal and comprehensive experimental evidence exists yet. Such an experimental setup would use two phytoplankton strains, first studied independently and later in competition experiments⁵. To ensure the observation of possible wide co-limitation zones, species need to show broad quota ranges, which must be determined along with the maximum values of growth and uptake rates. Next is the determination of the stationary values of the quotas when the input ratio is varied across a wide range of dilution rates including values close to μ_{\max} . These experiments should give a clear picture of the different limitation zones, as well as an estimate of the optimal ratio for each species (Figs. 5 and 6). If enough pairs ($w, [N_0] : [P_0]$) are explored, the behavior of the populations with changes in the dilution rate can be also reconstructed (Figs. 7 and 8). Complemented with, e.g. measurements of RNA, these experiments would suffice to provide a clear picture of the cellular allocation strategies under different environmental conditions. As a last step, experiments can be performed where the two populations compete in chemostats at one (or several) selected dilution rates, and for different input nutrient ratios (see Fig. 9). The observation of exclusion in an a priori expected coexistence zone would evidence the existence of a broad co-limitation region. On the other hand, optimal ratios matching $Q_{N_{\min}}/Q_{P_{\min}}$ and framing the co-existence regions would

⁵In order to observe competition, one species must be more efficient at nitrogen uptake, and less efficient at phosphorus uptake.

A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



confirm the classical hypothesis (Rhee, 1978). Thus, these competition experiments serve to confirm the results of the experiments performed previously, and to alleviate a possible lack of resolution in the exploration of the $(w, [N_0] : [P_0])$ space.

5 Conclusions

5 Phytoplankton stoichiometry is a dynamic characteristic of the cells. It is the result of the allocation strategy adopted by the organism under different environmental conditions. Our model for phytoplankton stoichiometry focuses on protein regulation as vehicle for cell acclimation to different environments. As nitrogen and phosphorus are major regulators of protein synthesis, a correct description of the cell's resource allocation is essential for our model. Considering allocation leads us to introduce the simple
10 expression and repression functions, which not only allow for the dynamic regulation of proteins but also introduce other essential features such as interactions between quotas and with growth. In addition, they yield realistic predictions of important behaviors for the cell. Nonetheless, our model represents a phytoplankton cell with a stoichiometric phenomenology more complicated than that inferred from the classic Redfield and Droop work, due to the plasticity of the cell in response to changing nutrient availability. However, the cell's physiological ranges (e.g. maximum and minimum values of the quotas), which are ultimately determined by evolutionary processes, constrain the plasticity of the cell. Thus, cells are limited in their ability to match the stoichiometry
15 of their environment, especially during blooms (cells growing close to their maximum growth rate) or in cases of nutrient scarcity (vanishing cell growth). Moreover, our dynamic equations for the number of uptake proteins allow our model to predict how cells respond to changes in the environment. Therefore, our model is suited to describe more realistic situations in which competition occurs under changing ocean conditions,
20 for instance due to diurnal or seasonal variation. Our approach may thus help understand the biotic regulations of oceanic nitrogen and phosphorus, as well as the role of stoichiometry in shaping phytoplankton communities. Nonetheless, the experimental

A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



test proposed here should be performed in order to further validate this model and significantly advance knowledge of phytoplankton stoichiometry.

Appendix A

Deductions and definitions

5 The relationship between the number of proteins devoted to the uptake of nutrient X , n_X , and the maximum uptake rate for that nutrient, $V_{X_{\max}}$, is easily obtained from the deduction of the Michaelis–Menten functional form for the uptake rate (e.g. (Bonachela et al., 2011)). The deduction, based on the analogy between the uptake process and an enzymatic reaction, states that:

$$10 V_{X_{\max}}(t) = k_{2X} n_X(t)/(B(t) N_A), \quad (\text{A1})$$

where k_{2X} is the nutrient ion handling rate and N_A is Avogadro's number. The same enzymatic analogy establishes that (Bonachela et al., 2011):

$$k_{2X} = 4D_X r_X K_X N_A, \quad (\text{A2})$$

15 where K_X is the half-saturation constant for nutrient X , D_X is its diffusivity in the medium where it is dissolved, and r_X is the radius of reactive part of the uptake protein (see Table 3 for units and values).

Thus, if ν_X is the maximum number of uptake proteins for nutrient X that the cell can synthesize per unit time, the maximum change in $V_{X_{\max}}$ per unit time is given by:

$$\tilde{\nu}_X = 4D_X r_X K_X \nu_X. \quad (\text{A3})$$

20 The handling constant also plays a role in the determination of the total absorbing area for the cell and, therefore, in the ratio absorbing : total area. Assuming, for

BGD

10, 3241–3279, 2013

A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



simplicity, that $r_N \sim r_P = r_S$:

$$A_{\text{rel}}(t) = \frac{(n_N(t)/k_{2N} + n_P(t)/k_{2P})r_S^2}{4B(t)r_C^2}, \quad (\text{A4})$$

which is the main argument of the “competition-for-space” term of Eqs. (9) and (10), represented by the Heaviside function:

$$H(x) = 1 \text{ for } x \geq 0 \text{ (and 0 elsewhere)}. \quad (\text{A5})$$

Deduction of the optimal ratio: at $[N_0] : [P_0]_{\text{opt}}$, the stationary concentrations of both nutrients in the chemostat are negligible. Thus, from Eqs. (13) and (14), $V_N/V_P = [N_0]/[P_0]$. On the other hand, the optimal ratio is the only point where this happens even for the maximum value of the growth rate. At those growth rates, the cell quotas reach their maximum values, and both V_N and V_P equal their respective maximum uptake rates, which in turn take their lower values, $V_{\text{max}_N}^{\text{lo}}$ and $V_{\text{max}_P}^{\text{lo}}$. By using the stationary solution to Eqs. (3) and (4), and assuming loss rates much smaller than the dilution rate (or, alternatively, $R_N \sim R_P$), we find that $[N_0]/[P_0]_{\text{opt}} = V_{\text{max}_N}^{\text{lo}}/V_{\text{max}_P}^{\text{lo}} = Q_{N_{\text{max}}}^{\text{eff}}/Q_{P_{\text{max}}}^{\text{eff}}$. In our case, $Q_{N_{\text{max}}}^{\text{eff}}/Q_{P_{\text{max}}}^{\text{eff}} \sim 26$, which is in accordance with the values deduced from Fig. 6b.

Acknowledgements. We thank Cecilia Batmalle, Renaud Berlemont, Edward DeLong, John Dunne, Caroline Farior, Patricia Glibert, Christopher Klausmeier, Elena Litchman, Michael Lomas, Jorge Sarmiento, and Charles Stock for helpful discussions and suggestions. We gratefully acknowledge support from NSF under grant OCE-1046001. Part of this work was funded by the Cooperative Institute for Climate Science (CICS) of Princeton University and the National Oceanographic and Atmospheric Administration’s (NOAA) Geophysical Fluid Dynamics Laboratory (GFDL).

References

- Ågren, G. I.: The C : N : P stoichiometry of autotrophs – theory and observations, *Ecol. Lett.*, 7, 185–191, 2004. 3244
- Alon, U.: *An Introduction to Systems Biology: Design Principles of Biological Circuits*, Chapman & Hall/CRC Press, Boca Raton, FL, 2007. 3248
- 5 Armstrong, R. A.: Nutrient uptake rate as a function of cell size and surface transporter density: a Michaelis-like approximation to the model of Pasciak and Gavis, *Deep-Sea Res. Pt. I*, 55, 1311–1317, 2008. 3247
- Bonachela, J. A., Raghieb, M., and Levin, S. A.: Dynamic model of flexible phytoplankton nutrient uptake, *P. Natl. Acad. Sci. USA*, 108, 20633–20638, 2011. 3247, 3248, 3255, 3262, 3270
- 10 Bougaran, G., Bernard, O., and Sciandra, A.: Modeling continuous cultures of microalgae colimited by nitrogen and phosphorus, *J. Theor. Biol.*, 265, 443–454, 2010. 3244, 3249, 3256, 3257
- Caperon, J.: Time lag in population growth response of *Isochrysis galbana* to a variable environment, *Ecology*, 50, 188–192, 1969. 3257
- 15 Droop, M. R.: Vitamin B₁₂ and marine ecology IV: the kinetics of uptake, growth and inhibition in *Monochrysis lutheri*, *J. Mar. Biol. Assoc. UK*, 48, 689–733, 1968. 3243
- Dyhrman, S. T. and Palenik, B.: A single-cell immunoassay for phosphate stress in the dinoflagellate *Prorocentrum minimum* (Dinophyceae), *J. Phycol.*, 37, 400–410, 2001. 3248
- 20 Elrifi, I. R. and Turpin, D. H.: Steady state luxury consumption and the concept of optimum nutrient ratios: a study with phosphate and nitrate limited *Selenastrum minutum* (Chlorophyta), *J. Phycol.*, 21, 592–602, 1985. 3243, 3256, 3258, 3268
- Flynn, K. J.: Use, abuse, misconceptions and insights from quota models – the Droop Cell Quota Model 40 years on, *Oceanogr. Mar. Biol.*, 46, 1–23, 2008. 3248
- 25 Flynn, K. J., Raven, J. A., Rees, T. A. V., Finkel, Z. V., Quigg, A., and Beardall, J.: Is the growth rate hypothesis applicable to microalgae?, *J. Phycol.*, 46, 1–12, 2010. 3270
- Geider, R. J. and La Roche, J.: Redfield revisited: variability of C : N : P in marine microalgae and its biochemical basis, *Eur. J. Phycol.*, 37, 1–17, 2002. 3242, 3249
- Geider, R. J., MacIntyre, H. L., and Kana, T. M.: A dynamic regulatory model of phytoplanktonic acclimation to light, nutrients, and temperature, *Limnol. Oceanogr.*, 43, 679–694, 1998. 3251
- 30 Goldman, J. C., McCarthy, J. J., and Peavey, D. G.: Growth rate influence on the chemical composition of phytoplankton in oceanic waters, *Nature*, 279, 210–215, 1979. 3242, 3268

A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Gotham, I. J. and Rhee, G. Y.: Comparative kinetic studies of nitrate-limited growth and nitrate uptake in phytoplankton in continuous culture, *J. Phycol.*, 17, 309–314, 1981a. 3244, 3248, 3268

Gotham, I. J. and Rhee, G. Y.: Comparative kinetic studies of phosphate-limited growth and phosphate uptake in phytoplankton in continuous culture, *J. Phycol.*, 17, 257–265, 1981b. 3244, 3248, 3268

Göthlich, L. and Oschlies, A.: Phytoplankton niche generation by interspecific stoichiometric variation, *Global Biogeochem. Cy.*, 26, GB2010, 1–8, 2012. 3243

Hall, S. R., Smith, V. H., Lytle, D. A., and Leibold, M. A.: Constraints on primary producer N : P stoichiometry along N : P supply ratio gradients, *Ecology*, 86, 1894–1904, 2005. 3256, 3269

Healey, F. P.: Interacting Effects of light and nutrient limitation on the growth rate of *Synechococcus linearis* (Cyanophyceae), *J. Phycol.*, 21, 134–146, 1985. 3256, 3258, 3268, 3270

Healey, F. P. and Hendzel, L. L.: Indicators of phosphorus and nitrogen deficiency in five algae in culture, *J. Fish. Res. Board Can.*, 36, 1364–1369, 1979. 3268

Hense, I. and Beckmann, A.: Towards a model of cyanobacteria life cycle-effects of growing and resting stages on bloom formation of N₂-fixing species, *Ecol. Model.*, 195, 205–218, 2006. 3270

Ikeya, T., Ohki, K., Takahashi, M., and Fujita, Y.: Study on phosphate uptake of the marine cyanophyte *Synechococcus sp.* NIBB 1071 in relation to oligotrophic environments in the open ocean, *Mar. Biol.*, 129, 195–202, 1997. 3270

Klausmeier, C. A., Litchman, E., Daufresne, T., and Levin, S. A.: Optimal nitrogen-to-phosphorus stoichiometry of phytoplankton, *Nature*, 429, 171–174, 2004a. 3243, 3245, 3256, 3258

Klausmeier, C. A., Litchman, E., and Levin, S. A.: Phytoplankton growth and stoichiometry under multiple nutrient limitation, *Limnol. Oceanogr.*, 49, 1463–1470, 2004b. 3243, 3256, 3257, 3258, 3269, 3272

Klausmeier, C. A., Litchman, E., and Levin, S. A.: A model of flexible uptake of two essential resources, *J. Theor. Biol.*, 246, 278–289, 2007. 3243, 3256, 3257, 3258

Kunikane, S., Kaneko, M., and Maehara, R.: Growth and nutrient uptake of green alga, *Scenedesmus dimorphus*, under a wide range of nitrogen/phosphorus ratio – I Experimental study, *Water Res.*, 18, 1299–1311, 1984. 3258, 3268, 3269

A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



- Legović, T. and Cruzado, A.: A model of phytoplankton growth on multiple nutrients based on the Michaelis–Menten–Monod uptake, Droop's growth and Liebig's law, *Ecol. Model.*, 99, 19–31, 1997. 3243
- 5 McCarthy, J. J. and Goldman, J. C.: Nitrogenous nutrition of marine phytoplankton in nutrient-depleted waters, *Science*, 203, 670–672, 1979. 3244, 3248
- Mierle, G.: Kinetics of phosphate transport by *Synechococcus leopoliensis* (Cyanophyta): evidence for diffusion limitation of phosphate uptake, *J. Phycol.*, 21, 177–181, 1985. 3247
- Morel, F. M. M.: Kinetics of nutrient uptake and growth in phytoplankton, *J. Phycol.*, 23, 137–150, 1987. 3244, 3248
- 10 Pahlow, M. and Oschlies, A.: Chain model of phytoplankton P, N and light co-limitation, *Mar. Ecol.-Prog. Ser.*, 376, 69–83, 2009. 3244, 3245, 3256, 3257, 3258, 3260, 3270
- Pasciak, W. J. and Gavis, J.: Transport limitation of nutrient uptake in phytoplankton, *Limnol. Oceanogr.*, 19, 881–888, 1974. 3247
- Rhee, G. Y.: A continuous culture study of phosphate uptake, growth rate and polyphosphate in *Scenedesmus Sp.*, *J. Phycol.*, 9, 495–506, 1973. 3268
- 15 Rhee, G. Y.: Effects of N : P atomic ratios and nitrate limitation on algal growth, cell compositions, and nitrate uptake, *Limnol. Oceanogr.*, 23, 10–25, 1978. 3242, 3243, 3253, 3255, 3256, 3259, 3261, 3269
- Rhee, G. Y. and Gotham, I. J.: Optimum N : P ratios and coexistence of planktonic algae, *J. Phycol.*, 16, 486–489, 1980. 3258
- 20 Riegman, R. and Mur, L. R.: Regulation of phosphate uptake kinetics in *Oscillatoria agardhii*, *Arch. Microbiol.*, 139, 28–32, 1984. 3248
- Roelke, D. L., Eldridge, P. M., and Cifuentes, L. A.: A model of phytoplankton competition for limiting and nonlimiting nutrient: implications for development of estuarine and nearshore management schemes, *Estuaries*, 22, 92–104, 1999. 3244
- 25 Saito, M. A., Goepfert, T. J., and Ritt, J. T.: Some thoughts on the concept of colimitation: three definitions and the importance of bioavailability, *Limnol. Oceanogr.*, 53, 276–290, 2008. 3244
- Smith, S. L. and Yamanaka, Y.: Optimization-based model of multinutrient uptake kinetics, *Limnol. Oceanogr.*, 52, 1545–1558, 2007. 3244, 3258
- 30 Song, B. and Ward, B. B.: Molecular cloning and characterization of high affinity nitrate transporters in marine phytoplankton, *J. Phycol.*, 43, 542–552, 2007. 3248, 3257
- Sterner, R. W. and Elser, J. J.: *Ecological Stoichiometry: The Biology of Elements from Molecules to the Biosphere*, Princeton University Press, Princeton, NJ, 2002. 3249

- Terry, K. L.: Nitrate and phosphate uptake interactions in a marine prymnesiophyte, *J. Phycol.*, 18, 79–86, 1982. 3244, 3256, 3268
- Terry, K., Laws, E. A., and Burns, D. J.: Growth rate variation in the N : P requirement ratio of phytoplankton, *J. Phycol.*, 21, 323–329, 1985. 3243, 3255, 3258, 3268
- 5 Tilman, D.: *Resource Competition and Community Structure*, Princeton University Press, Princeton, NJ, 1982. 3258, 3259

A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.

Table 1. Trend (increasing, \uparrow , or decreasing, \downarrow , function) and shape reported in the literature for the cell content (quota, Q) of nitrogen and phosphorus when the dilution rate w is varied in laboratory experiments.

Plot	Trend when N lim. (shape)	Trend when P lim. (shape)	Reference
μ vs Q_X	\uparrow with Q_N (hyp.)	\uparrow with Q_P (hyp.)	Rhee (1973); Gotham and Rhee (1981a,b); Elrifi and Turpin (1985)
Q_N vs w	\uparrow (line or curve)	\uparrow (curve)	Healey and Hendzel (1979); Terry (1982); Kunikane et al. (1984); Terry et al. (1985); Healey (1985); Elrifi and Turpin (1985)
Q_P vs w	\downarrow or no trend (line or curve)	\uparrow (curve)	Goldman et al. (1979); Healey and Hendzel (1979); Terry (1982); Kunikane et al. (1984); Healey (1985); Terry et al. (1985); Elrifi and Turpin (1985)
$Q_N : Q_P$ vs w	\uparrow or no trend (curve)	\downarrow (line or curve)	Goldman et al. (1979); Healey and Hendzel (1979); Terry (1982); Elrifi and Turpin (1985)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



BGD

10, 3241–3279, 2013

A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)**Table 2.** Trend and shape reported in experiments for Q when the input ratio $[N_0] : [P_0]$ is varied.

Plot	Trend (shape)	Ref.
Q_N vs $[N_0] : [P_0]$	Increasing (line or curve)	Kunikane et al. (1984); Rhee (1978)
Q_P vs $[N_0] : [P_0]$	Decreasing (curve)	Kunikane et al. (1984); Rhee (1978)
$Q_N : Q_P$ vs $[N_0] : [P_0]$	Increasing (line)	Klausmeier et al. (2004b), after Rhee (1978); Hall et al. (2005)



Table 3. List of variables and parameters used in this manuscript, representing a generic *Synechococcus* strain – values collected from Healey (1985); Hense and Beckmann (2006); Ikeya et al. (1997); Pahlow and Oschlies (2009); Flynn et al. (2010). For reasoning on the value (or deduction) of the remaining parameters, see text or (Bonachela et al., 2011).

Symbol	a.k.a.	Units	Value
[N]	Dissolved Inorganic Nitrogen Concentration	mol L ⁻¹	Variable
[P]	Dissolved Inorganic Phosphorus Concentration	mol L ⁻¹	Variable
[N ₀]	Dissolved Inorganic Nitrogen Supply Concentration	mol L ⁻¹	15 – 240 × 10 ⁻⁶
[P ₀]	Dissolved Inorganic Phosphorus Supply Concentration	mol L ⁻¹	3 × 10 ⁻⁶
D _N	Nitrogen Diffusion Constant in Water	m ² d ⁻¹	1.296 × 10 ⁻⁴
D _P	Phosphorus Diffusion Constant in Water	m ² d ⁻¹	8.64 × 10 ⁻⁵
w	Chemostat Dilution Rate	d ⁻¹	0.01 – 0.825
C	Population (Organic) Carbon Concentration	mol L ⁻¹	Variable
N	Population (Organic) Nitrogen Concentration	mol L ⁻¹	Variable
P	Population (Organic) Phosphorus Concentration	mol L ⁻¹	Variable
B	Number of Cells in Chemostat	cells	Variable
μ	Population Growth Rate	d ⁻¹	Variable
μ _{max}	Maximum Population Growth Rate	d ⁻¹	Emergent
V _N	Population N-Uptake Rate	mol L ⁻¹ d ⁻¹	Variable
V _P	Population P-Uptake Rate	mol L ⁻¹ d ⁻¹	Variable
V _{N,max}	Maximum cell N-Uptake Rate	mol cell ⁻¹ d ⁻¹	Variable
V _{P,max}	Maximum cell P-Uptake Rate	mol cell ⁻¹ d ⁻¹	Variable
V _{N,min}	Minimum value for max. N-Uptake Rate	mol cell ⁻¹ d ⁻¹	Emergent
V _{P,min}	Minimum value for max. P-Uptake Rate	mol cell ⁻¹ d ⁻¹	Emergent
K _N	Half-Saturation Constant for N	mol L ⁻¹	0.3 × 10 ⁻⁶
K _P	Half-Saturation Constant for P	mol L ⁻¹	10 × 10 ⁻⁹
K _N ^{eff}	Effective Half-Saturation Constant for N	mol L ⁻¹	Variable
K _P ^{eff}	Effective Half-Saturation Constant for P	mol L ⁻¹	Variable
Q _N	Nitrogen Cell Quota	mol cell ⁻¹	Variable
Q _P	Phosphorus Cell Quota	mol cell ⁻¹	Variable
Q _{N,max}	Maximum N Quota	mol cell ⁻¹	2.15 × 10 ⁻¹⁴
Q _{N,min}	Minimum N Quota	mol cell ⁻¹	6.14 × 10 ⁻¹⁵
Q _{P,max}	Maximum P Quota	mol cell ⁻¹	8.23 × 10 ⁻¹⁶
Q _{P,min}	Minimum P Quota	mol cell ⁻¹	1.11 × 10 ⁻¹⁶
Q _C	Carbon Content per Cell	mol cell ⁻¹	8.33 × 10 ⁻¹⁴
k _{S,N}	Handling Rate for N	d ⁻¹	From Eq. (A2)
k _{S,P}	Handling Rate for P	d ⁻¹	From Eq. (A2)
v _N	Max. N-uptake Sites Synthesis Rate	sites cell ⁻¹ d ⁻¹	10 ⁴
v _P	Max. P-uptake Sites Synthesis Rate	sites cell ⁻¹ d ⁻¹	10 ⁴
V̇ _N	Max. Change of V _{max,N} per Unit Time	mol cell ⁻¹ d ⁻²	From Eq. (A3)
V̇ _P	Max. Change of V _{max,P} per Unit Time	mol cell ⁻¹ d ⁻²	From Eq. (A3)
A _{rel}	Ratio Absorbing:Total Area	–	Variable
r _c	Cell Radius	m	0.82 × 10 ⁻⁶
r _s	Uptake Site Radius	m	2.5 × 10 ⁻⁹
P _{max}	Maximum Photosynthetic Rate	d ⁻¹	5
M _C	Uptake Maintenance Cost	–	2
R _C	Respiration Cost	d ⁻¹	0
R _N	Loss Rate	d ⁻¹	0.2
R _P	Loss Rate	d ⁻¹	0.2
F	Protein Expression Function	–	Variable
G	Protein Repression Function	–	Variable
H	Heaviside (or Step) Function	–	Variable
k _F	Sigmoid Slope Parameter for F	–	5
k _{G,1}	Sigmoid Slope Parameter for G	–	20
k _{G,2}	Sigmoid Shift Parameter for G	–	0.25
N _A	Avogadro Number	Units mol ⁻¹	6.02 × 10 ²³

A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.




Fig. 1. Patterns for growth rate μ and quotas Q observed in the experimental literature (see Tables 1 and 2). In panels **a** to **c**, when the behavior observed under N limitation is different from that under P limitation, two colors and line style are used, with green dashed lines indicating phosphorus limitation. **(a)** Saturating form for the growth rate dependence on limiting nutrients. **(b)** And **(c)** dependence of the quotas on dilution rate, with asymmetric behavior for N limitation. **(d)** Response of phytoplankton N : P ratio to changes in the input ratio; the black line represents $N : P = [N_0] : [P_0]$. See Tables 1 and 2 for references.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.




Fig. 2. Curves for growth rate μ , quotas Q , and N : P ratio obtained with a threshold model (Klausmeier et al., 2004b). **(a)** The growth rate depends only on Q_X in the X -limited regime, and the shape of the function (hyperbolic) does not depend on which element is limiting. **(b)** The use of an identical functional form for the dependence of growth on the limiting quota facilitates a similar behavior for Q_N and Q_P (symmetry between N and P quotas). **(c)** Phytoplankton N : P equals $[N_0] : [P_0]$ only when the growth rate is far from its maximum (small dilution rates). **(d)** Both nitrogen and phosphorus are essential resources, and as a consequence co-limitation is reached only at one point: the optimal ratio. For the parametrization used here, which corresponds to *Scenedesmus* sp., this point is $[N_0] : [P_0]_{opt} = 27.7$.

A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.




Fig. 3. Contribution of the nutrient quotas to the synthesis of nitrogen **(a)** and phosphorus **(b)** uptake proteins, summarized here by $\text{Prot}_N(Q_N, Q_P) = F \left(\frac{Q_{N,\max} - Q_N(t)}{Q_{N,\max} - Q_{N,\min}} \right) G \left(\frac{Q_P(t) - Q_{P,\min}}{Q_{P,\max} - Q_{P,\min}} \right)$ and $\text{Prot}_P(Q_P) = F \left(\frac{Q_{P,\max} - Q_P(t)}{Q_{P,\max} - Q_{P,\min}} \right) G \left(\frac{Q_P(t) - Q_{P,\min}}{Q_{P,\max} - Q_{P,\min}} \right)$.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.




Fig. 4. (a) Dependence of the photosynthetic term on the N and P quotas, summarized here by $\text{Photo}(Q_N, Q_P) = F \left(\frac{Q_N(t) - Q_{N\min}}{Q_{N\max} - Q_{N\min}} \right) G \left(\frac{Q_P(t) - Q_{P\min}}{Q_{P\max} - Q_{P\min}} \right)$. (b) Flow diagram describing the different physiological processes in the model: Inorganic nutrients ($[N]$ and $[P]$) are taken up at rates V_N and V_P , respectively, and assimilated as quotas, Q_N and Q_P . Quotas are the main influential factor for the synthesis of uptake and photosynthetic proteins, through the expression and repression functions, F and G . Photosynthetic proteins participate through photosynthesis in cell growth, which is also affected by the expenditure in the maintenance of the different apparatuses. In this model, we mainly consider uptake costs, through the terms V_N/N and V_P/P . Finally, changes in the number of uptake proteins (i.e. in V_{\max}) influence the uptake of the corresponding nutrient. See text for further details.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.




Fig. 5. Variation of N quota (main panel in **a**) and P quota (inset in panel **a**) with the supply ratio, for different values of the dilution rate. Effective maximum quota values are shown as dashed lines. **(b)** Phosphorus quota versus nitrogen quota for different dilution rates; single-limitation zones show constant values for the quota of the limiting nutrient.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.




Fig. 6. (a) N : P ratio for the cell as a function of the nutrient supply ratio for different chemostat dilution rates. (b) Identification of the optimal ratio as the input ratio for which the cell takes up nutrients at the supply ratio for any dilution rate; for the generic *Synechococcus* parametrization used here, $[N_0] : [P_0]_{opt} \sim 26$.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.




Fig. 7. Upper row: variation of the nitrogen and phosphorus quotas with dilution rate (dashed line: maximum effective values). **(a)** Q_N shows linear behavior or convexity for the different input ratios; for very high ratios, the quota saturates at its maximum value. **(b)** Q_P shows a similar behavior, but non-monotonicity is also possible for high relative phosphorus concentrations (low input ratio); that is the case of the $[N_0]:[P_0] = 16$ curve at high dilution rates, although the parametrization used here complicates the observation of that behavior (small values of the half-saturation constant, for instance, provide a more marked non-monotonicity, with clearer decline regimes for Q_P). Lower row: Dependence of the growth rate μ on nutrient quota; only the range of ratios for which Q_N **(c)** or Q_P **(d)** influence μ are shown in each panel (dashed lines: minimum value for the quotas, Q_{min}). Note the saturating shape of μ , reaching a maximum $\mu_{max} = 0.825 \pm 0.005$. For extreme input ratios, curves do not show any dependence on the input ratio (e.g. $[N_0]:[P_0] < 22$ in **c**). For intermediate input ratios, the value of the quotas at which μ reaches its minimum and maximum values change with the nutrient relative input concentration.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.




Fig. 8. Maximum uptake rate V_{\max} and *per-cell* uptake rate V/B for nitrogen **(a)** and phosphorus **(b)** as a function of the external nutrient concentration in the zone of co-limitation (axes in logarithmic scale). V_{\max} decreases with increasing nutrient availability due to the down-regulation of the uptake proteins in the population. V_{\max} and V/B converge for large nutrient concentrations (dashed lines, see text).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A model for variable phytoplankton stoichiometry

J. A. Bonachela et al.




Fig. 9. Different potential outcomes of competition experiments between two generic phytoplankton species, A and B. The labeled zones represent situations in which there is a priori opportunity for co-existence, while the non-labeled zones are those in which the outcome depends on each species' $\Delta = [(N : P)_{opt} - (N : P)] / (N : P)_{opt}$. **(a)** For non-interacting nutrients, each species' optimal ratio $[N_0] : [P_0]_{opt}$ matches the point where co-limitation occurs; the optimal ratios of the two species divide the phase diagram, delimiting the co-existence (grey) and exclusion (blank) zones. **(b)** In our model, the optimal ratio is one specific point within a wide co-limitation zone, which is delimited by the ratios at which quotas reach their effective maximum values (saturation points). This introduces exclusion in zones where threshold models predict co-existence (color patters, see above).

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

