

This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Review article

“Air-sea exchanges of CO₂ in world’s coastal seas”

C.-T. A. Chen^{1,2,3}, T.-H. Huang¹, Y.-C. Chen¹, Y. Bai⁴, X. He⁴, and Y. Kang⁴

¹Institute of Marine Geology and Chemistry, National Sun Yat-sen University,
Kaohsiung, Taiwan

²Institute of Marine Resources, Zhejiang University, Hangzhou, China

³The International Institute for Carbon-Neutral Energy Research (I²CNER), Kyushu University,
Fukuoka, Japan

⁴Second Institute of Oceanography, State Oceanic Administration, Hangzhou, China

Received: 28 January 2013 – Accepted: 2 February 2013 – Published: 13 March 2013

Correspondence to: C.-T. A. Chen (ctchen@mail.nsysu.edu.tw)

Published by Copernicus Publications on behalf of the European Geosciences Union.

BGD

10, 5041–5105, 2013

Air-sea Exchanges
of CO₂ in World’s¹
Coastal Seas

C.-T. A. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

The air-sea exchanges of CO₂ in the world's 165 estuaries and 87 continental shelves are evaluated. Generally and in all seasons, upper estuaries with salinities of less than two are strong sources of CO₂ ($39 \pm 56 \text{ mol C m}^{-2} \text{ yr}^{-1}$, negative flux indicates that the water is losing CO₂ to the atmosphere); mid-estuaries with salinities of between 2 and 25 are moderate sources ($17.5 \pm 34 \text{ mol C m}^{-2} \text{ yr}^{-1}$) and lower estuaries with salinities of more than 25 are weak sources ($8.4 \pm 14 \text{ mol C m}^{-2} \text{ yr}^{-1}$). With respect to latitude, estuaries between 23.5 and 50° N have the largest flux per unit area ($63 \pm 101 \text{ mmol C m}^{-2} \text{ d}^{-1}$); these are followed by mid-latitude estuaries (23.5–10° S: $44 \pm 29 \text{ mmol C m}^{-2} \text{ d}^{-1}$; 0–23.5° N: $39 \pm 55 \text{ mmol C m}^{-2} \text{ d}^{-1}$), and then regions north of 50° N ($36 \pm 91 \text{ mmol C m}^{-2} \text{ d}^{-1}$). Estuaries south of 50° S have the smallest flux per unit area ($9.5 \pm 12 \text{ mol C m}^{-2} \text{ d}^{-1}$). Mixing with low-*p*CO₂ shelf waters, water temperature, residence time and the complexity of the biogeochemistry are major factors that govern the *p*CO₂ in estuaries but wind speed, seldom discussed, is critical to controlling the air-water exchanges of CO₂. The total annual release of CO₂ from the world's estuaries is now estimated to be 0.10 PgC yr^{-1} , which is much lower than published values mainly because of the contribution of a considerable amount of heretofore unpublished or new data from Asia and the Arctic. The Asian data, although indicating high in *p*CO₂, are low in sea-to-air fluxes because the wind speeds are lower than previously determined values, which rely heavily on data from Europe and North America, where *p*CO₂ is lower but wind speeds are much higher, such that the CO₂ fluxes are higher than in Asia. Newly emerged CO₂ flux data in the Arctic reveal that estuaries there mostly absorb, rather than release CO₂.

Most continental shelves, and especially those at high latitude, are under-saturated in terms of CO₂ and absorb CO₂ from the atmosphere in all seasons. Shelves between 0° and 23.5° S are on average a weak source and have a small flux per unit area of CO₂ to the atmosphere. Water temperature, the spreading of river plumes, upwelling, and biological production seem to be the main factors in determining *p*CO₂ in the shelves.

BGD

10, 5041–5105, 2013

Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Wind speed, again, is critical because at high latitudes, the winds tend to be strong. Since the surface water $p\text{CO}_2$ values are low, the air-to-sea fluxes are high in regions above 50° N and below 50° S. At low latitudes, the winds tend to be weak, so the sea-to-air CO_2 flux is small. Overall, the world's continental shelves absorb 0.4 PgC yr^{-1} from the atmosphere.

5

1 Introduction

Carbon is arguably one of the most important elements on earth and understanding the global carbon cycle is fundamental to elucidating the effect of human activities in the Anthropocene era. The oceans are known to have an important role in regulating the climate on annual to millennial scales by absorbing CO_2 and exchanging carbon with various carbon-storing compartments, such as the atmosphere, the land, the biota and the fossil fuel carbon pool. Yet, despite the success of quantifying the air-sea CO_2 exchange and the uptake of anthropogenic CO_2 by the major oceans, effect of the land on these processes is still poorly understood and little discussed (Khatiwala et al., 10 2012; Le Quéré et al., 2012; Schuster et al., 2012; Wanninkhof et al., 2012).

15

Coastal waters link the land, the oceans, the atmosphere, biota and sediments. Although they constitute only a little over 7 % of the surface area of oceans and less than 0.5 % of the volume of the oceans, coastal oceans have a disproportionately large role in primary and new production, remineralization and sedimentation of organic matter (Walsh et al., 1981; Walsh, 1988, 1991; Kempe and Pegler, 1991; Mackenzie et al., 20 1991, 1998a, b; Chen, 1993; Wollast, 1993, 1998; Gattuso et al., 1998; Carrillo and Karl, 1999; Liu et al., 2000; de Haas et al., 2002; Elliott and McLusky, 2002; Muller-Karger et al., 2005; Thomas, 2010). Coastal waters receive large inputs of terrestrial material, such as suspended sediments and nutrients in solution or in particulate matter, in organic or inorganic forms and through river and groundwater discharge, as 25 well as by exchange with the atmosphere and the open ocean. They therefore tend to show greater temporal and spatial variability than open oceans, and are more affected

BGD

10, 5041–5105, 2013

Air-sea Exchanges of CO_2 in World's Coastal Seas

C.-T. A. Chen et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



by human activities (Cameron and Pritchard, 1963; Alongi, 1998; Chen and Tsunogai, 1998; Rabouille et al., 2001; Chen, 2002, 2003, 2004; Slomp and Van Cappellen, 2004; Beusen et al., 2005; Chavez et al., 2007; Doney et al., 2007; Radach and Patsch, 2007; Peng et al., 2008; Seitzinger et al., 2010; Durr et al., 2011; Jiang et al., 2012).

5 However, unlike the open oceans, in which millions of observations have been made and the air-sea exchanges of CO₂ have been valued using various developed models (such as by Khatiwala et al., 2012; Schuster et al., 2012; Wanninkhof et al., 2012), coastal waters have been relatively poorly examined.

Although estuaries are known to be generally sources of CO₂ (Frankignoulle et al., 10 Cai et al., 1999, 2000; Sarma et al., 2001, 2011; Abril et al., 2002; Borges et al., 2003; Dagg et al., 2005; Gao et al., 2005; Dai et al., 2008; Leinweber et al., 2009), only in the last few years have continental shelves been firmly established to absorb CO₂ from the atmosphere. (See, for example, Liu et al., 2000; Chen et al., 2003; Chen, 2004; Abril and Borges, 2005; Borges, 2005; Borges et al., 2005; Cai et al., 2006; Chen and Borges, 2009; Laruelle et al., 2010 and references therein.) Indeed, whether coastal seas are sources or sinks of CO₂ has remained an open question until only recently. The first report of the project on the Land Ocean Interaction in the Coastal Zone under the International Geosphere Biosphere Programme (IGBP) is entitled, "Coastal seas: a net source or sink of atmosphere carbon dioxide" (Kempe, 1995). The first report 15 of LOICZ did not provide any data concerning the air-sea exchanges of carbon in the continental margins, although it concluded that net carbon oxidation in the coastal zone is around 7×10^{12} mol yr⁻¹ (Crossland et al., 2005), implying that the coastal zone is a source of CO₂ to the atmosphere.

Unfortunately, Fasham et al. (2001), summarizing the work of the Joint Global Ocean 20 Flux Study (JGOFS, another IGBP project), concluded that there is a net sea-to-air CO₂ flux from continental margins of 0.5 PgC yr⁻¹. They drew this conclusion despite the fact that, at the time, the Joint JGOFS/LOICZ Continental Margins Task Team (Chen et al., 1994) had already gathered sufficient data to demonstrate that, the margins, rather than being a source of CO₂, are in fact a sink of CO₂. Indeed, in the same year, Fasham

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

published another paper that claimed that the continental shelves are actually a sink of CO₂ of the order of 0.6 PgC yr⁻¹ (Yool and Fasham, 2001). In 2003, the JGOFS also concluded that the shelves take up 0.3 PgC yr⁻¹ of atmospheric CO₂ (Chen et al., 2003). This view however, was not universally accepted (Cai et al., 2003; Cai and Dai, 2004) until more data, especially data obtained in the winter, became available. Many shelves that had been thought to be sources of CO₂ are now known to be sinks of CO₂ when winter data reveal severe under-saturation of CO₂ (Thomas et al., 2004; Cai et al., 2006; Schiettecatte et al., 2007; Jiang et al., 2008b).

Strangely, despite the fact that coastal waters play a major role in the livelihood of humans, and are strongly affected by human activities, our understanding of these waters is mostly semi-quantitative. For example, such basic information as the area of the continental shelf is uncertain. The most recent work of Y. Kang (private communication, 2013) yielded an area of 26.15×10^6 km² for waters shallower than 200 m. This value compares with 26.39×10^6 km² obtained by Laruelle et al. (2012), 24.72×10^6 km² presented by Laruelle et al. (2010), 30.16×10^6 km² obtained by Jahnke (2010), 26×10^6 km² presented by Chen and Borges (2009), 25.83×10^6 km² presented by Cai et al. (2006) and 36×10^6 km² presented by Liu et al. (2000). Merely comparing the total flux across various studies may not be very useful whereas comparing flux per unit area eliminates the problem of an uncertain global shelf area, which varies by as much as 50 % among studies. Even more strangely, despite the fact that rivers export approximately 1 PgC yr⁻¹ (Meybeck, 1982), or roughly half of the carbon that is absorbed by the open oceans each year, this value needs to be confirmed as it was based only on a few studies, and the well regarded study of Meybeck was based on a database of only 27 rivers.

The export of carbon by rivers comprises 40 % organic carbon (0.22 PgC yr⁻¹ of dissolved organic carbon (DOC) and 0.18 PgC yr⁻¹ of particulate organic carbon (POC)) and 60 % inorganic carbon (0.43 PgC yr⁻¹ of dissolved inorganic carbon (DIC) and 0.17 PgC yr⁻¹ of particulate inorganic carbon (PIC)) (Meybeck, 1982; Richey, 2004; IPCC, 2007; Schlunz and Schneider, 2000; Dai et al., 2012; Huang et al., 2012).

Hoever, estuarine filtering prevents some of the carbon that reaches the estuaries from also entering the oceans (Keil et al., 1997; Kemp et al., 1997; Middelburg and Herman, 2007; Chen et al., 2012; Dai et al., 2012). Whether speciation changes between the organic and inorganic or dissolved and particulate carbon in the estuaries, and how much of each of these forms of carbon actually enters the oceans are yet unknown (Woodwell et al., 1973; Raymond and Bauer, 2000; Wiegner and Seitzinger, 2001).

The above may be summarized by noting that nutrients from land, which may be transported by rivers or submarine groundwater discharge, or may be atmospheric fall-out, markedly affect estuaries and continental shelves (Ittekkot et al., 1991; Cole and Caraco, 2001; Neubauer and Anderson, 2003; Clark et al., 2004; Thomas et al., 2004; Gazeau et al., 2005; Hales et al., 2008; Jiang et al., 2012; Lauerwald et al., 2012). Consequently, estuaries and proximal continental shelves typically sustain high biological productivity (Walsh et al., 1981; Wollast, 1993, 1998; Cai, 2003) which may draw down CO₂. This phenomenon, however, may be more than counteracted by enhanced heterotrophic activity, supported by organic carbon input from rivers (Smith and Hollibaugh, 1993; Heip et al., 1995; Hedges and Keil, 1995; Hedges et al., 1997; Hansell and Carlson, 1998; Bouillon et al., 2006; Jiang et al., 2010). Additionally, direct inorganic carbon input from river water, submarine groundwater discharge and exchanges with tidal marshes and mangroves play an important role in increasing the pCO₂ of estuarine and shelf waters (Moran et al., 1991; Miller and Moran, 1997; Neal et al., 1998; Raymond et al., 2000; Raymond and Bauer, 2001; Borges et al., 2003, 2006; Cai et al., 2003; Wang and Cai, 2004; Jahnke et al., 2005; Bouillon et al., 2008; Jiang et al., 2008, 2010; Chen et al., 2012).

Since the above complex and conflicting factors influence the pCO₂ of estuarine and shelf waters, the air-sea exchanges of CO₂ in these waters can not yet be estimated by models, and so field data are required. Determinations of the air-sea flux of CO₂ in the world's estuaries and continental shelves, based on direct measurements, are presented below. Data from the literature and some unpublished data from CTA Chen are tabulated. Data for upper, mid and lower estuaries are compared. Seasonal and

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

|◀

▶|

◀

▶|

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



latitudinal variations are discussed and the global flux is presented. Data concerning continental shelves are also considered with reference to season and latitude before the global flux is determined.

BGD

10, 5041–5105, 2013

2 Air-sea CO₂ exchanges in Estuaries

- 5 Rivers are the main sources of carbon to the estuaries. Riverine organic carbon is supplied primarily by the erosion of soil organic matter or plant detritus (allochthonous) and by phytoplankton in water (autochthonous). The inorganic carbon is derived mainly from soil and rock erosion, and by the oxidation of organic matter mostly through microbial processes (Odum and Hoskin, 1958; Odum and Wilson, 1962; Probst et al., 1994;
- 10 Neal et al., 1998; Nelson et al., 1999; Pomeroy et al., 2000). These organic and inorganic forms of carbon in dissolved and particulate phases, reach the estuaries, which are typically wider than river channels. Therefore, particles tend to settle down and decompose, releasing carbon back into the water. Salt marshes, mangroves, and submarine groundwater discharge also export carbon to estuaries, increasing their $p\text{CO}_2$.
- 15 Rivers are the main sources of nutrients to estuaries. However, high turbidity and limited light cause nutrients rarely to be fully utilized for biological production in rivers or estuaries. Hence, the biological drawdown of CO₂ does not suffice to reduce the estuarine water $p\text{CO}_2$ to below saturation. Consequently, almost all estuaries are sources of CO₂ to the atmosphere. The influence of freshwater from large rivers frequently
- 20 extends hundreds of kilometers offshore. The enormous discharge of freshwater, sediments and the associated particulate and dissolved organic and inorganic carbon, nitrogen and phosphorus all greatly affect the biological and geochemical processes in the estuary, the plume and the adjacent continental shelf (Chen and Wang, 1999; Gong et al., 2000; Chen et al., 2003). A saline interface normally separates the plume water from the shelf water, with the width of the interface determined by interactions between river discharge and marine driving forces (Shen, 2001; Shen et al., 2003; Chen et al., 2008a). Complex biophysical and geochemical processes govern the direction of

CO_2 exchange between the plume-affected shelf area and the atmosphere (Kortzinger, 2003; Cooley et al., 2007), but in this investigation, river plumes outside of the estuaries are not considered.

Numerical data are gathered for 165 estuaries. They include 50 heretofore unpublished data from CTA Chen (Fig. 1, Table 1). The Wanninkhof (1992) quadratic equation is used to determine the CO_2 flux per unit area. That is, the flux is a function of the square of the wind speed. Figure 2 presents the $p\text{CO}_2$ and CO_2 fluxes per unit area in the upper, mid and lower estuaries worldwide. Almost all estuaries outside of the Arctic region except for only a few release CO_2 to the atmosphere. Unsurprisingly, upper estuaries, where the riverine effect is the strongest (Kempe, 1979, 1982; Chen et al., 2012), have the highest $p\text{CO}_2$ ($5026 \pm 6190 \mu\text{atm}$) and the highest sea-to-air CO_2 flux ($39.0 \pm 55.7 \text{ mol C m}^{-2} \text{ yr}^{-1}$, where the negative sign indicates that the seawater is losing CO_2); these are followed by the mid estuaries ($p\text{CO}_2 = 2230 \pm 2725 \mu\text{atm}$; flux = $17.5 \pm 34.2 \text{ mol C m}^{-2} \text{ yr}^{-1}$). Lower estuaries have the lowest $p\text{CO}_2$ ($723 \pm 957 \mu\text{atm}$) and CO_2 flux ($8.4 \pm 14.3 \text{ mol C m}^{-2} \text{ yr}^{-1}$). Except for those of the upper estuaries, these $p\text{CO}_2$ values compare favorably with those found by Chen et al. (2012), which were 3033, 2277, and $692 \mu\text{atm}$ for the upper, mid and lower estuaries, respectively. This study yields much higher $p\text{CO}_2$ values for upper estuaries mainly because new data from Asia are associated with high $p\text{CO}_2$ values. The fluxes obtained by Chen et al. (2012), however, are higher. Their values are 68.5, 37.4 and $9.92 \text{ mol C m}^{-2} \text{ yr}^{-1}$ for the upper, mid and lower estuaries, respectively. The seeming inconsistency among results is discussed below.

Figure 3 displays histograms of reported daily CO₂ fluxes per unit area in different seasons and the annual flux per unit area in the world's estuaries. Little seasonal-
 25
 ity is observed, except that the flux is lower in the winter when the *pCO₂* is usually lower, perhaps because the temperature is lower than other seasons. The flux is only marginally higher in summer than in spring or autumn. The numerical average annual flux per unit area is $16.5 \pm 27.7 \text{ mol C m}^{-2} \text{ yr}^{-1}$, which is significantly lower than that,
 $23.9 \pm 33.1 \text{ mmol C m}^{-2} \text{ d}^{-1}$, obtained by Chen et al. (2012). The numerical average



**Air-sea Exchanges
of CO₂ in World's
Coastal Seas**

C.-T. A. Chen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

annual flux per unit area, however, is not used to calculate the global release of CO₂ because small estuaries dominate the numerical average but they contribute relatively little to the total flux.

With respect to latitude, the flux per unit area is the highest, at $63.3 \pm 100.7 \text{ mmol C m}^{-2} \text{ d}^{-1}$ between 23.5° and 50° N, followed by $44.1 \pm 29.3 \text{ mmol C m}^{-2} \text{ d}^{-1}$ between 23.5° S and 0°, followed by $38.8 \pm 55.4 \text{ mmol C m}^{-2} \text{ d}^{-1}$ between 0° and 23.5° N, and $-35.9 \pm 91.2 \text{ mmol C m}^{-2} \text{ d}^{-1}$ north of 50° N. The flux south of 50° S ($9.5 \pm 11.7 \text{ mmol C m}^{-2} \text{ d}^{-1}$) is greatly lower but data are available for only two estuaries. By way of comparison, Chen et al. (2012) obtained $65.5 \pm 78.1 \text{ mmol C m}^{-2} \text{ d}^{-1}$ a slightly higher value than obtained in this study between 23.5° N and 23.5° S. The values of Chen et al. (2012), $67.4 \pm 108 \text{ mmol C m}^{-2} \text{ d}^{-1}$ between 23.5 and 50° N and $59.2 \pm 80 \text{ mmol C m}^{-2} \text{ d}^{-1}$ north of 50° N, are, however, significantly higher than those obtained herein. Notably, most other investigations have presented fluxes higher than those that were presented by Chen et al. (2012).

The fact that the annual average flux herein is lower than those reported previously, despite the fact the average *p*CO₂ is higher, warrants discussion. It follows mainly from the fact that many data from the low latitude bands in Asia have been added, and these areas are mostly areas of low wind energy. Figure 5 plots the wind energy potential which is, like the air-sea gas exchange rate, a quadratic function of wind speed. The areas of high wind energy at low latitudes are concentrated in the dry Middle East and northeastern, northern and northwestern Africa with few rivers, and therefore few estuaries. For example, the total area of the estuaries in the Red Sea region is almost zero (Table 3; Laruelle et al., 2012). Accordingly, the global average CO₂ flux herein is substantially affected by estuaries in areas of low wind energy, and therefore of low CO₂ flux.

The 50 newly considered estuaries in Taiwan, southern China and Southeast Asia, all at low latitudes, have lower fluxes than determined from previously obtained results (Table 1), which include many data for European rivers. For instance, only two of the

19 estuaries that were considered by Abril and Borges (2005), who published perhaps the first global study of CO₂ emissions from estuaries, are outside Europe and the eastern seaboard of the USA. Those authors found a global CO₂ flux per unit area of 35.7 mol C m⁻² yr⁻¹, which is more than triple the value obtained in this study. This finding does not imply that European rivers have higher *p*CO₂: they do not. Rather, Europe has more windy coasts than elsewhere in the world, and especially Asia. As mentioned above, the wind potential is a quadratic function of wind speed, as is the 1992 Wanninkhof air-sea CO₂ exchange equation. The mean *p*CO₂ of European estuaries is roughly 1600 µatm, whereas that of Asian estuaries is much higher, around 4000 µatm. Yet, the mean wind speed on European coasts is approximately 4 m s⁻¹, compared with about 1.6 m s⁻¹ on Asian coasts. The resulting CO₂ fluxes for European estuaries average about 16.9 mol C m⁻² yr⁻¹ vs. a much lower 8.1 mol C m⁻² yr⁻¹ for Asian estuaries (Table 3; Fig. 6) despite their higher *p*CO₂.

In the above calculation, the areas of estuaries are taken from Laruelle et al. (2012), who divided the world into 45 regions, and calculated a total estuarine area of 1.012×10^6 km², which is slightly smaller than the value of 1.067×10^6 km², given in Laruelle et al. (2010). Table 3 lists the total surface area in each of the 45 regions and the average CO₂ flux per unit area for each region. For regions without data, the mean flux for a similar region is used. Estuaries in North America have the largest total area but the lowest average flux per unit area among all continents, and therefore a low total flux of 10.8 TgC yr⁻¹. That is, a continent with 41 % of world's estuarine area accounts for only 12 % of the world's estuarine CO₂ release (Fig. 6). African, European and South American estuaries have similarly high fluxes per unit area but the areas of the estuaries are only moderate so they are responsible for only 16 % (14.7 TgC yr⁻¹), 26 % (24.1 TgC yr⁻¹) and 12 % (11.6 TgC yr⁻¹), respectively, of the global release. The largest contributor is Asia which has 31.5 % of the world's estuarine area and releases almost the same percentage of the world's estuarine released CO₂ (32 % or 30.6 TgC yr⁻¹; Fig. 6).

Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[|◀](#)

[▶|](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Largely on account of the distribution of data, which include data from high wind regions on both sides of the North Atlantic and around the Arabian Sea in the Indian Ocean, as well as those generally in the low wind regions around the Pacific Ocean, the mean CO₂ flux per unit area is the lowest for estuaries that flow into the Pacific Ocean, with a value of 10.5 mol C m⁻² yr⁻¹ (Fig. 7). This value compares with 12.4 mol C m⁻² yr⁻¹ for estuaries that flow into the Atlantic Ocean and 13.9 mol C m⁻² yr⁻¹ for estuaries that flow into the Indian Ocean. Because the total area of estuaries that enter the Atlantic Ocean exceeds the sum of areas of estuaries that enter the Pacific and Indian Oceans, the total flux of CO₂ released from the estuaries around the Atlantic (54.1 TgC yr⁻¹) exceeds the total flux from estuaries around the Pacific (30.8 TgC yr⁻¹) and the Indian (13.3 TgC yr⁻¹) Oceans. The total area of estuaries that enter the Arctic Ocean is substantial (324×10^3 km²), equaling the total areas of the estuaries around the Atlantic and Indian Oceans. Unfortunately, the relevant data are scarce and the available data seem to reveal that the Arctic estuaries absorb, rather than release CO₂. The flux per unit area and total flux are -1.1 mol C m⁻² yr⁻¹ and -4.2 TgC yr⁻¹, respectively. The global total release of 94 TgC yr⁻¹ is less than half of any previous estimates (Table 2). New data from low wind regions and the Arctic Ocean are responsible for this difference.

3 Air-Sea CO₂ exchanges in continental shelves

Data are available from 87 continental shelves (Table 3 and Fig. 8). Table 3 presents the fluxes in the 45 regions that were identified by Laruelle et al. (2010) and the areas of each region. Figure 9 displays a histogram of the reported daily CO₂ fluxes in different seasons and the annual flux for the world's continental shelves. Respiration rates are higher in summer and fall than in winter and spring (Hopkinson, 1985, 1988; Griffith et al., 1990; Hopkinson and Smith, 2005; Jiang et al., 2010). However, as with estuaries, no seasonality of flux per unit area on continental shelves is evident, and the values fall between -4.0 and -5.5 mmol C m⁻² d⁻¹, except in autumn, when

the flux is only $-0.5 \text{ mmol C m}^{-2} \text{ d}^{-1}$. The numerical annual mean air-to-sea flux is $-1.09 \pm 2.9 \text{ mol C m}^{-2} \text{ yr}^{-1}$. Multiplying this value by the total global area of the shelves yields a global flux of $-0.40 \text{ PgC yr}^{-1}$, which is slightly less than the published value (Table 5). A positive value indicates that the shelves absorb CO₂.

Figure 10 presents a histogram of the reported daily fluxes of CO₂ in different latitude bands. Most shelves absorb CO₂ from the atmosphere (negative fluxes) while shelves at low latitudes have a slight tendency to release CO₂ (positive fluxes). This finding is consistent with the work of Cai et al. (2006), who found that shelves at low latitudes between 30° N and 30° S are a source of CO₂ of the order of 0.11 PgC yr^{-1} , whereas those in temperate and high-latitude regions are sinks of CO₂ of the order of 0.33 PgC yr^{-1} . The CO₂ flux per unit area is highest on the South American shelves ($-3.6 \text{ mol C m}^{-2} \text{ yr}^{-1}$) but since their total area is moderate, the South American shelves absorb the second largest amount of CO₂ from the atmosphere annually at $-103.5 \text{ TgC yr}^{-1}$, or 26 % of the global shelf absorption.

Asian shelves have the highest total area but their flux per unit area ($-0.13 \text{ mol C m}^{-2} \text{ yr}^{-1}$) is the lowest of all, primarily because of the generally low wind speed and because some shelves release rather than absorb CO₂. The total annual flux from Asian shelves is only -22 TgC yr^{-1} , or 5 % of the global absorption by all shelves. North American shelves rank second in terms of both flux per unit area ($-2.1 \text{ mol C m}^{-2} \text{ yr}^{-1}$) and shelf area. Accordingly, North American shelves absorb the most CO₂ from the atmosphere at -156 TgC yr^{-1} , or 39 % of the global absorption by all shelves. Shelves around the Antarctica have the third highest flux per unit area ($-2.0 \text{ mol C m}^{-2} \text{ yr}^{-1}$) and the third largest total shelf area, resulting in the third highest total annual flux at -70 TgC yr^{-1} , or 18 % of the global absorption (Fig. 11). Unfortunately, data are available for only two such shelves.

Figure 12 shows the total CO₂ flux per unit area and the total CO₂ flux from shelves in different oceans. Two shelves in the Southern Ocean have the highest flux per unit area ($-2 \text{ mol C m}^{-2} \text{ yr}^{-1}$) with a total annual flux of -70 TgC yr^{-1} , or 18 % of the global shelf absorption. The second highest flux per unit area is that of shelves in the Arctic Ocean

Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



($-1.8 \text{ mol C m}^{-2} \text{ yr}^{-1}$), which also have the second highest total flux at -129 TgC yr^{-1} , or 33 % of the global absorption. Shelves in the Atlantic Ocean have the highest total absorption (-130 TgC yr^{-1} , or 33 % of the global absorption), the third highest flux per unit area ($-1.2 \text{ mol C m}^{-2} \text{ yr}^{-1}$) and the second highest total shelf area. The largest shelf area is that of the shelves around the Pacific Ocean but since the flux per unit area is low ($-0.4 \text{ mol C m}^{-2} \text{ yr}^{-1}$), their total absorption is only -49 TgC yr^{-1} , or 12 % of global absorption. Shelves around the Indian Ocean have the least total area and the second lowest flux per unit area ($-0.6 \text{ mol C m}^{-2} \text{ yr}^{-1}$), resulting in the lowest total flux (-18 TgC yr^{-1} , or only 4 % of the global absorption). In total, the world's shelves absorb 396 TgC yr^{-1} or $0.396 \text{ PgC yr}^{-1}$.

The shift from the sinking of CO_2 at higher latitudes to acting as a weak source at lower latitudes is explained by four major factors. The first is that waters on continental shelves are mostly dominated by the open oceans, as revealed simply by the salinity of most shelves, which is only slightly lower than that of the open ocean waters, except close to the estuaries. For example, for a shelf with 10 % input from rivers with a salinity of 0.5 and a $p\text{CO}_2$ of $1000 \mu\text{atm}$, and 90 % input from open oceans with a salinity of 35 and a $p\text{CO}_2$ of $300 \mu\text{atm}$, the resulting salinity (S) is 31.55. For the sake of argument, the $p\text{CO}_2$ of this $S = 31.55$ shelf water is approximately $370 \mu\text{atm}$, depending on the alkalinity of the river water. Restated, mixing with open ocean waters with low $p\text{CO}_2$ causes the $p\text{CO}_2$ of river water with high $p\text{CO}_2$ be reduced to below saturation. Notably, open ocean waters at high latitudes are frequently under-saturated and open ocean waters at low latitudes are frequently super-saturated (Takahashi et al., 2002; Kaltin et al., 2002; Kaltin and Anderson, 2005; Chen et al., 2006a, b, 2008a, b; Caias et al., 2008). Therefore, mixing with open ocean waters at high latitudes helps shelf waters become under-saturated whereas mixing with open ocean waters at low latitudes frequently yields shelf waters that are still super-saturated (Hidalgo-Gonzalez et al., 1997; Ito et al., 2005; Cai et al., 2006; Chen et al., 2008a, 2012).

The second factor that contributes to the super-saturation of shelf waters at low latitudes is temperature because $p\text{CO}_2$ increases by 4.3 % for an increase of a degree

Air-sea Exchanges of CO_2 in World's Coastal Seas

C.-T. A. Chen et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[|◀](#)

[▶|](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Celcuis (Bakker et al., 1999; Takahashi et al., 2002). Simply increasing by 15 °C the temperature of the shelf water with a $p\text{CO}_2$ of 370 µatm in the example given above would result in a $p\text{CO}_2$ of above 600 µatm, if all other factors are held constant. Notably, the difference between the temperatures of shelves at high latitude shelves and those at low latitude commonly exceeds 15 °C. Temperature similarly affects open ocean water so not only do warm temperatures increase the $p\text{CO}_2$ of shelf waters but also these waters also mix with open ocean waters with higher $p\text{CO}_2$, mainly because they are hotter than open oceans at high latitudes. The proximal shelves in waters with a depth of, say, less than 40 m, typically exhibit a greater seasonal range of water temperatures than the distal shelves with a water depth of between 40 and 200 m. The effect of temperature on the $p\text{CO}_2$ of proximal shelf waters is thus greater than that of distal shelf waters. Unfortunately, the readily available $p\text{CO}_2$ data do not suffice for a meaningful synthesis. As a matter of fact, for navigational or geopolitical reasons, $p\text{CO}_2$ is rarely measured along the coast or the data are frequently not disclosed.

The third factor in affecting $p\text{CO}_2$ on shelves is the fact that the discharge of organic matter by rivers is higher at lower latitudes. As much as 60 % of the riverine organic carbon discharge to the shelves occurs between 0°–30° (Walsh, 1988; Ludwig et al., 1996a, b; Borges et al., 2005). The total amount is approximately -0.3 PgC yr^{-1} , most of which is decomposed in the continental margins (McKee, 2003; Cai et al., 2006). Importantly, however, the cited studies did not identify the recipient of the riverine export. As indicated above, a significant fraction of the export is decomposed in the estuaries and does not reach the shelves.

The fourth factor that is responsible for the higher $p\text{CO}_2$ in shelves at lower latitudes involves lower biological productivity. Shelves at mid and high latitudes are generally highly productive whereas those at low latitudes, especially the non-upwelling shelves, are typically oligotrophic. The more effective biological pumping in the shelves at mid and high latitudes causes these shelves to have lower $p\text{CO}_2$.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

4 Temporal changes

Although efforts have been made to evaluate the parameters that affect the carbon cycle and to model past and future changes in carbon fluxes (Friederich et al., 2002; Fransson et al., 2006; Macpherson et al., 2008; Ver et al., 1999b, a; Mackenzie et al., 5 2000, 2004; Thomas et al., 2007; Borges, 2011), whether the size of the estuarine source and the continental shelf sink for CO₂ changes with time has not been determined because too few data are available. As stated above, the fact that the shelves are a sink rather than a source of CO₂ has only been established in the last few years. The conclusion of LOICZ that the coastal seas are a source of CO₂ (Crossland et al., 2005) 10 was not based on data. Rather, it was based on the reasoning that since rivers transport more organic carbon to the oceans than is buried in the sediments, the oceans must be releasing the remaining CO₂ back to the atmosphere (Smith and Mackenzie, 1987; Smith and Hollibaugh, 1993). However, the amount of organic carbon that is actually transported to the oceans after it passes through the estuaries is not clear, as 15 mentioned above.

Before Industrialization, more organic carbon may well have reached the oceans than accumulated in the sediments, such that the oceans were overall a source of CO₂. However, whether the CO₂ was released in the coastal oceans or in the open oceans remains unclear. Further, the increasing CO₂ in the atmosphere must have 20 reduced the difference between the $p\text{CO}_2$ of the atmosphere and that of the oceans, even if the oceans used to be supersaturated with CO₂ before Industrialization. In any case, the present data clearly demonstrate that the coastal oceans are a sink of CO₂. Whether a threshold has been crossed or whether the metabolism of the ecosystem has been changed cannot yet be determined (Mackenzie et al., 2004).

25 Interestingly, heterotrophic systems are commonly treated as CO₂ sources while autotrophic systems are considered to be CO₂ sinks (Walsh et al., 1981; Smith and Hollibaugh, 1993; Ducklow and McCallister, 2004). This assumption is usually true but must be applied with caution. For instance, if a shelf is regarded as a system, then

Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



it can be heterotrophic overall but its surface layer may still be under-saturated owing to cooling or biological production, and so absorbs CO₂ from the atmosphere. The deep layer may become super-saturated simply because it is warmer. Alternatively, the DOC or POC that is produced in the surface layer, or imported from rivers and submarine groundwater discharge, decomposes in the deep layer. As long as more CO₂ is generated in the deep layer than is taken up by the surface layer, the shelf as a whole will remain heterotrophic.

5 Future changes in carbon fluxes

Increasing air temperature (Belkin, 2009) tends to increase precipitation and continental runoff. These processes enhanced rock weathering during the last century (Probst et al., 1994). Intuitively, this fact suggests an increased export of carbon by rivers but whether the global river runoff has increased is uncertain (Dai et al., 2011; Syed et al., 2010). The construction of dams around the world has caused a substantial fraction of exported sediment to be impounded in recent decades (Chen, 2002; Syvitski et al., 2005). A related issue is that global warming is warming the oceans. The global mean sea surface temperature has reportedly risen by 0.67 °C over the last century (IPCC, 2007; Trenberth et al., 2007). The most rapid warming, two to four times the global average of 0.177 °C per decade between 1981 and 2005, has been observed in the land-locked or semi-enclosed European and East Asian Seas, including the Baltic, North, Black, Japan and East China Seas as well as over the Newfoundland-Labrador Shelf (Belkin, 2009). The thermodynamics of seawater dictate that for each °C rise in temperature, the pCO₂ increases by 4 %, or approximately 14 µatm. This fact would compensate for some of the increase in CO₂ in the atmosphere, which is of the order of 18 µatm per decade.

BGD

10, 5041–5105, 2013

Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



6 Conclusions

Data from 165 estuaries and 87 continental shelves around the globe have been evaluated to show that world's estuaries release $0.094 \text{ PgC yr}^{-1}$ to the atmosphere. This value is substantially lower than any published values mainly because of the newly available data in Asia which has low wind speed in general, hence low per unit area flux. In addition, new data in the polar regions indicate that estuaries there may absorb, instead of release CO_2 . Overall, the world's continental shelves absorb 0.4 PgC yr^{-1} from the atmosphere, which is in line with published data.

Acknowledgements. The National Science Council of Taiwan (NSC 101-2611-M-110-010-MY3, 98-2621-M-110-001-MY3) and the Aim for the Top University Program (01 C0302 04) provided financial assistance.

References

- 15 Abril, G. and Borges, A.: Carbon dioxide and methane emissions from estuaries, in: Greenhouse Gases Emissions from Natural Environments and Hydroelectric Reservoirs: Fluxes and Processes, Environ. Sci. Ser., edited by: Tremblay, A., Varfalvy, L., Roehm, C., and Garneau, M., Springer-Verlag Berlin, New York, 187–207, 2005.
- 20 Abril, G., Nogueira, M., Etcheber, H., Cabecadas, G., Lemaire, E., and Brogueira, M. J.: Behaviour of organic carbon in nine contrasting European estuaries, Estuarine Coastal. Shelf Sci., 54, 241–262, doi:10.1006/ecss.2001.0844, 2002.
- 25 Abril, G., Etcheber, H., Delille, B., Frankignoulle, M., and Borges, A. V.: Carbonate dissolution in the turbid and eutrophic Loire estuary, Mar. Ecol. Prog. Ser., 259, 129–138, doi:10.3354/meps259129, 2003.
- Algesten, G., Wikner, J., Sobek, S., Tranvik, L. J., and Jansson, M.: Seasonal variation of CO_2 saturation in the Gulf of Bothnia: Indications of marine net heterotrophy, Glob. Biogeochem. Cy., 18, Gb4021, doi:10.1029/2004gb002232, 2004.
- Alongi, D. M.: Coastal ecosystem processes, Marine science series, CRC Press, Boca Raton, 448 pp., 1998.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Álvarez-Salgado, X. A., Doval, M. D., and Pérez, F. F.: Dissolved organic matter in shelf waters off the Ría de Vigo (NW Iberian upwelling system), *J. Mar. Syst.*, 18, 383–394, doi:10.1016/s0924-7963(97)00114-0, 1999.

5 Bakker, D. C. E., de Baar, H. J. W., and de Jong, E.: The dependence on temperature and salinity of dissolved inorganic carbon in East Atlantic surface waters, *Mar. Chem.*, 65, 263–280, doi:10.1016/s0304-4203(99)00017-1, 1999.

Bates, N. R.: Air-sea CO₂ fluxes and the continental shelf pump of carbon in the Chukchi Sea adjacent to the Arctic Ocean, *J. Geophys. Res.*, 111, C10013, doi:10.1029/2005jc003083, 2006.

10 Bates, N. R., Hansell, D. A., Carlson, C. A., and Gordon, L. I.: Distribution of CO₂ species, estimates of net community production, and air-sea CO₂ exchange in the Ross Sea polynya, *J. Geophys. Res.*, 103, 2883–2896, doi:10.1029/97jc02473, 1998.

Belkin, I. M.: Rapid warming of large marine ecosystems, *Prog. Oceanogr.*, 81, 207–213, doi:10.1016/j.pocean.2009.04.011, 2009.

15 Beusen, A. H. W., Dekkers, A. L. M., Bouwman, A. F., Ludwig, W., and Harrison, J.: Estimation of global river transport of sediments and associated particulate C, N, and P, *Glob. Biogeochem. Cy.*, 19, Gb4s05, doi:10.1029/2005gb002453, 2005.

Bianchi, T. S. and Allison, M. A.: Large-river delta-front estuaries as natural “recorders” of global environmental change, *Proc. Natl. Acad. Sci. USA*, 106, 8085–8092, doi:10.1073/pnas.0812878106, 2009.

Boehme, S. E., Sabine, C. L., and Reimers, C. E.: CO₂ fluxes from a coastal transect: a time-series approach, *Mar. Chem.*, 63, 49–67, doi:10.1016/s0304-4203(98)00050-4, 1998.

Borges, A. V.: Do we have enough pieces of the jigsaw to integrate CO₂ fluxes in the coastal ocean?, *Estuaries*, 28, 3–27, doi:10.1007/bf02732750, 2005.

25 Borges, A. V.: Present day carbon dioxide fluxes in the coastal ocean and possible feedbacks under global change, in: *Oceans and the atmospheric carbon content* edited by: Duarte, P. M. d. S., and Casiano, J. M. S., 47–77, 2011.

Borges, A. V. and Frankignoulle, M.: Distribution of surface carbon dioxide and air-sea exchange in the English Channel and adjacent areas, *J. Geophys. Res.*, 108, 3140, doi:10.1029/2000jc000571, 2003.

30 Borges, A. V., Djennidi, S., Lacroix, G., Theate, J., Delille, B., and Frankignoulle, M.: Atmospheric CO₂ flux from mangrove surrounding waters, *Geophys. Res. Lett.*, 30, 1558, doi:10.1029/2003gl017143, 2003.

BGD

10, 5041–5105, 2013

Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



**Air-sea Exchanges
of CO₂ in World's
Coastal Seas**

C.-T. A. Chen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Borges, A. V., Delille, B., and Frankignoulle, M.: Budgeting sinks and sources of CO₂ in the coastal ocean: Diversity of ecosystems counts, *Geophys. Res. Lett.*, 32, L14601, doi:10.1029/2005gl023053, 2005.

Borges, A. V., Schiettecatte, L. S., Abril, G., Delille, B., and Gazeau, E.: Carbon dioxide in European coastal waters, *Estur. Coast. Shelf Sci.*, 70, 375–387, doi:10.1016/j.ecss.2006.05.046, 2006.

Bouillon, S., Frankignoulle, M., Dehairs, F., Velimirov, B., Eiler, A., Abril, G., Etcheber, H., and Borges, A. V.: Inorganic and organic carbon biogeochemistry in the Gautami Godavari estuary (Andhra Pradesh, India) during pre-monsoon: The local impact of extensive mangrove forests, *Glob. Biogeochem. Cy.*, 17, 1114, doi:10.1029/2002gb002026, 2003.

Bouillon, S., Korntheuer, M., Baeyens, W., and Dehairs, F.: A new automated setup for stable isotope analysis of dissolved organic carbon, *Limnol. Oceanogr. Methods*, 4, 216–226, 2006.

Bouillon, S., Dehairs, F., Schiettecatte, L. S., and Borges, A. V.: Biogeochemistry of the Tana estuary and delta (northern Kenya), *Limnol. Oceanogr.*, 52, 46–59, 2007a.

Bouillon, S., Dehairs, F., Velimirov, B., Abril, G., and Borges, A. V.: Dynamics of organic and inorganic carbon across contiguous mangrove and seagrass systems (Gazi Bay, Kenya), *J. Geophys. Res.*, 112, G02018, doi:10.1029/2006jg000325, 2007b.

Bouillon, S., Middelburg, J. J., Dehairs, F., Borges, A. V., Abril, G., Flindt, M. R., Ulomi, S., and Kristensen, E.: Importance of intertidal sediment processes and porewater exchange on the water column biogeochemistry in a pristine mangrove creek (Ras Dege, Tanzania), *Biogeosciences*, 4, 311–322, doi:10.5194/bg-4-311-2007, 2007c.

Bouillon, S., Connolly, R. M., and Lee, S. Y.: Organic matter exchange and cycling in mangrove ecosystems: Recent insights from stable isotope studies, *J. Sea Res.*, 59, 44–58, doi:10.1016/j.seares.2007.05.001, 2008.

Cai, W. J.: Riverine inorganic carbon flux and rate of biological uptake in the Mississippi River plume, *Geophys. Res. Lett.*, 30, 1032, doi:10.1029/2002gl016312, 2003.

Cai, W. J.: Estuarine and coastal ocean carbon paradox: CO₂ sinks or sites of terrestrial carbon incineration?, *Annu. Rev. Mar. Sci.*, 3, 123–145, doi:10.1146/annurev-marine-120709-142723, 2011.

Cai, W. J. and Dai, M.: Comment on “Enhanced open ocean storage of CO₂ from shelf sea pumping”, *Science*, 306, 1477–1477, 2004.

Cai, W. J. and Wang, Y.: The chemistry, fluxes, and sources of carbon dioxide in the estuarine waters of the Satilla and Altamaha Rivers, Georgia, *Limnol. Oceanogr.*, 43, 6570–668, 1998.

**Air-sea Exchanges
of CO₂ in World's
Coastal Seas**

C.-T. A. Chen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Cai, W. J., Pomeroy, L. R., Moran, M. A., and Wang, Y. C.: Oxygen and carbon dioxide mass balance for the estuarine-intertidal marsh complex of five rivers in the southeastern US, *Limnol. Oceanogr.*, 44, 639–649, 1999.

Cai, W. J., Wiebe, W. J., Wang, Y. C., and Sheldon, J. E.: Intertidal marsh as a source of dissolved inorganic carbon and a sink of nitrate in the Satilla River-estuarine complex in the southeastern US, *Limnol. Oceanogr.*, 45, 1743–1752, 2000.

Cai, W. J., Wang, Y. C., Krest, J., and Moore, W. S.: The geochemistry of dissolved inorganic carbon in a surficial groundwater aquifer in North Inlet, South Carolina, and the carbon fluxes to the coastal ocean, *Geochim. Cosmochim. Acta*, 67, 631–639, doi:10.1016/s0016-7037(02)01167-5, 2003.

Cai, W. J., Dai, M. H., and Wang, Y. C.: Air-sea exchange of carbon dioxide in ocean margins: A province-based synthesis, *Geophys. Res. Lett.*, 33, L12603, doi:10.1029/2006gl026219, 2006.

Cameron, W. M. and Pritchard, D. W.: Estuaries, in: *The Sea*, edited by: Hill, M. N., John Wiley and Sons, New York, 306–324, 1963.

Carrillo, C. J. and Karl, D. M.: Dissolved inorganic carbon pool dynamics in northern Gerlache Strait, Antarctica, *J. Geophys. Res.*, 104, 15873–15884, doi:10.1029/1999jc900110, 1999.

Chavez, F. P., Takahashi, T., Cai, W.-J., Friederich, G., Hales, B., Wanninkhof, R., and Feely, R. A.: Coastal Oceans, in: *The first state of the carbon cycle report (SOCCR): The north American carbon budget and implications for the global carbon cycle*, edited by: King, A. W., Dilling, L., Zimmerman, G. P., Fairman, D. M., Houghton, R. A., Marland, G., Rose, A. Z., and Wilbanks, T. J., National Oceanic and Atmospheric Administration, National Climatic Data Center, Washington, DC, USA, 157–166, 2007.

Chen, C. T. A.: The oceanic anthropogenic CO₂ sink, *Chemosphere*, 27, 1041–1064, 1993.

Chen, C. T. A.: Dams may impact fisheries even beyond the estuaries, *Netherlands Institute for Sea Research*, The Netherlands, 177–178, 2002.

Chen, C. T. A.: New vs. export production on the continental shelf, *Deep Sea Res.*, 50, 1327–1333, doi:10.1016/S0967-0645(03)00026-2, 2003.

Chen, C. T. A.: Exchanges of carbon in the coastal seas, in: *The global carbon cycle:integrating humans, climate, and the natural world*, edited by: Field, C. B. and Raupach, M. R., Island Press, 341–351, 2004.

Chen, C. T. A. and Borges, A. V.: Reconciling opposing views on carbon cycling in the coastal ocean: Continental shelves as sinks and near-shore ecosystems as sources of atmospheric CO₂, *Deep-Sea Res.*, 56, 578–590, doi:10.1016/j.dsr2.2009.01.001, 2009.

5 Chen, C. T. A. and Tsunogai, S.: Carbon and nutrients in the ocean, in: *Asian Change in the Context of Global Change*, edited by: Galloway, J. N. and Melillo, J. M., Cambridge University Press, Cambridge, UK, 271–307, 1998.

Chen, C. T. A. and Wang, S. L.: Carbon, alkalinity and nutrient budgets on the East China Sea continental shelf, *J. Geophys. Res.*, 104, 20675–20686, doi:10.1029/1999jc900055, 1999.

10 Chen, C. T. A., Holligan, P., Hong, H. S., Iseki, K., Krishnaswami, S., Wollast, R., and Yoder, J.: Land–ocean interactions in the coastal zone, 20 pp., 1994.

Chen, C. T. A., Liu, K. K., and MacDonald, R.: Continental margin exchanges, in: *Ocean Biogeochemistry: a JGOFS Synthesis*, edited by: Fasham, M. J. R., Springer, Berlin, 53–97, 2003.

15 Chen, C. T. A., Hou, W. P., Gamo, T., and Wang, S. L.: Carbonate-related parameters of subsurface waters in the West Philippine, South China and Sulu Seas, *Mar. Chem.*, 99, 151–161, doi:10.1016/j.marchem.2005.05.008, 2006a.

Chen, C. T. A., Wang, S. L., Chou, W. C., and Sheu, D. D.: Carbonate chemistry and projected future changes in pH and CaCO₃ saturation state of the South China Sea, *Mar. Chem.*, 101, 277–305, 2006b.

20 Chen, C. T. A., Wang, S. L., Lu, X. X., Zhang, S. R., Lui, H. K., Tseng, H. C., Wang, B. J., and Huang, H. I.: Hydrogeochemistry and greenhouse gases of the Pearl River, its estuary and beyond, *Quatern. Int.*, 186, 79–90, doi:10.1016/j.quaint.2007.08.024, 2008a.

25 Chen, C. T. A., Zhai, W., and Dai, M.: Riverine input and air-sea CO₂ exchanges near the Changjiang (Yangtze River) Estuary: Status quo and implication on possible future changes in metabolic status, *Cont. Shelf Res.*, 28, 1476–1482, doi:10.1016/j.csr.2007.10.013, 2008b.

Chen, C. T. A., Huang, T. H., Fu, Y. H., Bai, Y., and He, X.: Strong sources of CO₂ in upper estuaries become sinks of CO₂ in large river plumes, *Curr. Opin. Env. Sust.*, 4, 179–185, 2012.

30 Ciais, P., Borges, A. V., Abril, G., Meybeck, M., Folberth, G., Hauglustaine, D., and Janssens, I. A.: The impact of lateral carbon fluxes on the European carbon balance, *Biogeosciences*, 5, 1259–1271, doi:10.5194/bg-5-1259-2008, 2008.

Clark, D. B., Taylor, C. M., and Thorpe, A. J.: Feedback between the land surface and rainfall at convective length scales, *J. Hydrometeorol.*, 5, 625–639, 2004.

Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.

- Codispoti, L. A., Friederich, G. E., and Hood, D. W.: Variability in the inorganic carbon system over the southeastern Bering Sea shelf during spring 1980 and spring – summer 1981, Cont. Shelf Res., 5, 133–160, 1986.
- Cole, J. J. and Caraco, N. F.: Carbon in catchments: connecting terrestrial carbon losses with aquatic metabolism, Mar. Freshwater Res., 52, 101–110, 2001.
- Cooley, S. R., Coles, V. J., Subramaniam, A., and Yager, P. L.: Seasonal variations in the Amazon plume-related atmospheric carbon sink, Glob. Biogeochem. Cy., 21, GB3014, doi:10.1029/2006gb002831, 2007.
- Crossland, C. J., Kremer, H. H., Lindeboom, H. J., Crossland, J. I. M., and Tissier, M. D. A. L.: Coastal fluxes in the anthropocene: the land-ocean interactions in the coastal zone project of the international geosphere-biosphere programme, Springer, Berlin, 232 pp., 2005.
- Dagg, M. J., Bianchi, T. S., Breed, G. A., Cai, W. J., Duan, S., Liu, H., McKee, B. A., Powell, R. T., and Stewart, C. M.: Biogeochemical characteristics of the lower Mississippi River, USA, during June 2003, Estuaries, 28, 664–674, doi:10.1007/bf02732905, 2005.
- Dai, M., Yin, Z., Meng, F., Liu, Q., and Cai, W.-J.: Spatial distribution of riverine DOC Inputs to the ocean: An updated global synthesis, Curr. Opin. Env. Sust., 4, 170–178, 2012.
- Dai, M. H., Zhai, W. D., Cai, W. J., Callahan, J., Huang, B. Q., Shang, S. L., Huang, T., Li, X. L., Lu, Z. M., Chen, W. F., and Chen, Z. Z.: Effects of an estuarine plume-associated bloom on the carbonate system in the lower reaches of the Pearl River estuary and the coastal zone of the northern South China Sea, Cont. Shelf Res., 28, 1416–1423, doi:10.1016/j.csr.2007.04.018, 2008.
- Dai, M. H., Lu, Z. M., Zhai, W. D., Chen, B. S., Cao, Z. M., Zhou, K. B., Cai, W. J., and Chen, C. T. A.: Diurnal variations of surface seawater $p\text{CO}_2$ in contrasting coastal environments, Limnol. Oceanogr., 54, 735–745, doi:10.4319/lo.2009.54.3.0735, 2009.
- Dai, Z. J., Du, J. Z., Zhang, X. L., Su, N., and Li, J. F.: Variation of Riverine Material Loads and Environmental Consequences on the Changjiang (Yangtze) Estuary in Recent Decades (1955–2008), Environ. Sci. Technol., 45, 223–227, doi:10.1021/es103026a, 2011.
- de Haas, H., van Weering, T. C. E., and de Stieger, H.: Organic carbon in shelf seas: sinks or sources, processes and products, Cont. Shelf Res., 22, 691–717, doi:10.1016/s0278-4343(01)00093-0, 2002.
- de la Paz, M., Gomez-Parra, A., and Forja, J.: Inorganic carbon dynamic and air-water CO₂ exchange in the Guadalquivir Estuary (SW Iberian peninsula), J. Mar. Syst., 68, 265–277, doi:10.1016/j.jmarsys.2006.11.011, 2007.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

1

1

1

▶

Back

Close

Full Screen / Esc

[Printer-friendly Version](#)

Interactive Discussion



- Ferron, S., Ortega, T., Gomez-Parra, A., and Forja, J. M.: Seasonal study of dissolved CH₄, CO₂ and N₂O in a shallow tidal system of the bay of Cadiz (SW Spain), *J. Mar. Syst.*, 66, 244–257, doi:10.1016/j.jmarsys.2006.03.021, 2007.
- Frankignoulle, M. and Borges, A. V.: European continental shelf as a significant sink for atmospheric carbon dioxide, *Glob. Biogeochem. Cy.*, 15, 569–576, 2001.
- Frankignoulle, M., Gattuso, J. P., Biondo, R., Bourge, I., CopinMontegut, G., and Pichon, M.: Carbon fluxes in coral reefs 2, Eulerian study of inorganic carbon dynamics and measurement of air-sea CO₂ exchanges, *Mar. Ecol. Prog. Ser.*, 145, 123–132, doi:10.3354/meps145123, 1996.
- Frankignoulle, M., Abril, G., Borges, A., Bourge, I., Canon, C., DeLille, B., Libert, E., and Theate, J. M.: Carbon dioxide emission from European estuaries, *Science*, 282, 434–436, doi:10.1126/science.282.5388.434, 1998.
- Fransson, A., Chierici, M., Anderson, L. C., Bussmann, I., Kattnar, G., Jones, E. P., and Swift, J. H.: The importance of shelf processes for the modification of chemical constituents in the waters of the Eurasian Arctic Ocean: implication for carbon fluxes, *Cont. Shelf Res.*, 21, 225–242, doi:10.1016/s0278-4343(00)00088-1, 2001.
- Fransson, A., Chierici, M. and Nojiri, Y.: Increased net CO₂ outgassing in the upwelling region of the southern Bering Sea in a period of variable marine climate between 1995 and 2001, *J. Geophys. Res.*, 111, C08008, doi:10.1029/2004jc002759, 2006.
- Friederich, G. E., Walz, P. M., Burczynski, M. G., and Chavez, F. P.: Inorganic carbon in the central California upwelling system during the 1997–1999 El Niño-La Niña event, *Prog. Oceanogr.*, 54, 185–203, doi:10.1016/s0079-6611(02)00049-6, 2002.
- Gao, X. L., Song, J. M., Li, X. G., Yuan, H. M., and Li, N.: Spatial and temporal variations in pH and total alkalinity at the beginning of the rainy season in the Changjiang Estuary, China, *Acta. Oceanol. Sin.*, 24, 68–77, 2005.
- Gattuso, J. P., Pichon, M., Delesalle, B., and Frankignoulle, M.: Community metabolism and air-sea CO₂ fluxes in a coral-reef ecosystem (Moorea, French-Polynesia), *Mar. Ecol. Prog. Ser.*, 96, 259–267, doi:10.3354/meps096259, 1993.
- Gattuso, J. P., Frankignoulle, M., and Wollast, R.: Carbon and carbonate metabolism in coastal aquatic ecosystems, *Annu. Rev. Ecol. Syst.*, 29, 405–434, doi:10.1146/annurev.ecolsys.29.1.405, 1998.
- Gazeau, F., Borges, A. V., Barron, C., Duarte, C. M., Iversen, N., Middelburg, J. J., Delille, B., Pizay, M. D., Frankignoulle, M., and Gattuso, J. P.: Net ecosystem metabolism in a micro-

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tidal estuary (Randers Fjord, Denmark): evaluation of methods, Mar. Ecol. Prog. Ser., 301, 23–41, doi:10.3354/meps301023, 2005.

Ghosh, S., Jana, T. K., Singh, B. N., and Choudhury, A.: Comparative study of carbon dioxide system in virgin and reclaimed mangrove waters of Sundarbans during freshet, Mahasagar-Bulletin of the National Institute of Oceanography, 20, 155–161, 1987.

Gibson, J. A. E. and Trull, T. W.: Annual cycle of fCO₂ under sea-ice and in open water in Prydz Bay, East Antarctica, Mar. Chem., 66, 187–200, doi:10.1016/s0304-4203(99)00040-7, 1999.

Gong, G. C., Shiah, F. K., Liu, K. K., Wen, Y. H., and Liang, M. H.: Spatial and temporal variation of chlorophyll a, primary productivity and chemical hydrography in the southern East China Sea, Cont. Shelf Res., 20, 411–436, doi:10.1016/s0278-4343(99)00079-5, 2000.

Goyet, C., Millero, F. J., O'Sullivan, D. W., Eischeid, G., McCue, S. J., and Bellerby, R. G. J.: Temporal variations of pCO₂ in surface seawater of the Arabian Sea in 1995, Deep Sea Res., 45, 609–623, doi:10.1016/s0967-0637(97)00085-x, 1998.

Griffith, P. C., Douglas, D. J., and Wainright, S. C.: Metabolic-activity of size-fractionated microbial plankton in estuarine, nearshore, and continental-shelf waters of Georgia, Mar. Ecol. Prog. Ser., 59, 263–270, doi:10.3354/meps059263, 1990.

Guilderson, T. P., Schrag, D. P., Fallon, S. J., Dunbar, R. B., Kilbourne, K. H., and Prouty, N. G.: Surface water radiocarbon ($\Delta^{14}\text{C}$) reconstructed from reef-building zooxanthellate corals from 1751–2004, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, Tennessee, 2005.

Guo, X. H., Dai, M. H., Zhai, W. D., Cai, W. J., and Chen, B. S.: CO₂ flux and seasonal variability in a large subtropical estuarine system, the Pearl River Estuary, China, J. Geophys. Res., 114, GB3013, doi:10.1029/2008jg000905, 2009.

Gupta, G. V. M., Sarma, V., Robin, R. S., Raman, A. V., Kumar, M. J., Rakesh, M., and Subramanian, B. R.: Influence of net ecosystem metabolism in transferring riverine organic carbon to atmospheric CO₂ in a tropical coastal lagoon (Chilka Lake, India), Biogeochemistry, 87, 265–285, doi:10.1007/s10533-008-9183-x, 2008.

Hales, B., Takahashi, T., and Bandstra, L.: Atmospheric CO₂ uptake by a coastal upwelling system, Glob. Biogeochem. Cy., 19, Gb1009, doi:10.1029/2004gb002295, 2005.

Hales, B., Cai, W. J., Mitchell, B. G., Sabine, C. L., and Schofield, O.: North American continental margins: a synthesis and planning workshop: report of the North American Continental Margins Working Group for the US, Carbon Cycle Scientific Steering Group and Interagency Working Group, US, Carbon Cycle Science Program, Washington, DC, 110 pp., 2008.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Air-sea Exchanges
of CO₂ in World's
Coastal Seas**

C.-T. A. Chen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Air-sea Exchanges
of CO₂ in World's
Coastal Seas

C.-T. A. Chen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Johnson, D. R., Boyer, T. P., Garcia, H. E., Locarnini, R. A., Baranova, O. K., and Zweng, M. M.: World Ocean Database 2009 Documentation, NOAA Printing Office Edn., NODC Internal Report 20, Silver Spring, MD, 2009.

Kaltin, S. and Anderson, L. G.: Uptake of atmospheric carbon dioxide in Arctic shelf seas: evaluation of the relative importance of processes that influence $p\text{CO}_2$ in water transported over the Bering-Chukchi Sea shelf, *Mar. Chem.*, 94, 67–79, doi:10.1016/j.marchem.2004.07.010, 2005.

Kaltin, S., Anderson, L. G., Olsson, K., Fransson, A., and Chierici, M.: Uptake of atmospheric carbon dioxide in the Barents Sea, *J. Mar. Syst.*, 38, 31–45, doi:10.1016/s0924-7963(02)00168-9, 2002.

Kang, Y., Pan, D., Bai, Y., He, X., Chen, X., Chen, C. T. A., and Wang, D.: Areas of the global major river plumes, *Acta. Oceanol. Sin.*, 32, 79–88, 2013.

Kawahata, H., Suzuki, A., and Goto, K.: Coral reefs as sources of atmospheric CO₂ – Spatial distribution of $p\text{CO}_2$ in Majuro Atoll, *Geochem. J.*, 33, 295–303, 1999.

Kayanne, H., Hata, H., Kudo, S., Yamano, H., Watanabe, A., Ikeda, Y., Nozaki, K., Kato, K., Negishi, A., and Saito, H.: Seasonal and bleaching-induced changes in coral reef metabolism and CO₂ flux, *Glob. Biogeochem. Cy.*, 19, GB3015, doi:10.1029/2004gb002400, 2005.

Keil, R. G., Mayer, L. M., Quay, P. D., Richey, J. E., and Hedges, J. I.: Loss of organic matter from riverine particles in deltas, *Geochim. Cosmochim. Acta*, 61, 1507–1511, doi:10.1016/s0016-7037(97)00044-6, 1997.

Kelley, J. J. and Hood, D. W.: Carbon dioxide in the surface water of the ice-covered Bering Sea, *Nature*, 229, 37–39, 1971.

Kemp, W. M., Smith, E. M., MarvinDiPasquale, M., and Boynton, W. R.: Organic carbon balance and net ecosystem metabolism in Chesapeake Bay, *Mar. Ecol. Prog. Ser.*, 150, 229–248, doi:10.3354/meps150229, 1997.

Kempe, S.: Carbon in the freshwater cycle, in: The global carbon cycle, edited by: Bolin, B., Degens, E. T., Kempe, S., and Ketner, P., Scope 13, Wiley, Chichester, UK, 317–342, 1979.

Kempe, S.: Long-term records of CO₂ pressure fluctuations in fresh waters, *Mitteilungen aus dem Geologisch Palaeontologischen Institut der Universität Hamburg*, 52, 91–332, 1982.

Kempe, S.: Coastal seas : a net source or sink of atmospheric carbon dioxide?, LOICZ Core Project, Netherlands Institute for Sea Research, Texel, The Netherlands, 1995.

Kempe, S. and Pegler, K.: Sinks and sources of CO₂ in coastal seas – the North-Sea, *Tellus B*, 43, 224–235, doi:10.1034/j.1600-0889.1991.00015.x, 1991.

Khatiwala, S., Tanhua, T., Mikaloff Fletcher, S., Gerber, M., Doney, S. C., Graven, H. D., Gruber, N., McKinley, G. A., Murata, A., Ríos, A. F., Sabine, C. L., and Sarmiento, J. L.: Global ocean storage of anthropogenic carbon, *Biogeosciences Discuss.*, 9, 8931–8988, doi:10.5194/bgd-9-8931-2012, 2012.

- 5 Kone, Y. J. M. and Borges, A. V.: Dissolved inorganic carbon dynamics in the waters surrounding forested mangroves of the Ca Mau Province (Vietnam), *Estuar. Coast. Shelf Sci.*, 77, 409–421, doi:10.1016/j.ecss.2007.10.001, 2008.
- 10 Kone, Y. J. M., Abril, G., Kouadio, K. N., Delille, B., and Borges, A. V.: Seasonal variability of carbon dioxide in the rivers and lagoons of Ivory Coast (West Africa), *Estuar. Coast.*, 32, 246–260, doi:10.1007/s12237-008-9121-0, 2009.
- Kortzinger, A.: A significant CO₂ sink in the tropical Atlantic Ocean associated with the Amazon River plume, *Geophys. Res. Lett.*, 30, 2287, doi:10.1029/2003gl018841, 2003.
- 15 Kristensen, E., Flindt, M. R., Ulomi, S., Borges, A. V., Abril, G., and Bouillon, S.: Emission of CO₂ and CH₄ to the atmosphere by sediments and open waters in two Tanzanian mangrove forests, *Mar. Ecol. Prog. Ser.*, 370, 53–67, doi:10.3354/meps07642, 2008.
- 20 Kuss, J., Roeder, W., Wlost, K. P., and DeGrandpre, M. D.: Time-series of surface water CO₂ and oxygen measurements on a platform in the central Arkona Sea (Baltic Sea): Seasonality of uptake and release, *Mar. Chem.*, 101, 220–232, doi:10.1016/j.marchem.2006.03.004, 2006.
- 25 Laruelle, G. G., Dürr, H. H., Slomp, C. P., and Borges, A. V.: Evaluation of sinks and sources of CO₂ in the global coastal ocean using a spatially-explicit typology of estuaries and continental shelves, *Geophys. Res. Lett.*, 37, L15607, doi:10.1029/2010gl043691, 2010.
- Laruelle, G. G., Dürr, H. H., Lauerwald, R., Hartmann, J., Slomp, C. P., and Regnier, P. A. G.: Global multi-scale segmentation of continental and coastal waters from the watersheds to the continental margins, *Hydrol. Earth Syst. Sci. Discuss.*, 9, 11319–11361, doi:10.5194/hessd-9-11319-2012, 2012.
- 30 Laruelle, R., Hartmann, J., Ludwig, W., and Moosdorf, N.: Assessing the non-conservative fluvial fluxes of dissolved organic carbon in North America, *J. Geophys. Res.*, 117, G01027, doi:10.1029/2011JG001820, 2012.
- Le Quéré, C., Andres, R. J., Boden, T., Conway, T., Houghton, R. A., House, J. I., Marland, G., Peters, G. P., van der Werf, G., Ahlström, A., Andrew, R. M., Bopp, L., Canadell, J. G., Ciais, P., Doney, S. C., Enright, C., Friedlingstein, P., Huntingford, C., Jain, A. K., Jourdain, C., Kato, E., Keeling, R. F., Klein Goldewijk, K., Levis, S., Levy, P., Lomas, M., Poulter, B.,

BGD
10, 5041–5105, 2013

Air-sea Exchanges
of CO₂ in World's
Coastal Seas

C.-T. A. Chen et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



**Air-sea Exchanges
of CO₂ in World's
Coastal Seas**

C.-T. A. Chen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

1

▶

Bac

Close

Full Screen / Esc

Muller-Karger, F. E., Varela, R., Thunell, R., Luerssen, R., Hu, C. M., and Walsh, J. J.: The importance of continental margins in the global carbon cycle, Geophys. Res. Lett., 32, L01602, doi:10.1029/2004gl021346, 2005.

Murata, A. and Takizawa, T.: Summertime CO₂ sinks in shelf and slope waters of the western Arctic Ocean, *Cont. Shelf Res.*, 23, 753–776, doi:10.1016/S0278-4343(03)00046-3, 2003.

Nakayama, N., Watanabe, S., and Tsunogai, S.: Difference in O₂ and CO₂ gas transfer velocities in Funka Bay, Mar. Chem., 72, 115–129, doi:10.1016/s0304-4203(00)00077-3, 2000.

Neal, C., House, W. A., Jarvie, H. P., and Etherall, A.: The significance of dissolved carbon dioxide in major lowland rivers entering the North Sea, *Sci. Total Environ.*, 210, 187–203, doi:10.1016/s0048-9697(98)00012-6, 1998.

Nedashkovsky, A. P., Sapozhnikov, V. V., Sagalaev, S. G., Isaeva, A. A., and Shevtsova, O. V.: Variability of the carbonate system components and inorganic carbon dynamics in the western Bering Sea in summer, in: Multiple investigations of the Bering Sea ecosystem, edited by: Kotenев, B. N. and Sapozhnikov, V. V., Moskva Russia Izd., Vniro, 155–162, 1995.

Nelson, J. R., Eckman, J. E., Robertson, C. Y., Marinelli, R. L., and Jahnke, R. A.: Benthic microalgal biomass and irradiance at the sea floor on the continental shelf of the South Atlantic Bight: Spatial and temporal variability and storm effects, *Cont. Shelf Res.*, 19, 477–505, doi:10.1016/s0278-4343(98)00092-2, 1999.

Neubauer, S. C. and Anderson, I. C.: Transport of dissolved inorganic carbon from a tidal freshwater marsh to the York River estuary, Limnol. Oceanogr., 48, 299–307, 2003.

Odum, H. T. and Hoskin, C. M.: Comparative studies of the metabolism of Texas Bays, Publications of the Institute of Marine Science, University of Texas, 5, 16-46, 1958.

Odum, H. T. and Wilson, R.: Further studies on the reaeration and metabolism of Texas Bays, Publications of the Institute of Marine Science, University of Texas, 8, 23-55, 1962.

Oh, D. C., Park, M. K., and Kim, K. R.: CO₂ exchange at air-sea interface in the Huanghai Sea, *Acta. Oceanol. Sin.*, 19, 79–89, 2000.

Ortega, T., Ponce, R., Forja, J., and Gomez-Parra, A.: Fluxes of dissolved inorganic carbon in three estuarine systems of the Cantabrian Sea (north of Spain), *J. Mar. Syst.*, 53, 125–142, doi:10.1016/j.marsys.2004.06.006, 2005.

Otsuki, A. S., Watanabe, S., and Tsunogai, S.: Absorption of atmospheric CO₂ and its transport to the intermediate layer in the Okhotsk Sea, *J. Oceanogr.*, 59, 709–717, doi:10.1023/b:joce.0000009599.94380.30, 2003.

**Air-sea Exchanges
of CO₂ in World's
Coastal Seas**

C.-T. A. Chen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Ovalle, A. R. C., Rezende, C. E., Lacerda, L. D., and Silva, C. A. R.: Factors affecting the hydrochemistry of a mangrove tidal creek, sepetiba bay, Brazil, *Estuar. Coast. Shelf Sci.*, 31, 639–650, doi:10.1016/0272-7714(90)90017-I, 1990.

Paquay, F. S., Mackenzie, F. T., and Borges, A. V.: Carbon dioxide dynamics in rivers and coastal waters of the “Big Island” of Hawaii, USA, during baseline and heavy rain conditions, *Aquat. Geochem.*, 13, 1–18, doi:10.1007/s10498-006-9005-5, 2007.

Peng, T. R., Chen, C. T. A., Wang, C. H., Zhanc, J., and Lin, Y. J.: Assessment of terrestrial factors controlling the submarine groundwater discharge in water shortage and highly deformed island of Taiwan, western Pacific Ocean, *J. Oceanogr.*, 64, 323–337, doi:10.1007/s10872-008-0026-0, 2008.

Peterson, D. H.: Sources and sinks of biologically reactive oxygen, carbon, nitrogen, and silica in Northern San Francisco Bay, 1979.

Pfeil, B., Olsen, A., Bakker, D. C. E., Hankin, S., Koyuk, H., Kozyr, A., Malczyk, J., Manke, A., Metzl, N., Sabine, C. L., Akl, J., Alin, S. R., Bellerby, R. G. J., Borges, A., Boutin, J., Brown, P. J., Cai, W.-J., Chavez, F. P., Chen, A., Cosca, C., Fassbender, A. J., Feely, R. A., González-Dávila, M., Goyet, C., Hardman-Mountford, N., Heinze, C., Hood, M., Hoppeima, M., Hunt, C. W., Hydes, D., Ishii, M., Johannessen, T., Jones, S. D., Key, R. M., Kötzinger, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lenton, A., Lourantou, A., Merlivat, L., Midorikawa, T., Mintrop, L., Miyazaki, C., Murata, A., Nakadate, A., Nakano, Y., Nakaoka, S., Nojiri, Y., Omar, A. M., Padin, X. A., Park, G.-H., Paterson, K., Perez, F. F., Pierrot, D., Poisson, A., Ríos, A. F., Santana-Casiano, J. M., Salisbury, J., Sarma, V. V. S. S., Schlitzer, R., Schneider, B., Schuster, U., Sieger, R., Skjelvan, I., Steinhoff, T., Suzuki, T., Takahashi, T., Tedesco, K., Telszewski, M., Thomas, H., Tilbrook, B., Tjiputra, J., Vandemark, D., Veness, T., Wanninkhof, R., Watson, A. J., Weiss, R., Wong, C. S., and Yoshikawa-Inoue, H.: A uniform, quality controlled Surface Ocean CO₂ Atlas (SOCAT), *Earth Syst. Sci. Data Discuss.*, 5, 735–780, doi:10.5194/essdd-5-735-2012, 2012.

Pomeroy, L. R., Sheldon, J. E., Sheldon, W. M., Blanton, J. O., Amft, J., and Peters, F.: Seasonal changes in microbial processes in estuarine and continental shelf waters of the southeastern USA, *Estuar. Coast. Shelf Sci.*, 51, 415–428, doi:10.1006/ecss.2000.0690, 2000.

Probst, J. L., Mortatti, J., and Tardy, Y.: Carbon river fluxes and weathering CO₂ consumption in the Congo and Amazon river basins, *Appl. Geochem.*, 9, 1–13, doi:10.1016/0883-2927(94)90047-7, 1994.

Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

1

A small red triangle icon, likely a play button or a navigation arrow.

1

Back

Close

Full Screen / Esc

[Printer-friendly Version](#)

Interactive Discussion



- Rabouille, C., Mackenzie, F. T., and Ver, L. M.: Influence of the human perturbation on carbon, nitrogen, and oxygen biogeochemical cycles in the global coastal ocean, *Geochim. Cosmochim. Acta*, 65, 3615–3641, doi:10.1016/s0016-7037(01)00760-8, 2001.

Radach, G. and Patsch, J.: Variability of continental riverine freshwater and nutrient inputs into the North Sea for the years 1977–2000 and its consequences for the assessment of eutrophication, *Estuar. Coast.*, 30, 66–81, 2007.

Ralison, O. H., Borges, A. V., Dehairs, F., Middelburg, J. J., and Bouillon, S.: Carbon biogeochemistry of the Betsiboka estuary (north-western Madagascar), *Org. Geochem.*, 39, 1649–1658, doi:10.1016/j.orggeochem.2008.01.010, 2008.

Raymond, P. A. and Bauer, J. E.: Bacterial consumption of DOC during transport through a temperate estuary, *Aquat. Microb. Ecol.*, 22, 1–12, doi:10.3354/ame022001, 2000.

Raymond, P. A. and Bauer, J. E.: Use of C-14 and C-13 natural abundances for evaluating riverine, estuarine, and coastal DOC and POC sources and cycling: a review and synthesis, *Org. Geochem.*, 32, 469–485, doi:10.1016/s0146-6380(00)00190-x, 2001.

Raymond, P. A. and Hopkinson, C. S.: Ecosystem modulation of dissolved carbon age in a temperate marsh-dominated estuary, *Ecosystems*, 6, 694–705, doi:10.1007/s10021-002-0213-6, 2003.

Raymond, P. A., Caraco, N. F., and Cole, J. J.: Carbon dioxide concentration and atmospheric flux in the Hudson River, *Estuaries*, 20, 381–390, doi:10.2307/1352351, 1997.

Raymond, P. A., Bauer, J. E., and Cole, J. J.: Atmospheric CO₂ evasion, dissolved inorganic carbon production, and net heterotrophy in the York River estuary, *Limnol. Oceanogr.*, 45, 1707–1717, 2000.

Ribas-Ribas, M., Gomez-Parra, A., and Forja, J. M.: Air-sea CO₂ fluxes in the north-eastern shelf of the Gulf of Cadiz (southwest Iberian Peninsula), *Mar. Chem.*, 123, 56–66, doi:10.1016/j.marchem.2010.09.005, 2011.

Richey, J. E.: Pathways of atmospheric CO₂ through fluvial systems, in: The global carbon cycle, integrating humans, climate, and the natural world, edited by: Field, C. B. and Raupach, M. R., Island Press, Washington, 329–340, 2004.

Ruiz-Halpern, S., Sejr, M. K., Duarte, C. M., Krause-Jensen, D., Dalsgaard, T., Dachs, J., and Rysgaard, S.: Air-water exchange and vertical profiles of organic carbon in a subarctic fjord, *Limnol. Oceanogr.*, 55, 1733–1740, doi:10.4319/lo.2010.55.4.1733, 2010.

Rysgaard, S., Mortensen, J., Juul-Pedersen, T., Sorensen, L. L., Lennert, K., Sogaard, D. H., Arendt, K. E., Blicher, M. E., Sejr, M. K., and Bendtsen, J.: High air-sea CO₂ uptake rates

in nearshore and shelf areas of Southern Greenland: Temporal and spatial variability, Mar. Chem., 128, 26–33, doi:10.1016/j.marchem.2011.11.002, 2012.

Sabine, C., Maenner, S., and Sutton., A.: High-resolution ocean and atmosphere $p\text{CO}_2$ time-series measurements from mooring LaPush 125° W 48° N., Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, Tennessee, 2011.

Sabine, C., Maenner, S., and Sutton., A.: High-resolution ocean and atmosphere $p\text{CO}_2$ time-series measurements from mooring CoastalMS 88° W, 30° N, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, Tennessee, 2012.

Sakamoto, A., Watanabe, Y. W., Osawa, M., Kido, K., and Noriki, S.: Time series of carbonate system variables off Otaru coast in Hokkaido, Japan, Estuar. Coast. Shelf Sci., 79, 377–386, doi:10.1016/j.ecss.2008.04.013, 2008.

Sarma, V., Kumar, M. D., and Manerikar, M.: Emission of carbon dioxide from a tropical estuarine system, Goa, India, Geophys. Res. Lett., 28, 1239–1242, doi:10.1029/2000gl006114, 2001.

Sarma, V., Kumar, N. A., Prasad, V. R., Venkataramana, V., Appalanaidu, S., Sridevi, B., Kumar, B. S. K., Bharati, M. D., Subbaiah, C. V., Acharyya, T., Rao, G. D., Viswanadham, R., Gawade, L., Manjary, D. T., Kumar, P. P., Rajeev, K., Reddy, N. P. C., Sarma, V. V., Kumar, M. D., Sadhuram, Y., and Murty, T. V. R.: High CO_2 emissions from the tropical Godavari estuary (India) associated with monsoon river discharges, Geophys. Res. Lett., 38, L08601, doi:10.1029/2011gl046928, 2011.

Sarma, V., Viswanadham, R., Rao, G. D., Prasad, V. R., Kumar, B. S. K., Naidu, S. A., Kumar, N. A., Rao, D. B., Sridevi, T., Krishna, M. S., Reddy, N. P. C., Sadhuram, Y., and Murty, T. V. R.: Carbon dioxide emissions from Indian monsoonal estuaries, Geophys. Res. Lett., 39, L03602, doi:10.1029/2011gl050709, 2012.

Schiettecatte, L. S., Thomas, H., Bozec, Y., and Borges, A. V.: High temporal coverage of carbon dioxide measurements in the Southern Bight of the North Sea, Mar. Chem., 106, 161–173, doi:10.1016/j.marchem.2007.01.001, 2007.

Schlunz, B. and Schneider, R. R.: Transport of terrestrial organic carbon to the oceans by rivers: re-estimating flux- and burial rates, Int. J. Earth Sci., 88, 599–606, doi:10.1007/s005310050290, 2000.

BGD

10, 5041–5105, 2013

Air-sea Exchanges of CO_2 in World's Coastal Seas

C.-T. A. Chen et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



- Schuster, U., McKinley, G. A., Bates, N., Chevallier, F., Doney, S. C., Fay, A. R., Gonzalez-Davila, M., Gruber, N., Jones, S., Krijnen, J., Landschutze, P., Lefevre, N., Manizza, M., Mathis, J., Metzl, N., Olsen, A., Rios, A. F., Rodenbeck, C., Santana-Casiano, J. M., Taka-hashi, T., Wanninkhof, R., and Watson, A. J.: Atlantic and Arctic sea-air CO₂ fluxes, 1990–2009, *Biogeosci. Disc.*, 9, 10669–10724, doi:10.5194/bgd-9-10669-2012, 2012.
- Seitzinger, S. P., Mayorga, E., Bouwman, A. F., Kroese, C., Beusen, A. H. W., Billen, G., Van Drecht, G., Dumont, E., Fekete, B. M., Garnier, J., and Harrison, J. A.: Global river nutrient export: A scenario analysis of past and future trends, *Glob. Biogeochem. Cy.*, 24, GB0a08 doi:10.1029/2009gb003587, 2010.
- Shadwick, E. H., Thomas, H., Azetsu-Scott, K., Greenan, B. J. W., Head, E., and Horne, E.: Seasonal variability of dissolved inorganic carbon and surface water *pCO₂* in the Scotian Shelf region of the Northwestern Atlantic, *Mar. Chem.*, 124, 23–37, doi:10.1016/j.marchem.2010.11.004, 2011.
- Shen, Z. L.: Historical changes in nutrient structure and its influences on phytoplankton composition in Jiaozhou Bay, *Estuar. Coast. Shelf Sci.*, 52, 211–224, doi:10.1006/ecss.2000.0736, 2001.
- Shen, Z. L., Liu, Q., and Zhang, S.: Distribution, variation and removal patterns of inorganic nitrogen in the Changjiang River, *Oceanol. Limnol. Sin.*, 34, 355–363, 2003.
- Shim, J., Kim, D., Kang, Y. C., Lee, J. H., Jang, S. T., and Kim, C. H.: Seasonal variations in *pCO₂* and its controlling factors in surface seawater of the northern East China Sea, *Cont. Shelf Res.*, 27, 2623–2636, doi:10.1016/j.csr.2007.07.005, 2007.
- Silvennoinen, H., Liikanen, A., Rintala, J., and Martikainen, P. J.: Greenhouse gas fluxes from the eutrophic Temmesjoki River and its Estuary in the Liminganlahti Bay (the Baltic Sea), *Biogeochemistry*, 90, 193–208, doi:10.1007/s10533-008-9244-1, 2008.
- Slomp, C. P. and Van Cappellen, P.: Nutrient inputs to the coastal ocean through submarine groundwater discharge: controls and potential impact, *J. Hydrol.*, 295, 64–86, doi:10.1016/j.jhydrol.2004.02.018, 2004.
- Smith, S. V. and Hollibaugh, J. T.: Coastal metabolism and the oceanic organic-carbon balance, *Rev. Geophys.*, 31, 75–89, doi:10.1029/92rg02584, 1993.
- Smith, S. V. and Mackenzie, F. T.: The ocean as a net heterotrophic system: implications from the carbon biogeochemical cycle, *Glob. Biogeochem. Cy.*, 1, 187–198, doi:10.1029/GB001i003p00187, 1987.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

1

A small red triangle icon, likely a play button or a navigation arrow.

1

1

ANSWER

Digitized by srujanika@gmail.com

Interactive Discussion

- Souza, M., Gomes, V., Freitas, S., Andrade, R., and Knoppers, B.: Net ecosystem metabolism and nonconservative fluxes of organic matter in a tropical mangrove estuary, Piauí River (NE of Brazil), *Estuar. Coast.*, 32, 111–122, doi:10.1007/s12237-008-9104-1, 2009.

Stainton, E. C.: Air-water carbon dioxide exchange in relation to chemical and physical characteristics of the Churchill River and estuary, southwestern Hudson Bay region Master of Science, Department of Environment and Geography University of Manitoba, Winnipeg, 123 pp., 2009.

Sweeney, C.: The annual cycle of surface CO₂ and O₂ in the Ross Sea: A model for gas exchange on the continental shelves of Antarctica, *Antarct. Res. Ser.*, 78, 295–312, 2003.

Syed, T. H., Famiglietti, J. S., Chambers, D. P., Willis, J. K., and Hilburn, K.: Satellite-based global-ocean mass balance estimates of interannual variability and emerging trends in continental freshwater discharge, *Proc. Natl. Acad. Sci. USA*, 107, 17916–17921, doi:10.1073/pnas.1003292107, 2010.

Svitshi, J. P., Vorosmarty, C. J., Kettner, A. J., and Green, P.: Impact of humans on the flux of terrestrial sediment to the global coastal ocean, *Science*, 308, 376–380, doi:10.1126/science.1109454, 2005.

Takahashi, T., Sutherland, S. C., Sweeney, C., Poisson, A., Metzl, N., Tilbrook, B., Bates, N., Wanninkhof, R., Feely, R. A., Sabine, C., Olafsson, J., and Nojiri, Y.: Global sea-air CO₂ flux based on climatological surface ocean pCO₂, and seasonal biological and temperature effects, 3rd International Symposium on Climatic Changes and the Cycle of Carbon, Brest, France, ISI:000175992000002, 1601–1622, 2002

Takahashi, T., Sutherland, S. C., and Kozyr, A.: Global ocean surface water partial pressure of CO₂ database: Measurements performed during 1957–2011 (Version 2011), Oak Ridge, Tennessee, 2012.

Ternon, J. F., Oudot, C., Dessier, A., and Diverres, D.: A seasonal tropical sink for atmospheric CO₂ in the Atlantic ocean: the role of the Amazon River discharge, *Mar. Chem.*, 68, 183–201, doi:10.1016/s0304-4203(99)00077-8, 2000.

Thomas, H., and Schneider, B.: The seasonal cycle of carbon dioxide in Baltic Sea surface waters, *J. Mar. Syst.*, 22, 53–67, doi:10.1016/s0924-7963(99)00030-5, 1999.

Thomas, H., Bozec, Y., Elkalay, K., and de Baar, H. J.: Enhanced open ocean storage of CO₂ from shelf sea pumping, *Science*, 304, 1005–1008, doi:10.1126/science.1095491, 2004.

Thomas, H., Prowe, A. E. F., van Heuven, S., Bozec, Y., de Baar, H. J. W., Schiettecatte, L. S., Suykens, K., Kone, M., Borges, A. V., Lima, I. D., and Doney, S. C.: Rapid decline of the



Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



**Air-sea Exchanges
of CO₂ in World's
Coastal Seas**

C.-T. A. Chen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶|](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Wollast, R.: Evaluation and comparison of the global carbon cycle in the coastal zone and in the open ocean, in: *The Global Coastal Ocean*, edited by: Brink, K. H. and Robinson, A. R., Wiley & Sons, New York, 213–252, 1998.

5 Woodwell, G. M., Houghton, R. A., and Tempel, N. R.: Atmospheric CO₂ at Brookhaven, long-island, New York – patterns of variation up to 125 meters, *J. Geophys. Res.*, 78, 932–940, doi:10.1029/JC078i006p00932, 1973.

Xue, L., Zhang, L. J., Cai, W. J., and Jiang, L. Q.: Air-sea CO₂ fluxes in the southern Yellow Sea: An examination of the continental shelf pump hypothesis, *Cont. Shelf Res.*, 31, 1904–1914, doi:10.1016/j.csr.2011.09.002, 2011.

10 Xue, L., Xue, M., Zhang, L. J., Sun, T. M., Guo, Z., and Wang, J. J.: Surface partial pressure of CO₂ and air-sea exchange in the northern Yellow Sea, *J. Mar. Syst.*, 105, 194–206, doi:10.1016/j.jmarsys.2012.08.006, 2012.

Yool, A. and Fasham, M. J. R.: An examination of the “continental shelf pump” in an open ocean general circulation model, *Glob. Biogeochem. Cy.*, 15, 831–844, doi:10.1029/2000gb001359, 2001.

15 Zemmelink, H. J., Slagter, H. A., van Slooten, C., Snoek, J., Heusinkveld, B., Elbers, J., Bink, N. J., Klaassen, W., Philippart, C. J. M., and de Baar, H. J. W.: Primary production and eddy correlation measurements of CO₂ exchange over an intertidal estuary, *Geophys. Res. Lett.*, 36, L19606 doi:10.1029/2009gl039285, 2009.

20 Zeng, F. W., Masiello, C. A., and Hockaday, W. C.: Controls on the origin and cycling of riverine dissolved inorganic carbon in the Brazos River, Texas, *Biogeochemistry*, 104, 275–291, doi:10.1007/s10533-010-9501-y, 2011.

Zhai, W. D. and Dai, M. H.: On the seasonal variation of air-sea CO₂ fluxes in the outer Changjiang (Yangtze River) Estuary, East China Sea, *Mar. Chem.*, 117, 2–10, doi:10.1016/j.marchem.2009.02.008, 2009.

25 Zhai, W., Dai, M., and Guo, X.: Carbonate system and CO₂ degassing fluxes in the inner estuary of Changjiang (Yangtze) River, China, *Mar. Chem.*, 107, 342–356, doi:10.1016/j.marchem.2007.02.011, 2007.

30 Zhai, W. D., Dai, M. H., Cai, W. J., Wang, Y. C., and Wang, Z. H.: High partial pressure of CO₂ and its maintaining mechanism in a subtropical estuary: the Pearl River estuary, China, *Mar. Chem.*, 93, 21–32, doi:10.1016/j.marchem.2004.07.003, 2005.

Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.

Title Page

Abstrac

Introduction

Conclusions

References:

Tables

Figures



Back

Close

Full Screen / Esc

[Printer-friendly Version](#)

Interactive Discussion



Table 1. Seasonal and annual air-to-sea fluxes of CO₂ in world's estuaries.

type	lon.(°)	lat.(°)	spring flux ^c (mmol C m ⁻² d ⁻¹)	summer flux (mmol C m ⁻² d ⁻¹)	autumn flux (mmol C m ⁻² d ⁻¹)	winter flux (mmol C m ⁻² d ⁻¹)	annual flux (mol C m ⁻² yr ⁻¹)	References
1-1 (fjord) (US) ^b	-152.5	57.7	-1.8		1.8		0.001	Takahashi et al. (2012) (LDED database)
11-1 (fjord) (CA)	-55.8	52.3		-2.1			-0.8	Takahashi et al. (2012) (LDED database)
14-1 (fjord) (IC)	-23.2	66.2	-0.7	-7.0	-12.9	-4.8	-2.3	Takahashi et al. (2012) (LDED database)
14-2 (fjord) (IC)	-23.6	66.1		-7.7			-2.8	Takahashi et al. (2012) (LDED database)
14-3 (fjord) (IC)	-23.7	65.7			5.4		2.0	Takahashi et al. (2012) (LDED database)
14-4 (fjord) (IC)	-24.1	65.6	-0.3				-0.1	Takahashi et al. (2012) (LDED database)
14-5 (fjord) (IC)	-18.6	66.0	-48.2	-7.8	-11.2	-9.0	-7.0	Takahashi et al. (2012) (LDED database)
Aby lagoon (CI)	-3.3	5.4	-10.1	1.2	-11.3	-4.1	-2.7	Kone et al. (2009)
Altamaha Sound (US)	-81.3	31.3	57.8	127.0	79.7	28.5	26.8	Jiang et al. (2008a)
Ambalaayaar (IN)	79.3	10.0		-0.02			-0.007	Sarma et al., (2012)
Amur River (RU)	141.1	52.9		0.1	1.5		0.3	Johnson et al. (2009) (WOD09 database)
Ason (ES)	-3.5	43.3		-3.0			-1.1	Ortega et al. (2005)
Aveiro lagoon (PT)	-8.7	40.7					12.4	Borges and Frankignoulle (unpublished)
Baitarani (IN)	86.9	20.5		20.7			7.6	Sarma et al. (2012)
Bancal (PH)	115.0	5.0	2.2				0.8	Chen (unpublished)
Bebar River (MY)	103.4	3.1					6.5	Chen (unpublished)
Bellamy (US)	-70.9	43.1	-11.0	43.0	6.0		4.6	Hunt et al. (2011)
Betsiboka (MG)	46.3	-15.7					3.3	Raison et al. (2008)
Bharatkaluzza (IN)	76.0	11.2		11.7			4.3	Sarma et al. (2012)
Bohniian Bay (FI)	21.0	63.0					3.5	Algesten et al. (2004)
Brazos River (US)	-95.4	28.9					0.033	Zeng et al. (2011)
Brunei River (BN)	96.4	16.5		53.7			19.6	Chen (unpublished)
Cauvery (IN)	79.89	11.26		2.23			0.8	Sarma et al. (2012)
Chalakudi (IN)	76.18	10.69		12.86			4.70	Sarma et al. (2012)
Changjiang (Yantze) (CN)	120.5	31.5	23.5	65.5	33.7	37.8	14.6	Zhai et al. (2007)
Chi Shui River (TW)	120.11	23.29		176		68.5	44.6	Chen (unpublished)
Chilka (lagoon) (IN)	85.5	19.1	9.8	141.0			27.5	Gupta et al. (2008)
Cho Shui River (TW)	120.3	23.9	651.0	13.4			121.0	Chen (unpublished)
Chung Kang River (TW)	120.8	24.7	45.8	53.4	28.8	144.0	24.8	Chen (unpublished)
Churchill River (CA)	-94.2	58.8		1.2	-3.6		-0.4	Stainton (2009)
Citanduy-Manganit (ID)	108.8	-7.7	25.7 ^d				9.4	Chen (unpublished)
Ciujung-Kragilan (ID)	106.4	-6.0	36.9 ^d				13.5	Chen (unpublished)
Cocheo (US)	-70.8	43.1	2.0	26.0	2.0		3.7	Hunt et al. (2011)
Cochin (IN)	76.0	9.5			267.0	65.0	60.6	Gupta et al. (2008)
Cross Sound (fjord) (US)	-134.1	56.6		-0.2	45.1		8.2	Takahashi et al. (2012)
(LDED database)								

Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Continued.

type	lon.(°)	lat.(°)	spring flux ^c (mmol C m ⁻² d ⁻¹)	summer flux (mmol C m ⁻² d ⁻¹)	autumn flux (mmol C m ⁻² d ⁻¹)	winter flux (mmol C m ⁻² d ⁻¹)	annual flux (mol C m ⁻² yr ⁻¹)	References
Doboy Sound (US)	-81.3	31.4	15.2	47.4	51.0	16.0	11.9	Jiang et al. (2008a)
Douro (PT)	-8.7	41.1			240.0		87.6	Frankignoulle et al. (1998)
Duplin River (US)	-81.3	31.5	53.4	83.0	73.2	23.4	21.3	Wang and Cai (2004)
Ebrí lagoon (CI)	-4.3	5.5	56.4	109.0	61.9	47.9	26.6	Kone et al. (2009)
Elbe (DE)	8.8	53.9	180.0				65.7	Frankignoulle et al. (1998)
Ems (DE)	6.9	53.4		110.0			40.2	Frankignoulle et al. (1998)
Endau River (MY)	103.6	2.7			1.0		0.4	Chen (unpublished)
Erh Jen River (TW)	120.2	22.9	68.5	11.1		26.5	12.9	Chen (unpublished)
Florida Bay (US)	-80.8	25.0					1.7	Miller et al. (2001)
Fong Kang River (TW)	120.7	22.2	6.7	-17.9		18.0	0.8	Chen (unpublished)
Gaderu creek (IN)	82.3	16.8		56.0			20.4	Borges et al. (2003)
Gironde (FR)	-1.1	45.6	110.0	110.0	65.0	50.0	30.6	Frankignoulle et al. (1998)
Godavari (IN)	82.3	16.7					8.0	Bouillon et al. (2003), Sarma et al. (2012)
Godthåbsfjord (GL) ^e	-51.9	64.1					-7.25	Rysgaard et al. (2012)
Golfo Almirante Montt (fjord) (CL)	-72.0	-52.1			-17.7 ^d		-6.5	Takahashi et al. (2012) (LDED database)
Great Bay (US)	-70.9	43.1					3.6	Hunt et al. (2011)
Guadaluquivir (ES)	-6.0	37.4		104.0			37.9	de la Paz et al. (2007)
Haldia (IN)	88.2	21.9		12.3			4.5	Sarma et al. (2012)
Hanjiang (CN)	116.8	23.4				0.9	0.3	Chen (unpublished)
Ho Ping River (TW)	121.8	24.3	5.3	22.0		68.5	11.7	Chen (unpublished)
Hooghly (IN)	88.0	22.0	31.8	-1.1	16.7	2.5	4.9	Mukhopadhyay et al. (2002)
Hou Lung River (TW)	120.8	24.6	72.9	9.3	7.6	21.0	10.1	Chen (unpublished)
Hsiu Ku Luan River (TW)	121.5	23.5	26.5	41.9		19.2	10.7	Chen (unpublished)
Hua Lien River (TW)	121.6	23.9	93.4	75.3		4.8	21.1	Chen (unpublished)
Hudson River Estuary (US)	-74.0	40.7					5.9	Raymond et al. (1997)
Isla Gordon (fjord) (CL)	-68.9	-55.2			-1.2 ^d		-0.4	Takahashi et al. (2012) (LDED database)
Itacuraca creek Sepetiba (Bay)(BR)	-44.0	-23.0					41.4	Ovalle et al. (1990), Borges et al. (2003)
Jiulong Jiang (Xiamen Bay) (CN)	118.1	24.5					0.5	Dai et al. (2009)
Jiulongjiang (CN)	118.0	24.5				4.3	1.6	Chen (unpublished)
Johor River (MY)	104.0	1.5			2.3		0.8	Chen (unpublished)
Kakinada Bay (IN)	82.3	16.7					3.0	Bouillon et al. (2003)
Kali (IN)	74.2	14.8		3.2			1.2	Sarma et al. (2012)
Kaneohe Bay and stream (US)	-157.8	21.5					1.5	Fagan and Mackenzie (2007)
Kao Ping River (TW)	120.4	22.5	98.1	51.8	30.5	12.4	17.6	Chen (unpublished)
Kapuas River (ID)	109.1	0.1				148.3	54.1	Chen (unpublished)

Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Table 1. Continued.

type	lon.(°)	lat.(°)	spring flux ^c (mmol C m ⁻² d ⁻¹)	summer flux (mmol C m ⁻² d ⁻¹)	autumn flux (mmol C m ⁻² d ⁻¹)	winter flux (mmol C m ⁻² d ⁻¹)	annual flux (mol C m ⁻² yr ⁻¹)	References
Kennebec River (US)	-69.8	43.8		22.5	22.0	-0.2	-49.6	-0.5
Khura River estuary(TH)	98.3	9.2						35.7
Kidogoweni creek (Gazi Bay) (KE)	39.5	-4.4						Miyajima et al. (2009)
Kien Vang creeks(VN)	105.1	8.7		32.2		154.7		21.8
Klang River (MY)	101.4	3.0				7.7		Bouillon et al. (2007b)
Kobbe fjord (GL)	-51.5	64.2		-2.7		-136.6		34.2
Kochi back waters (IN)	76.4	10.0			8.1			Chen (unpublished)
Kola Bay (RU)	33.4	69.1		-2.5	-0.2	-3.5		2.8
Krishna (IN)	81.1	15.8			6.8			Sarma et al. (2012)
Lan Yang River (TW)	121.8	24.7		65.5	66.0			2.9
Limingenlahti Bay (FI)	25.4	64.9			-0.9			Johnson et al. (2009)
Lin Pien River (TW)	120.5	22.4		44.4	54.5			(WOD09 database)
Little Bay (US)	-70.9	43.1		-5.1	33.9	3.9		
Loire (FR)	-2.2	47.2				155.0		
Luohu (CN)	115.6	22.9					0.1	
Mahanadi (IN)	86.6	20.0			3.1			0.022
Mahisagar (IN)	72.6	22.1						Chen (unpublished)
Mandovi (IN)	73.8	15.7			18.1			1.1
Mandovi-Zuari (IN)	73.5	15.3						Sarma et al. (2012)
Matolo creek (KE)	40.1	-2.1						3.7
Mekong (VN)	106.5	10.0						Sarma et al. (2012)
Mempawah River (ID)	89.0	22.0			23.2			6.6
Moni (TZ)	39.3	-6.9						Sarma et al. (2012)
Nagada creek (Papua New Guinea) (ID)	145.8	-5.2				43.6 ^d		14.2
Nagavalil (IN)	84.0	18.2			0.2			Sarma et al. (2001)
Nalongne (CN)	112.0	21.8				10.1		Bouillon et al. (2007a)
Narmada (IN)	73.0	20.2			8.8			21.2
Netravathi (IN)	75.0	12.7			70.7			Borges (unpublished)
Norman's Pond (BS)	-76.1	23.8					30.8	
Orinoco River (VE)	-62.3	8.6		31.8				Chen (unpublished)
Oyster (US)	-70.9	43.1		-17.2	51.5	2.5		8.5
Pa Chang River (TW)	120.1	23.3		29.9	94.2			Kristensen et al. (2008)
Patang River (MY)	103.5	3.5				34.8		2.4
Palau lagoon (PW)	134.5	7.5		0.03		3.5		Borges et al. (2003)
Parker River estuary (US)	-70.8	42.8			3.2	-1.0		15.9
Hopkinson (2003)								0.1
Pei Kang River (TW)	120.2	23.5		27.3	80.0	35.1		Sarma et al. (2012)
Pei Nan River (TW)	121.2	22.8		155.0	96.2			3.7
Penna (IN)	80.2	14.4				5.2		3.2
Piaui River estuary (BR)	-37.5	-11.5						3.2

Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Continued.

type	lon.(°)	lat.(°)	spring flux ^c (mmol C m ⁻² d ⁻¹)	summer flux (mmol C m ⁻² d ⁻¹)	autumn flux (mmol C m ⁻² d ⁻¹)	winter flux (mmol C m ⁻² d ⁻¹)	annual flux (mol C m ⁻² yr ⁻¹)	References
Po Tzu River (TW)	120.1	23.4		85.5		89.9	32.0	Chen (unpublished)
Ponnyaar (IN)	80.3	12.4		96.3			35.2	Chen (unpublished)
Potou lagoon (CI)	-3.8	5.6	40.3	186.0	45.5	82.7	36.8	Kone et al. (2009)
Qiantang River (CN)	122.0	30.1					0.1	Chen (unpublished)
Qinjiang (CN)	108.6	21.7				6.3	2.3	Chen (unpublished)
Rajang River (MY)	115.5	2.1			7.1		2.6	Chen (unpublished)
Randers Fjord (DK)	10.3	56.6	-5.0	52.9			8.7	Gazeau et al. (2005)
Ras Dege creek (TZ)	39.3	-6.9					12.0	Kristensen et al. (2008), Bouillon et al. (2007c)
Rhine (NL)	4.1	52.0					21.9	Frankignoulle et al. (1998)
Ria de Vigo (FR)	8.6	42.1	-0.1		-0.5	0.5	-0.1	Álvarez-Salgado et al. (1999)
Rio San Pedro (ES)	-6.1	36.4					39.4	Ferron et al. (2007)
Rompin River (MY)	103.5	2.8			1.5		0.6	Chen (unpublished)
Rongjiang (CN)	116.7	23.3				14.0	5.1	Chen (unpublished)
Rushikulya (IN)	85.2	19.3		-0.02			-0.01	Sarma et al. (2012)
S. Muar (MY)	102.6	2.0			3.2		1.2	Chen (unpublished)
Sabarmathi (IN)	72.8	21.6		13.8			5.1	Sarma et al. (2012)
Sado (PT)	-8.9	38.5			396.0		145.0	Frankignoulle et al. (1998)
Saja-Besaya (ES)	-4.0	43.4		446.0			163.0	Ortega et al. (2005)
São Francisco Estuary (US)	-122.3	37.7			1.8	0.5	0.4	Peterson (1979)
Sapelo Sound (US)	-81.3	31.6	19.1		41.1	47.1	16.8	Jiang et al. (2008a)
Saptamukhi creek (IN)	89.0	22.0					20.7	Ghosh et al. (1987), Borges et al. (2003)
Satilla River (US)	-81.5	31.0			116.0		42.5	Cai and Wang (1998)
Scheldt (BE/NL)	3.5	51.4	175.0	233.0	326.0	240.0	94.1	Frankignoulle et al. (1998)
Sedili Besar (MY)	104.1	1.9			12.6		4.6	Chen (unpublished)
Sentosa River (MY)	104.1	1.9			17.2		6.3	Chen (unpublished)
Sharavathi (IN)	74.5	14.4					3.7	Sarma et al. (2012)
Shark River (US)	-81.1	25.2					16.0	Kone and Borges (2008)
Skagit River (US)	-130.1	53.9			65.6		23.9	Takahashi et al. (2012) (LDED database)
Subarnalekha (IN)	87.6	21.5		0.03			0.01	Sarma et al. (2012)
Szu Chung River (TW)	120.7	22.1	12.9	50.4			7.6	Chen (unpublished)
Ta An River (TW)	120.6	24.4	-0.4	3.4	27.3	17.0	4.3	Chen (unpublished)
Ta Chia River (TW)	120.6	24.3	-6.3	25.3	-29.2		-1.2	Chen (unpublished)
Tagba lagoon (CI)	-5.0	5.4	18.1	114.0	28.5	13.2	18.5	Kone et al. (2009)
Tam Giang creeks (VN)	105.2	8.8	141.5		128.5		49.3	Kone and Borges (2008)
Tamar (UK)	-4.2	50.4	90.1	120.0			38.3	Frankignoulle et al. (1998)
Tan Shui River (TW)	121.5	25.1	168.0	160.0	214.0	3.3	49.8	Chen (unpublished)
Tana (KE)	40.1	-2.1					21.2	Bouillon et al. (2007a)
Tapti (IN)	72.7	21.1		362.5			132.4	Sarma et al (2012)

Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Continued.

type	lon.(°)	lat.(°)	spring flux ^c (mmol C m ⁻² d ⁻¹)	summer flux (mmol C m ⁻² d ⁻¹)	autumn flux (mmol C m ⁻² d ⁻¹)	winter flux (mmol C m ⁻² d ⁻¹)	annual flux (mol C m ⁻² yr ⁻¹)	References
Tendo laggon (CI)	-3.2	5.3	-17.7	75.6	-4.9	-3.0	7.0	Kone et al. (2009)
Thames (UK)	0.9	51.5			250.0		91.3	Frankignoulle et al. (1998)
Tou Chien River (TW)	120.9	24.8	55.9	10.5	7.2	46.6	11.0	Chen (unpublished)
Trang River estuary (TH)	99.4	7.2					30.9	Miyajima et al. (2009)
Tseng Wen River (TW)	120.1	23.1	93.2	-1.8	12.4		34.6	Chen (unpublished)
Tung Kang River (TW)	120.4	22.5	114.0	160.0	48.9	121.0	40.5	Chen (unpublished)
Urdabai (ES)	2.7	43.4		22.8			8.3	Ortega et al. (2005)
Vaigai (IN)	78.9	9.3		0.2			0.1	Sarma et al. (2012)
Vamsadhara (IN)	84.7	18.9		0.4			0.1	Sarma et al. (2012)
Vellar (IN)	79.9	11.7		17.0			6.2	Sarma et al. (2012)
Wadden Sea estuary (NL)	4.8	53.0	-160.0				-58.4	Zemmelink et al. (2009)
Waialoa river-estuary (US) ^e	-159.5	22.2	1032		422	607	251	Paquay et al. (2007)
Wailuku River (US)	-155.08	19.72	5.73				5.73	Paquay et al. (2007)
Wu River (TW)	120.5	24.2	44.4	92.1			24.9	Chen (unpublished)
Yangon (MM)	121.8	31.3				5.4	2.0	Chen (unpublished)
Yen Shui River (TW)	120.2	23.0	50.1	125.0		14.4	23.1	Chen (unpublished)
Yenisey (RU)	82.7	71.8	29.7	16.7	3.5	27.5	7.1	Johnson et al. (2009) (WOD09 database)
York River (US)	-78.4	37.2	10.0	29.0	16.7	6.5	5.6	Raymond et al. (2000)
Zhujiang (Pearl River) (CN)	113.5	22.5	60.2	70.7	47.0	22.2	6.9	Guo et al. (2009)
Zuari (IN)	74.0	-15.3		6.4			2.3	Sarma et al. (2012)

^a Positive fluxes indicate an emission of CO₂ from water to the atmosphere;

^b BE: Belgium; BN: Brunei; BR: Brazil; BS: Bahamas; CI: Côte d'Ivoire, CL: Chile; CN: China; DE: Germany; DK: Denmark; ES: Spain; FI: Finland; FR: France; GL: Greenland; IC: Iceland; ID: Indonesia; IN: India; KE: Kenya; MG: Repoblikan'i Madagasikara; MM: Myanmar; MY: Malaysia; NL: Netherlands; PH: Philippines; PT: Portugal; PW: Palau; RU: Russia; TH: Thailand; TW: Taiwan; TZ: Tunisia; UK: United Kingdom; US: United States; VE: Venezuela; VN: Vietnam;

^c Spring: March–May; Summer: June–August; Fall: September–November; Winter: December–February;

^d Austral seasons;

^e not used in the calculation.

Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Table 2. Summary of reported total air-to-sea fluxes of CO₂ in world's estuaries.

	Unit area flux (mol C m ⁻² yr ⁻¹)	Area (10 ⁶ km ²)	Total flux (PgCyr ⁻¹)	References
Estuaries (<i>n</i> = 19)	35.71	1.40	0.60	Abril and Borges (2005)
Estuaries (<i>n</i> = 16)	38.12	0.94	0.43	
Non-estuarine salt marshes (<i>n</i> = 1)	23.45	0.14	0.04	
Mangroves	13.66	0.20	0.04	Borges (2005)
Ave/Total	33.20	1.28	0.51	
Estuaries (<i>n</i> = 16)	28.62	0.94	0.32	
Non-estuarine salt marshes	21.40	0.14	0.036	
Mangroves	18.66	0.15	0.033	Borges et al. (2005)
Ave/Total	26.42	1.23	0.39	
Estuaries (<i>n</i> = 32)	32.10	0.943	0.36	
Non-estuarine salt marshes	30.40	0.384	0.09	
Mangroves	27.10	0.147	0.05	Chen and Borges (2009)
Ave/Total	28.27	1.474	0.50	
Small deltas and estuaries	25.7 ± 15.8	0.084	0.026 ± 0.016	
Tidal systems and embayments	28.5 ± 24.9	0.276	0.094 ± 0.082	
Lagoons	17.3 ± 16.6	0.252	0.052 ± 0.050	Laruelle et al. (2010)
Fjords and fjords	17.5 ± 14.0	0.456	0.096 ± 0.077	
Ave/Total (<i>n</i> =60)	21.0 ± 17.6	1.067	0.268 ± 0.225	
Estuaries (including both river-dominated and nonriverine coastal lagoons)	20.83	1.05	0.25	Cai (2011)
Estuaries (<i>n</i> = 106)	23.9 ± 33.1	1.07	0.26	Chen et al. (2012)
Estuaries (<i>n</i> = 165)	7.74	1.01	0.094	This study

MARCATS Segments Number	continent	ocean	System Name	Class	Estuarine Surface (10^3 km^2)	average CO ₂ flux (mol C m ⁻² yr ⁻¹)	CO ₂ flux (TgCyr ⁻¹)	Shelf Surface (10^3 km^2)	average CO ₂ flux (mol C m ⁻² yr ⁻¹)	CO ₂ flux (TgCyr ⁻¹)
1	NA	PA	North Eastern Pacific	Subpolar	33.9	10.71 ($n = 3$)	4.36	461	-3.51 ($n = 3$)	-19.40
2	NA/OC	PA	Californian Current	EBC	8.9	0.93 ($n = 3$)	0.10	214	-2.22 ($n = 3$)	-5.69
3	NA	PA	Tropical Eastern Pacific	Tropical	6.2	14.47	1.08	198	-0.05 ($n = 2$)	-0.13
4	SA	PA	Peruvian Upwelling Current	EBC	4.2	34.71	1.75	143	-0.62 ($n = 4$)	-1.07
5	SA	AT	Southern America	Subpolar	22	-3.46 ($n = 2$)	-0.91	1230	-3.25 ($n = 3$)	-47.98
6	SA	AT	Brazilian Current	WBC	26.3	28.20 ($n = 2$)	8.90	521	3.97 ($n = 1$)	24.81
7	SA	AT	Tropical Western Atlantic	Tropical	13.4	11.60 ($n = 1$)	1.87	517	-12.78 ($n = 1$)	-79.26
8	NA	AT	Caribbean Sea	Tropical	26.2	5.04 ($n = 1$)	1.58	344	0.66 ($n = 1$)	2.74
9	NA	AT	Gulf of Mexico	Marginal Sea	31.9	8.02 ($n = 2$)	3.07	544	-0.19 ($n = 2$)	-1.26
10	NA	AT	Florida Upwelling	WBC	34	9.81 ($n = 15$)	4.00	858	-1.10 ($n = 4$)	-11.27
11	NA	AT	Sea of Labrador	Subpolar	36.1	-0.76 ($n = 1$)	-0.33	395	-2.11 ($n = 1$)	-10.02
12	NA	AT	Hudson Bay	Marginal Sea	39	-0.44 ($n = 1$)	-0.20	1064	0.84 ($n = 1$)	10.73
13	NA	AR	Canadian Archipelagos	Polar	163.7	-1.08	-2.11	1177	-4.06 ($n = 2$)	-57.34
14	NA	AR	Northern Greenland	Polar	24.1	-2.05 ($n = 5$)	-0.59	614	-6.14 ($n = 1$)	-45.20
15	NA	AR	Southern Greenland	Polar	8.8	-1.08	-0.11	270	-5.95 ($n = 1$)	-19.29
16	EU	AR	Norwegian Basin	Polar	17	-17.30 ($n = 1$)	-3.53	171	-3.63 ($n = 1$)	-7.45
17	EU	AT	North Eastern Atlantic	Marginal Sea	37.6	37.73 ($n=8$)	17.02	1112	-1.04 ($n = 2$)	-13.88
18	EU	AT	Baltic Sea	Marginal Sea	26.3	1.28 ($n = 2$)	0.40	383	-1.95 ($n = 1$)	-8.96
19	EU	AT	Iberian Upwelling	EBC	12.7	58.75 ($n = 10$)	8.95	283	-1.33 ($n = 5$)	-4.51
20	EU	AT	Mediterranean Sea	Marginal Sea	15.1	-0.06 ($n = 1$)	-0.01	580	1.47 ($n = 3$)	10.21
21	EU	AT	Black Sea	Marginal Sea	10.3	10.00	1.24	172	-0.79	-1.63
22	AF	AT	Moroccan Upwelling	EBC	5.6	34.71	2.33	225	3.02 ($n = 1$)	8.15
23	AF	AT	Tropical Eastern Atlantic	Tropical	26.6	17.25 ($n = 5$)	5.51	284	0.29 ($n = 1$)	0.99
24	AF	AT	Southern Western Africa	EBC	1.7	34.71	0.71	308	-2.41 ($n = 1$)	-8.91
25	AF	IN	Aguilhas Current	WBC	28.4	14.52	4.95	254	-4.03 ($n = 1$)	-12.28
26	AF	IN	Tropical Western Indian	Tropical	5.8	15.73 ($n = 5$)	1.09	72	1.03 ($n = 1$)	0.89
27	AF	IN	Western Arabian Sea	Indian Margins	2	3.32 ($n = 1$)	0.08	102	-0.32 ($n = 2$)	-0.40
28	AF	IN	Red Sea	Marginal Sea	0.04	10.00	0.005	190	0.12 ($n = 2$)	0.28
29	AS	IN	Persian Gulf	Marginal Sea	2.3	10.00	0.28	233	-0.79	-2.20
30	AS	IN	Eastern Arabian Sea	Indian Margins	14.5	9.02 ($n = 25$)	1.57	342	0.01 ($n = 1$)	0.06
31	AS	IN	Bay of Bengal	Indian Margins	10.1	19.82 ($n = 10$)	2.40	230	-0.22 ($n = 1$)	-0.60
32	AS	IN	Tropical Eastern Indian	Indian Margins	16.2	13.73 ($n = 6$)	2.67	809	-0.28 ($n = 4$)	-2.74

Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.

Title Page

Abstract

Introduction

Conclusion

References

Tables

Figures



Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.

Table 3. Areas and air-sea fluxes of CO₂ in estuaries and continental shelves by biogeochemical provinces.

MARCATS Segments Number	continent	ocean	System Name	Class	Estuarine Surface (10 ³ km ²)	average CO ₂ flux (mol C m ⁻² yr ⁻¹)	CO ₂ flux (TgCyr ⁻¹)	Shelf Surface (10 ³ km ²)	average CO ₂ flux (mol C m ⁻² yr ⁻¹)	CO ₂ flux (TgCyr ⁻¹)
33	OC	IN	Leeuwin Current	EBC	0.6	34.71	0.25	118	-0.58 (n = 1)	-0.82
34	OC	PA	Southern Australia	Subpolar	13.1	2.82	0.44	452	-0.94 (n = 1)	-5.12
35	OC	PA	Eastern Australian Current	WBC	7.9	14.52	1.38	139	-0.19 (n = 3)	-0.31
36	OC	PA	New Zealand	Subpolar	7.3	2.82	0.25	283	-0.17 (n = 1)	-0.58
37	AS	PA	Northern Australia	Tropical	40.5	15.90 (n = 1)	7.73	2463	0.11 (n = 3)	3.35
38	AS	PA	South East Asia	Tropical	45.6	17.70 (n = 49)	9.68	2318	0.86 (n = 1)	23.92
39	AS	PA	East China Sea and Kuroshio	WBC	27.8	7.33 (n = 2)	2.44	1299	1.04 (n = 8)	16.26
40	AS	PA	Sea of Japan	Marginal Sea	6.7	10.00	0.80	277	-3.89 (n = 2)	-12.93
41	AS	PA	Sea of Okhotsk	Marginal Sea	19.7	0.30 (n = 1)	0.07	992	-1.67 (n = 1)	-19.82
42	AS	PA	North Western Pacific	Subpolar	22.3	2.82	0.76	1082	-2.12 (n = 2)	-27.56
43	AS	AR	Siberian Shelves	Polar	37.8	-1.08	-0.49	1918	0.01 (n = 1)	0.25
44	AS	AR	Barent and Kara Seas	Polar	72.2	3.07 (n = 2)	2.66	1727	0.01 (n = 1)	0.23
45	AN	AN	Antarctic Shelves	Polar	—			2952	-1.98 (n = 2)	-69.96
Total					1012.44	7.74	94.08	30320	-1.09	-395.71

* Shaded area is regions without data and data from a similar region is given.





Table 4. Seasonal and annual air-sea fluxes of CO₂ in world's continental shelves.

type	Lon.(°)	Lat.(°)	spring fluxes ^b (mmol C m ⁻² d ⁻¹)	summer fluxes (mmol C m ⁻² d ⁻¹)	autumn fluxes (mmol C m ⁻² d ⁻¹)	winter fluxes (mmol C m ⁻² d ⁻¹)	annual flux (mol C m ⁻² yr ⁻¹)	References
1 NEP	-155.9	56.4	-9.31	-12.22	-10.30		-3.87	Pfeil et al. (2012) (SOCAT database)
10T	-1.7	4.5	0.80				0.29	Takahashi et al. (2012) (LDED database)
11EBC	20.1	-35.7		-5.79 ^c	-7.42 ^c		-2.41	Takahashi et al. (2012) (LDED database)
11LAB	-53.4	47.0	-9.77	0.46	-5.91	-7.95	-2.11	Pfeil et al. (2012) (SOCAT database)
14 NGR	-14.0	69.2	-14.98	-18.64			-6.14	Pfeil et al. (2012) (SOCAT database)
14WBC	26.2	-34.1		-11.04 ^c			-4.03	Takahashi et al. (2012) (LDED database)
15 SGR	-28.5	63.8	-16.61	-40.20	-6.07	-2.37	-5.95	Pfeil et al. (2012) (SOCAT database)
16 NOR	14.1	66.9	-11.65	-6.41	-12.77	-8.93	-3.63	Pfeil et al. (2012) (SOCAT database)
17EBC	114.8	-29.9	-3.23 ^c		-3.94 ^c		-1.31	Pfeil et al. (2012) (SOCAT database)
20 MED	3.1	39.5	-0.31	0.06	0.33	-2.28	-0.20	Pfeil et al. (2012) (SOCAT database)
22 MOR	-16.7	18.8	20.22	4.00	0.59		3.02	Pfeil et al. (2012) (SOCAT database)
26 TWI	46.4	-12.4			2.83 ^c		1.03	Pfeil et al. (2012) (SOCAT database)
27WAS	57.1	25.3	0.71	0.66			0.25	Pfeil et al. (2012) (SOCAT database)
28 RED	32.7	29.2		0.23	1.06		0.23	Pfeil et al. (2012) (SOCAT database)
3 TEP	-82.5	8.8	1.99	-1.29	-1.28	-0.39	-0.09	Pfeil et al. (2012) (SOCAT database)
31 BEN	92.4	19.7				-0.59	-0.22	Pfeil et al. (2012) (SOCAT database)
33 LEE	113.5	-27.4	-4.27 ^c	-0.39 ^c	-0.10 ^c		-0.58	Pfeil et al. (2012) (SOCAT database)
34 SAU	146.8	-42.1	-4.21 ^c	-1.39 ^c	-1.42 ^c	-3.34 ^c	-0.94	Pfeil et al. (2012) (SOCAT database)
39 CSK	150.2	45.6				20.05	7.32	Pfeil et al. (2012) (SOCAT database)
4 HUM-1	-77.8	-11.6			-0.14 ^c		-0.05	Takahashi et al. (2012) (LDED database)
4 HUM-2	-80.0	-26.3		-0.91 ^c			-0.33	Takahashi et al. (2012) (LDED database)
4 HUM-3	-73.1	-36.1		-11.82 ^c			-4.32	Takahashi et al. (2012) (LDED database)
42 NWP	-169.1	60.4	-33.81	-13.73	-3.41	6.61	-4.05	Pfeil et al. (2012) (SOCAT database)
4HUM-4	-72.7	-36.7	-2.14 ^c	9.74 ^c	10.53 ^c		2.20	Pfeil et al. (2012) (SOCAT database)
5 SAM	-71.8	-50.5		-18.15 ^c			-6.62	Pfeil et al. (2012) (SOCAT database)
6WBC-1	-56.5	-37.8		-4.56 ^c	-5.19 ^c		-1.78	Takahashi et al. (2012) (LDED database)
6WBC-2	-47.4	-25.7		10.87 ^c			3.97	Takahashi et al. (2012) (LDED database)
8 CAR	-68.0	17.3	0.86	3.76	2.58	0.07	0.66	Pfeil et al. (2012) (SOCAT database)
Amazon River Plume	-52.5	6					-12.78	Ternon et al. (2000), Kortzinger (2003)
Arafura Sea	136.3	-9.9	-0.02 ^c				-0.01	Hydes et al. (2012)
Atlantic Bight (Middle)	-74.5	38.5					-1.8	DeGrandpre et al. (2002)
Atlantic Bight (Southern)	-80.6	31	-0.44	-0.22	-0.24	-0.26	-0.48	Jiang et al. (2008b)

Table 4. Continued.

type	Lon.(°)	Lat.(°)	spring fluxes ^b (mmol C m ⁻² d ⁻¹)	summer fluxes (mmol C m ⁻² d ⁻¹)	autumn fluxes (mmol C m ⁻² d ⁻¹)	winter fluxes (mmol C m ⁻² d ⁻¹)	annual flux (mol C m ⁻² yr ⁻¹)	References
Baltic Sea	20; 13.9	57; 54.9	-92.9	-66.5	-3.6	-34.4	-1.95	Thomas and Schneider (1999), Kuss et al. (2006)
Bass Strait	148.0	-38.8		-0.11 ^c	-0.73 ^c		-0.15	Hydes et al. (2012)
Bay of Biscay (Northern)	-7.9	49					-0.8	Borges et al. (2006)
Bay of Biscay (Southern)	-3.5	46.5					-2.65	de la Paz et al. (2010)
Beaufort Shelves	-155	72		-2.81			-2.79	Murata and Takizawa (2003), Cai et al. (2006)
Bering Sea Shelf	-165	57	-1.2	-0.66			-6.15	Nedashkovsky et al. (1995), Codispoti et al. (1986), Walsh and Dieterle (1994)
Bering Sea Shelf	-165.4	56.7					-8	Codispoti et al. (1986)
Bristol Bay	-164	58					-0.2	Borges et al. (2005), Kelley and Hood (1971), Codispoti et al. (1986), Chen (1993), Murata and Takizawa (2003)
Canterbury Bight	170.7	-45.8	-0.64 ^c	-0.43 ^c	-0.41 ^c	-0.37 ^c	-0.17	Guilderson et al. (2005)
Chukchi Sea	-165	72.5	-0.05	-2.3	-2.47	-0.04	-5.33	Bates (2006)
Coastal Calif. (Monterey Bay)	-121.9	36.9					0.05	Friederich et al. (2002)
East China Sea (Middle)	124	31	-8.8	-4.9	2.9	-10.4	-1.9	Zhai and Dai (2009)
East China Sea(Northern)	126	33	-5.04	-2.52	1.9		-0.79	Shim et al. (2007)
East China Sea (Southern Eastern)	125	30	-4.87	-3.32	-5.14	-8.57	-1.45	Wang et al. (2000)
English Channel	-1.2	50.2					-0.15	Borges and Frankignoulle (2003), Thomas et al. (2007)
Funka Bay	140.6	42.3					-7	Nakayama et al. (2000)
Gray's Reef	-80.9	31.4	0.28	-0.35	-0.01	-1.72	-0.16	Sabine et al. (2012)
Great Barrier Reef	145.5	-15					0.33	Kawahata et al. (1999)
Gulf of Biscay	-6.5	49	-6.98	-15.08	-1.43	0.94	-2.88	Frankignoulle and Borges (2001)
Gulf of Cadiz	-6.5	36.75	-0.85	1.45	-0.4	-1.75	-0.16	Ribas-Ribas et al. (2011), Huertas et al. (2006) ^a
Gulf of Lion	4	43					7.1	de Madron et al. (2010)
Gulf of Mexico Shelf (Northwest)	-88.6	30.0	-1.35	-0.16	-0.31		-0.22	Sabine et al. (2012)
Gulf of Nicoya	-84.9	9.6			-0.05			
Gulf of Trieste	13.6	45.7					-0.02	Pfeil et al. (2012) (SOCAT database)
Hudson Bay	-85	59		5.43	0.77		-2.5	Turk et al. (2010)
Ishigaki Island ^d	124.3	24.4	-27	55	25		0.84	Else et al. (2008)
Java Sea	112.9	-5.6	0.26 ^c	-0.01 ^c	0.07 ^c	0.23 ^c	6.45	Kayanne et al. (2005)
Jiaozhou Bay	120.3	36.15	4.14	19.47	17.07	-0.15	0.05	Hydes et al. (2012)
Kameohe Bay	-157.8	21.5					3.7	Li et al. (2007)
Kara Sea	74.0	74.0					1.45	Fagan and Mackenzie (2007)
La Push	-125.0	48.0		-1.62	0.14		0.01	Fransson et al. (2001)
Laptev Sea	130.0	74.0					-0.27	Sabine et al. (2011)
Malacca Strait	101.6	2.4	-0.10		0.63		0.01	Fransson et al. (2001)
Moorea ^e	-149.9	-17.5		1.5 ^c		-1.2 ^c	0.10	Hydes et al. (2012)
New Jersey Coast	-74.2	39.4					0.05	Frankignoulle et al. (1996), Gattuso et al. (1993)
							-0.68	Boehme et al. (1998)

Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 4. Continued.

type	Lon.(°)	Lat.(°)	spring fluxes ^b (mmol C m ⁻² d ⁻¹)	summer fluxes (mmol C m ⁻² d ⁻¹)	autumn fluxes (mmol C m ⁻² d ⁻¹)	winter fluxes (mmol C m ⁻² d ⁻¹)	annual flux (mol C m ⁻² yr ⁻¹)	References
North Coastal of California	-123.8	39.0		2.52			0.92	Pfeil et al. (2012) (SOCAT database)
North Sea (Northern and Middle)	2.6	56.7					-1.38	Thomas et al. (2004)
North Sea (Southern)	2.5	52.0	-12.47	6.8	4.35	-0.35	-0.7	Schiettecatte et al. (2007), Hoppeema (1991) ^a
Northeast Coastal of Australia	151.5	-23.5	-0.16 ^c	0.28 ^c	0.04 ^c		0.02	Takahashi et al. (2012) (LDED database)
NorthEast Sunda Shelf	105.7	0.7	-0.04		0.28	0.01	0.03	Hydes et al. (2012), Chen (unpublished)
Okhotsk Sea	143.5	44.5		-4.1			-1.67	Chen et al. (2003), Otsuki et al. (2003), Wakita et al. (2003)
Omani coast	59.0	20.0	0.75	-7.13	-0.95	-1.17	-0.9	Goyet et al. (1998)
Oregon Coast	-124.5	44.5		-20			-7.3	Hales et al. (2005)
Otaru Bay	141.0	43.3	-8.8	-6.9	7.4	-6.9	-0.78	Sakamoto et al. (2008)
Palau Islands ^d	134.4	7.4		33	49		15.0	Kayanne et al. (2005)
Patagonian shelf	-65.0	-45.0	-7 ^c	-3.8 ^c	-2.9 ^c	-1 ^c	-1.35	Bianchi and Allison (2009)
Prydz Bay	78.9	-68.6				-75 ^c	-2.45	Gibson and Trull (1999), Borges et al. (2005), Wang et al. (1998)
Red Sea	42.8	13.4			0.04		0.01	Hydes et al. (2012)
Ross Sea	180.0	-75.0				-13 ^c	-1.5	Sweeney (2003), Wang et al. (1998), Bates et al. (1998)
Scotian shelf	-63.0	44.0	1.6	-3.1	-5.9	-8.3	-1.42	Shadwick et al. (2011)
Southeast Coastal of Australia	152.4	-32.8	-0.74 ^c	-0.57 ^c			-0.24	Takahashi et al. (2012) (LDED database)
South China Sea (Northern)	116.0	22.0	2.7	7.5	1.4		0.86	Zhai et al. (2005, 2007)
Sydney Coast (Port Hacking time series station)	151.2	-34.1					-0.17	McNeil (2010)
Taiwan St. ^e	120.3	25.0	-17.6				-6.4	Ma et al. (1999)
Vancouver Is. Coast	-126.0	49.0					-0.5	Ianson and Allen (2002)
West Coastal of India	74.0	14.1	0.06		0.03	0.03	0.01	Hydes et al. (2012)
Yellow Sea	122.0	35.5	-4.4	1.8	-4.4	-13	-2.2	Oh et al. (2000), Wang et al. (2001)
Yellow Sea (Northern)	122.5	38.5	1.88	3.38	1.39	0.24	1.68	Xue et al. (2012)
Yellow Sea (Southern)	122.0	34.5	4.47	1.56	4.85		1.99	Xue et al. (2011)

^a Positive fluxes indicate an emission of CO₂ from water to the atmosphere;^b Spring: March–May; Summer: June–August; Fall: September–November; Winter: December–February;^c Austral seasons;^d not used in the calculation.

Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 5. Summary of reported annual global air-sea CO₂ fluxes in world's continental shelves.

CO ₂ sink in the coastal ocean (PgC yr ⁻¹)	References
-1.00	Tsunogai et al. (1999)
-0.10	Liu et al. (2000)
0.50	Fasham et al. (2001)
-0.60	Yool and Fasham (2001)
-0.24	Rabouille et al. (2001)
-0.30	Chen et al. (2003)
-0.36	Chen (2004)
-0.40	Thomas et al. (2004)
-0.90	Ducklow and McCallister (2004)
-0.37	Borges (2005)
-0.45	Borges et al. (2005)
-0.22	Cai et al. (2006)
-0.33 to -0.36	Chen and Borges (2009)
-0.21	Laruelle et al. (2010)
-0.25	Cai (2011)
-0.40	This study

**Air-sea Exchanges
of CO₂ in World's
Coastal Seas**

C.-T. A. Chen et al.

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)



Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.




Fig. 1. Distribution of estuaries studied.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.




Fig. 2. CO₂ flux in (a) upper, (b) mid and (c) lower estuaries.

**Air-sea Exchanges
of CO₂ in World's
Coastal Seas**

C.-T. A. Chen et al.




Fig. 3. Histogram of reported daily CO₂ fluxes per unit area in different seasons and annual flux of world's estuaries.

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)



Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.




Fig. 4. Histogram of reported annual CO₂ fluxes of world's estuaries in various latitude bands.

Title Page

Abstract

Introduction

Conclusion:

References

Tables

Figures



Back

Close

Full Screen / Esc

[Printer-friendly Version](#)

Interactive Discussion

Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.




Fig. 5. Global wind potential (courtesy Zhou et al., 2012).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

[Printer-friendly Version](#)

Interactive Discussion



Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.





Fig. 6. Total annual CO₂ flux and CO₂ flux per unit area from estuaries in each continent.

**Air-sea Exchanges
of CO₂ in World's
Coastal Seas**

C.-T. A. Chen et al.

**Fig. 7.** Total annual CO₂ flux and CO₂ flux per unit area from estuaries around each ocean.

Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.




Fig. 8. Distribution of continental shelves studied.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[|◀](#)

[▶|](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.




Fig. 9. Histogram of reported daily CO₂ fluxes in different seasons and annual flux on world's continental shelves.

Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.




Fig. 10. Histogram of reported annual CO₂ fluxes of continental shelves in various latitude bands.

Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.





Fig. 11. Total annual CO₂ flux and CO₂ flux per unit area of continental shelves in different continents.

Air-sea Exchanges of CO₂ in World's Coastal Seas

C.-T. A. Chen et al.




Fig. 12. Total annual CO₂ flux and CO₂ flux per unit area of continental shelves in different oceans.