Biogeosciences Discuss., 10, C3394–C3396, 2013 www.biogeosciences-discuss.net/10/C3394/2013/

© Author(s) 2013. This work is distributed under the Creative Commons Attribute 3.0 License.



BGD

10, C3394-C3396, 2013

Interactive Comment

Interactive comment on "Microbial activity and carbonate isotope signatures as a tool for identification of spatial differences in methane advection: a case study at the Pacific Costa Rican margin" by S. Krause et al.

Anonymous Referee #1

Received and published: 10 July 2013

Review "Microbial activity and carbonate isotope signatures as a tool for identification of spatial differences in methane advection: A case study at the Pacific Costa Rican margin" by Krause et al.

Krause and co-workers present measurements of microbial anaerobic oxidation of methane and sulfate reduction rates from sediments at the Pacific Coasta Rican Margin. In addition, they discuss the analysis of carbonate $^{18}O,\,\delta^{13}C,$ and $^{87}Sr/^{86}Sr$ signatures. They combine these measurements with numerical modelling to constrain

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



the advection velocities and the origin of methane fluxes at these mounds. Results indicate pronounced differences between the two settings and thus reveal a strong temporal and spatial variability of methane charged fluid flow at the Pacific Costa Rican margin. The authors represent an interesting, multi-disciplinary analysis that integrates observations and numerical modelling techniques. The manuscript is generally well structured and well written. However, the authors need to emphasise the original aspects of their work. They should explain the differences between their study and previous studies. What is new? What distinguishes this study from the results presented in, for instance, Hensen et al., 2004; Mavromatis et al., 2012? How do their results fit into a regional/broader context? In addition, the description and rational for the chosen model approach could be improved. In particular, simplifying assumptions (e.g. CaCO3 treatment, FeS precipitation, pH dynamics, TA) should be better justified and critically discussed.

The presented manuscript could be considered for publication if the authors address these points and emphasis the originality of the presented research.

Specific Comments

p.2, l.24: a^{-1}

p.3, I.67: dissociate

p.9, I.228: remove brackets around Berner, 1980

p.9, I.228: The equation is only formulated for dissolved species. What about solid species?

p. 10, I. 250: What about bioturbation?

p. 10, l. 258: add with p. 10, l. 261: mixed

p. 11, l. 271: justify the use of this equation for iron sulphide precipitation. Why don't you consider precipitation as a two step reversible process? What about the oxidation

BGD

10, C3394-C3396, 2013

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



of FeS by O2 and the oxidation of H2S by Fe(OH)3?

p. 11, l. 279: explain/justify why you use this simplified approach. Why is pH and the carbonate system not explicitly resolved? What are the errors associated with the chosen approach?

p.11, I. 289: how do you quantify HCO3- HS- and CO32- concentrations without resolving pH dynamics? For instance, the speciation of DIC and TH2S released during AOM will depend on ambient pH and will also influence ambient pH.

p.11, I.290: B(OH)4- is an important component of TA in the marine environment.

p. 12, I. 293: What are the boundary conditions for Fe2O3 and CaCO3?

p. 13, Result section: reference the figures more often

p.19-20, I. 495 onwards and I. 507: what is new?

Interactive comment on Biogeosciences Discuss., 10, 8159, 2013.

BGD

10, C3394-C3396, 2013

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

