

Agricultural peat lands; towards a greenhouse gas sink – a synthesis of a Dutch landscape study

A.P. Schrier-Uijl^{*1}, P.S. Kroon², D.M.D. Hendriks³, A. Hensen², J.C. Van Huissteden⁴, F. Berendse¹, E.M. Veenendaal¹.

[1] Nature Conservation and Plant Ecology, Wageningen University, Droevendaalse steeg 3a, 6708 PD Wageningen, The Netherlands.

[2] Energy research Centre of the Netherlands (ECN), Department of Air Quality and Climate Change, 1755 LE Petten, The Netherlands

[3] Deltares, dept. Soil and Groundwater Systems, Princetonlaan 6, 3584 CB Utrecht, The Netherlands.

[4] Hydrology and Geo-Environmental Sciences, Faculty of Earth and Life Sciences, Vrije Universiteit Amsterdam, De Boelelaan 1085, 1081 HV Amsterdam, The Netherlands.

[*] Correspondence to: Arina P. Schrier-Uijl (arina.schrier@wetlands.org).

Gewijzigde veldcode

Abstract

It is generally known that managed, drained peatlands act as carbon sources. In this study we examined how mitigation through the reduction of [the intensity of land](#) management and through rewetting may affect the greenhouse gas (GHG) emission and the carbon balance of intensively managed, drained, agricultural peatlands. Carbon and GHG balances were determined for three peatlands in the western part of the Netherlands from 2005 to 2008 by considering spatial and temporal variability of emissions (CO_2 , CH_4 and N_2O). One area (Oukoop) is an intensively managed grass-on-peatland, including a dairy farm, with the ground water level at an average annual depth of 0.55 m below the soil surface. The second area (Stein) is an extensively managed grass-on-peatland, formerly intensively managed, with a dynamic ground water level at an average annual depth of 0.45 m below the soil surface. The third area is an (since 1998) rewetted former agricultural peatland (Horstermeer), close to Oukoop and Stein, with the average annual ground water level at a depth of 0.2 m below the soil surface. During the measurement campaigns we found that both agriculturally managed sites acted as carbon and GHG sources but the rewetted agricultural peatland acted as a carbon and GHG sink. The terrestrial GHG source strength was $1.4 \text{ kg CO}_2\text{-eq m}^{-2} \text{ yr}^{-1}$ for the intensively managed area and $1.0 \text{ kg CO}_2\text{-eq m}^{-2} \text{ yr}^{-1}$ for the extensively managed area; the unmanaged area acted as a GHG sink of $0.7 \text{ kg CO}_2\text{-eq m}^{-2} \text{ yr}^{-1}$. Water bodies contributed significantly to the terrestrial GHG balance because of a high release of CH_4 and the loss of DOC only played only a minor role. Adding the farm-based CO_2 and CH_4 emissions increased the source strength for the managed sites to $2.7 \text{ kg CO}_2\text{-eq m}^{-2} \text{ yr}^{-1}$ for Oukoop and $2.1 \text{ kg CO}_2\text{-eq m}^{-2} \text{ yr}^{-1}$ for Stein. Shifting from intensively managed to extensively managed grass-on-peat reduced GHG emissions mainly because N_2O emission and farm-based CH_4 emissions decreased. Overall, this study suggests that managed peatlands are large sources of GHG and carbon, but, if appropriate measures are taken they can be turned back into GHG and carbon sinks within 15 years of abandonment and rewetting.

Keywords: GHG balance, carbon balance, restoration, management reduction, water bodies, agricultural peatland, rewetting

Introduction

Although peatlands cover only 6% of the earth surface, they play a central role in the global carbon cycle (Gorham et al., 2012). In their natural state, peatlands capture carbon as carbon dioxide (CO₂) with a long term average uptake rate of 25 g C m⁻² yr⁻¹ (Borren et al., 2006). Natural peatlands emit methane (CH₄) as a result of anaerobic conditions that lead to methanogenesis. The total balance between CO₂ uptake and CH₄ release is in most cases negative (sequestration of carbon) and is dependent on moisture conditions, temperature, vegetation composition, availability of degradable substrates and microbial activity (e.g. Hendriks et al., 2009). Generally, nitrous oxide (N₂O) does not play a significant role in the greenhouse gas (GHG) budgets of natural peatlands. While natural peatlands act as sinks for carbon, agricultural peatlands commonly act as sources for carbon and GHGs.

In Europe, 50% of all peatlands are subject to various sorts of agricultural practices (Joosten and Clarke, 2002), often associated with drainage resulting in oxidation of peat and release of CO₂ to the atmosphere. In the Netherlands about 270 000 ha (7 % of the land surface) is peatland. Since the industrial period these peat soils were heavily drained and fertilized and they turned into carbon sources (e.g. Langeveld et al., 1997; Veenendaal et al., 2007). As a result, peat subsidence rates in the Netherlands are up to 10 mm yr⁻¹ and already 20% of the peat soils have disappeared and classified as mineral soil in the last 30 years (Kempen et al., 2009). In wet peatlands however, CH₄ is commonly released (e.g. Carter et al., 2012; Teh et al., 2011; Schrier-Uijl et al., 2010; Hendriks et al., 2007). Also water bodies in peat ecosystems are important contributors to the GHG balance and have therefore to be considered when calculating GHG budgets (Schrier-Uijl et al., 2011; Billet and Harvey, 2012). Emissions from wetlands, water bodies and grasslands are currently not (or only partly) included in national emission inventories (Maas et al., 2008, Nol et al., 2008). While it does not play a significant role in the GHG budgets of natural peatlands, in intensively managed peatlands high inputs of chemical fertiliser and manure lead to increased N₂O emissions.

Since there are still few comprehensive studies that report on the effects of restoration activities on the total GHG emission balance, the main goal of this paper is to analyze the long term effects of restoration through reducing the management and decreasing the ground water table depth on the GHG balance of intensively managed peatlands in the Netherlands.

Here we present the synthesis of a landscape scale experiment that has been performed in three temperate peatlands in the Netherlands under different management regimes (intensively managed, extensively managed and a rewetted, former agricultural peatland (unmanaged)). To obtain spatially and temporally explicit GHG sources and sinks, three to four years of simultaneous measurements were conducted with chamber and micrometeorological techniques to cover different spatial and temporal scales and the entire suite of biogenic key GHGs (CO₂, CH₄ and N₂O). The studies that published on the first results of the separate sites are Schrier-Uijl et al (2009, 2010, 2011), Kroon et al (2010a,b,c), Hendriks et al (2007) and Veenendaal et al (2007). In this paper results of later years have been analysed and the three sites have been compared. For full accounting of the carbon and GHG balance also carbon import and export by management has been analysed and water bodies are included in the calculations.

Verwijderd: under restoration

Materials and methods

At three low land peat sites in the Mid West of the Netherlands, the exchange of CO₂, CH₄ and N₂O between the soil-plant continuum and the atmosphere was measured in detail at various scales, using various measurement techniques. Measurements have been cross-checked to robustly investigate the effect of restoration on the GHG balance. The three research sites (Oukoop, Stein and Horstermeer) are located below sea level. The peat soils at all sites are (fibric) eutric histosols, the mean annual air temperature is 9.8 °C and the mean annual rainfall is 797 mm. In Fig. 1 the average monthly water table (WT) and soil temperature (T_{soil}) is given for the three sites. Table 1 gives an overview of the main site characteristics and the management per site.

Verwijderd: in a moderate climate

Verwijderd: All sites

Verwijderd: In this publication a general description of materials and methods is given; the reader is referred to previous publications (Schrier-Uijl et al., 2009, 2010, 2011; Kroon et al., 2010a,b,c; Nol et al., 2008; Hendriks et al., 2007; Veenendaal et al., 2007) for more detailed information for the specific sites and methods used.

Verwijderd: ¶

Figure 1. Monthly averages for water table and soil temperature in Oukoop (Ou), Stein (St) and Horstermeer (Ho).

Measurement sites

The Oukoop site (Ou) is an intensively managed grassland polder on peat in the west of the Netherlands (lat. 52° 02'N, long. 4° 47'E, altitude – 1.8 m a.s.l.). The site is part of a dairy farm and grass is regularly mowed to feed cows that are kept on the farm. Manure and fertilizers are applied four or five times annually in the period February to September. The area has a clayey peat or peaty clay top layer of about 0.25 m thickness on a 12 m thick peat layer on a mineral subsoil. Sixteen percent of the total polder is open water (drainage ditches, small ponds, shallow lakes), 5% are bordering (water saturated) edges and the remaining part consists of relatively dry fields with a dynamic water table of mean annual depth of 0.55 m. The dominant grass species are *Lolium perenne* and *Poa trivialis*.

Verwijderd:

Verwijderd: annually

Verwijderd: with dynamic mean annual ground water tables at

Verwijderd:

Verwijderd: below field level.

Verwijderd:

Verwijderd: are

The Stein site (St) is an extensively managed grassland polder on peat 4 km South-West of the Oukoop site (lat. 52° 01'N, long. 4° 46'E, altitude – 1.7 m a.s.l.). The area has been managed as a meadow bird reserve since 2001 which implies that no manure or artificial fertilizers have been applied ever since and that management only comprises the removal of above ground biomass three times a year. The polder was intensively used for grass production in the same way as the Oukoop polder before it gradually became a meadow bird reserve. The area has the same soil characteristics as the Oukoop site and land use history was similar before Stein was taken out of production. The Stein site has a dynamic water table since 2006 with high water tables in winter and low water tables in summer; the ground water table is on average 0.45 m below field level. The proportions of land and water are similar to the Oukoop site (16% open water; 5% water saturated borders; 79% relatively dry land). The dominating plant species were *Lolium perenne* and *Poa trivialis*, but over time *Anthoxanthum odoratum* and *Rumex acetosa* have become more abundant.

Verwijderd: The ground water table is

The Horstermeer site (Ho) is a grassland/wetland polder on peat in a former intensively used dairy farm area in the centre of the Netherlands (lat 52° 02'N, 5° 04'E, altitude -2.2 m a.s.l.) located about 40 km NE from Oukoop and Stein. The site has been abandoned in 1998 and has not been exploited agriculturally ever since. The management was about similar to that of Stein and Oukoop until abandonment. After abandonment, the ditch water table has been raised to approximately 0.10 m below the land surface. The vegetation has developed

towards a semi-natural grassland. Five percent of the area is open water (ditches), 10% is year-round saturated soil (mostly along the ditches), 25% is relatively wet soils and 60% is relatively dry land with a fluctuating ground water table (between 0 – 0.40 m below the soils surface, 0.20 m annual average) with a dry top-layer during the largest part of the year. No management takes places, except for ditch water table regulation. Dominant species are *Holcus lanatus*, *Phalaris arundinacea*, *Glyceria fluitans*, reeds and high forbs.

Verwijderd: an aerated

Table 1. Site descriptions, land use and management for each peat site

Verwijderd: per

Landscape scale flux measurements

Landscape scale flux measurements of CO₂, CH₄ and N₂O were performed with an averaging time of 30 minutes from 10 Hz data using the EC flux technique. EC flux systems for CO₂ at the three sites consisted of a sonic anemometer (Oukoop and Stein: Campbell CSAT C3 Sonic Anemometer (Campbell Scientific, Utah, USA), Horstermeer: Windmaster Pro 3 Axis Ultra Sonic Anemometer (Gill Instruments, Lymington, UK) and a fast response CO₂-H₂O open path gas analyzer (all sites: LICOR 7500 (LICOR, Lincoln, NE, USA)) placed immediately below (Oukoop & Stein) or next too (Horstermeer) the Sonic Anemometer.

Met opmaak: Niet Markeren

Met opmaak: Niet Markeren

Met opmaak: Niet Markeren

The EC flux systems were installed 4.3 meter above the soil surface in Horstermeer and 3.05 meter above the soil surface in Oukoop and Stein. Oukoop was also equipped with an EC flux system for CH₄ and N₂O in the period 2006-2008 consisting of a Sonic Anemometer (model WMPRO from October 2007- July 2008 and model R3 for the rest of the measurement period, (Gill instruments, Lymington, UK) and a Quantum cascade laser spectrometer (model QCL-TILDAS-76, Aerodyne Research Inc., Billerica MA, USA).

Met opmaak: Niet Markeren

Met opmaak: Niet Markeren

The EUROFLUX protocol for quality control (including filtering for spikes, frequency losses and webb-corrections) and data correction was applied to determine the fluxes on a thirty minute basis (Aubinet et al., 2000). Footprint analysis was performed according the Kormann and Meixner method (Kormann and Meixner, 2001) and Schuepp et al., (1990). The footprint of the masts included fields, ditches and ditches edges but excluded farms and other GHG hotspots.

Met opmaak: Niet Markeren

Met opmaak: Niet Markeren

Met opmaak: Niet Markeren

Soil respiration in the three sites was described as a function of soil temperature by using the Arrhenius relation (Lloyd and Taylor, 2004). For CO₂, data coverage was over 60% for each site. Annual balances for CH₄ at Oukoop were derived from a site specific multivariate regression model, including soil temperature and wind velocity. Annual balances for N₂O were calculated from four contributions: background emissions, event emissions due to fertilizer application, leaching and deposition. Background emissions and event emissions were derived from the EC flux data (see table 5). For EC measurements of N₂O and CH₄, data coverage was 48%. For more details about EC flux measurements in the three sites see Veenendaal et al., 2007; Hendriks et al., 2007; Kroon et al., 2009, 2010).

Met opmaak: Niet Markeren

Small-scale flux measurements

For all sites, small-scale flux measurements were performed on dry fields, saturated ditch edges and open water (ditches) for CO₂ and CH₄. A Photo Acoustic Field Gas Monitor was used (Oukoop and Stein: Innova type 1412 and Horstermeer: type 1312, Innova AirTech Instruments, Ballerup, Denmark). The gas analyser was connected with Teflon tubes to

Met opmaak: Niet Markeren

closed opaque chambers (height 0.2 m, diameter 0.3 m). For each gas measurement, five samples were taken from the headspace of a closed cylindrical, non-transparent chamber over a period of 5-10 minutes from which the flux (height of the chamber multiplied by dC/dt) was calculated. Linearity was checked using the intercept method (Kroon et al., 2008). Fans were installed in the chamber to homogenize the inside air and two external filters were added: a soda lime filter for CO_2 (when measuring CH_4) and a silica gel filter for water vapor. To avoid disturbances, vegetation was not removed prior to the flux measurements. Because the relations with explanatory variables were non-linear, multiple non-linear regression was used to calculate annual emissions. Landscape scale fluxes were upscaled from the emissions from the different landscape elements (fields, saturated edges and ditches) relative to their proportion in the landscape.

Additional to the three years measurements on ditches, in the summer of 2009 an intensive measurement campaign was performed on 6 large shallow lakes and 14 drainage ditches in peatlands (Schrier-Uijl et al (2011)) to compare fluxes from water bodies that were different in depth, size and nutrient status. Cross-checks of emission values were performed by comparing EC measurements to upscaled chamber measurements within the footprint of the EC systems (Schrier-Uijl et al (2009); Hendriks et al (2009)).

Additional measurements and analyses

Meteorological measurements and soil measurements

In addition to GHG measurements global radiation, net radiation, air temperature, vapor pressure, wind speed, wind direction and precipitation were measured at each site using the sonic anemometers. Soil measurement sensors included soil heat flux plates (HPF01, Campbell Scientific, Utah, USA), soil temperature sensors at depths of 0.02, 0.04, 0.08, 0.16 and 0.32 m (Campbell Scientific, Utah, USA) and soil moisture probes to measure volumetric moisture contents at depths of 0.10, 0.20 and 0.30 m (Theta probes ML 2x; Delta T devices Burwell, UK). These systems provided 30 minute values for soil heat fluxes, soil temperature, soil moisture and water table.

Analyses

At the start and the end of the experiments (2005 and 2008), soils were sampled and analyzed for C and N content, organic matter, NO_3^- , NH_4^+ , PO_4^{3-} and pH. Water from drainage ditches was sampled for pH, C content (not for Horstermeer), N content (not for Horstermeer), organic matter, NO_3^- , NH_4^+ , PO_4^{3-} , SO_4^{2-} , Fe^{2+} , dissolved CH_4 , oxygen saturation and electrical conductivity. Well-stirred samples of slurry manure were sampled just before manure application in the Oukoop site and were analyzed for dry matter and C content. Vegetation height was measured every three to four weeks with a disk pasture meter (Eijkelkamp Giesbeek, The Netherlands) to estimate above ground biomass and biomass removal using a site specific empirical linear relationship between vegetation height and biomass weight (Dry Biomass (g) = $29.1 \times$ Disk height (cm) + 50.2; $R^2 = 0.84$; $n = 51$) (Veenendaal et al. 2007). The calculated biomass production data were found to be in agreement with the grass-production data provided by the farmer.

Estimates of GHG balances and carbon balances

The total GHG balance and carbon balance of the sites consist of 1) ecosystem sinks and sources (including fluxes from fields, saturated parts of the land and drainage ditches and

Met opmaak: Niet Markeren

Met opmaak: Niet Markeren

Met opmaak: Niet Markeren

Verwijderd: Instrumentation and methodology ¶

¶
To capture temporal variability of fluxes, all three sites were equipped with eddy covariance (EC) systems for CO_2 since the end of 2004 for four years (for details of the systems see Veenendaal et al., 2007; for Oukoop and Stein and Hendriks et al., 2007 for Horstermeer). In addition, Oukoop was equipped with an EC system for CH_4 and N_2O since the beginning of 2006 for three years (details in Kroon et al., 2007, 2010b). In April 2006 the Horstermeer site was equipped with an EC system for CH_4 (Hendriks et al., 2008, 2009). At the Oukoop and Stein sites chamber measurements were performed from January 2006 to December 2008 (Schrier-Uijl et al 2010) and at the Horstermeer site from January 2005 to December 2008 (Hendriks et al., 2007). ¶
For the three sites, similar measurement procedure(... [1]

Met opmaak: Niet Markeren

Met opmaak: Subscript

Met opmaak: Subscript

Met opmaak: Lettertype: 10 pt

Met opmaak: Lettertype: 10 pt

Met opmaak: Lettertype: 10 pt, Subscript

Met opmaak: Lettertype: 10 pt

Met opmaak: Lettertype: 10 pt, Subscript

Met opmaak: Lettertype: 10 pt

Met opmaak: Lettertype: 10 pt, Subscript

Met opmaak: Lettertype: 10 pt

Met opmaak: Lettertype: 10 pt

Met opmaak: Lettertype: 10 pt

Met opmaak: Lettertype: Cursief

Met opmaak: Lettertype: Cursief

Met opmaak: Superscript

release of dissolved organic carbon (DOC) through water bodies) and in the managed sites 2) sinks and sources related to farm activities such as carbon that is lost or gained through mowing of plant biomass, animal body mass and milk production and slurry and fertilizer application. An overview of the GHG fluxes and C fluxes that have been considered in the calculation of balances in this study is given in Fig. 2.

Verwijderd: ure

Verwijderd: 1

The ecosystem GHG balance of each experimental site was calculated for three years by summation of the net ecosystem exchange of CO₂, CH₄ and N₂O using the global warming potential (GWP) of each gas at the 100 years time horizon (IPCC, 2007). Thus

$$NEE_{GHG} = NEE_{CO_2} + 25NEE_{CH_4} + 298NEE_{N_2O} \quad (1)$$

Verwijderd:

where 25 and 298 are the global warming potentials of CH₄ and N₂O for a 100-year time horizon.

Figure 2. Ecosystem and farm-based GHG fluxes (CO₂ respiration (R_{CO_2}), CO₂ gross primary production or photosynthesis (GPP_{CO_2}), CH₄ and N₂O) and carbon fluxes (CO₂-C, CH₄-C, manure and fertilizer-C, biomass-C) that are being considered in the current study for Oukoop, Stein and Horstermeer. White arrows are farm-related fluxes and dark grey arrows are ecosystem fluxes.

Verwijderd: 1

A more detailed overview of carbon flows in the intensively managed peat area Oukoop is given in Fig. 3. The dashed box represents the boundary of the total polder system.

Verwijderd: 2

Figure 3. System boundaries and fluxes of the intensively managed area Oukoop. Black arrows are C flows, thick dashed arrows are CH₄ fluxes and dashed arrows are CO₂ fluxes (autotrophic and heterotrophic respiration; $R_{CO_2\text{auto}}$ and R_{CO_2} , respectively and photosynthesis(GEP_{CO_2})).

Verwijderd: .

Verwijderd: 2

External carbon inputs from imported feeds and outputs through milk and meat and dissolved organic carbon losses (arrows 12, 13 and 14, Fig. 3) in Oukoop have been considered negligible relative to the other sources and sinks (e.g. Nieveen et al., 2005; Lovett et al., 2008; Wells, 2001). The farm-based N₂O source strength was estimated by using the farm measurements of Hensen et al. (2006). The ecosystem N₂O fluxes which are shown in Fig. 2. were measured in the Oukoop site with Eddy covariance, whereas for the Stein and Horstermeer sites these components have been estimated from Velthof et al. (1997) who estimated peat N₂O emissions from 2 years of measurements on similar peat soils in the same region.

Verwijderd: 2

Verwijderd: 2005

Verwijderd: 1

Verwijderd: ¶

Farm-based CO₂ emissions (arrows 1 and 3, Fig. 3) were estimated from the amount of biomass-C imported into the farm subtracted by the amount of manure-C added on the fields and the amount of C emitted as CH₄. A production efficiency (the amount of energy intake that is transferred into meat or milk) of 7% for large mammals, is used (Van Raamsdonk et al., 2007; Nieveen et al., 2005; Guinand-Flament et al., 2007).

Verwijderd: ¶

Verwijderd: 2

Farm-based CH₄ emissions (arrows 2 and 4, Fig. 3) from the cattle and the stable were estimated following the emission factor approach described by Hensen et al. (2006)

Verwijderd: 2

$$E_{CH_4\text{farm}} = N_{\text{dairy}}E_d + N_{\text{heifer}}E_y + N_{\text{calves}}E_c + A_{\text{manure}}E_m + A_{\text{FYME}}E_f \quad (2)$$

Met opmaak: Lettertype: Cursief

with N the number of animals (number of dairy cows = 65, number of heifers = 20, number of calves = 10), and the amount of manure or farmyard manure (volume of manure storage = 780 m³) and with emission factors for dairy cows (E_d), heifers (E_y), calves (E_c), manure in storages (E_m) and farmyard manure (E_f). The emission parameters were 274 g CH₄ day⁻¹ animal⁻¹ for cows 170 g CH₄ day⁻¹ animal⁻¹ for heifers, 48 g CH₄ day⁻¹ animal⁻¹ for calves, 53 g CH₄ day⁻¹ m⁻³ for fertiliser and 40 g CH₄ day⁻¹ m⁻³ for farmyard manure – all ±50% (Sneath et al., 2006; Van Amstel et al., 2003).

Verwijderd: A

For the fluxes that are considered in the Horstermeer site, the reader is referred to Hendriks et al (2007). Indirect emissions due to leaching and run-off were measured in the Horstermeer site, but were not directly measured in the Oukoop and Stein site. In Kroon et al (2010b) an estimate of leaching and runoff based on the annual amount of synthetic fertilizer and the annual amount of applied cow manure (IPCC 2006) is given for the Oukoop site.

Emissions from large water bodies such as shallow lakes were measured in the summer season, and thus no annual values have been presented for these ecosystems in this paper. Summer emissions from water bodies within peat areas and emissions from peatland were therefore compared using June and July data only. These summer fluxes have been shown to represent around 70% of the annual CH₄ emission from drainage ditches (Schrier-Uijl et al., 2010).

Up-scaling of fluxes to regional scale

To be able to scale the site fluxes up to the entire western peatland area (115,000 ha) of the Netherlands, a detailed database was compiled using the topographic vector-based Top10Vector database (TDN, 2006), a field inventory (Nol et al., 2008), and databases of Dutch natural peatlands (Natuurmonumenten, Staatsbosbeheer, Provinciale landschappen). The resulting database distinguishes between intensively and extensively managed peatland and ditches and ditch edges within these peatlands. Under the assumption that fluxes measured in the intensively managed area and the extensively managed area in this study were representative for the Dutch western peatland area, the emissions have been extrapolated to a larger area. Table 2 shows the areas of the land use and landscape elements.

Table 2. Landscape elements in the Dutch peatlands.

Results

Carbon dioxide balance

Fig. 4 shows the daily measured CO₂ NEE for all three sites for the period 2005-2008 and the cumulative NEE for each year. The unmanaged site was a CO₂ sink in all years (range -1034 to -1939 g CO₂ m⁻² yr⁻¹), with periods of a net (small) release in late winter/early spring periods and with net uptake in the rest of the year. The managed sites appeared to be sources of CO₂ in all years, except for Oukoop being a marginal sink for CO₂ in 2007 which was a relatively wet and cold year and Stein being a marginal sink of CO₂ in 2005 (range -173 to +747 and -88 to +790 g CO₂ m⁻² yr⁻¹ for Oukoop and Stein, respectively).

Verwijderd: ure

Verwijderd: 3

Figure 4. Daily NEE CO₂ for the Oukoop (intensively managed), Stein (extensively managed) and Horstermeer (unmanaged) peatlands measured by the eddy covariance flux technique. The black line represents the cumulative NEE for each year separately (y-axis on the right) and the grey line represents the temporal variability of NEE on the time scale of a day (y-axis on the left).

Verwijderd: 3

Temporal variability of the annual net ecosystem exchange (NEE) (excluding management related fluxes) was high, but NEE of CO₂ in the three areas showed a clear difference between the managed (Oukoop and Stein, net release) and the unmanaged (Horstermeer, net uptake) peatlands (Table 3). The unmanaged peatland (Horstermeer) had an average CO₂ NEE uptake of 1.4 kg CO₂ m⁻² yr⁻¹, while the two managed peatlands (Stein and Oukoop) had an average release of 0.4 kg CO₂ m⁻² yr⁻¹ over a four years period. Inter-annual variability was high, but seasonal trends were the same for each year (Fig. 4). In the years 2006 and 2008 the managed systems had the highest release of CO₂ in, while the unmanaged system had the highest uptake in the year 2007.

Verwijderd: Figure

Verwijderd: 3

Table 3. Terrestrial carbon dioxide flux estimates (kg CO₂ m⁻²) measured by eddy covariance in the period 2005-2008 for the intensively managed production grassland on peat (Ou), the extensively managed hayfield on peat (St) and the unmanaged former agricultural peatland (Ho). Fluxes from shallow lakes are measured in the summer of 2009. Fluxes from removed biomass are not included here.

Monthly CO₂ NEE values show (Fig. 5) that the difference between the managed areas with low water tables and the unmanaged area with high water table is largest in the growing season. The former agricultural peatland, Horstermeer, with its unmanaged vegetation is a large sink in this period while Oukoop and Stein are only minor sinks.

Verwijderd: Figure

Verwijderd: 4

Figure 5. Monthly NEE CO₂ values for the three experimental sites. The CO₂ NEE is given on the y-axes in g CO₂ m⁻² d⁻¹ and the month numbers are given from 2006-2008 on the x-axes.

Verwijderd: 4

Farm-based CO₂ emissions were calculated from the biomass-C fed to the cows on the farm and the transformation to manure-C. With an assimilation efficiency of 7% for large mammals, and 142 g C m⁻² yr⁻¹ exported as manure to the fields and a farm based CO₂ emission of 57 g C m⁻² yr⁻¹, the total farm based CO₂ release is estimated at 0.7 kg CO₂ m⁻² yr⁻¹. The total (ecosystem + farm based) CO₂ emission is than 2.7, 2.1 and -1.4 kg CO₂ m⁻² yr⁻¹ for Oukoop, Stein and Horstermeer, respectively.

Methane balance

For all three sites, soil and water temperature were the most significant predictors of CH₄ emissions and temperature is therefore used as explanatory variable to determine annual balances (Schrier-Uijl et al., 2010; Hendriks et al., 2007); for all sites the CH₄ flux is calculated as e^{a+bT} with T = soil temperature and factors a and b specific for the site and landform (field, ditch and ditch edge). Annual emissions for Oukoop and Stein for 2006 - 2008 and for Horstermeer from 2005 - 2008 are given in Table 4.

Met opmaak: Subscript

Met opmaak: Lettertype: Cursief

Gewijzigde veldcode

Verwijderd: .

Verwijderd: of

Table 4. Yearly CH₄ fluxes (g CH₄ m⁻² yr⁻¹) 1) measured by the chamber method and calculated by using landscape element weighted predictive relationships for Oukoop, Stein and Horstermeer and 2) measured by eddy covariance in the intensively managed site and modelled by using predictive relationships. Farm based emissions are not included in this table.

Fig. 6 shows the temporal variability (daily values) and the cumulative terrestrial NEE_{CH₄} (including fields, saturated land and open water) over three years for all three sites, calculated from chamber measurements based regressions. For the Oukoop site also three years of eddy covariance measurements are shown. Modelled emissions based on chamber measurements are less detailed compared to the eddy covariance measurements because only temperature is used as predictive variable. Annual cumulative CH₄ values are similar for Oukoop and Stein and are higher for the Horstermeer site. The CH₄ emissions varied widely with the season, reaching highest levels during summer. Spatial variability was found to be high between landscape elements within an ecosystem (Schrier-Uijl et al., 2010; Hendriks et al., 2007, 2009) and between ecosystems (this study).

Verwijderd: Figure

Verwijderd: 5

Figure 6. Temporal variability of terrestrial CH₄ fluxes for the three experimental sites modelled by predictive relationships based on chamber measurements and additionally for Oukoop measured by eddy covariance. The right-hand y-axes represents the cumulative CH₄ flux over the three years. Fluxes are weighted by the contribution of each landscape element.

Verwijderd: 5

Schrier-Uijl et al. (2010) reported additional farm-based emissions of 17 and 26 g CH₄ m⁻² yr⁻¹ for Stein and Oukoop, respectively, for the years 2006, 2007 and 2008. The sum of terrestrial CH₄ emissions and farm-based CH₄ emissions amounted to 43.0 g CH₄ m⁻² yr⁻¹ for Oukoop, 33.7 g CH₄ m⁻² yr⁻¹ for Stein, and 19.2 g CH₄ m⁻² yr⁻¹ for Horstermeer.

Carbon balance

The terrestrial CO₂ source estimates of 0.4 kg CO₂ m⁻² yr⁻¹ on average over four years for Oukoop and Stein and -1.4 kg CO₂ m⁻² yr⁻¹ for Horstermeer result in an average carbon source strength of 1091 kg C ha⁻¹ yr⁻¹ for Oukoop and Stein and 4515 kg C ha⁻¹ yr⁻¹ for Horstermeer.

Met opmaak: Superscript

Met opmaak: Superscript

The average CH₄ emission estimates of 17.0 and 16.7 g CH₄ m⁻² yr⁻¹ for Oukoop and Stein and 19.2 for Horstermeer (Table 4) result in an average carbon source strength of 127.5, 125.3 and 144.0 kg C ha⁻¹ yr⁻¹, respectively.

Met opmaak: Superscript

The average annual remittal of C through manure into the field was estimated at 157 g C m⁻² on average over the period 2005-2008; 64.5 m³ ha⁻¹ manure was applied during this period of which 310 kg N ha⁻¹. Fertiliser application amounted to 88 kg N ha⁻¹ on average for the four years.

Removal of biomass in Oukoop was estimated at 8.1 t dry matter on average for 2005, 2006 and 2008, respectively or on average a loss of 400 g C m⁻² yr⁻¹. In Stein carbon loss by biomass removal was similar in all years estimated at 420 g C m⁻² yr⁻¹. The removed biomass was fed to the dairy cattle in Oukoop and is transformed to manure-C. With a assimilation efficiency of 7% for large mammals, 142 g C m⁻² yr⁻¹ exported to the fields as manure and a farm based CH₄ emission of 57 g C m⁻² yr⁻¹, the farm based C release will be around 0.19 kg m⁻² yr⁻¹.

Fig. 7 shows the total carbon balance for the three sites, taking into account the emissions of CO₂ and CH₄, manure application and biomass export as described above. The total C-release in Oukoop and Stein is 5.9 and 7.4 Mg C ha⁻¹ yr⁻¹, respectively and the total C-uptake in Horstermeer is 4.4 Mg C ha⁻¹ yr⁻¹.

Verwijderd: Figure

Verwijderd: 6

Figure 7. Summary of all considered carbon fluxes in the research areas Horstermeer (Ho), Stein (St) and Oukoop (Ou) averaged over 2005, 2006, 2007 and 2008. The annual carbon balance is presented in Mg C ha⁻¹ yr⁻¹, (+) is release and (-) is uptake, and consists of fluxes due to GHG emissions (NEE CO₂ and NEE CH₄) and fluxes due to management (farm based fluxes, manure application and biomass removal).

Verwijderd: 6

Nitrous oxide balance

Measured cumulative NEE N₂O was previously determined over a period of three years at the intensively managed site (Kroon et al., 2010b). Emissions have been separated in 1) background emissions, 2) fertilizer related indirect (peak) emission and 3) emissions due to atmospheric deposition. In Oukoop also farm-based N₂O emissions from manure storages (estimated at 1.5 10⁻² g N₂O m⁻² yr⁻¹) were added to the total N₂O balance, although not significant. Nitrous oxide emissions in the extensively managed site (Stein) and unmanaged site (Horstermeer) were estimated based on (Velthof et al., 1997), since the used chamber set-up was not sufficient to detect the low N₂O emissions in these sites. Table 5 shows all N₂O flux estimates.

Verwijderd: period was in

Verwijderd: (Oukoop) previously determined (

Table 5. Nitrous oxide flux estimates (kg N₂O ha⁻¹ yr⁻¹) and their uncertainties (u) for the intensively managed site (Oukoop), extensively managed site (Stein) and the unmanaged peatland (Horstermeer).

Total GHG balance

Fig. 8 shows the total GHG balance of the three sites in terms of **global** warming potential.

Verwijderd: Figure

Verwijderd: 7

Figure **8**. The GHG balances including CO₂, CH₄ and N₂O for the three sites: intensive (Oukoop), extensive (Stein) and unmanaged (Horstermeer). On the left excluding farm-based CH₄ and CO₂ emissions and on the right including farm-based CH₄ and CO₂ emissions, averaged over 2006, 2007 and 2008.

Verwijderd: 7

Methane emissions from drainage ditches and saturated soil edges along ditches were significantly higher compared to fluxes from the relatively dry land (Schrier et al., 2010; Hendriks et al., 2007). The CH₄ component in the GHG balance in the studied sites consists of outgoing fluxes only and N₂O emission in the intensively managed site consists of emissions originating from fertilizer events and from background emission.

Overall, the managed peatlands acted as terrestrial GHG sources of 1.4 and 1.0 kg CO₂-eq m⁻² yr⁻¹ (including CO₂, CH₄ and N₂O), respectively for Oukoop and Stein and the unmanaged site acted as a GHG sink of 0.8 kg CO₂-eq m⁻² yr⁻¹. Ecosystem N₂O emissions were dominant in the intensively managed peatland, while CO₂ and CH₄ dominated the ecosystem GHG balance of the extensively managed peatland. In the unmanaged peatland CO₂ was the dominating ecosystem GHG. Adding the farm based CH₄ and CO₂ emissions decreased the relative importance of N₂O in the total GHG balance of the intensively managed peatland. The difference in total source strength between the intensively managed peatland and the extensively managed peatland was mainly attributed to the higher N₂O emission and the higher farm-based CH₄ emissions in the intensively managed site.

Upscaling GHG emissions from Dutch peatlands areas

In the western Dutch peat area 68% is intensively managed grassland, 8% is extensively managed grassland or unmanaged grassland, 6% is water (Table 6) and the remaining part is road, farm or has other land use. With the emission values found in this study for intensively and extensively managed peatland and the total area for both of these land uses, emission estimates are performed for the total intensively managed grassland and extensively managed/unmanaged grasslands in the western peatland. The total emission, estimated using a time-horizon of 100 years from the western peatlands is approximately 1210 Gg CO₂-eq (=kton CO₂-eq). In lakes the annual CO₂ emissions are estimated from summer measurements only, however, these fluxes have to be verified by performing year round measurements and by including all three GHG's in the balance.

Table 6. Estimated area and annual GHG release for the area of intensively managed and extensively managed (mown only) or unmanaged grasslands on peat within the total western peatland region of the Netherlands. Farm-based emissions are not included.

Discussion

Balances

Long term emission values of the GHGs and carbon fluxes were compared for peatlands under different management: a drained intensively managed grass-on-peatland with application of fertilizer and biomass export, a drained extensively managed grass-on-peatland with biomass export only and a shallow drained former agricultural peatland **that has been restored** since 1998. Significant differences in GHG emissions have previously been reported between landscape elements within these three sites: CH₄ emissions from drainage ditches and saturated soil were significantly higher compared to CH₄ emissions from the relatively dry land, (Schrier-Uijl et al., 2009; Hendriks et al., 2007) and CH₄ fluxes from shallow lakes in the peat area contribute significantly to the GHG balance (Schrier-Uijl et al., 2010). Emissions originating from the operating farm in the intensively managed peat were found to be important contributors to the GHG balance **as well as to the carbon balance** (Schrier-Uijl et al., 2009; Veenendaal et al., 2007).

Verwijderd: under restoration

Verwijderd: and

Both managed experimental grass-on-peat-areas, Oukoop and Stein, acted as CO₂ emissions sources. Variation in **CO₂ emissions** in Oukoop and Stein was mainly a result of management: Oukoop has the most variable NEE which is a result of the very high frequency of mowing, grazing and manure application. Sharp decreases in NEE are a direct result of mowing events in Oukoop. In Stein management, and therefore variability in NEE, showed less variability with the first biomass removal on the 15th of June and the second biomass removal in September of each year. Both areas had the same history in terms of management, and only during the past 20 years the Stein site has gradually become a meadow bird reserve. This change has not resulted in a significant difference of annual NEE compared to the intensively managed site. The unmanaged site, Horstermeer, **acted as a CO₂ sink**. The cumulative NEE shows a stable pattern with high uptake rates in spring and summer. The Horstermeer site is still **being restored** and despite its abandonment since 1998 soil conditions have remained eutrophic because of influx of eutrophic ground water from the surrounding area. The continuing nutrient-rich conditions generate high plant productivity and microbial activity, resulting in high carbon fluxes (both uptake and emissions) and more organic matter is accumulated than oxidized (Hendriks et al., 2009). However, **development of the nutrient rich, formerly managed system to a more nutrient poor system with natural peat vegetation, may imply a reduction of the NEE** in the future.

Verwijderd: NEE

Verwijderd: the site under restoration,

Verwijderd: under restoration

Verwijderd: as succession may move

Verwijderd: this

Verwijderd: ,

Verwijderd: NEE (and thus uptake)

Verwijderd: reduce

Verwijderd: Temporal variability (daily, seasonally and annually) was high at all three sites, but in the two managed sites no diurnal cycles were observed after correction for temperature (Schrier-Uijl et al., 2011). In the Horstermeer site clear diurnal cycles were found during all seasons synchronous to incoming radiation (Hendriks et al., 2009). They attribute this to the stomatal opening and/or pressurized convective gas transport through the vascular plants with highest emissions in the late afternoon and lowest emissions during the night, as is also reported for other swamp areas (e.g. Whiting and Chanton, 1993; Hirota et al., 2004).

Verwijderd: under restoration

Verwijderd: but

Verwijderd: for succession

All three sites, Oukoop, Stein and Horstermeer, acted as sources for CH₄. Differences in ecosystem CH₄ emissions between sites were not significant. Farm practices in Oukoop and Stein caused an estimated additional CH₄ emission of 26 and 17 g CH₄ m⁻².

The total source strength (ecosystem + farm-based emissions) decreases when management intensity decreases. The Horstermeer is **an unmanaged polder, and the end stage of restoration will depend on management intensity of the surrounding area** affecting groundwater supply and nutrient input. Even if influx of nutrient rich water from surrounding areas and atmospheric nutrient deposition stops, it may be necessary to remove the strongly eutrophic top layer **to reach a near-natural system as e.g. studied by Van den Pol-van Dasselaar et al. (1999)**. Van den Pol-van Dasselaar et al. (1999) studied the emission of CH₄ over two years in three near-natural peatlands in a Dutch nature preserve with narrow

grasslands (mown once a year), reed fields and ditches and ground water level at 18 cm below field level. Soils were similar to the soils of the sites in this research. The average field-CH₄ fluxes were 13.3, 20.4 and 7.9 g CH₄ m⁻² yr⁻¹ and ditch fluxes were 11.3 g m⁻² yr⁻¹ on average. After weighing the contributions of water and land CH₄ emissions were on average significantly lower than the emissions measured in this study. Reported CH₄ emissions from undisturbed peatlands are highly variable. For example a natural peatland in Quebec, Canada showed CH₄ emissions of 9.8 g m⁻² yr⁻¹ (Moore and Knowles, 1990), and Nykänen et al (1995) reported CH₄ emissions of 34.7 g CH₄ m⁻² yr⁻¹ for Scandinavian undisturbed peat lands.

Summer CH₄ emissions from the lakes were significantly higher compared to the emissions from the managed ecosystems. In a study of Schrier-Uijl et al. (2011) the emissions from lakes appeared to be smaller than the emission from drainage ditches within the managed and unmanaged ecosystems. Comparison of 'polder-ecosystem emissions' to emissions from large shallow fresh water lakes shows that water bodies are important contributors to the CH₄ balance (Fig. 9).

Verwijderd: Summer CH₄ emissions from the three peatlands were compared to emissions from 5 large shallow fresh water lakes located in peatlands

Verwijderd: Figure

Verwijderd: 8

Verwijderd: 8

Figure 9. Comparison of summer (June/July) CH₄ fluxes between different ecosystems in peatlands. Fluxes are averaged over three years for the polder Oukoop (intensive), the polder Stein (extensive) and the polder Horstermeer (unmanaged). The lake fluxes are measured in one summer (June/July, n=97) and are averaged over 5 large shallow lakes; error bars represent the standard deviation of the mean.

Calculation of the total CH₄ polder balances for the three sites is based on the current classification of the landscape with 16% open water in Oukoop and Stein and 5% in Horstermeer. Changing the contribution of water and/or saturated land in the landscape by reclassification will cause large changes in the CH₄ balance because these landscape elements together are responsible for over 50% of the total flux. In conclusion, large scale spatial differences in CH₄ emission depend on the combination of management and water table and the presence or absence of water bodies. Drainage ditches, large shallow lakes and saturated land are CH₄ hotspots and therefore spatial differences greatly depend on the proportion of these landscape elements in the landscape. Temporal variability within sites was largely driven by temperature.

Verwijderd: ¶

The carbon balance considered in this study consisted of CO₂-C, CH₄-C, biomass removal and manure and fertilizer application. The two managed sites acted as C-sources and the unmanaged site acted as a C-sink. In the two managed sites, the CO₂ emission (farm based + terrestrial) and the biomass removal accounted for the largest part of the C-release. Because in Oukoop carbon was added through manure and fertilizer inputs, the total C-release turned out to be smaller compared to Stein. In the unmanaged site Horstermeer, the C-balance was dominated by the uptake of CO₂-C. Except for the small release of C through CH₄, no other C-sources or sinks were involved in this undisturbed system. Release of C through ditch water was marginal (Hendriks et al., 2007). Measurements at the three contrasting sites show that an intensively managed fen meadow area can shift from a carbon source towards a carbon sink when the water table is raised and when land management is reduced to zero. It has to be noted that possible CO₂ and/or CH₄ spikes after rewetting are not included in the calculations. These possible spikes may cause an initial increase in emissions in the first years after rewetting. Estimates of these 'spike emissions' are currently uncertain, more research is needed to get robust emissions factors for this emission after rewetting.

The higher background emission in Oukoop compared to Stein is likely the result of the build-up of easily decomposable organic materials in the soil due to manure application 5 times a year. In Stein this application stopped almost 20 years ago. N_2O in these types of ecosystems is produced during nitrification and/or denitrification of NO_3^- . Nitrate is released during mineralization of soil organic N. Leaching is considered to be negligible, but this is not measured. Eddy covariance measurements in Oukoop showed a typical pattern of long periods with low emissions (background emissions) followed by short periods of high emissions (peak-emissions) around manure application (Kroon et al., 2010b). Emissions due to manure application accounted for 25% of the total annual N_2O emissions in Oukoop. Hensen et al. (2006) show that manure based emissions from storages around the farm can cause an additional emission of $14.8 \text{ mg } N_2O \text{ m}^{-3} \text{ manure d}^{-1}$. With the 700 m^{-3} slurry stored around the Oukoop farm this would result in an extra (marginal) emission of $3.8 \text{ kg } N_2O \text{ yr}^{-1}$ over 50 ha or $0.08 \text{ kg } N_2O \text{ ha}^{-1} \text{ yr}^{-1}$. Emission factors around manure application were calculated by subtracting the background emission from the total emissions measured by the EC system. The emission factor range was from 1.2 to 2.8% which is higher than the IPCC default emission factor of 1% (Kroon et al., 2010b).

Verwijderd: has

Combining all incoming and outgoing GHG fluxes shows that Oukoop is the largest GHG source in terms of warming potential. N_2O dominated the emissions in Oukoop and CO_2 and CH_4 contributed equally. In Stein N_2O was the least contributing GHG and the total emission was lower compared to Oukoop. The Horstermeer appeared to be a GHG sink with a release of CH_4 and N_2O from the system, but a large uptake of CO_2 . It suggests that changing the management from intensively to extensively and further to unmanaged may change the total GHG balance from release to uptake. Water bodies were large contributors to the GHG balance when considering summer emissions of CO_2 and CH_4 (Schrier-Uijl et al., 2009, 2010).

Verwijderd: Upscaling GHG emissions from Netherlands peatlands areas¶

In the western Dutch peat area 68% is intensively managed grassland, 8% is extensively managed grassland or unmanaged grassland, 6% is water (Table 6) and the remaining part is road, farm or has other land use. With the emission values found in this study for intensively and extensively managed peatland and the total area for both of these land uses, emission estimates are performed for the total intensively managed grassland and extensively managed/unmanaged grasslands in the western peatland. The total emission, estimated using a time-horizon of 100 years from the western peatlands is approximately $1210 \text{ Gg } CO_2\text{-eq}$ (=kton $CO_2\text{-eq}$). In lakes the annual CO_2 emissions are estimated from summer measurements only, however, these fluxes have to be verified by performing year round measurements and by including all three GHG's in the balance. ¶

¶ Table 6. Estimated area and annual GHG release for the area of intensively managed and extensively managed (mown only) or unmanaged grasslands on peat within the total western peatland region of the Netherlands. Farm-based emissions are not included. ¶

Verwijderd: to stop

Verwijderd: Inverse drainage systems may for this purpose be applied.

Potential ways for mitigation

Mitigation of CO_2 , CH_4 and N_2O in peat areas is important for two reasons: 1) to maintain this ecosystem, stop the degradation of peat soils and soil subsidence and 2) to reduce GHG emissions from drained peatlands. Strategies to reduce GHG emissions from these areas and to increase carbon uptake may be oriented toward rewetting of intensively cultivated peatlands combined with reducing farm-based fluxes and decreasing management intensity.

This study shows that rewetting of agricultural peatland may turn areas of carbon release into areas of carbon uptake; the GHG balance switched from GHG source to sink. The effect might be even stronger in peat soils that lack a clay layer on top of the peat. These peat soils are extremely vulnerable to oxidation (Schothorst, 1977) and to subsidence. The dynamic water tables in the extensively managed polder (high water tables in winter and low water tables in summer) resulted in only a small reduction in GHG emission mainly due to a decrease in farm-based CH_4 emissions and a reduction in N_2O emissions because no fertilizer is applied. High water tables both in winter and in summer will likely reduce emissions. The long-term duration for the sink strength in the unmanaged polder may slow down at centennial timescales due to a decrease in nutrient availability and thus a decreased growth of vegetation.

Farm-based emissions have not been studied separately. Sommer et al. (2009) studied farm-based emissions in Sweden, Denmark, France and Italy. The results showed that shortening the in-house manure storage and decreasing storage temperatures reduced GHG emissions from manure by 0-40% depending on current management and climatic conditions. Large GHG reductions were obtained with slurry separation in a liquid phase and a solid,

organic phase in combination with the early application of the liquid fraction compared to the solid fraction.

Summer emissions on large shallow lakes are higher than the emissions from the intensively and extensively managed polder ecosystems, but lower than the emissions from drainage ditches within the polders. It has been shown that summer emissions from water bodies can contribute significantly to the summer release in the fen meadow area. In the establishment of emission factors for the peatlands these landscape elements should be included in further inventories (Schrier-Uijl et al., 2009, 2010; Kankaala et al., 2007; Billet et al., 2012). Reduction of inputs of organic material and nutrients from the surroundings will likely reduce emissions from these water bodies (Schrier-Uijl et al., 2009, 2010).

Conclusions

This study strongly suggests that intensively managed, drained, agricultural peat soils, which are large GHG and carbon sources, can be turned into sinks in the long term if appropriate mitigation measures are taken. Appropriate mitigation measures are decreasing the water table depth in combination with reducing the management intensity. The switch from an intensively managed peatland to an extensively managed peatland may not significantly alter the ecosystem GHG balance, however, if farm-based emissions are zero, the total (ecosystem + farm-based) emissions decrease significantly. In addition, when implementing mitigation strategies to reduce emissions from one source, GHG emissions from other sources might also be reduced. For example, when the input of nutrient-rich (ground) water in lakes and drainage ditches will be reduced by reducing management in the surrounding catchments, also emissions from water bodies will likely be decreased. Conclusions related to the effect of management on GHG emissions from peatlands are summarized in Table 7.

Table 7. Overview of the expected effects of different mitigation strategies on the total GHG balance (including emissions due to management). The effect on the GHG balance has been determined for the three research sites Oukoop (intensively managed), Stein (extensively managed) and Horstermeer (unmanaged) and are not including the expected future temperature rise. (-) = decrease in emission, (+) = increase in emission, (0) = neutral effect, (?) = effect unknown, (x) = not relevant.

References

- Aubinet, M., Grelle, A., Ibrom, A., Rannik, U., Moncrieff, J., Foken, T., Kowalski, A.S., Martin, P.H., Berbigier, P., Bernhofer, C., Clement, R., Elbers, J., Granier, A., Grunwald, T., Morgenstern, K., Pilegaard, K., Rebmann, C., Snijders, W., Valentini, R. and Vesala T.: Estimates for the annual net carbon and water exchange of forests. The EUROFLUX methodology. *Adv. in Ecol. Res.*; **30**, 113-175, 2000.
- Berendse, F., Oomes, M.J.M., Altena, H.J., Visser, W. : A Comparative Study of Nitrogen Flows in Two Similar Meadows Affected by Different Groundwater Levels. *Journ. Of Applied Ecol.* 31, pp. 40-48, 1994.
- Billet, M.F and Harvey, F.H: Measurements of CO₂ and CH₄ evasion from UK peatland headwater streams. *Biogeochemistry DOI* 10.1007/S105 33-012-9798-9, 2012.
- Burgerhart, N. in Dutch: Mogelijkheden voor koolstofopslag in Nederlandse ecosystemen. Dutch National Inventory Report for LULUCF sector, category 5D1, 2001.
- Carter, M. S., Larsen, K. S., Emmett, B., Estiarte, M., Field, C., Leith, I. D., Lund, M., Meijide, A., Mills, R. T. E., Niinemets, Ü., Peñuelas, J., Portillo-Estrada, M., Schmidt, I. K., Selsted, M. B., Sheppard, L. J., Sowerby, A., Tietema, A., and Beier, C: Synthesizing greenhouse gas fluxes across nine European peatlands and shrublands – responses to climatic and environmental changes. *Biogeosciences*, 9, 3739–3755, 2012.
- Christensen, T. R., Ekberg, A., Strom, L., Mastepanov, M., Panikov, N., Mats, O., Svensson, B. H., Nykanen, H., Martikainen, P. J., and Oskarsson, H.: Factors controlling large scale variations in methane emissions from wetlands, *Geophysical Research Letters*, 30, 2003.
- Dalal, R.C., Allen, D.E., Livesley, S.J.: Magnitude and biophysical regulators of methane emission and consumption in the Australian Agricultural, forest, and submerged landscapes: a review. *Plant and Soil*, doi:10.1007/s11104-007-9446-7, 2007.
- Falge, E., Baldocchi, D., Olson, R., Anthoni, P., Aubinet, M., Bernhofer, C., Burba, G., Ceulemans, R., Clement, R., Dolman, H., Granier, A., Gross, P., Grunwald, T., Hollinger, D., Jensen, N. O., Katul, G., Keronen, P., Kowalski, A., Lai, C. T., Law, B. E., Meyers, T., Moncrieff, H., Moors, E., Munger, J. W., Pilegaard, K., Rannik, U., Rebmann, C., Suyker, A., Tenhunen, J., Tu, K., Verma, S., Vesala, T., Wilson, K., and Wofsy, S.: Gap filling strategies for defensible annual sums of net ecosystem exchange, *Agr. Forest Meteorol.*, 107, pp. 43–69, 2001.
- Gorham, E., Lehman, C., Dyke, A., Clymo, D., Janssens, J: Long-term carbon sequestration in North American peatlands. *Quaternary Science Reviews* 58 (2012) 77-82, 2012.
- Guinand-Flment, F., Delamaire, E., Lambertson, P., Peyraud, J.L.: Adaptations of mammary uptake and nutrient use to once-daily milking and feed restriction in dairy cows, *J. Dairy Sci.* 90, pp 5062-5072, 2007.
- Hendriks, D. M. D., van Huissteden, J., Dolman, A. J., and van der Molen, M. K.: The full greenhouse gas balance of an abandoned peat meadow, *Biogeosciences*, 4, 411-424, 2007.
- Hendriks, D.M.D., van Huissteden, J., Dolman, A.J.. Multi-technique assessment of spatial and temporal variability of methane fluxes in a peat meadow. *Ag. and Forest Meteorol.*, doi:10.1016/j.agrformet.2009.06.017, 2009a.
- Hendriks, D.M.D. thesis: Integrated observations of greenhouse gas budgets at the ecosystem level – changing environment and management practives in peat meadows, 2009b.
- Hensen, A., Groot, T.T., van den Bulk, W.C.M: Dairy farm emissions, from one square meter to the full farm scale. *Agriculture, Ecosystems and Environment* 112: Issues 2-3, 146-152, 2006.

Met opmaak: Lettertype: Arial, 10 pt, Engels (V.S.)

Met opmaak: Inspringing: Links: 0 cm, Afstand Na: 0 pt

Met opmaak: Lettertype: Arial, 10 pt, Engels (Groot-Brittannië)

Met opmaak: Lettertype: 10 pt, Engels (Groot-Brittannië)

Met opmaak: Lettertype: Arial, 10 pt, Engels (Groot-Brittannië)

Met opmaak: Lettertype: Arial, 10 pt, Engels (V.S.)

Met opmaak: Lettertype: Arial, 10 pt, Engels (V.S.)

Met opmaak: Lettertype: Arial, 10 pt, Engels (V.S.)

Met opmaak: Lettertype: Arial, 10 pt, Engels (V.S.)

Verwijderd: ¶

Verwijderd: ¶

Verwijderd: ¶

Verwijderd: ¶

- Hirota, M., Tang, Y.H., Hu, Q.W., Hirata, S., Kato, T., Mo, W.H., Cao, G.M., Mariko, S.: Methane emissions from different vegetation zones in a Qinghai-Tibetan Plateau wetland. *Soil Biology and Biochemistry* 36, pp737-748, 2004.
- Howden, S.M. and Reyenga, P.J.: Methane emissions from Australian livestock: implications of the Kyoto Protocol. *Australian Journ. Of Agr. Res.* 50, pp. 1285-1291, 1999.
- Iepema, G., Baars, T., Bolk, L. in Dutch: Afgewenteld grondgebruik op melkveebedrijven: externe hectares, Rapport 10 Bioveem., 2005
- IPCC 2006: *Guidelines for National Greenhouse Gas Inventories*. Prepared by the National Greenhouse Gas Inventories Programme, Eggleston, H.S., Buendia, L., Miwa, K., Ngara T. & Tanabe, K., (eds), IGES, Japan, 2006.
- IPCC 2007: *The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2007.
- Janssens, I. A., Medlyn, B., Gielen, B., Laureysens, I., Jach, M. E., Van Hove, D. & Ceulemans, R.: Carbon budget of *Pinus sylvestris* saplings after four years of exposure to elevated atmospheric carbon dioxide concentration. *Tree Physiology* 25: 325-337, 2005.
- Joosten, H and Clarke, D.: *Wise use of mires and peatlands – Background and principles including a framework for decision-making*. International Mire Conservation Group / International Peat Society, 304 pp, 2002.
- Kankaala P, Taipale S, Nykanen H et al: Oxidation, efflux, and isotopic fractionation of methane during autumnal turnover in a polyhumic, boreal lake. *J Geoph Res-Biogeosc.* 112 G2 G02003, 2007.
- Kempen, B., Brus, D.J., Heuvelink, G.B.M., Stoorvogel, J.J.: Updating the 1:50,000 Dutch soil map using legacy soil data: A multinomial logistic regression approach. *Geoderma* 151, pp. 311-326, 2009.
- Kormann, R., Meixner, F.X.: *An analytical footprint model for neutral stratification*. *Boudary-Layer Meteorology*: 99, 207-224, 2001.
- Kroon, P.S., Hensen, A., Jonker, H.J.J., Zahniser, M.S., van 't Veen, W.H., Vermeulen, A.T.: Suitability of quantum cascade laser spectroscopy for CH₄ and N₂O eddy covariance flux measurements, *BG*, 4, 715-728, 2007.
- Kroon, P.S., Hensen, A., van den Bulk, W.C.M., Jongejan, P.A.C., Vermeulen, A.T.: The importance of reducing the systematic error due to non-linearity in N₂O flux measurements by static chambers. *Nutr Cycl Agroecosyst* 82:175-186, 2008.
- Kroon, P.S., Hensen, A., Jonker, H.J.J., Ouwensloot, H.G. Vermeulen, A.T. and Bosveld, F.C.: Uncertainties in eddy covariance flux measurements assessed from CH₄ and N₂O observations. *Agric. For. Meteorol.*, doi:10.1016/j.agrformet. 2009.08.008, 2010a.
- Kroon, P.S., Schrier-Uijl, A.P., Hensen, A., Veenendaal., E.M., Jonker, H.J.J.: Annual balances of CH₄ and N₂O from a managed fen meadow using eddy covariance flux measurements. *Eur. J. Soil Sci*, 2010b.
- Kroon, P.S, Thesis: *Eddy covariance observations of methane and nitrous oxide emissions - towards more accurate estimates from ecosystems*. Technische Universiteit Delft, ISBN: 978-90-9025388-6, 2010c.
- Langeveld, C.A., Segers, R., Dirks, B.O.M., Van den Pol-van Dasselaar, A., Velthof, G. C. & Hensen, A.: Emissions of CO₂, CH₄ and N₂O from pasture on drained peat soils in the Netherlands. *Eur. J. Agron.*, 7(1-3), 25-42, 1997.
- Limpens, J., Berendse, F., Blodau, C., Canadell, J.G., Freeman, C., Holdens, Roulet, N., Rydin, H., and Scheapman-Strub, G.: Peatlands and the carbon cycle: from local processes to global implications – a synthesis. *Biogeosciences*, 5, pp. 1475–1491, 2008.
- Lloyd, J., Taylor, J.A.: On the temperature dependency of soils. *Functional Ecology*: 8, 315-323, 1994.

Verwijderd: ¶

Verwijderd:

Met opmaak: Lettertype: Arial, 10 pt, Engels (Groot-Brittannië)

Met opmaak: Inspringing: Links: 0 cm

Met opmaak: Lettertype: Arial, 10 pt, Engels (Groot-Brittannië), Subscript

Met opmaak: Lettertype: Arial, 10 pt, Engels (Groot-Brittannië)

Met opmaak: Lettertype: Arial, 10 pt, Engels (Groot-Brittannië)

Met opmaak: Lettertype: Arial, 10 pt, Engels (Groot-Brittannië)

Met opmaak: Lettertype: Arial, 10 pt, Engels (Groot-Brittannië)

Met opmaak: Lettertype: Arial, 10 pt, Engels (Groot-Brittannië)

Verwijderd: ¶

- Lloyd, C.R.: Annual carbon balance of a managed wetland meadow in the Somerset Levels, UK. *Agricultural and Forest Meteorology* 138, pp. 168-179, 2006.
- Maas, C.W.M., Coenen, P.W.H.G., Ruysenaars, P.G., Vreuls, H.H.J., Brandes, L.J., Baas, K., Van den Berghe, G., Van den Born, G. J.: Greenhouse gas emissions in the Netherlands 1990-2006, National inventory report 2008. MNP report 500080009, Netherlands Environmental Assessment Agency (MNP), 2008.
- Moore, T.R and Knowles, R: Methane emissions from fen, bog and swamp peatlands in Quebec. *BIOGEOCHEMISTRY* 11, pp. 45-61, 1990.
- Nieveen, J.P., Campbell, D.I., Schipper, L.A., Blair, I.J.: Carbon exchange of grazed pasture on a drained peat soil, *Global Change Biology* 11, pp 607-618, 2005.
- Nol, L., Verburg, P.H., Heuvelink, G.B.M., Molenaar, K.: Effect of land cover data on nitrous oxide inventory in fen meadows. *J. of Envir. Qual.*: 37, pp. 1209-1219, 2008.
- Martikainen, P.J., Nykanen, H., Alm, J., Silvola, J: Change in fluxes of carbon-dioxide, methane and nitrous oxide due to forest drainage of mire sites of different trophic. *Plant and Soil* 168, pp. 571-577, 1995.
- Parmentier, F.J.W., van der Molen, M.K., de Jeu, R.A.M., Hedriks, D.M.D., Dolman, A.J.: CO₂ fluxes and evaporation on a peatland in the Netherlands appear not affected by water table fluctuations. *Agricultural and Forest Meteorology*, 149, pp. 1201-1208, 2008.
- Raamsdonk, L.W.D., Kan, C.A., Meijer, G.A.L., Kemme, P.A. in Dutch: Kengetallen van enkele landbouwhuisdieren en hun consumptiepatronen, Rapport 2007.010, RIKILT – Instituut voor voedselveiligheid, 2007.
- Ruiter, J.F., Van Pul, W.A.J., Van Jaarsveld, J.A., & Buijsman, E.: Zuur- en stikstofdepositie in Nederland in de periode 1981-2002. [RIVM rapport 500037005/2006 \(In Dutch\)](#), 2006.
- Schrier-Uijl, A.P., Kroon, P.S., Hensen, A., Leffelaar, P.A., Berendse, F. & Veenendaal, E.M.: Comparison of chamber and eddy covariance based CO₂ and CH₄ emission estimates in a heterogeneous grass ecosystem on peat, *Agric. For. Meteorol.*, doi:10.1016/j.agrformet.2009.11.007, 2009.
- Schrier-Uijl, A.P., Veenendaal, E.M., Leffelaar, P.A., van Huissteden, J.C., Berendse, F.: Methane emissions in two drained peat agro-ecosystems with high and low agricultural intensity. *Plant Soil*, doi:10.1007/s11104-009-0180-1, 2010.
- Schrier-Uijl, A.P., Veraart, A.J., Leffelaar, P.A., Berendse, F., Veenendaal, E.M: Release of CO₂ and CH₄ from lakes and drainage ditches in temperate wetlands. *Biogeochemistry*, doi:10.1007/s10533-010-9440-7, 2011.
- [Schuepp, P. H., Leclerc, M. Y., McPherson, J. I., and Desjardins, R. L.: Footprint prediction of scalar fluxes from analytical solutions of the diffusion equation, *Boundary-Layer Meteorol.*, 50, 355–374, 1990.](#)
- Schulp, C.J.E., Nabuurs, G.J., Verburg, P.H.: [Future carbon sequestration in Europe – Effects of land use change. *Ecosyst. And Environm.* 127, pp. 251-264, 2008.](#)
- Sneath, W, Beline F, Hilhorst A, Peu P: Monitoring GHG from manure stores on organic and conventional dairy farms, *Agric. Ecosyst. Environ.* 112, pp. 122–128, 2006.
- Sommer, S.G., Olesen, J.E., Petersen, S.O., Weisberg, M.R., Vallis, L., Rodhe, L., Beline, F.: Region-specific assessment of greenhouse gas mitigation with different manure management strategies in four agro ecological zones. *Global Change Biology* 15, pp. 2825-2837, 2009.
- Soussana, J.F., Loiseau, P., Vuichard, N., Ceschia, E., Balesdent, J., Chevallier, T., Arrouays, D.: Carbon cycling and sequestration opportunities in temperate grasslands, *Soil Use and Management* 20, pp. 219-230, 2004.
- Soussana, J.F., Maire, V., Gross, N., Reinhold, T., Dahring, H., Pontes, L.S., Picon-Cochard, C., Fontaine, S., Pages, L., Wirth, C.: Modeling the relationships between the diversity and the functioning of pasture swards with a complex floristic composition, *Fourrages* 195, pp. 259-274, 2008.

Met opmaak: Engels (V.S.)

Met opmaak: Engels (V.S.)

Met opmaak: Lettertype: Arial, 10 pt, Engels (V.S.)

Met opmaak: Inspringing: Links: 0 cm

Met opmaak: Lettertype: Arial, 10 pt, Engels (V.S.)

Verwijderd: ¶

Met opmaak: Lettertype: 10 pt, Engels (V.S.)

Met opmaak: Lettertype: 10 pt, Engels (V.S.)

Met opmaak: Inspringing: Links: 0 cm, Afstand Na: 0 pt

Stouthamer, E. and ten Brinke, W., in Dutch: Fluvial sedimentology, *Geologie en mijnbouw* 86, pp. 1-1, 2008.

TDN. TOP10vector Product information (in Dutch): Topografische Dienst Kadaster Emmen the Netherlands. Available at <http://www.tdn.nl/> (verified 6 Dec 2007), 2006.

Teh, Y.A., Silver, W.L., Sonnentag, O., Detto, M., Kelly, M and Baldocchi, D.D.: Large Greenhouse Gas Emissions from a Temperate Peatland Pasture. *Ecosystems* 14: 311–325. DOI: 10.1007/s10021-011-9411-4, 2011

Van Amstel A., Kroeze C., van Eerd M., Dumont M., Both D.: National emissions of methane and nitrous oxide from agriculture in The Netherlands: current estimates and future trends, Workshop on Inventories and projections of greenhouse gas and ammonia emissions from agriculture, 27–28 February 2003, European Environmental Agency, Copenhagen, 2003.

Van den Pol-van Dasselaar, A., Van Beusichem, M.L., Oenema, O.: Effects of nitrogen input and grazing on methane fluxes of extensively and intensively managed grasslands in the Netherlands. *Biol. Fertil. Soils*, 29, pp. 24-30, 1999.

Veenendaal, E.M., Kolle, O., Leffelaar, P.S. Schrier-Uijl, A.P., Van Huissteden, J., Van Walsem, J., Möller, F. & Berends, F.: CO₂ exchange and the carbon balance in two grassland sites on eutrophic drained peat soils. *Biogeosciences*, 4, pp. 1027-1040, 2007.

Velthof, G.L. and Oenema, O.: Nitrous oxide emission from dairy farming systems in the Netherlands. *Netherlands Journal of Agricultural Science*, 45, pp. 347-360, 1997.

Whiting, G.J. and Chanton, J.P.: Primary Production Control of Methane Emission from wetlands. *Nature*. 364, pp. 794-795, 1993.

Met opmaak: Engels (Groot-Brittannië)

Met opmaak: Engels (Groot-Brittannië)

Met opmaak: Engels (Groot-Brittannië)

Met opmaak: Engels (Groot-Brittannië)

Met opmaak: Engels (Groot-Brittannië)

Met opmaak: Engels (Groot-Brittannië)

Met opmaak: Engels (Groot-Brittannië)

Table 1. Site descriptions, land use and management per peat site

Loc.	Mean annual WT depth (m)	Peat depth (m)	Land elements				Land use	Grazing ²	Biomass Removal ²	Cow manure Applied ²	Fertiliser Applied ²
			Dry land	Wet Land	Saturated land	Water					
	(m)	%	%	%	%			(ton ha ⁻¹ yr ⁻¹)	(kgN ha ⁻¹ yr ⁻¹)	(kgN ha ⁻¹ yr ⁻¹)	
Ou ₁	0.55	12	79		5	16	intensively managed grassland	2005 and 2006 by some cows	12	300	88
St ¹	0.45	12	79		5	16	extensively managed hayfield	young cattle few days per year	10	0	0
Ho ₁	0.25	2.1	60	25	10	5	former managed area under restoration	None	0	0	0

¹Ou= Oukoop, St= Stein, Ho= Horstermeer.

²Values related to management are averaged over the years 2006, 2007 and 2008.

Table 2. Landscape elements in the Dutch peatlands.

Landscape element/land use		Surface area (ha)	Surface area (%)
Grassland/managed	intensively	78,375	68%
Grassland/managed	extensively	8,786	8%
Water		6,717	6%
Urban area (incl. greenhouses)		983	1%
Roads		4,490	4%
Forest		2,716	2%
Cropland		1,818	2%
Other land use		11,258	10%
Total		115,142	100%

¹n.e. = not estimated

Kadaster, 2010. Product information TOP10vector (in Dutch). Topografische Dienst Kadaster, Emmen, the Netherlands. Available at <http://www.tdn.nl/> (verified 3 May 2010).

Met opmaak: Engels (Groot-Brittannië)

Table 3. Terrestrial carbon dioxide flux estimates ($\text{kg CO}_2 \text{ m}^{-2}$) measured by eddy covariance in the period 2005-2008 for the intensively managed production grassland on peat (Ou), the extensively managed hayfield on peat (St) and the unmanaged former agricultural peatland (Ho). Fluxes from shallow lakes are measured in the summer of 2009. Fluxes from removed biomass are not included in this table.

Site	Carbon dioxide NEE ¹ per year ($\text{kg CO}_2 \text{ m}^{-2} \text{ yr}^{-1}$)				Average
	2005	2006	2007	2008	
Ou	0.4	0.7	-0.2	0.7	0.4
St	-0.09	0.8	0.3	0.8	0.4
Ho	-1.0	-1.0	-1.9	-1.5	-1.4

¹NEE= Net Ecosystem Exchange

² Fluxes are measured in the summer (June/July) of 2009 in 5 large shallow lakes located in peatlands.

Table 4. Yearly methane fluxes ($\text{g CH}_4 \text{ m}^{-2} \text{ yr}^{-1}$) 1) measured by the chamber method and calculated by using landscape element weighted predictive relationships for Oukoop, Stein and Horstermeer and 2) measured by eddy covariance in the intensively managed site and modelled by using predictive relationships. Uncertainties are given within brackets and are averaged over the three or four years. Farm based emissions are not included in this table.

Ecosystem	Annual methane fluxes				Average emission
	$(\text{g CH}_4 \text{ m}^{-2} \text{ yr}^{-1})$				$(\text{g CH}_4 \text{ m}^{-2} \text{ yr}^{-1})$
Spatially weighted for contribution of fields, ditches and edges	2005	2006	2007	2008	Average
Ou (chamber-method)	NA ¹	20.3	16.2	14.6	17.0 ($\pm 56\%$)
Ou (eddy covariance-method)	NA ¹	17.2	16.6	15.5	16.4 ($\pm 37\%$)
St (chamber-method)	NA ¹	15.7	18.0	16.3	16.7 ($\pm 59\%$)
Ho (chamber-method)	19.1	20.5	19.8	17.6	19.2 ($\pm 65\%$)

¹NA = not available.

Table 5. Nitrous oxide flux estimates ($\text{kg N}_2\text{O ha}^{-1} \text{ yr}^{-1}$) and their uncertainties (u) for the intensively managed site (Oukoop), extensively managed site (Stein) and the unmanaged peatland (Horstermeer).

Site	Source	Reference	Total emission ($\text{kg N}_2\text{O ha}^{-1} \text{ yr}^{-1}$) ¹
Oukoop	Background emission ¹	Kroon et al., 2010	24 ($\pm 28\%$)
	due to fertilizers ²	Kroon et al., 2010	
	due to leaching and run-off	Kroon et al., 2010; IPCC, 2006	
	due to deposition	Kroon et al., 2010; IPCC, 2006	
Stein	Total emission	Velthof et al., 1997; IPCC, 2006	8 ($\pm 100\%$)
Horstermeer	Total emission	Velthof et al., 1997; IPCC, 2006	8 ($\pm 100\%$)

¹ Background emissions are determined by a multivariate regression model based on EC flux data excluding EC fluxes measured around a management event.

² Emissions due to fertilizer application have been determined by subtraction the background emission from the total measured N_2O emission around fertilizer application. The IPCC default value of 1% is used for the missing fertilizing events.

Tabel 6. Estimated area and annual GHG release for the areal of intensively managed and extensively managed (mown only) or unmanaged grasslands on peat within the total western peatland region of the Netherlands. Farm-based emissions are not included.

Ecosystem type	Area in western peatland		Total N ₂ O emission	Total CH ₄ emission	Total CO ₂ emission
	(ha)	(% of total)	10 ³ kg N ₂ O yr ⁻¹	10 ³ kg CH ₄ yr ⁻¹	10 ³ kg CO ₂ yr ⁻¹
Intensively managed grassland	78,375	68%	1,653	12,853	313,498
Extensively managed/ unmanaged grassland	8,786	8%	43	1,577	35,145
Shallow water bodies	87	6%	unknown	unknown	33,583 ¹

¹An annual emission of 0.5 kg CO₂ m⁻² yr⁻² was assumed (Table 3).

Tabel 7. Overview of the expected effects of different mitigation strategies on the total GHG balance (ecosystem + farm based emissions). The effect on the GHG balance has been determined for the three research sites Oukoop (intensively managed), Stein (extensively managed) and Horstermeer (unmanaged) and are not including the expected future temperature rise. (-) = decrease in emission, (+) = increase in emission, (0) = neutral effect, (?) = effect unknown, (x) = not relevant.

	Rewetting+management reduction			Management reduction towards extensively managed			Increase in % open water with no reduction of management in the catchment		
	CO ₂	CH ₄	N ₂ O	CO ₂	CH ₄	N ₂ O	CO ₂	CH ₄	N ₂ O
Intensive management	-	-	-	0	-	-	-	+	-
Extensive management	-	-	-	x	x	x	-	+	?
Unmanaged/rewettered	x	x	x	x	x	x	-	+	?
Open water*	-	-	-	-	-	-	x	x	x

* The influence of rewetting and land management reduction in the surrounding area on emissions from 'open water'.


Met opmaak: Engels (Groot-Brittannië)

Met opmaak: Engels (Groot-Brittannië)

Met opmaak: Engels (Groot-Brittannië)

Met opmaak: Engels (Groot-Brittannië)

Figure 1.




Met opmaak: Engels
(Groot-Brittannië)

Met opmaak: Engels
(Groot-Brittannië)

Figure 1. Monthly averages for water table and soil temperature in Oukoop (Ou), Stein (St) and Horstermeer (Ho).

Figure 2.



Met opmaak: Engels (Groot-Brittannië)

Met opmaak: Engels (Groot-Brittannië)

Verwijderd: 1

Met opmaak: Engels (Groot-Brittannië)

Met opmaak: Engels (Groot-Brittannië)

Figure 2. Ecosystem and farm-based GHG fluxes (CO_2 respiration (R_{CO_2}), CO_2 gross primary production or photosynthesis (GPP_{CO_2}), CH_4 and N_2O) and carbon fluxes ($\text{CO}_2\text{-C}$, $\text{CH}_4\text{-C}$, manure and fertilizer-C, biomass-C) that are being considered in the current study for Oukoop, Stein and Horstermeer. White arrows are farm-related fluxes and dark grey arrows are ecosystem fluxes.

Verwijderd: 1

Figure 3.





Fig. 3. System boundaries of the intensively managed area Oukoop. Black arrows are C flows, thick dashed arrows are CH₄ fluxes and dashed arrows are CO₂ fluxes (autotrophic and heterotrophic respiration; R_{CO₂auto} and R_{CO₂}, respectively and photosynthesis(GEP_{CO₂})).

Verwijderd: 3

Figure 4.




Verwijderd:

Figure 4. Daily NEE CO₂ for the Oukoop (intensively managed), Stein (extensively managed) and Horstermeer (unmanaged) peatlands measured by the eddy covariance flux technique. The black line represents the cumulative NEE for each year separately (y-axis on the right) and the grey line represents the temporal variability of NEE on the time scale of a day (y-axis on the left). The x-axis represents the day number since 1 January 2005.

Verwijderd: 3

Verwijderd: 4

Figure 5.



Verwijderd: 4






Figure 5. Monthly NEE CO₂ values for the three experimental sites. The CO₂ NEE is given on the y-axis in g CO₂ m⁻² d⁻¹ and the month numbers are given from 2006-2008 on the x-axis.

Verwijderd: 4

Figure 6.

Verwijerd: 5




Verwijerd:

Figure 6. Temporal variability of terrestrial CH₄ fluxes for the three experimental sites modelled by predictive relationships based on chamber measurements and additionally for Oukoop measured by continuous eddy covariance. The right-hand y-axis represents the cumulative CH₄ flux over the three years. Fluxes are weighted by the contribution of each landscape element.

Verwijerd: 5

Verwijderd: 6

Figure 7.





Verwijderd: 6

Figure 7. Summary of all considered carbon fluxes in the research areas Horstermeer (Ho), Stein (St) and Oukoop (Ou) averaged over 2005, 2006, 2007 and 2008. The annual carbon balance is presented in kg C ha⁻¹ yr⁻¹, (+) is release and (-) is uptake, and consists of fluxes due to GHG emissions (NEE CO₂ and NEE CH₄) and fluxes due to management (farm based fluxes, manure application and biomass removal).

Figure 8.

Verwijderd: 7





Verwijderd:

Figure 8. The GHG balances including CO₂, CH₄ and N₂O for the three sites: intensive (Oukoop), extensive (Stein) and unmanaged (Horstermeer). On the left excluding farm-based CH₄ and CO₂ emissions and on the right including farm-based CH₄ and CO₂ emissions, averaged over 2006, 2007 and 2008 (fluxes are given in warming potentials, kg CO₂-equivalents m⁻² yr⁻¹).

Verwijderd: 7

Figure 9.

Verwijderd: 8



Verwijderd:

Figure 9. Comparison of summer (June/July) CH₄ fluxes between different ecosystems in peatlands. Fluxes are averaged over three years for the polder Oukoop (intensive), the polder Stein (extensive) and the polder Horstermeer (unmanaged). The lake fluxes are measured in one summer (June/July, n=97) season and are averaged over 5 large shallow lakes; error bars represent the standard deviation of the mean.

Verwijderd: 8

Instrumentation and methodology

To capture temporal variability of fluxes, all three sites were equipped with eddy covariance (EC) systems for CO₂ since the end of 2004 for four years (for details of the systems see Veenendaal et al., 2007; for Oukoop and Stein and Hendriks et al., 2007 for Horstermeer). In addition, Oukoop was equipped with an EC system for CH₄ and N₂O since the beginning of 2006 for three years (details in Kroon et al., 2007, 2010b). In April 2006 the Horstermeer site was equipped with an EC system for CH₄ (Hendriks et al., 2008, 2009). At the Oukoop and Stein sites chamber measurements were performed from January 2006 to December 2008 (Schrier-Uijl et al 2010) and at the Horstermeer site from January 2005 to December 2008 (Hendriks et al., 2007).

For the three sites, similar measurement procedures, flux calculation methods, gap filling techniques and up-scaling methods were used to calculate annual GHG fluxes. Procedures to calculate annual fluxes are explained in more detail in Schrier-Uijl et al. (2009, 2010); Kroon et al. (2007, 2010b); Hendriks et al. (2009, 2007) and Veenendaal et al., (2007). For spatial up-scaling of chamber measurements, the landscape elements (ditches, saturated soil, dry soil) that contributed significantly different to the GHG balance were taken into account proportionally.

Landscape scale flux measurements

Landscape scale flux measurements of CO₂, CH₄ and N₂O were performed with an averaging time of 30 minutes from 10 Hz data using the EC flux technique. EC flux systems for CO₂ at the three sites consisted of a sonic anemometer (Oukoop and Stein: Campbell CSAT C3 Sonic Anemometer (Campbell Scientific, Utah, USA) Horstermeer: Windmaster Pro 3 Axis Ultra Sonic Anemometer (Gill Instruments Limited, UK)) and a fast response CO₂-H₂O analyzer (all sites: LICOR 7500 open path infrared gas analyser (LICOR, Lincoln, NE, USA).

The EC flux systems were installed 4.3 meter above the soil surface in Horstermeer and 3.05 meter above the soil surface in Oukoop and Stein. Oukoop was also equipped with an EC flux system for CH₄ and N₂O in the period 2006-2008 consisting of a sonic anemometer (model WMPRO from October 2007- July 2008 and model R3 the rest of the measurement period, Gill instruments, Lymington, UK in 2008) and a Quantum cascade laser spectrometer (model QCL-TILDAS-76, Aerodyne Research Inc., Billerica MA, USA).. The EUROFLUX methodology for quality control (including filtering for spikes, frequency losses and webb-corrections) and data correction was applied to determine the fluxes on a thirty minute basis (Aubinet et al., 2000). Footprints of the fluxes were calculated by using the Kormann and Meixner method (Kormann and Meixner, 2001) and Schuepp et al., 1990). The footprint included fields, ditches and ditches edges but excluding farms and other GHG hotspots. Soil respiration in the three sites was described as a function of soil temperature by using the Arrhenius relation (Lloyd and Taylor, 2004). For CO₂, data coverage was over 60% for each site. Annual balances for CH₄ at Oukoop were derived using a multivariate regression model, including soil temperature and wind velocity. Annual balances for N₂O were calculated from four contributions: background emissions, event emissions due to fertilizer application, leaching and deposition. Background emission was derived from the EC flux data by using a multivariate regression model including soil temperature and wind velocity. Also the fertilizer event emissions are separately derive from EX flux data (see table 5). For N₂O and CH₄ data coverage was 48%. For more details about EC flux measurements see Veenendaal et al., 2007; Hendriks et al., 2007; Kroon et al., 2009; Kroon et al., 2010).

Small-scale flux measurements

For all sites, small-scale flux measurements were performed on dry fields, saturated ditch edges and open water (ditches) for CO₂ and CH₄ using a Photo Acoustic Field Gas Monitor (Oukoop and Stein: type 1412 and Horstermeer type 1312,2, Innova AirTech Instruments, Ballerup, Denmark) connected with Teflon tubes to closed opaque chambers (height 0.2 m, diameter 0.3 m). For each gas measurement, five samples were taken from the headspace of a closed cylindrical, non-transparent chamber over a period of 5-10 minutes from which the flux (height of the chamber multiplied by dC/dt) was calculated. Linearity was checked using the intercept method (Kroon et al., 2008). A fan was installed in the chamber to homogenize the inside air and two external filters were added: a soda lime filter for CO₂ (when measuring CH₄) and a silica gel filter for water vapour. To avoid disturbances, vegetation was not removed prior to the flux measurements. Concentration build-up in the chamber headspace was measured for 6 minutes at one minute intervals. Because the relations with explanatory variables were non-linear, multiple non-linear regression was used to calculate annual emissions, and the landscape elements that contributed significantly differently to the GHG balance were taken into account. Landscape scale fluxes were upscaled from the emissions from the different landscape elements (fields, saturated edges and ditches) by using a weight factor relative to their proportion in the landscape. Additional to the three years measurements on ditches, in the summer of 2009 an intensive measurement campaign was performed on 6 large shallow lakes and 14 drainage ditches in peatlands (Schrier-Uijl et al (2011)) to compare fluxes from water bodies that were different in depth, size and nutrient status. Cross-checks of emission values were performed by comparing eddy covariance measurements to upscaled chamber measurements within the footprint of the eddy covariance systems (Schrier-Uijl et al (2009); Hendriks et al (2009)).

Landscape scale flux measurements

Landscape scale flux measurements of CO₂, CH₄ and N₂O were performed using the EC flux technique. Footprints of all EC flux towers were over the entire landscape, including fields, ditches and ditch edges but excluding farms and other GHG hotspots. EC flux systems for CO₂ at the three sites consisted of a sonic anemometer and a fast response CO₂-H₂O analyzer. Open path infrared gas analyzers (LI-COR Lincoln, NE, USA) were used 4.3 meter above the soil surface in Horstermeer and 3.05 meter above the soil surface in Oukoop and Stein. Oukoop was also equipped with an EC flux system for CH₄ and N₂O in the period 2006-2008 consisting of a sonic anemometer and a Quantum cascade laser spectrometer (model QCL-TILDAS-76, Aerodyne Research Inc., Billerica MA, USA). In all three sites additional micrometeorological measurements were performed.

Additional measurements and analyses

Meteorological measurements and soil measurements

In addition to EC devices and chamber measurements, GHG measurements, towers with meteorological instruments were installed at each site which provided 30-minute averages of global radiation (R_{in}), net radiation (R_{net}), air temperature (T_{air}), vapour pressure (P_{vap}), wind speed (U), wind direction (D) and precipitation (P) were measured at each site using the sonic anemometers.

Soil measurement sensors included soil heat flux plates (HPF01, Campbell Scientific, USA), soil temperature sensors at depths of 0.02, 0.04, 0.08, 0.16 and 0.32 m (Campbell Scientific, USA) and soil moisture probes to measure volumetric moisture contents at depths of 0.10, 0.20 and 0.30 m (Theta probes ML 2x; Delta T devices Burwell, UK). These systems provided 30 minute values for soil heat fluxes, soil temperature, soil moisture and water table. At the beginning and the end of the experiments (2005 and 2008), soils were sampled and analyzed for C and N content, organic matter, NO_3^- , NH_4^+ , PO_4^{3-} and pH.

Analyses

Water from drainage ditches was sampled for pH, C content (not for Horstermeer), N content (not for Horstermeer), organic matter, NO_3^- , NH_4^+ , PO_4^{3-} , SO_4^{2-} , Fe^{2+} , dissolved CH_4 , oxygen saturation and electrical conductivity. Well-stirred samples of slurry manure were sampled just before manure application in the Oukoop site and were analyzed for dry matter and C content. To estimate the above ground biomass and biomass removal, vegetation height was measured every three to four weeks to estimate above ground biomass and biomass removal using the relationship between vegetation height and biomass weight ($R^2 = 0.84$; $n = 51$) as discussed in Veenendaal et al. (2007) .