

Interactive comment on “A model of potential carbon dioxide efflux from surface water across England and Wales using headwater stream survey data and landscape predictors” by B. G. Rawlins et al.

Anonymous Referee #2

Received and published: 18 December 2013

General Comment

1. The authors present a model containing 10 variables which predicts 24% of the spatial/temporal variation in pCO₂. This suggests it has limited value. I believe that one of the main problems with this approach is that the model is predicting pCO₂ values which have been modelled themselves.
2. There is a significant N American, European (inc. UK) literature on pCO₂ and CO₂ evasion, which is poorly represented in the manuscript.

C7468

Specific Comments

16455, L1: Several papers (see Kling et al. 1991 Science; Hope et al. 2001 L&O) already show that CO₂ evasion is likely to account for much, much more than 10% of NEE.

16456, L16-17: it would be better to turn this into a testable hypothesis.

16457, L3-6: what is the justification for this statement or assumption? These are the areas of the catchment (headwaters) where evasion rates are known (and have been measured) to be highest.

16457, L11-14: the authors are making a huge assumption here, that all free CO₂ evades downstream so there is no need to calculate k values. There is a significant body of literature to show that CO₂ concentrations (even in many large rivers systems) never reach equilibrium with the atmosphere. It is also unclear what the authors mean by “limited downstream changes in water chemistry”. The statement needs to be clarified, particularly as rivers typically show significant spatial changes in water chemistry.

16457, L21: this is a misrepresentation of Dinsmore et al. (2010) – I believe their work is based on one headwater site only.

16459, L19: “theoretical pCO₂” is being modeled by the authors, from several variables, but it is unclear which ones (pH? temperature? DIC?). The authors need to clearly state how they are doing this in their model, so readers can evaluate its usefulness for themselves. It would also be appropriate (like most models) to validate it against real pCO₂ data.

16461, L6: the authors need to define “dominant” land cover class. Does this mean >50% coverage?

16461, L16: what is the BFIHOST value?

16464, L8-10: It's not clear why the authors expect to find higher pCO₂ in agricultural

C7469

streams; pCO₂ is not just based on soil productivity. These systems typically contain well-aerated, highly managed soils, which lose soil CO₂ rapidly to the atmosphere. Poorly drained, organic-rich natural systems are where highest pCO₂ occur because of their poor drainage and their ability to accumulate significant sub-surface CO₂ stores which connect to the aquatic pathway.

16467, L2-3: CO₂ evasion = pCO₂ x flow is a gross over-simplification, because there are so many factors which influence CO₂ concentration in the aquatic system.

Figure 7: Is the data presented in this figure realistic? The suggestion is that the highest evasion rates occur in the higher pH soils and not in the organic-rich upland areas of the UK. High pH soils produce circum-neutral or high pH streamwater, which based on the carbonate equilibrium, contains little or no free CO₂. I suggest the authors revisit their model as it appears to calculate pCO₂ incorrectly. Further underlying data or variables like pH and DIC, would help the reader sense-check these regional differences.

Interactive comment on Biogeosciences Discuss., 10, 16453, 2013.