

## ***Interactive comment on “Comparison of different methods to determine the degree of peat decomposition in peat bogs” by H. Biester et al.***

**H. Biester et al.**

h.biester@tu-bs.de

Received and published: 20 March 2014

Bg-2013-444 Final Response to referee comments:

We thank the reviewers for their constructive comments and suggestions which we believe will improve the manuscript.

Referee #1 This paper aims to evaluate peat decomposition proxies as indicators for decomposition processes in two ombrotrophic mires. They characterize and compare several measures of decomposition proxies in depth profiles of these mires, including pyrolysis-GC-MS, C/N ratios, stable isotopes of C and N, FTIR spectroscopy, Rock Eval oxygen and hydrogen indices, and humic acid UV adsorption. This study is an impressive and comprehensive comparison of different methods used to quantify OM de-

C9328

composition. The authors highlight that depth profiles of many of these indices are both affected by decomposition processes, vegetation changes, and possibly other external and internal factors (e.g., N deposition and recycling). A particular strength of this study is the use of molecular analysis using GC-MS to attempt to differentiate between these processes and discuss the performance of each of the decomposition proxies in regards to molecular tracers. As mentioned, this is a very comprehensive characterization of biogeochemical decomposition measures in peat bogs with high relevance for climate reconstruction and characterization of historic deposition inputs of trace elements. The analyses, although I am not fully familiar with all of the techniques used, seem very sound, are well described, and seem to have appropriate quality control measures. Overall, this is a highly suitable paper for publication in Biogeosciences. The introduction is very well written, clearly lays out the issues with many of these decomposition proxies, and is very well referenced – in short, it does an excellent job to highlight the need and importance of this study.

My major concern with the paper is the current structure of the results and discussion sections which makes it very difficult for the reader to understand the analysis, comparisons, and their respective interpretations. One reason for this may be that the paper is extremely dense and comprehensive, and as such presents a vast amount of data and chemical analysis which are difficult to fully absorb. However, I had to read the results and discussion sections many times in order to understand the direction and interpretation of all analysis performed. I strongly suggest that these two sections need to be comprehensively restructured for the readers to be able to follow the arguments made by the authors – see section below for some suggestions. In particular section 4.3. and the following subsections (4.3.1 through 4.3.5) are extremely cumbersome to follow. I think the main reason is the introduction of the Principal Component Analysis (PCA) and the two factors F1 and F2 that represent water table heights and degradation of vascular plants based on analysis of GC-MS data. While this is very elegant analysis to compare decomposition measures to molecular tracers, the authors need to prepare the reader much earlier and more directly for this approach. For example, they should

C9329

make it a specific goal in the introduction that their evaluation of decomposition proxies is based in large parts on molecular analysis and PCA factors, and clearly state that the factors allow to differentiate between processes contributing to changes in decomposition proxies (i.e., water table height affecting species abundance and decomposition, and decomposition of vascular plants).

A: We agree with the reviewer that the manuscript contains a lot of data and high information density and that some restructuring as proposed by the reviewer might help to improve the readability of the manuscript. We restructured the manuscript following the reviewer's suggestions. First we described at the end of the introduction section that peat structural information obtained by pyrolysis-GC-MS was used as the basis for the interpretation of the peat decomposition indicators. We also added a sentence on how PCA was used to decipher biogeochemical processes related to peat decomposition and vegetation changes. We have also moved the description of pyrolysis-GC-MS results to the beginning of the result section to give the molecular composition of the peat first.

R: If this is clearly laid out in the introduction, and the results and discussion section also put an emphasis on this approach, it will be much easier to follow comparison of other decomposition proxies to this approach. Similarly, figures should be re-arranged in away to help this restructuring. For example, Figure 4 introducing the two main factors should be very early in the manuscript and be discussed early in results and discussion sections. Currently, figure 3 and many text sections show and discuss PCA factors F1 and F2 before they are clearly discussed and interpreted.

A: We have now moved the interpretation and discussion of the pyrolysis-GC-MS results to the beginning of the discussion section so that this information is now available for further discussion of the other decomposition proxies. According to this new structure, Fig.1 is now showing the results (depth profiles) of pyrolysis-GC-MS, followed by two figures showing the results (depth profiles) of the decomposition proxies. The discussion section is starting now with the figure showing the interpretation of the

C9330

pyrolysis-GC-MS results (distribution of factor loading of proxies and pyrolysis products).

R: Then, I suggest that a more structured comparison of decomposition proxies to these factors should be done; for example, in Figure 5 only one regression is shown relating decomposition proxies to F1 (i.e., for C/N), but large parts of sections 4.3.1 through 4.3.5. specifically discuss relationships to F1 and F2 to decomposition proxies without showing these relationships. To help this, the correlation matrix between factors F1/F2 and decomposition proxies currently shown in Suppl. Table S3 could be moved to the main text. Other figures of the manuscript (e.g., Fig. 1, Fig. 2, and possibly Figure 3) could possibly be moved from the main manuscript to the supplemental information.

A: We agree that some more of the relationships (correlations) between decomposition proxies and F1 and F2 factors should be shown. We decided to add this information to Figure 5 because we think that a correlation matrix with large amounts of proxies in the main text is even more difficult to follow.

R: While I do not insist on restructuring the manuscript exactly the way I mentioned above, I need to re-emphasize the need to restructure the results and discussion section. This is really a great dataset with unique and elegant analysis and interpretations, but I think in its current form many readers will be overwhelmed and not be able to follow the reasoning of the authors without repeated reading.

A: We agree on that, however, we doubt that it is possible to thoroughly understand a data set like this without repeated reading, whatever the structure might be.

R: Other comments: Page 17352: Abstract: what is missing in the abstract is the entire approach using PCA analysis and PCA factors based on molecular characterization and their relationships to decomposition proxies. This is one of the main aspects in discussion and should be clearly mentioned and described in the abstract. Has been added

C9331

Page 17352, line 8: “if not misleading” – clarify what is meant here or reword, why/what is misleading?

Has been removed

Page 17352, line 17: define abbreviation “UV-ABS” when first mentioned

Has been added

Page 17352, line 25: “showed less variation” – less than what, do they mean they were stable through the profile?

Has been added

Page 17352, line 28: “reasonable information despite their bulk nature” clarify what is meant and reword

Has been changed

Page 17353, line 21: “has been reported recently” delete recently as these studies are over 10 years ago now.

Has been removed

Page 17354, line 16: add reference

Has been added

Page 17354, line 17: write out versus

Has been changed

Page 17355, line 11: add reference about the impact of fire incidence.

Has been changed

Page 17355, line 19: “more narrow” I assume they mean lower C/N ratios indicative of more decomposed peat material?

C9332

Has been changed

Page 17357, line 4: reword the entire sentence starting with “Unless the fact that. . .”

Has been changed

Page 17357, line 20, the authors use H/C and O/C indices earlier in the manuscript and here use HI and OI (I guess they mean hydrogen and oxygen indices) – they should be consistent with the terminology through the manuscript and in figures.

A: This was correct because with mentioning the H/C and O/C ratios we intend to indicate what the HI and OI indices actually show.

Page 17359, line 22: define GC-TCD

Has been added

Page 17362: Results: a very important component of the discussion is based on PCA and comparison of decomposition proxies to factors F1 and F2. This should be presented in the results section, which will make the discussion easier to follow.

Has been restructured

Page 17362, line 12-18; and Table 1 and 2: the authors should show stdev in Table 1, and discuss if  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  are statistically different instead of stating “slightly lighter”

Has been changed

Page 17362, line 23: “a similar trend of increasing values. . .”: the authors should quantify the similarities of trends, e.g., by referring to the correlation matrix or perform consistent scatterplots/regressions between different proxies.

A: We did not quantify the trend, but referred to the correlation matrix now

Page 17363, line 2: “seem to show inverse relationship. . .” same as above, please quantify.

C9333

A: We referred to the correlation matrix now.

Page 17363, line 24: “a remarkable sharp decrease. . .” this was also observed in the KK profile at about 43 cm depth, why not mention and discuss that also for the KK profile?

Has been added

Page 17363, line 12: “do not correspond to those of any other decomposition proxy” – as mentioned above, the correlations/correlation matrix should be done for all proxy-comparison to quantify similarities/differences between proxies.

A: This information is in the correlation matrix, reference added.

Page 17364, line 3: for that reason, molecular component were only determined in the KK core” I have no problem that molecular composition was only determined in onecore, but I would remove the justification given – as mentioned above, some of the traditional decomposition proxies show differences between the cores.

A: We think we should keep this explanation, otherwise the reader will probably not understand why the KRB core has not been analyzed.

Page 17365 and 17366: section 3.2.1. and 3.2.2. The reason for presenting these vegetation markers and molecular characterization becomes only clear later in the discussion section and the use of these markers in Figure 4. It would help tremendously if the reader is prepared for the use of these markers in the PCA analysis and their factors, maybe make a separate results section how the markers are used to develop factors F1 and F2 which will then be compared systematically to traditional decomposition proxies.

A: As described above, we changed the structure of the manuscript to make the role of the pyrolysis-GC-MS data and the PCA clearer. We think that PCA should not be a part of the results, but we moved it to the beginning of the discussion section, so that the information of F1 and F2 is now available for discussing the other proxies.

C9334

Page 17366 and following: The discussion should start with a detailed description of the use of PCA and Factors 1 and 2, and their interpretation, and then systematically interpret all traditional decomposition proxies in regards to this analysis. This would greatly help to follow the manuscript.

Has been restructured (see comments above).

Page 17366 section 4.1; This section would be much clearer to follow if analysis of molecular tracers was done first and the authors already discuss relationships to factors F1 and F2 although it is not clear yet what these mean.

Has been restructured (see comments above).

Page 17373, lines 4: “the fact that in our KK peat not lignin. . .” reword

Has been changed

Page 17374, line 12: A comparison of the FTIR intensity. . . is given in Suppl Fig S4). This should probably be mentioned earlier in this discussion section, why introduce new results at the end of a discussion section?

Has been changed

Section 4.3.3. Be consistent with the use of O/C and H/C and OI and HI.

Has been modified

Page 17378, line 7 reflecting change in mass loss and ARE related. . .

Sentence has been corrected

Page 17378, line 10 : change “sand” to and

Has been corrected

Page 17378, line 18: change “despites” to despite

Has been corrected

C9335

Page 17379, line 13: change "correlates" with correlate

Has been corrected

Page 17379, line 22, remove comma before changes

Has been changed

Referee #2 General Comments This manuscript has interest because it makes comparison between several techniques that might be used to investigate past changes in peat decomposition. It is based on studies on two peat cores from one locality – Harz Mountains – and the authors' general conclusions are drawn inductively from this particular location. However, whether it is possible to extrapolate from these specific findings to wider general application elsewhere is not known (and to an extent, this is acknowledged in the Conclusions). Indeed, because of the constituents of the peat from this location, the choice of mires for making these comparisons between techniques may limit applicability of the findings. For example, it is somewhat unfortunate that the authors chose mires in which *Eriophorum* remains are present in the peat, because this taxon is known to produce erratic and potentially erroneous values using one of the techniques reviewed (the colorimetric method), owing to the strong contrast in relative decay between unhumified *Eriophorum* fibres and the more humified peat matrix. Markedly different results can be obtained from the same analysed horizon in the peat, depending upon which of these fractions happens to predominate in any 0.2 g sub-sample (Chambers et al., 2011). So, it is perhaps not surprising that the UV-ABS method used on these Harz Mountain peats produced results that are different from some of the other techniques applied (this is partly admitted on p. 17377, lines 16–19, but the specific problem of *Eriophorum* is not mentioned).

A: We consider the choice of a peatland that has contributions from *Eriophorum* as representative, as most *Sphagnum*-dominated peatlands have contributions from this graminoid. Thus, the fact that the colorimetric method is influenced by vascular plants, such as *Eriophorum*, says more about the shortcomings of the method than the selec-

C9336

tion of the peatland. Our data do not indicate that the proposed mechanism influenced the method.

Specific Comments: The review of the literature is not complete. In places, these omissions have led inadvertently to inaccuracies. For example, on p. 17355 the authors state that "Blackford and Chambers (1993) introduced an alkaline extraction (NaOH) procedure combined with UV absorption measurements to determine differences in the degree of peat humification based on the leachate's color intensity." It would be more complete to say that "Aaby and Tauber (1975) used Bahnson's (1968) alkaline extraction (NaOH) procedure combined with UV absorption measurements to determine differences in the degree of peat humification based on the leachate's color intensity, and related this to bog surface wetness at the time of peat formation. Blackford and Chambers (1993) compared this colorimetric 'determination of peat humification' for reconstructing past bog surface wetness with various other simple methods (such as fibre content; von Post visual humification scale, etc.), and considered fibre content and the colorimetric technique to be superior. However, because 'percentage peat humification' is a dubious concept, they recommended that results from colorimetry should instead be reported using percentage light transmission values. A revised protocol for this colorimetric method was published recently by Chambers et al. (2011)."

Aaby, B. & Tauber, H. (1975) Rates of peat formation in relation to degree of humification and local environment, as shown by studies of a raised bog in Denmark. *Boreas*, 4, 1–14.

Bahnson, H. (1968) Kolorimetrisk bestemmelse af humificeringstal i højmosetør v fra Fuglsø mose på Djursland (Colorimetric determination of humification for bog peat from Fuglsø Mire in Jutland). *Meddelelser fra Dansk Geologisk Forening*, 18, 55–63.

Chambers, F.M., Beilman, D.W. & Yu, Z. (2011) Methods for determining peat humification and for quantifying peat bulk density, organic matter and carbon content for-

C9337

palaeostudies of climate and peatland carbon dynamics. *Mires and Peat*, 7, Article 07,1–10.

A: We appreciate this comment and have re-written this section according to the reviewer's suggestion. We also included the suggested references.

p. 17377, lines 22–55. There is no mention here of the work of Morgan et al. (2005), who used size-exclusion chromatography and showed, inter alia, that humic and fulvic acid compounds were being extracted from peats through alkali digestion; there were implications about their structures and molecular masses. Morgan, T.J., Herod, A.A.,

Brain, S.A., Chambers, F.M. and Kandiyoti, R. (2005) Examination of soil contaminated by coal-liquids by size-exclusion chromatography in 1-methyl-2-pyrrolidinone solution to evaluate interference from humic and fulvic acids and extracts from peat. *Journal of Chromatography*, 1095, 81–88

A: We suggest avoiding these references as they are focused solely on the chemical characteristics of NaOH extracted humic and fulvic acids, which is not the topic of our manuscript.

Nowhere in the manuscript are there cited any of the several relevant papers on proxy-climate methods from peats that were published in *Mires and Peat*, vol 7: <http://www.mires-and-peat.net/mpj3.html#Vol7> It is important to cite some of these because they represent state-of-the-art laboratory protocols for various proxy-climate methods from peats, including those of testate amoebae, peat geochemistry, pollen and non-pollen microfossils, peat humification, etc., and they are more up-to-date than other papers regarding specific techniques (e.g. analysis of testate amoebae) that are cited by the authors. Some of the papers (in that volume of *Mires and Peat*) were published in 2010; others in 2011. It is unclear also whether the field-sampling protocol of De Vleeschouwer et al. (2010; in that same volume) was followed by the authors when they sampled the Harz mires.

C9338

A: We have cited two references from *Mires and Peat* (Chambers, Klavins) which we think are relevant for this manuscript on peat decomposition. Our study does not include testate amoebae, pollen or non-pollen microfossils which are climate proxies but no specific proxies for peat decomposition; we therefore decide not to cite them here. Our peat cores were extracted in 2008 prior to the publication of the field-sampling protocol of De Vleeschouwer et al. (2010). We also think that this protocol is not crucial here, as it is predominately designed for trace element analysis and not validated for decomposition studies. We also believe that we gained sufficient experience during the past 15 years to extract representative peat samples.

p.17367 The comment that “Changes in humic acids may thus reflect more a signal of vegetation changes than of changes in humification or decomposition alone” is not a new finding. That there is a ‘species’ signal in peat humification records using the NaOH colorimetric technique was recognised by Chambers et al. (1995). This finding was also mentioned by Yeloff & Mauquoy, but the first recognition was in the following: Chambers, F.M., Barber, K.E., Maddy, D. and Brew, J. (1997) A 5500-year proxy climate and vegetational record from blanket mire at Talla Moss, Peebleshire, Scotland. *The Holocene*, 7, 391–399.

References has been added

Technical (text-drafting) corrections Several times, sentences begin ‘Due to’. In all these instances, what was meant was ‘Owing to’. There might also be occasions when ‘due to’ (= caused by) was used within sentences when ‘owing to’ (= because) might be better. See advice in Booth, V. *Communication in Science: Writing and Speaking*, CUP, Cambridge, p.

Has been changed according to the reviewer's suggestion.

17352, line 20, p. 17353, line 10, 17357, line 16 and p. 17368, line 3: change ‘extend’ to ‘extent’

C9339

Has been changed

p. 17358, line 8: change 'present day' to 'present-day'

Has been changed

p. 17371, line 6: change 'dryer' to 'drier'

Has been changed

p. 17361 lines 17-18 I wonder whether 'was adopted by Blackford and Chambers(1993)' should instead be 'of Blackford and Chambers (1993) was adopted' [Note also that this paper has been superseded by Chambers et al. (2011).]

p. 17362, line 15: hyphenate 'peat forming': 'peat-forming'

Has been changed

p. 17374, line 9. Write 'FTIR data, however, confirm' [not 'confirms']

Has been changed

p. 17375, line 14. Re-word the end of this sentence: 'discussed controversial' does not read well.

Has been changed

p. 17376, lines 17-18 'present day increase in grasses on the bogs surface' There are two things to change here: (i) hyphenate 'present day' (to 'present-day'); (ii) do something about the missing apostrophe in 'bogs surface'. It is not clear whether this sentence refers to one or to two bogs. If two, then place an apostrophe after the sin bogs. Or, can avoid an apostrophe altogether by writing as follows: 'present-day increase in grasses on the bog surface'

Has been changed

p. 17377, lines 16-17. Change 'due to the fact that' to 'because'

C9340

Has been changed

p. 17379. Change 'Moreover, the amount of UV-absorbing aromatic compounds which are e.g. abundant in lignin also depend on changes in vegetation, which does not necessarily correlate with' to 'Moreover, the amount of UV-absorbing aromatic compounds that are abundant in lignin, for example, also depend on changes in vegetation, which do not necessarily correlate with'.

Has been changed

---

Interactive comment on Biogeosciences Discuss., 10, 17351, 2013.

C9341