

# 1 Multi-factor controls on terrestrial 2 carbon dynamics in urbanised areas

---

3

4 C. Zhang<sup>1,2</sup>, H. Tian<sup>2</sup>, S. Pan<sup>2</sup>, G. Lockaby<sup>2</sup>, A. Chappelka<sup>2</sup>

5 1, Xinjiang Institute of Ecology and Geography, Chinese Academy of Sciences

6 2, International Center for Climate and Global Change Research, School of Forestry

7 and Wildlife Sciences, Auburn University, Auburn AL, 36849, USA

8

9

10 **Corresponding author:** Hanqin Tian. Phone: 334-844-1059. Fax: 334-844-1084.

11 Email: tianhanq@gmail.com

12

13

14

## Abstract

As urban land cover and populations continue rapidly increasing across the globe, much concern has been raised that urbanization may alter terrestrial carbon dynamics. Urbanization involves complex changes in land structure and multiple environmental factors. Relative contribution of these and their interactive effects need be quantified to better understand urbanization effects on regional carbon (C) dynamics as well as assess the effectiveness of C sequestration policies focusing on urban green space development. This study developed a comprehensive analysis framework for the factors (and their interactions) that control urbanization effect on ecosystem C dynamics, and proposed a factorial analysis scheme to analyze their relative contributions. In total fifteen factors belonging to five categories were identified. Some of them, like the interactive effects between global changes and urban land conversion, have not been studied before. Guided by the factorial analysis scheme, 24 number experiments were designed to systematically isolate and quantify the relative contribution of individual factors. A case study was conducted in the southern United States (SUS) to test this newly developed factorial analysis scheme. We found the impact of land conversion was far larger than the other factors. The urban management and the interactive effects among the multiple factors were also important. The mechanisms underlying the complex interactive effects were analyzed. Our findings provide valuable information for regional C management in the urbanised areas of SUS: (1) it is important to preserve pre-urban C pools rather than to rely on the C sink in urban ecosystem to compensate for the lost C during land

37 conversion. (2) In forested area, it is recommendable to improve landscape design  
38 (by arrange green spaces close to the city center) to maximize the urbanization-  
39 induced effect on C sequestration; while in the arid shrubland regions, urban  
40 managements will be a more effective way for C sequestration. (3) Lawn  
41 managements could create strong C sink even when the fossil fuel C cost are taken  
42 into account. (4) Protecting urban forests from disturbances such as logging and  
43 wildfire could be an effective way to enhance urban C sink. In general, the proposed  
44 factorial analysis scheme provides a useful tool for quantifying the complex  
45 mechanisms controlling C dynamics, and defining best development practices in  
46 urbanised areas.

47

48 **Keywords:** Urbanization, Urban ecosystems, Land use change, Carbon cycle, Global  
49 Change, Dynamic Land Ecosystem Model (DLEM)

50

## 1. Introduction

Urbanization, the aggregation of population in cities and transformation of rural areas into urban/developed land-use, became a dominant demographic trend and important land transformation process in recent decades (Seto *et al.*, 2010; Pickett *et al.*, 2011). At present about 3-5% of global land area has been converted to urban and developed land-use (hereafter refer to as urban) (Svirejeva-Hopkins & Schellnhuber, 2008; Seto *et al.*, 2010), 13-17% of which were intensively developed (Schneider *et al.*, 2010). Urban areas in the United States increased about 130% between 1960 and 2000 (www.census.gov, last accessed in Jul. 2012). Global urban areas could increase by about one million km<sup>2</sup> over the next 25 years (McDonald, 2008). The spatial prominence of urban areas and fast urban land conversion rate is reason enough to study its environmental impacts (Zipperer and Pickett 2012). A major finding of urban ecological research in the past decade is that urban ecosystems play an important role in both local and regional biogeochemical cycles (Imhoff *et al.*, 2000; Pataki *et al.*, 2003; Grimm *et al.*, 2008), esp. urban ecosystems account for a significant portion of terrestrial carbon (C) storage (Nowak & Crane, 2002; Pataki *et al.*, 2006; Pouyat *et al.*, 2006; Churkina *et al.*, 2010; Davies *et al.*, 2011; Hutyra *et al.*, 2011; Edmondson *et al.*, 2012). Zhang *et al.* (2012) estimated that urban and developed land accounts for about 6.7–7.6% of total ecosystem C storage within the Southern United States (SUS), larger than the pool size of shrubland. The potential for C sequestration in urban vegetation (McPherson *et al.*, 1997) and soil (Pouyat *et al.*, 2008) has drawn attention from both ecologists and decision makers (Poudyal *et al.*,

2010). Municipal interest in climate change mitigation through C offset trading has increased as many cities have established substantial programs, such as tree planting, to increase ecological services of urban ecosystems (Nowak, 2006; Tratalos *et al.*, 2007; Young, 2010). A management strategy for urban and peri-urban land, as suggested by the Intergovernmental Panel on Climate Change (IPCC, 2001), including tree planting, improved waste management and wood production could lead to a C sink of  $0.3 \text{ t C ha}^{-1} \text{ a}^{-1}$ . Escobedo *et al.* (2010) indicated that urban forest management can create moderate carbon sink in the southeastern US.

However, the ecological consequence of urbanization is highly complex (Pickett *et al.*, 2011), not only because of the strong spatial heterogeneity of urban ecosystems, which is composed by land cover types with distinct biogeochemical characteristics (Cannell *et al.*, 1999; Alberti, 2005; Buyantuyev *et al.*, 2010), but also because urbanization usually results in significant changes in many interacting environmental factors that affects ecosystem C processes, such as land conversion from rural to urban land-use (Schaldach & Alcamo, 2007), shifts in disturbance and management regimes (Kissling *et al.*, 2009; Fissore *et al.*, 2012), and urban-induced climate and atmospheric changes (Koerner & Klopatek, 2002; Fenn *et al.*, 2003; Kuttler, 2011). Furthermore, the legacy effect of pre-urban land-use changes (Ramalho & Hobbs, 2012) and influences from global climate changes (McCarthy *et al.*, 2010) could also modify ecosystem's responses to the urbanization-induced environmental changes. Analyzing the impacts of these changes and their interactive effects will help in our understanding of how regional C cycles are affected by

urbanization, quantifying the impacts of various environmental stresses, and identifying the major factors that control C dynamics of developed areas. Such knowledge can be valuable for policy makers and managers to predict the long-term ecological consequences of urbanization, to elucidate where management efforts should focus, and to formulate meaningful guidelines and tailor strategies for urban C managements.

Despite its importance and complexity, urbanization is an often-missing component in global change studies (Kaye *et al.*, 2005; Pouyat *et al.*, 2006). There are several remote sensing analyses that addressed urbanization effect on net primary productivity (NPP) (Imhoff *et al.*, 2000; Milesi *et al.*, 2003; Lu *et al.*, 2010). With an empirical inventory approach, Cannell *et al.* (1999) roughly estimated the effects of urbanization on the C budget of the United Kingdom. Only a few modeling studies have analyzed the responses of regional C dynamics to the environmental changes induced by urbanization. Many studies suggested that urban land conversion could have strong negative impact on regional to global C storage (Schaldach & Alcamo, 2007; Svirejeva-Hopkins & Schellnhuber, 2008; Zhang *et al.*, 2008; Eigenbrod *et al.*, 2011). Trusilova and Churkina (2008) compared the impacts of different urban-induced environmental changes on the C cycle in Europe, and found strong C sequestration due to urbanization-induced atmospheric changes. Milesi *et al.* (2005) assessed effects of different management practices on the C storage of US urban lawn. Zhang *et al.* (2012) found that pre-urbanization vegetation type and time since land conversion were closely related to the extent of urbanization effects on C dynamics of

Southern US during the last six decades. Despite these efforts, a comprehensive study that investigates the dominant environmental changes and addresses their relative importance on regional C dynamics is still not available, although it has been repeatedly suggested that due to the complex interactions among multiple involving factors, the ecological consequences of urbanization could not be fully understood without a full set of controlling drivers and their interactions being addressed (Hutyra *et al.*, 2011; Pickett *et al.*, 2011; Ramalho & Hobbs, 2012).

In this study, we first comprehensively analyzed the factors that may control the urbanization effect on ecosystem C dynamic (Fig. 1), and proposed a numeric experimental scheme, i.e. scenarios design, to conduct factorial analysis on the effects of different factors in section 2. Then as a case study of the newly developed analysis scheme, the dynamic land ecosystem model (DLEM, Tian *et al.* 2011a) was applied to quantify the urbanization effect on the C dynamics of the SUS from 1945-2007, and to analyze the relative contributions from each environmental factors and their interactive effects (Zhang *et al.*, 2012). SUS was selected as the study area because it was identified as the region with the most rapid urbanization in the US, where about one-third of the developed area has been added in the last 15 years of the 20th century (Alig *et al.*, 2004). Our study only considered the C dynamics of ecosystem (i.e. vegetation and soil) in urban. Fossil fuel emissions unrelated to urban managements were out of the scope of this study (Townsend-Small & Czimczik, 2010; Bartlett & James, 2011).

## 2. Factors controlling urbanization effects

To study the effects of urbanization on regional C balance, Zhang *et al.* (2012) compared the model simulation results of the urbanization scenario (or the “business as usual scenario”) against the results from a non-urbanization scenario, in which urbanization process was controlled and all lands remained in pre-urban land types. They found the urbanization from 1945-2007 resulted in a regional C loss of 0.21 Pg C in the SUS. The study, like others (McCarthy *et al.*, 2010), also indicated that urbanization is not a simple C release process, but involves complex changes in land structure and multiple environment factors, whose effects should not be treated independently. Whenever an ecosystem component is modified by one environmental stress, the ecosystem’s responses to other factors could also be altered due to the non-linear interactions among the coupled ecosystem components and processes (Wu, 1999). For example, elevated CO<sub>2</sub> in urban areas could be particularly important in relieving water stress induced by urban heat island effect (Groffman *et al.*, 2006). Therefore, it is important to consider all the major environmental factors and their interactive effects on C processes when studying the urbanization effects on regional C balance.

In Fig. 1, we generalize the factors that may control the urbanization effects (UBNZ; descriptions for the abbreviations are found in Fig. 1): (1) Urban vegetation is intensively managed. Irrigation, fertilization, and weed/disease controls improve lawn productivity (Milesi *et al.*, 2005). Remnant ecosystems in urban areas are generally protected from intensive disturbances such as agricultural soil tillage, wild



160 fire, and commercial logging (Raciti *et al.*, 2011). All these urban managements  
161 (UBMG) could result in high C density in urban ecosystems as observed in former  
162 studies (Nowak & Crane, 2002; Hutyra *et al.*, 2011; Edmondson *et al.*, 2012). (2)  
163 Urbanization-induced environmental changes (UIEC), such as urban heat island  
164 (UHI), elevated CO<sub>2</sub> (UCO<sub>2</sub>) and N deposition (UNDP), and reduced solar radiation  
165 due to aerosol pollutions (UDIM) could affect plant growth, succession and soil  
166 respiration in urban (Lovett *et al.* 2000, Awal *et al.* 2010, Zipperer 2011). According  
167 to Shen *et al.* (2008), the interactive effects among these UIEC factors (IT\_UIEC)  
168 should not be ignored. (3) Urban land conversion (UBNC) alters the landscape  
169 structure, where pre-urban land-covers are replaced by impervious surfaces and  
170 artificial green spaces such as urban lawns. During the process, vegetation biomass is  
171 removed, soils are disturbed, and large amount of C are released from the ecosystem  
172 (Schaldach & Alcamo, 2007; Zhang *et al.*, 2008). (4) Global changes (GLBC) in  
173 climate (CLM), land use (LUC), and atmosphere (e.g., CO<sub>2</sub>, N deposition (NDP), and  
174 O<sub>3</sub>) have different effects on different vegetation/land-cover types. In this study, LUC  
175 only refers to pre-urban land-use change. Because UBNC alters the vegetation/land-  
176 cover type, it also indirectly affects ecosystem's responses to global changes. For  
177 example, the legacy effects of pre-urban land-use history could explain the spatial  
178 pattern and temporal dynamic of ecosystem C pools in urban and developed areas  
179 (Golubiewski, 2006; Jenerette *et al.*, 2006). Therefore, the interactive effects between  
180 GLBC and UBNC (GLBC-UBNC) should not be ignored when investigating  
181 urbanization effects. Furthermore, the interactive effects among global changes

182 (IT\_GLBC) could have important ecological impacts (McMurtrie *et al.*, 2008). (5)

183 Finally, the interactive effects among the above four major type of urban controls

184 (IT\_OTHER in Fig. 1) should not be overlooked (Wu, 1999).

185 Numeric experiments and factorial analyses can be conducted to quantify the

186 effects of each of the above factors on carbon balance. For this purpose, a model

187 scenario scheme is presented in Table 1. Based on these scenario outputs, factorial

188 analyses can be conducted to isolate the effect of individual factor and their

189 interactive effects. According to Fig. 1, we have

$$190 \quad \text{UBNZ} = \text{UBNC} + \text{UBMG} + \text{UIEC} + \text{GLB-UBNC} + \text{IT\_OTHER} = S_{\text{UBNZ}} - S_{\text{GLBC}}$$

$$191 \quad \Rightarrow \text{IT\_OTHER} = (S_{\text{UBNZ}} - S_{\text{GLBC}}) - (\text{UBNC} + \text{UBMG} + \text{UIEC} + \text{GLB-UBNC}) \quad (1)$$

192 Where  $S_{\text{UBNZ}}$  is the urbanization scenario (or the “business as usual scenario”),

193 and  $S_{\text{GLBC}}$  is the control scenario, in which no urbanization takes place (Table 1). The

194 difference indicates the overall urbanization effect on C balance (Zhang et al., 2012).

195 UBNC is estimated with the  $S_{\text{UBNC}}$  scenario, in which only urban land conversion

196 occurs.

$$197 \quad \text{UBMG} = \text{LWN} + \text{UFM}$$

$$198 \quad \text{LWN} = S_{\text{LWN\&UBNC}} - S_{\text{UBNC}} \quad (2)$$

$$199 \quad \text{UFM} = S_{\text{UFM\&UBNC}} - S_{\text{UBNC}}$$

200  $S_{\text{LWN\&UBNC}}$  and  $S_{\text{UFM\&UBNC}}$  simulate the C balance in managed grass (lawn)

201 and urban forests in (converted) urban areas, respectively. It should be noted that it is

impossible to simulate urban land management without also simulate the urban land conversion. Their results are compared against the UBNC to isolate the effects of lawn (LWN) and urban forest (UFM) management.

$$UIEC = UHI + UCO2 + UNDP + UDIM + IT\_UIEC = S_{UIEC\&UBNC} - S_{UBNC}$$

$$\Rightarrow IT\_UIEC = (S_{UIEC\&UBNC} - S_{UBNC}) - (UHI + UCO2 + UNDP + UDIM) \quad (3)$$

$S_{UIEC\&UBNC}$  simulates the combination effects of multiple urban induced environmental changes and urban land conversion. We cannot simulate urban induced environmental changes without also simulating urban land conversion (land use change). Therefore, the effects of UHI, UCO2, UNDP, and UDIM are calculated similarly to the LWN and UFM:

$$\left. \begin{aligned} UHI &= S_{UHI\&UBNC} - S_{UBNC} \\ UCO2 &= S_{UCO2\&UBNC} - S_{UBNC} \\ UNDP &= S_{UNDP\&UBNC} - S_{UBNC} \\ UDIM &= S_{UDIM\&UBNC} - S_{UBNC} \end{aligned} \right\} \quad (4)$$

Finally, the interactive effects between global changes and urban land conversion can be derived as:

$$GLB-UBNC = LUC-UBNC + NDP-UBNC + O3-UBNC + CO2-UBNC + CLM-UBNC + IT\_GLBC = S_{GLBC\&UBNC} - (S_{GLBC} + S_{UBNC})$$

$$\Rightarrow IT\_GLBC = S_{GLBC\&UBNC} - (S_{GLBC} + S_{UBNC}) - (LUC-UBNC + NDP-UBNC + O3-UBNC + CO2-UBNC + CLM-UBNC) \quad (5)$$

Where  $S_{GLBC}$  simulates the global change effects; and  $S_{GLBC\&UBNC}$  simulates the combined effects of global change and urban land conversion. The difference between the result from the combined scenario and the sum of the GLBC and UBNC scenarios, i.e.  $S_{GLBC}$  and  $S_{UBNC}$ , shows the interactive effects between the two factors. Similarly,

$$LUC-UBNC = S_{LUC\&UBNC} - (S_{LUC} + S_{UBNC}) \quad (6)$$

$$NDP-UBNC = S_{NDP\&UBNC} - (S_{NDP} + S_{UBNC}) \quad (7)$$

$$O3-UBNC = S_{O3\&UBNC} - (S_{O3} + S_{UBNC}) \quad (8)$$

$$CO2-UBNC = S_{CO2\&UBNC} - (S_{CO2} + S_{UBNC}) \quad (9)$$

$$CLM-UBNC = S_{CLM\&UBNC} - (S_{CLM} + S_{UBNC}) \quad (10)$$

Detailed information about scenario design can be found in Table 1. Based on the work reported by Zhang et al. (2012), we conducted two additional scenarios to simulate urban C storage under extreme conditions:  $S_{CMAX}$ ,  $S_{CMIN}$ , to assess the uncertainties related to model parameters. For  $S_{CMAX}$ , parameters were selected to maximize C sequestration capacity of urban ecosystem, while for  $S_{CMIN}$ , parameters were selected to provide a conservative estimation.

### 3. Materials and methods in the case study

The DLEM is a process-based model that integrates the biophysical, biogeochemical, and hydrological processes to simulate impacts of environmental changes on water, C, and N cycles (see Fig. S1a in the supplementary material). The model has been parameterized and validated against intensively studied natural sites, and has been applied in multiple regional C dynamic studies (Tian *et al.*, 2011ab, 2012). Zhang *et al.* (2012) have developed an urbanization module for DLEM to assess the impacts of urbanization on long-term C dynamics in the SUS. Their study, however, only focused on the overall effects of urbanization without investigating the relative contribution from individual factor. In this current study, by conducting factorial analysis, we examined the relative contribution of different environmental controls and their interactive effects on regional C dynamics during urbanization. Here, we briefly introduce the study area, model structure, and the development of model inputs including the background global-change datasets (Supplementary Table S1) as well as the parameters for human-induced changes in urban. More detailed descriptions are found in Zhang *et al.* (2012).

#### 3.1. Study area

Because an ecological understanding of urban effect must include the suburban areas and settled villages as well as city cores (Pickett *et al.*, 2011), the “urban” areas refer to all the urban and developed lands in the SUS in this study. This study focuses on the  $1.2 \times 10^5 \text{ km}^2$  urban lands in the SUS (red areas in Fig. S2 in the supplementary material). Following Zhang *et al.* (2012), this study focuses on the

impacts of urbanization from 1945-2007 on regional net carbon exchange (NCE). NCE quantifies the C balance (with positive value indicating C sequestration) of the ecosystems in response to environmental change in a certain period (Tian et al., 2003, 2011).

### 3.2. Model description

Urban landscapes are composed by two major land functional types – urban impervious surface (UIS), and urban vegetation. Stearns (1971) identified three urban vegetation types - ruderal, residual, and managed. For simplification, ruderal and residual are merged into the dominant/potential local vegetation type in urban (UVG), and the managed vegetation is represented by urban lawn (ULW), an important characteristic of urban land-use conversion with respect to the C cycle (Kaye *et al.*, 2005; Golubiewski, 2006). Therefore, an urban landscape is treated as a mosaic of UIS, UVG, and ULW in DLEM. The development of UIS and ULW land typically includes the clearing of existing vegetation, massive movements of soil. DLEM not only models the disturbances on vegetation and soil during land clearing, but also tracks the fate of removed biomasses following the study of Houghton (1999) and Nowak and Crane (2002). Converting agricultural land to UVG will result in cropland abandonment and regeneration of potential vegetation (Dwyer et al. 2000). Otherwise, UVG conversion will not directly disturb pre-urban ecosystem. The disturbance regimes in UVG land, however, change after urbanization in DLEM. Urban forest and other residual ecosystems are protected from wildfire and commercial logging (Campbell *et al.*, 2007; Defosse *et al.*, 2011), disturbances that are responsible for the

low biomass density in the SUS forest (Birdsey 1992). Taking the disturbances' effect into account, the overall turnover rate of rural forest (Tian et al., 2012) is about 10% higher than that of urban forest, whose annual mortality was set to 2.2% to 3.5% according to Nowak (1986, 1994).

### 3.3. Model inputs

In model simulation, the background climate, atmosphere, and land use drivers were modified by urbanization-induced environmental changes, the values of which were estimated based on literature reviews (Table 2). The background environmental drivers provide global change information because they are transient datasets that changed annually or daily from 1945 to 2007. To control a certain global change driver, we fixed its value to the year 1945. For example, in the climate only scenario ( $S_{CLM}$  in Table 1), only climate data changed from 1945-2007, the values of other drivers ( $CO_2$ , N deposition,  $O_3$ , and land use) were fixed to the value of 1945. If a certain urbanization-induced environmental change factor is considered in the simulation, the corresponding background value will be modified by its parameter from the Table 2.

#### 3.3.1. The background climate, atmosphere, and land use dataset

We reconstructed an 8 km resolution daily climate dataset of the entire SUS from 1895 to 2005 (Fig. 2a,b,c) by integrating the daily climate pattern of the North American Regional Reanalysis (NARR; 32 km resolution) dataset (Mesinger et al., 2006) into the monthly PRISM (Parameter-elevation Regressions on Independent Slopes Model; 4 km resolution; 1895-present) climate data (Daly et al., 2008;

[prism.oregonstate.edu/](http://prism.oregonstate.edu/)). A detailed description of the method is found in Zhang (2008). PRISM is a knowledge-based system to interpolate climate elements under the assumption that for a localized region, elevation is the most important factor in the distribution of temperature and precipitation. To make predictions, PRISM dynamically calculates a linear climate–elevation relationship for each DEM grid cell using a moving-window, a procedure that smooth out signals of urbanization-induced climate changes (Daly et al., 2008). Like most reanalysis data, the surface temperatures of NARR were estimated from the atmospheric values by regional climate modeling, thus were not sensitive to changes in land surface (Kanamitsu et al., 2002). Therefore, the reanalysis datasets were used to provide information of background climate change in former UHI studies (Si *et al.*, 2012). The climate change data from 1895-2005 reconstructed based on the PRISM and NARR datasets were used to address the background global climate change (Fig. 1; GLBC-CLM) in this study.

Ozone AOT40 data (Fig. 2d) were retrieved from a global dataset developed by Felzer et al. (2005). EDGAR-HYDE 1.3 nitrogen emission data (Van Aardenne et al., 2001) were used to interpolate three maps from Dentener (2006) to generate a time-varying annual nitrogen deposition dataset (Fig. 2e). Both data sources had coarse spatial resolutions ( $0.5^{\circ}$ - $1^{\circ}$ ), and were downscaled to  $8 \times 8 \text{ km}^2$  using bilinear interpolation. Due to their coarse resolutions and because the atmospheric models that were used to generate these datasets did not consider the local urbanization effects (Felzer et al., 2005; Dentener, 2006), the AOT40 and nitrogen deposition datasets



represented the background global atmospheric changes in this study. The background global annual CO<sub>2</sub> concentration was obtained from the National Oceanic and Atmospheric Administration (NOAA) ([www.esrl.noaa.gov](http://www.esrl.noaa.gov)).

To simulate the land-use changes, DLEM requires annual urban and cropland maps (1 represents urban or croplands; 0 for natural vegetation). Distribution maps for cropland and urban/developed lands from 1895 to 2007 (Fig. 2f) were reconstructed by combining the contemporary land-use map that was derived from NLCD2001 (Homer et al., 2007) with historical census dataset for cropland, urban, and population (Waisanen and Bliss, 2002). A detailed description can be found in Zhang et al. (2012).

Sources of other inputs including the base maps (potential vegetation, soil properties, and topographic characteristics, etc.) and cropland management (irrigation and fertilization) datasets can be found in the Supplementary Table S1. Detailed description of the data development methodologies are found in Zhang (2008).

### **3.3.2. Urban-induced environmental changes**

DLEM further models the effect of urban-induced environmental changes (i.e. UHI, aerosol pollutions, and increased CO<sub>2</sub> and N deposition) on urban ecosystem, which (except for the aerosol pollutions) generally enhance the growth and biomass accumulation rate of urban vegetation (Ziska *et al.*, 2004). Based on literature review, we estimated the parameters controlling urban-induced environmental changes (Table 2). To evaluate the effects of parameterization uncertainties on the model simulations,

we designed two additional scenarios to simulate urban C storage under extreme conditions: UBNZ\_Cmin and UBNZ\_Cmax (Table 1). Parameters of UBNZ\_Cmin were set so that carbon sequestration were minimized while carbon loss was maximized during urbanization; UBNZ\_Cmax was the contrary (Table 2).

#### (1) The urban heat island effect

DLEM estimates the elevated temperature in urban (i.e. UHI, unit: °C) with the regression model developed by Karl et al. (1988):

$$UHI = \alpha \times (p)^{0.45} \quad (11)$$

where  $\alpha$  is a regression coefficient that varies with seasons and size of urban population ( $p$ ). Based on the climate records from 1219 stations in the US, Karl et al. (1988) determined the values of  $\alpha$  for the maximum, minimum, and average temperature for each season in three different urban sizes ( $p < 10000$ ;  $p \in [10000, 100000]$ ;  $p > 10000$ ). To develop 8-km resolution urban population dataset from 1945-2007, county-level urban population data developed by Goldewijk (2005) was divided by the area of urban/developed land to calculate the mean urban population density of each county. Then, the urban population map for each year was developed by multiplying the area of each urban region/patch in the NLCD 2001 land-use map with the urban population density of the local county.

## (2) The elevated atmospheric CO<sub>2</sub> concentration in urban/developed lands

Rural-urban CO<sub>2</sub> gradient is highly variable depending on time, location, wind direction, and distance from traffic, etc. (Idso et al., 1998, 2001, 2002; Vogt et al. 2006). The reported daily urban-rural CO<sub>2</sub> gradient ranged from 5 ppmv (Berry and Colls, 1990) to 66 ppmv (George et al., 2007). However, the daytime CO<sub>2</sub> gradient that determines the CO<sub>2</sub> fertilization effect on urban ecosystem is usually much smaller than the daily average due to solar-induced convective mixing (Idso et al., 2002) and the C uptake by plants (Kordowski & Kuttler, 2010). Day et al. (2002) reported that the daytime CO<sub>2</sub> concentration of the vegetated area in the center of Phoenix, AZ was only 8 ppmv higher than the background value. According to the measurements by Clark-Thorne and Yapp (2003), the daytime CO<sub>2</sub> concentration of urban interior was averagely 5-7 ppmv higher than the rural CO<sub>2</sub> in Dallas, TX. Angeles Garcia *et al.* (2012) found the daytime suburban CO<sub>2</sub> concentrations were 6-16 ppmv higher than rural levels in Northern Spain Based on these and other reports (Berry and Colls, 1990; (Li *et al.*, 2010; Rice & Bostrom, 2011), we assumed that the atmospheric CO<sub>2</sub> concentration of an urban vegetated area is 10 ppmv higher than the background value.

## (3) Urbanization-induced air pollution

Urban atmospheres have higher concentrations of nitrogen and aerosols than the rural region (Lovett et al., 2000; Azimi et al., 2005). In general, urban boundary layer pollutants are believed to reduce solar irradiance by 0-10% in North American Cities

(Oke, 1979, 1982; Peterson and Stoffel, 1980; Estournel et al., 1983). DLEM assumed that aerosol pollutions reduce the urban solar radiation by 5%. Urban air pollution generates high concentrations of both ozone precursors and ozone scavengers. Gregg et al. (2003) found the detrimental effects of tropospheric ozone were lower in urban than in suburban. Due to the uncertainties in urban ozone (Trusilova and Churkina, 2008), we did not consider the urbanization-induced ozone change in this study. Like atmospheric CO<sub>2</sub>, the temporal and spatial patterns of urban nitrogen deposition are highly variable. Previous studies indicate that the daytime atmospheric NO<sub>2</sub> concentration of urban and developed land usually ranges from 0.03 to 0.06 ppmv, an order higher than the value measured in rural ecosystem (Hanson et al. 1989). In this study, we used the mean value of 0.045 ppmv as the elevated urban NO<sub>2</sub> concentration.

It should be noted that these UIEC were not static but changed through time. In this study, we assumed that the rise of CO<sub>2</sub> and air pollutants in urban areas were positive correlated to the historical per capita fossil fuel emissions:

$$PL_i = PL_{mean} \times f\_EMS_i \quad (12)$$

where  $PL_i$  is the urbanization-induced atmospheric change (i.e., elevated NO<sub>2</sub>, CO<sub>2</sub>, and aerosol) in year  $i$ ;  $PL_{mean}$  refers to the mean urbanization-induced air pollution according to recent (since 1980) studies (parameters in Table 2);  $f\_EMS_i$  is the normalized fossil fuel emission factor for year  $i$ :

$$f\_EMS_i = \frac{EMS_i}{EMS_{1980\_2000}} \quad (13)$$

where  $EMS_i$  is the per capita annual fossil fuel emission of US in year  $i$ ;  
 $EMS_{1980\_2000}$  denotes the mean value between 1980 and 2000. To calculate the annual  
per capita fossil fuel emissions, we obtained the annual national fossil fuel emission  
data compiled by Marland et al. (2008) and the historical US population data from the  
US Census Bureau (<http://www.census.gov/population/www/popclockus.html>).

### 3.3.3. Urban managements

#### (1) Lawn managements

Urban lawns are irrigated, fertilized, and clipped. In DLEM, urban lawns are  
irrigated whenever the soil water content is lower than 50% of the field capacity.  
Since many of the US lawns are irrigated excessively (Milesi et al., 2005), we may  
underestimated the water use in irrigation. This uncertainty in irrigated water did not  
significantly affect the predicted C dynamics in urban lawn.

Based on the values provided by several reports in the literature, e.g., 8 - 9 g N  
 $m^{-2} yr^{-1}$  (Rockwell, 1929), 5 - 10 g N  $m^{-2} yr^{-1}$  (Thompson, 1961), 10 g N  $m^{-2} yr^{-1}$   
(Qian et al., 2003), 2.4 - 15 g N  $m^{-2} yr^{-1}$  (Osmond and Hardy, 2004), 9 g N  $m^{-2} yr^{-1}$   
(Law et al., 2004), and 9.7 g N  $m^{-2} yr^{-1}$  (Zhou et al., 2008), DLEM assumes that 10 g  
N  $m^{-2} yr^{-1}$  will be the N fertilization rate for the professionally managed lawns. This  
value is close to the 10.9 g N  $m^{-2} yr^{-1}$  fertilization rate for the managed lawns in US as  
estimated by Zirkle et al. (2011). In reality, however, the rates of N fertilization to  
lawns varied significantly from household to household in the US (Augustin, 2007).  
The professionally managed lawn only account for half of the US lawn area (Grounds

Maintenance, 1996). Only half of the home lawns are fertilized in a given year (Augustin, 2007). Only 25% of the fertilized home lawns are professionally managed. The remaining 75% are managed by the home owners with fertilization rates ranging from 4 - 9 g N m<sup>-2</sup> yr<sup>-1</sup>. Combining all these information, we deduced that the annual N fertilization rates to US lawns vary from 6.4 g N m<sup>-2</sup> yr<sup>-1</sup> (when 4 g N m<sup>-2</sup> yr<sup>-1</sup> is applied by home owners) to 7.3 g N m<sup>-2</sup> yr<sup>-1</sup> (when 9 g N m<sup>-2</sup> yr<sup>-1</sup> is applied by home owners). In the simulation, DLEM used the average value of 6.8 g N m<sup>-2</sup> yr<sup>-1</sup> for the urban lawn.

Urban lawns are usually clipped every 0.5 to 2 weeks in the US (Milesi et al., 2005; Kaye et al., 2005). In this study, we assumed an averaged mowing cycle of 10 days in SUS. This estimation agrees with a 900-person survey in Illinois, which reported an average mowing rate of 30 per year (Zirkle et al., 2011). Following Milesi et al. (2005), a lawn will only be mowed if its leaf area index exceeds the threshold value of 1.5. After mowing, 20% of the vegetation biomass will be removed. The belowground biomass will enter the soil litter pool, while the aboveground portion will enter the product pool and decay in one year. All clipped biomass will enter the product pool and decay in one year.

## (2) Mortality and management of urban forest

We assumed that urban trees were protected from commercial logging, and thus could grow very old. Large uncertainties exist in the mortality rate of urban trees. Field measurements revealed that street trees could have various mortality rates

450 depending on their size - 2.1% to 3.0% for trees whose DBH < 77 cm; and 5.4% for  
451 larger trees (Nowak, 1986). Nowak (1994) assumed an annual mortality rate of 2.6%  
452 in their urban forest modeling study. In DLEM, the annual background mortality rate  
453 of urban trees ranged from 2.2% to 3.5%, positively correlated with tree size (Nowak,  
454 1986). Following Sitch et al. (2003), the background mortality is modified by light  
455 competition at the stand level. In DLEM, forest die back will take place to maintain  
456 the foliage-projected coverage under 95%. Urban forests have relatively open canopy  
457 compared to rural forests, providing it an advantage to suppress light competition and  
458 support bigger trees. DLEM calculates the foliage-projected coverage of urban forest  
459 based on the total land area of the urban to simulate the open canopy effect.

460       Like lawns, urban forests may be managed, such as pruning and litter raking. It  
461 was found that intensive pruning might reduce the biomass of urban trees by as much  
462 as 25% (Nowak, 1994; Nowak et al., 2002; Escobedo et al., 2010). Unlike lawn,  
463 however, intensively managed trees, such as street trees that account for about 62% of  
464 the managed urban forest in the US (Kielbaso, 2008), only contribute to a small  
465 fraction (e.g., 2-4% in Oakland, CA and Chicago) of urban forests (Dwyer et al.,  
466 2000). Furthermore, a national survey revealed that more than 60% of US cities do  
467 not have urban forest management programs (Kielbaso, 2008). Even if all cities in the  
468 SUS have forest management program, and 10% of urban forest is street tree that  
469 accounts for 50% of the managed forest, managed tree will only account for 20% of  
470 urban forest. Under this assumption, about  $20\% \times 25\% = 5\%$  of the forest biomass  
471 was removed by pruning (Nowak, 1994; Nowak et al., 2002) (Table 2).

In some managed urban forests, a fraction of the litter (such as the litter from the pruned trees) will be removed and disposed of in a landfill. Nowak et al. (2002) assumed that only 3.7% of the removed carbon would be released during the first 5 years, and the remaining would be permanently locked up in a landfill. Accordingly, DLEM simulated the process of litter removal by allocating 1.85%, 1.85%, and 96.3% of the removed litter to 1-, 10-, 100-yr product pools that have turnover rates of 1 year, 10 years, and 100 years, respectively. No information about the patterns of litter management is currently available for the urban/developed land in the SUS. Since the fraction of intensively managed urban forest is quite low (Dwyer et al., 2000), we assumed that only 10% of litter will be removed and disposed in a landfill (Table 2).

#### **4. Case study results**

The temporal pattern of carbon dynamic during urbanization was controlled by the UBNC, which was estimated to result in about 0.37 Pg C loss from 1945-2007 (Fig. 3). In contrast, the urban managements (i.e. UBMG) and UIEC enhanced C storage by about 0.12 Pg and 0.03 Pg, respectively. Factorial analysis based on numeric experiments indicated the interactive effects between global changes and urban land conversion has negative effect on C storage, causing the study area to lose about 0.02 Pg C from 1945-2007. The complex interactive effects (i.e. IT\_OTHER) among the four major types of environmental changes, urban land conversion, urban managements, UIEC, and GLBC-UBNC, resulted in C sequestration of 0.04 Pg, comparable to the effects of UIEC and GLBC-UBNC.



The effects of UEIC, urban managements, and GLBC-UBNC can be further broken down to reflect the effect of individual factors (Fig. 1). From 1945-2007, urban lawn management (i.e. LWN) enhanced C storage by 489.9 g m<sup>-2</sup> (the SUS subgroup in Table 3) or 63.6 Tg in the SUS (Fig. 1), having the strongest C sequestration effect among all factors. Urban forest managements (i.e. UFM), including direct management (Table 2) and indirect effects from altered disturbance regimes (e.g., protection from commercial logging and wildfire), also resulted in large C sequestration of 396.3 g m<sup>-2</sup> or 51.5 Tg. Other factors that have significant positive effects on C sequestration included the increased N deposition (248.9 g m<sup>-2</sup> and 32.3 Tg in the SUS) and CO<sub>2</sub> (220.5 g m<sup>-2</sup> and 28.6 Tg in the SUS) in urban. In comparison, UHI and interactive effects among UIEC factors caused 15.6 Tg (120.3 g m<sup>-2</sup>) and 16.0 Tg (123.2 g m<sup>-2</sup>) C loss from the SUS, respectively. The interactive effect between UBNC and global change factors were smaller than other controls. While its interactions with global O<sub>3</sub> and climate change may enhance C sequestration, interactions between UBNC and other global changes (pre-urban land-use change, atmospheric CO<sub>2</sub> and N deposition change, and the interactive effects among the global change factors) caused C loss (Fig. 1).

Because the juxtaposition of land use and ecotypes strongly influences regional patterns of urban ecosystem functions (Nowak et al., 1996), we further analyze the impacts of urbanization on ecosystem C density based on the dominant/potential local vegetation type (i.e. UVG; Table 3). The results indicated that urbanization had strong negative effect on C density (-2084 g m<sup>-2</sup>) in forest area, only slight negative effect on

C density ( $-95 \text{ g m}^{-2}$ ) in grasslands, and positive effect on C density ( $390 \text{ g m}^{-2}$ ) in shrubland/desert (Table 3). The C sequestration effects of UIEC and forest managements were strongest in forest area, followed by grassland and shrubland/desert areas. The interactive effects between global change and urban land conversion had negative effect ( $-276 \text{ g m}^{-2}$ ) on C density in forest area and positive effect ( $168 \text{ g m}^{-2}$ ) on C density in grassland area. Because of the large forest area in the SUS and because of the relatively strong responses of forest C dynamics to land conversion and urban induced changes, forest area determined the pattern of regional C dynamics in response to urbanization from 1945-2007(Fig. 1; Table 3).

## 5. Discussion

### 5.1. Importance of the ability to quantify the relative contributions from multiple controls on urban C dynamics

Nowhere is ecological complexity more apparent than in urban areas (Kaye *et al.*, 2006). The aggregated effects of urbanization on land–atmosphere exchange processes remains highly uncertain despite decades of study on components of the problem (Pickett *et al.*, 2011). Only considering certain aspects of urbanization, former studies drew contradictory conclusions about urbanization effect. Those focusing on urban land conversion concluded that urbanization have negative effects on ecosystem productivity and C storage (e.g., Imphoff *et al.*, 2000; Schaldach and Alcamo, 2007; Zhang *et al.*, 2008). Those focusing on urbanization-induced environmental changes found positive effects on carbon sequestration from the

elevated CO<sub>2</sub>, N deposition, and prolonged growth season in urban (e.g., Trusilova and Churkina, 2008). Other studies emphasized on the effects of urban managements on promoting C sequestration in urban ecosystem (e.g., Qian et al., 2010; Milesi et al., 2005). Still others suggested that the interactive effects among different factors should not be overlooked (e.g., Shen et al., 2008). So, which factor is more important?

To answer this question, we first identified 15 factors that may affect ecosystem carbon dynamics during urbanization and organized them into five major categories under an urban analysis framework that shows the relationship among the factors (section 2, Fig. 1). Although it may not include all factors, the framework provides the most comprehensive analysis on the dominant factors so far. Among the five major categories in the Fig. 1, GLBC-UBNC (i.e., the interactive effect between global changes and urban land conversion) and IT\_OTHER (i.e., the overall interactive effects among urban land conversion, urban management, UIEC, and GLBC-UBNC) are newly identified controls that have never been addressed before. Our case study in the SUS showed the IT\_OTHER even had a larger impact on C dynamics than the UIEC. Furthermore, we proposed that the high carbon density of urban forest could be explained by a new mechanism – the low turnover rate of trees due to the suppressed disturbances (flooding, wild fire, pest, commercial logging, etc.) in urban (see the following discussion in the section 5.3). Considering their potentially important effects, field observations and experiments should be setup to evaluate the importance of these newly identified factors. Other factors, such as the urban land conversion, urban managements, and urbanization-induced environmental changes have been

individually analyzed in former studies (Milesi et al., 2005; Schaldach and Alcamo, 2007; Trusilova and Churkina, 2008), but their effects on urban ecosystem carbon dynamics have never been compared before this study.

Then, we designed 24 numeric experiments, based on the result of which a factorial analysis scheme was developed to systematically isolate and quantify the relative contribution of individual factors (section 2; Table 1). This is an important contribution to urban and global change researches. Global urban area is large and increasing rapidly. Urbanization effects on C cycle and climate have become the focus of global change research (Grimm et al., 2008). Reforestation projects have been initiated in many cities for C sequestration and climate regulation (Young, 2010). Of particular importance to climate-change policy and carbon management is the ability to quantify the relative contributions of multiple environmental factors to net carbon source and sink behavior (Heimann and Reichstein, 2008). Our study, for the first time, provides the ability to quantify the relative contribution of the factors that controls C dynamics during urbanization. Guided by this factorial analysis scheme, our case study in the SUS found that the urban land conversion, urban managements, and overall interactive effects among major factors (IT\_OTHER in Fig 1) were the first, second, and third most important controls on the ecosystem carbon dynamics from 1945-2007 (Fig 1). The impact of land conversion was far larger than the other factors. Beside its theoretical merit, this finding is also important for carbon management in urban. It not only shows the big potential of carbon sequestration by improving urban managements, but also highlights the large uncertainties related to

the complex interactive effects among multiple environmental changes (see section 5.2 in the Discussion). Although the findings in our case study only reflect the urban carbon dynamics in the SUS from 1945-2007, our framework of urban environmental controls and the factorial analysis scheme can be applied in other regions.

## **5.2. Complex interactive effects among factors**

One of the major uncertainties in urban C dynamic is from the interactive effects among environmental factors, which could have strong ecological impacts, sometimes even determined the direction of the overall ecosystem C balance (Shen et al., 2008; Tian et al., 2011b). However, many former urban studies overlooked the interactive effects and assumed the effects of multiple factors to be additive (e.g., Zirkle et al., 2011). Our case study found the overall interactive effects of the major control factors could increase C sequestration in the SUS by about 39.9 Tg, larger than the effect of urbanization-induced environmental changes (29 Tg) (Fig. 1). This C sink mainly located in the forested areas, which in average gained  $411 \text{ gC m}^{-2}$  due to the overall interactive effects of urbanization from 1945-2007 (Table 3). Compared to the pre-urban forests, urban trees in general had higher biomass and productivity, because they were protected (by human managements) from disturbances (such as commercial logging) that caused the high turnover rates and low biomass of the rural forest in SUS (Birdsey 1992). Our simulations shows that these larger trees are more responsive to urbanization-induced environmental changes and can fix more C, a phenomenon confirmed by recent observations from Escobedo et al. (2010) and Stephenson et al. (2014). The underlying mechanism is related to the relatively large

total leaf area of big trees. According to the Pipe model (Shinozaki et al., 1964) that controls Photosynthate allocation in woody plant, total tree leaf mass increases as the square of trunk diameter. A typical tree that experiences a tenfold increase in diameter will therefore undergo a roughly 100-fold increase in total leaf mass. Larger leaf mass means the tree has higher growth potential if not limited by water and nutrient availability. Therefore, bigger trees are more sensitive to elevated CO<sub>2</sub> and N deposition in urban. In rural forest stand, the high C sequestration rate of large, old trees could be offset by intensified mortality related to light competition. The urban forest, however, has relatively open canopy, and are able to support large trees (see section 3.3.3). Therefore, when a rural forest became a remnant forest in urban, its trees could grow bigger, faster, and were more sensitive to the increased urban CO<sub>2</sub> and N deposition because of the urban management effect that suppressed disturbances (commercial logging) and light competition.

We also found strong interactive effect (-16 Tg) among the UIEC factors (UHI, CO<sub>2</sub> dome, and elevated N deposition), comparable to the negative effects of UHI (-15.6 Tg) (Fig. 1). Unlike Trusilova and Churkina (2008), who found the UIEC interactive effect increased C sequestration in Europe, we found it suppress the urban C sink in the SUS (Fig. 1). This is mainly because the two regions experienced different urbanization-induced climate changes. In our simulation, urbanization will increase local surface temperature in the SUS, but Trusilova and Churkina (2008)'s data indicated significant reduction of temperature by 0.73-1.26 °C followed the urbanization in Europe. We found the UHI effect increased potential

evapotranspiration and exacerbated the water stress in the warm temperate ecosystems of the urban areas in the SUS. Like Shen et al. (2008), our simulation indicated that increase water stress suppressed elevated CO<sub>2</sub> and N deposition effects on ecosystem C sequestration. Trusilova and Churkina (2008)'s data, in contrast, indicated reduced temperature and increased precipitation in urbanised areas in Europe, both climate changes improved water availability and magnified elevated CO<sub>2</sub> and N deposition effects.

Shen *et al.* (2008) suggested that the effect of urbanization-induced changes are difficult to predict due to the influenced from other factors such as global climate change. Guided by the factorial analysis scheme developed in this study, for the first time, we found a way to separate the global change effects (i.e. GLBC) from the urban land conversion (i.e. UBNC) and quantify their interactions (i.e. GLBC-UBNC) (Fig. 1). We found GLBC-UBNC had negative effects on regional C storage (-24 Tg), almost offset the C sink induced by UEIC (29 Tg) (Fig. 1). Such an important mechanism, however, had been overlooked in former studies. The interaction between UBNC and different global change factors had different effects on C dynamics. In general, GLBC-UBNC would have negative impact on C storage if the global change factor enhances ecosystem C sequestration. This is because the lands converted to impervious surface are no more responsive to global change. For example, elevated CO<sub>2</sub> and N deposition in atmosphere stimulate C sequestration. After a pre-urban ecosystem is converted to impervious surface, the related C sinks (in response to CO<sub>2</sub> fertilization) disappear. Therefore, the interactive effects between urban land

conversion and changes in global CO<sub>2</sub> and N deposition seem to have negative effect on C sequestration (Fig. 1).

### 5.3. Implications for urban management

Many cities and regional governments are taking significant steps to reduce and offset their carbon emission and increase ecological services of urban ecosystems (Nowak, 2006; Tratalos *et al.*, 2007; Young, 2010). Our findings provide valuable information for regional C management in the urbanised areas of SUS: First, we found the C loss caused by urban land conversion dominated the carbon sink induced by all other factors from 1945-2007 in the SUS (Fig. 1). This finding highlights the importance to preserve pre-urban C pools during land development, probably by reducing soil disturbances or reserving large areas of remnant green space, rather than relying on the carbon sink in urban ecosystems to compensate for the C loss during land conversion. Escobedo *et al.* (2010) suggested that preserving large trees had a larger C benefit than planting young trees in the urbanised areas of the SUS. This is especially important for the forested regions, which when converted to urban could release nine times more C than the shrubland (Table 3). Our analysis shows about 77% of urban and developed areas in the SUS were converted from forest, becoming a primary threat to the C sequestration in the forested area of the SUS (Wear, 2002).

Second, our study, as well as others (Ziska *et al.*, 2004; Trusilova & Churkina, 2008), indicated the urban-induced environmental changes possibly promote NPP and C sequestration in urban ecosystems. Because these UIEC factors generally have a “dome” pattern that peaks at the city center and gradually levels off along urban-rural



gradient (Idso et al., 1998), it is advisable to arrange green spaces close to the city center to maximize their C sequestration capacity. The distinct responses from different vegetation types to urbanization should also be taken into consideration. Shen et al. (2008) suggested grass to be more sensitive to urban CO<sub>2</sub> dome effect than desert shrub. We found the carbon sink effect of UEIC decreased, while the carbon sink effect of lawn managements increased, in the sequence of forest, grass, and shrub areas (Table 3). Therefore, in the forested areas, it is recommendable to improve landscape design (such as arrange green spaces close to the city center) to maximize the UIEC effect; while in the arid shrubland areas, the focus should be put on improving urban managements to enhance C sequestration.

Third, our study indicates managements could create strong C sinks in urban vegetation (115.1 Tg C from 1945-2007 in SUS, about 55% of which was induced by lawn managements). Grasslands receiving fertilizer produced 7% to 298% more dry biomass than unfertilized grassland (Zirkle et al., 2011). Qian *et al.* (2010) observed that irrigation increased SOC of turfgrass, which could sequester 32-78 g C m<sup>-2</sup> a<sup>-1</sup>. Zirkle et al. (2011) estimated that N fertilization and irrigate together could increase the SOC of US home lawn (or the so-called “Do It Yourself lawn”) by 78.5-79.5 gC m<sup>-2</sup> a<sup>-1</sup>. However, high CO<sub>2</sub> uptake in lawns is not without a “carbon cost” from fossil fuel CO<sub>2</sub> emitted during maintenance (Townsend-Small and Czimczik, 2010). In a literature review on US lawn managements, Zirkle et al. (2011) estimated the mean hidden carbon cost of N fertilizer, pesticide, and irrigation to be about 10.1-20.4 gC m<sup>-2</sup> a<sup>-1</sup>, 0.4-2.6 gC m<sup>-2</sup> a<sup>-1</sup>, and 0.1-0.3 gC m<sup>-2</sup> a<sup>-1</sup>, respectively for home lawn. In

690 total, lawn maintenance could result in about  $23.6\text{--}43.9 \text{ gC m}^{-2} \text{ a}^{-1}$  fossil fuel carbon  
691 emission, equaling to 30%-56% of the carbon sink ( $\sim 79 \text{ gC m}^{-2} \text{ a}^{-1}$ ) induced by N  
692 fertilization and irrigation (Zirkle et al., 2011). This translates to about 25.7-47.8 Tg  
693 hidden carbon cost in lawn maintenance in the US from 1945-2007. In another word,  
694 about 40%-75% of the 63.6 Tg C sink induced by lawn management was offsetting by  
695 the related hidden carbon cost. Previous also studies indicated that if carbon-based  
696 maintenance is performed, urban forest will eventually become a C source (Nowak et  
697 al., 2002), or weak sink (Escobedo et al., 2010). Unlike lawn, however, intensively  
698 managed trees only account for a small fraction of the urban forest in US. A national  
699 survey revealed that less than 40% of US cities have urban forest managements  
700 programs, and 62% of the managed urban forest is street tree (Kielbaso, 2008).  
701 Considering the fact that street tree usually contribute to a small very fraction (e.g., 2-  
702 4% in Oakland, CA and Chicago, IL) of urban forests (Dwyer et al., 2000), hidden  
703 carbon cost from urban tree maintenance should be relatively small at regional scale.  
704 Furthermore, there is substantial scope in reducing management-related  $\text{CO}_2$  emission,  
705 because different equipment and maintenance techniques may have distinct carbon  
706 emission rate (Reid et al., 2010). For example, walk-behind lawnmower produces far  
707 less carbon emission than the riding mowers. It was estimated that half of the  
708 lawnmowers used by US homeowner belong to riding mower (Quigley, 2001), which  
709 has far larger C emission than walk-behind mower (Zirkle et al., 2011). By improving  
710 the efficiency of riding mowers or choosing to use the walk-behind mower, the  
711 maintenance carbon emission of urban vegetation could be significantly reduced.

Another possibility is to collect and utilize the 164 Tg dry biomass of lawn clippings and pruned tree twigs/limbs produced annually in the managed urban ecosystem in US for bioenergy production (Springer, 2012). Finally, well-managed urban vegetation can also indirectly reduce the C emission with its shading and cooling effects (Akbari et al., 1992; Taha et al., 1996). These factors and processes will be considered and integrated in the DLEM in our future urban studies.

Fourth, we found that altered disturbance regime might explain the observed C sink in urban forest. Urban forest was reported to have higher C density and growth rate than the average rural forest (Nowak and Crane, 2002; Golubiewski, 2006). This phenomenon was attributed to urbanization-induced environmental changes (Pouyat et al., 2007) and reduced light competition in urban forest due to its open canopy (Nowak, 1994). We propose another possible mechanism – the altered disturbance regime after urbanization may enhance C sink in urban forest. Most city trees are protected from frequent tillage, wildfire, and commercial logging, leading to suppressed soil disturbances and increased tree age (lower mortality and higher biomass) (Hutyra et al. 2011). In comparison, the influence of more intensive management of plantations and natural forests resulted in a mosaic of different age classes and an averaged low biomass density (Birdsey 1992). The risk of C emission through catastrophic wildfire burning is also considerably reduced by shifted fire regime (Guilden et al. 1990; Pickett et al. 2011) or fire control/management in urban areas (Campbell et al. 2007, Defosse et al. 2011). Based on a literature review (see section 3.2 and 3.3.3), we estimated that the overall turnover rate (considering both

the mortality rate and the disturbance effect) of rural forest is about 10% higher than that of urban forest (assuming zero disturbance from fire or commercial logging in urban) in SUS. Our simulation showed the effect of management and altered disturbance regimes together resulted in a C sink of 51.5 Tg in the SUS urban forest. Because direct management such as pruning negatively affected C storage, the C sink can be attributed to the altered disturbance regime; a potentially important mechanism that should be further investigated in future study.

#### 5.4. Uncertainties

Modeled C balance is essentially a result of the interactions among model assumptions, empirical and mechanistic relationships, and model parameterization. Urban ecosystem modelling is especially difficult and bound to large uncertainties (Churkina, 2008). Urbanization has complex effects on local climate, atmosphere, and disturbance regimes. Urban environment conditions and land management vary from place to place and from time to time (Alberti, 2005). Because the ecophysical and socioeconomic mechanisms underlying these urban-induced environmental changes are largely unclear (Pouyat et al., 2007; Pickett et al., 2011), we have to rely on an empirical parameterization approach to address the multiple controls on urban C dynamics in this study. As described in section 3.3, based on extensive literature review and academic reasoning, we derived the model parameters to approximate the average urban-induced environmental changes in the study region (Table 2). To evaluate the effects of parameterization uncertainties on the simulation results, we designed two additional scenarios to simulate urban C storage under extreme

conditions: UBNZ\_Cmin and UBNZ\_Cmax (Table 1). The simulation results indicated that uncertainties related to parameterization of urban-induced environmental changes amounted to -2% to 3% of the urban-induced C dynamics (Fig. 3).

Previously, we validated the DLEM simulated C and water fluxes, nitrogen cycle, and soil processes and trace gas emission against intensively studied ecological research sites (Tian et al., 2011a,b). Because urbanization does not change genetic characteristics of plants or fundamental mechanisms of ecological processes (Niemela, 1999), former validation results indicated that DLEM can correctly simulate ecosystem's responses to multiple environmental stresses in urbanised areas. To evaluate DLEM performance for simulating C processes in urban ecosystems, we further compared model predictions with 16 field observations (including VEGC, SOC, and NPP) from 12 studies that were located in or close to the SUS (Table 4). For those studies with sample variance, all of our model predictions fall in the range of one standard error.

## **6. Conclusion**

Urbanization involves complex changes in land structure and multiple environment factors, whose effects should not be treated independently. As urban land cover and human population continue to increase across the globe rapidly, it is important to investigate the individual effects of and complex interactions among multiple factors on the ecosystem structure and processes in urbanised lands. Our case

777 study revealed how the C dynamics in the  $1.2 \times 10^5 \text{ km}^2$  urbanised areas of the SUS  
778 were influenced by multiple environmental factors from 1945-2007, but the numeric  
779 experimental design and the factorial analysis schemes proposed in this study could  
780 be applied in other regions. Such efforts as the one reported not only improves our  
781 understanding of the complex effects of urbanization on regional C dynamics, but also  
782 provides a quantitative approach for assessing the effectiveness of policies and  
783 defines best development practices.

784

## 785    **Acknowledgements**

786            This study has been supported by the NSFC program (#31170347), the “Hundred  
787   Talents Program” of the Chinese Academy of Sciences, US Department of Energy  
788   NICCR Program, NASA IDS Program, AAES and the Center for Forest  
789   Sustainability at Auburn University.

## References

- Akbari, H., S. Davis, S. Dorsano, J. Huang, and S. Winnett (editors). 1992. Cooling Our Communities: A Guidebook on Tree Planting and Light-Colored Surfacing, U. S. Environmental Protection Agency, Office of Policy Analysis, Climate Change Division.
- Alberti, M. 2005. The effects of urban patterns on ecosystem function. *International Regional Science Review* 28:168-192.
- Alig, R. J., J. D. Kline, and M. Lichtenstein. 2004. Urbanization on the US landscape: looking ahead in the 21st century. *Landscape and Urban Planning* 69:219-234.
- Awal, M. A., T. Ohta, K. Matsumoto, T. Toba, K. Daikoku, S. Hattori, T. Hiyama, and H. Park. 2010. Comparing the carbon sequestration capacity of temperate deciduous forests between urban and rural landscapes in central Japan. *Urban Forestry & Urban Greening* 9:261-270.
- Bartlett, M. D. and I. T. James. 2011. A model of greenhouse gas emissions from the management of turf on two golf courses. *Science of the Total Environment* 409:1357-1367.
- Berry, R.D., Colls, J.J., 1990. Atmospheric carbon dioxide and sulphur dioxide on an urban/rural transect-I. Continuous measurements at the transect ends. *Atmospheric Environment* 24A, 2681–2688.
- Bettencourt, L. and G. West. 2010. A unified theory of urban living. *Nature* 467:912-913.
- Birdsey, R. A.: Carbon storage for major forest types and regions in the conterminous United States, in: *Forests and Global Change, Volume Z: Forest Management*



812 Opportunities for Mitigating Carbon Emissions, edited by: Sampson, R.L. and Hair,  
813 D., American Forests, Washington, DC, 1–26, 1996.

814 Birdsey, R. A.: Carbon Storage and Accumulation in United States Forest Ecosystems,  
815 Gen. 20 Tech. Rep. WO-59, US Department of Agriculture, Forest Service,  
816 Washington Office, Washington, DC, 51 pp., 1992.

817 Buyantuyev, A., Wu, J., Gries, C., and Adger, W. N.: Multiscale analysis of the  
818 urbanization pattern of the Phoenix metropolitan landscape of USA: time, space and  
819 thematic resolution, *Landscape Urban Plan.*, 94, 206–217, 2010.

820 Campbell, J., Donato, D. C., Azuma, D., and Law, B.: Pyrogenic carbon emission from a  
821 large wildfire in Oregon, United States, *J. Geophys. Res.-Biogeo.*, 112, G04014,  
822 doi:10.1029/2007JG000451, 2007.

823 Cannell, M. G. R., R. Milne, K. J. Hargreaves, T. A. W. Brown, M. M. Cruickshank, R. I.  
824 Bradley, T. Spencer, D. Hope, M. F. Billett, W. N. Adger, and S. Subak. 1999.  
825 National inventories of terrestrial carbon sources and sinks: The UK experience.  
826 *Climatic Change* 42:505-530.

827 Chen, G., H. Tian, C. Zhang, M. Liu, W. Ren, W. Zhu, A. Chappelka, S. Prior and G.  
828 Lockaby. 2012. Drought in the Southern United States over the 20th century:  
829 Variability and its impacts on terrestrial ecosystem productivity and carbon storage.  
830 *Climatic Change* 114: 379-397.

831 Churkina, G., D. G. Brown, and G. Keoleian. 2010. Carbon stored in human settlements:  
832 the conterminous United States. *Global Change Biology* 16:135-143.

833 Daly, C., Halbleib, M., Smith, J.I., Gibson, W.P., Doggett, M.K., Taylor, G.H., Curtis, J.,  
834 and Pasteris, P.A. 2008. Physiographically-sensitive mapping of temperature and

835 precipitation across the conterminous United States. *International Journal of*  
836 *Climatology*, 28: 2031-2064.

837 Davies, Z. G., J. L. Edmondson, A. Heinemeyer, J. R. Leake, and K. J. Gaston. 2011.  
838 Mapping an urban ecosystem service: quantifying above-ground carbon storage at a  
839 city-wide scale. *Journal of Applied Ecology* 48:1125-1134.

840 Defosse, G. E., G. Loguercio, F. J. Oddi, J. C. Molina, and P. D. Kraus. 2011. Potential  
841 CO<sub>2</sub> emissions mitigation through forest prescribed burning: A case study in  
842 Patagonia, Argentina. *Forest Ecology and Management* 261:2243-2254.

843 Dentener, F.J. 2006. Global maps of atmospheric nitrogen deposition, 1860, 1993, and  
844 2050. Available online <http://daac.ornl.gov/> from Oak Ridge National Laboratory  
845 Distributed Active Archive Center, Oak Ridge, Tennessee, USA.

846 Dwyer, J.F.; Nowak, D.J.; Noble, M.H.; Sisinni, S.M. 2000. Connecting people with  
847 ecosystems in the 21st century: an assessment of our nation's urban forests. Gen.  
848 Tech. Rep. PNW-GTR-490. Portland, OR: U.S. Department of Agriculture, Forest  
849 Service, Pacific Northwest Research Station. 483 p.

850 Edmondson, J. L., Davies, Z. G., McHugh, N., Gaston, K. J., and Leake, J. R.: Organic  
851 carbon hidden in urban ecosystems, *Scientific Reports*, 2, 963,  
852 doi:10.1038/srep00963, 2012.

853 Eigenbrod, F., V. A. Bell, H. N. Davies, A. Heinemeyer, P. R. Armsworth, and K. J.  
854 Gaston. 2011. The impact of projected increases in urbanization on ecosystem  
855 services. *Proceedings of the Royal Society B-Biological Sciences* 278:3201-3208.

856 Escobedo, F., S. Varela, M. Zhao, J. E. Wagner, and W. Zipperer. 2010. Analyzing the  
857 efficacy of subtropical urban forests in offsetting carbon emissions from cities.  
858 *Environmental Science & Policy* 13:362-372

859 Felzer, B., Kicklighter, D.W., Melillo, J.M., Wang, C., Zhuang, Q., Prinn, R. 2004.  
860 Effects of ozone on net primary production and carbon sequestration in the  
861 Conterminous United States using a biogeochemistry model. *Tellus* 56B:230–48.

862 Fenn, M. E., R. Haeuber, G. S. Tonnesen, J. S. Baron, S. Grossman-Clarke, D. Hope, D.  
863 A. Jaffe, S. Copeland, L. Geiser, and H. M. Rueth. 2003. Nitrogen emissions,  
864 deposition, and monitoring in the western United States. *BioScience* 53:391-403.

865 Fissore, C., S. E. Hobbie, J. Y. King, J. P. McFadden, K. C. Nelson, and L. A. Baker.  
866 2012. The residential landscape: fluxes of elements and the role of household  
867 decisions. *Urban Ecosystems* 15:1-18.

868 Folke, C., A. Jansson, J. Larsson, and R. Costanza. 1997. Ecosystem appropriation by  
869 cities. *Ambio* 26:167-172.

870 George, K., L. H. Ziska, J. A. Bunce, and B. Quebedeaux. 2007. Elevated atmospheric  
871 CO<sub>2</sub> concentration and temperature across an urban-rural transect. *Atmospheric*  
872 *Environment* 41:7654-7665.

873 Goldewijk, K., 2005. Three Centuries of Global Population Growth: A Spatial  
874 Referenced Population (Density) Database for 1700 – 2000. *Population and*  
875 *Environment* 26 (5), 343-367.

876 Golubiewski, N. E. 2006. Urbanization increases grassland carbon pools: Effects of  
877 landscaping in Colorado's front range. *Ecological Applications* 16:555-571.

878 Grimm, N. B., S. H. Faeth, N. E. Golubiewski, C. L. Redman, J. Wu, X. Bai, and J. M.  
879 Briggs. 2008. Global change and the ecology of cities. *science* 319:756-760.

880 Groffman, P. M., R. V. Pouyat, M. L. Cadenasso, W. C. Zipperer, K. Szlavecz, I. D.  
881 Yesilonis, L. E. Band, and G. S. Brush. 2006. Land use context and natural soil  
882 controls on plant community composition and soil nitrogen and carbon dynamics in  
883 urban and rural forests. *Forest Ecology and Management* 236:177-192.

884 Guilden, J.M., Smith, J.R., Thompson, L., 1990. Stand structure of an old-growth upland  
885 hardwood forest in Overton Park, Memphis, Tennessee. In: Mitchell, R.S., Shevial,  
886 C.J., Leopold, D.J. (Eds.), *Ecosystem Management: Rare Species and Significant*  
887 *Habitats*. New York State Museum, Albany, pp. 61-66.

888 Hart JF. 1980. Land use change in a piedmont county. *Ann Assoc Am Geogr* 70:492–525.

889 Homer, C., Dewitz, J., Fry, J., Coan, M., Hossain, N., Larson, C., Herold, N., McKerrow,  
890 A., Van Driel., J.N., Wickham, J. 2007. Completion of the 2001 National Land  
891 Cover Database for the conterminous United States. *Photogrammetric Engineering*  
892 *and Remote Sensing*, 73: 337-341.

893 Hutyra, L. R., B. Yoon, and M. Alberti. 2011. Terrestrial carbon stocks across a gradient  
894 of urbanization: a study of the Seattle, WA region. *Global Change Biology* 17:783-  
895 797.

896 Idso, C. D., Idso, S. B., and Balling, R. C. J.: The urban CO<sub>2</sub> dome of Phoenix, Arizona,  
897 *Phys. Geogr.*, 19, 95–108, 1998.

898 Imhoff, M. L., C. J. Tucker, W. T. Lawrence, and D. C. Stutzer. 2000. The use of  
899 multisource satellite and geospatial data to study the effect of urbanization on

900 primary productivity in the United States. *Ieee Transactions on Geoscience and*  
 901 *Remote Sensing* 38:2549-2556.

902 IPCC (2001) *Land Use, Land-use Change and Forestry*. Cambridge, UK: Cambridge  
 903 University Press.

904 Jenerette, G. D., J. Wu, N. B. Grimm, and D. Hope. 2006. Points, patches, and regions:  
 905 scaling soil biogeochemical patterns in an urbanised arid ecosystem. *Global Change*  
 906 *Biology* 12:1532-1544.

907 Ji, HX., Shi, Y., Zhu, YM., Wen, JS., Tang, YL., Ge, Y., Chang, J. 2011. Tree growth  
 908 and carbon sequestration in different land-use types in Hangzhou City. *Shengtaixue*  
 909 *Zazhi*, 30: 2405-2412.

910 Kanamitsu M, Ebisuzaki W, Woollen J et al., 2002. NCEP-DOE AMIP- II  
 911 REANALYSIS (R-2). *Bulletin of the American Meteorological Society* 83(11):  
 912 1631–1643.

913 Kaye, J. P., P. M. Groffman, N. B. Grimm, L. A. Baker, and R. V. Pouyat. 2006. A  
 914 distinct urban biogeochemistry? *Trends in Ecology & Evolution* 21:192-199.

915 Kaye, J. P., R. L. McCulley, and I. C. Burke. 2005. Carbon fluxes, nitrogen cycling, and  
 916 soil microbial communities in adjacent urban, native and agricultural ecosystems.  
 917 *Global Change Biology* 11:575-587.

918 Kissling, M., K. T. Hegetschweiler, H.-P. Rusterholz, and B. Baur. 2009. Short-term and  
 919 long-term effects of human trampling on above-ground vegetation, soil density, soil  
 920 organic matter and soil microbial processes in suburban beech forests. *Applied Soil*  
 921 *Ecology* 42:303-314.

922 Koerner, B. and J. Klopatek. 2002. Anthropogenic and natural CO<sub>2</sub> emission sources in  
 923 an arid urban environment. *Environmental Pollution* 116:S45-S51.  
 924 Kuttler, W.: Climate change in urban areas, Part 1, Effects, *Environmen. Sci. Eur.*, 23, 11,  
 925 doi:10.1186/2190-4715-23-11, 2011.  
 926 Lovett, G. M., M. M. Traynor, R. V. Pouyat, M. M. Carreiro, W. X. Zhu, and J. W.  
 927 Baxter. 2000. Atmospheric deposition to oak forests along an urban-rural gradient.  
 928 *Environmental Science & Technology* 34:4294-4300.  
 929 Lu, D., X. Xu, H. Tian, E. Moran, M. Zhao, and S. Running. 2010. The Effects of  
 930 Urbanization on Net Primary Productivity in Southeastern China. *Environmental*  
 931 *Management* 46:404-410.  
 932 McCarthy, M. P., Best, M. J., and Betts, R. A.: Climate change in cities due to global  
 933 warming and urban effects, *Geophys. Res. Lett.*, 37, L09705,  
 934 doi:10.1029/2010GL042845, 2010.  
 935 McDonald, R. I. 2008. Global urbanization: can ecologists identify a sustainable way  
 936 forward? *Frontiers in Ecology and the Environment* 6:99-104.  
 937 McHale, M., Baker, L., Koerner, B., Li, K., Hall, S., and Grimm, N.: Impacts of  
 938 urbanization on carbon cycling: a complete carbon budget of the Phoenix  
 939 metropolitan area, in: *Ecological Society of America Annual Meeting*, Albuquerque,  
 940 NM, COS-46-2, 2009.  
 941 McMurtrie, R. E., R. J. Norby, D. Ellsworth, and D. T. Tissue. 2008. Why is plant-  
 942 growth response to elevated CO<sub>2</sub> amplified when water is limiting but reduced when  
 943 nitrogen is limiting? A growth-optimisation hypothesis. *Journal Name: Functional*  
 944 *Plant Biology*; *Journal Volume*: 35; *Journal Issue*: 6; *Medium*: X; *Size*: 521-534.

945 McPherson, E. G., D. Nowak, G. Heisler, S. Grimmond, C. Souch, R. Grant, and R.  
 946 Rowntree. 1997. Quantifying urban forest structure, function, and value: the  
 947 Chicago Urban Forest Climate Project. *Urban Ecosystems* 1:49-61.

948 Mesinger, F., DiMego, G., Kalnay, E., Mitchell, K., Shafran, P. C., Ebisuzaki, W., Jovic,  
 949 D., Woollen, J., Rogers, E., Berbery, E. H., Ek, M. B., Fan, Y., Grumbine, R.,  
 950 Higgins, W., Li, H., Lin, Y., Manikin, G., Parrish, D., and Shi, W.: North American  
 951 regional reanalysis, *B. Am. Meteorol. Soc.*, 87, 343–360, 2006.

952 Milesi, C., C. D. Elvidge, R. R. Nemani, and S. W. Running. 2003. Assessing the impact  
 953 of urban land development on net primary productivity in the southeastern United  
 954 States. *Remote Sensing of Environment* 86:401-410.

955 Milesi, C., S. W. Running, C. D. Elvidge, J. B. Dietz, B. T. Tuttle, and R. R. Nemani.  
 956 2005. Mapping and modeling the biogeochemical cycling of turf grasses in the  
 957 United States. *Environmental Management* 36:426-438.

958 Newman, P. 2006. The environmental impact of cities. *Environment and Urbanization*  
 959 18:275-295.

960 Nowak, D. J. 2006. Institutionalizing urban forestry as a “biotechnology” to improve  
 961 environmental quality. *Urban Forestry & Urban Greening* 5:93-100.

962 Nowak, D. J. and Crane, D. E.: Carbon storage and sequestration by urban trees in the  
 963 USA, *Environ. Pollut.*, 116, 381–389, 2002.

964 Nowak, D. J., J. C. Stevens, S. M. Sisinni, and C. J. Luley. 2002. Effects of urban tree  
 965 management and species selection on atmospheric carbon dioxide. *Journal of*  
 966 *Arboriculture* 28:113-122.

967 Pataki, D. E., Bowling, D. R., and Ehleringer, J. R.: Seasonal cycle of carbon dioxide and  
 968 its isotopic composition in an urban atmosphere: anthropogenic and biogenic effects,  
 969 J. Geophys. Res.-Atmos., 108, 4735, doi:10.1029/2003JD003865, 2003.

970 Pataki, D. E., R. J. Alig, A. S. Fung, N. E. Golubiewski, C. A. Kennedy, E. G.  
 971 McPherson, D. J. Nowak, R. V. Pouyat, and P. R. Lankao. 2006. Urban ecosystems  
 972 and the North American carbon cycle. *Global Change Biology* 12:2092-2102.

973 Pickett, S. T. A., M. L. Cadenasso, J. M. Grove, C. G. Boone, P. M. Groffman, E. Irwin,  
 974 S. S. Kaushal, V. Marshall, B. P. McGrath, C. H. Nilon, R. V. Pouyat, K. Szlavecz,  
 975 A. Troy, and P. Warren. 2011. Urban ecological systems: Scientific foundations and  
 976 a decade of progress. *Journal of Environmental Management* 92:331-362.

977 Poudyal, N. C., J. P. Siry, and J. M. Bowker. 2010. Urban forests' potential to supply  
 978 marketable carbon emission offsets: A survey of municipal governments in the  
 979 United States. *Forest Policy and Economics* 12:432-438.

980 Pouyat, R. V., D. E. Pataki, K. T. Belt, P. M. Groffman, J. Hom, and L. E. Band. 2007.  
 981 Effects of urban land-use change on biogeochemical cycles. *Terrestrial ecosystems*  
 982 *in a changing world*:45-58.

983 Pouyat, R. V., I. D. Yesilonis, and D. J. Nowak. 2006. Carbon storage by urban soils in  
 984 the United States. *Journal of Environmental Quality* 35:1566-1575.

985 Pouyat, R. V., I. D. Yesilonis, and N. E. Golubiewski. 2009. A comparison of soil  
 986 organic carbon stocks between residential turf grass and native soil. *Urban Ecosyst.*  
 987 12: 45-62.



988 Pouyat, R. V., I. D. Yesilonis, K. Szlavecz, C. Csuzdi, E. Hornung, Z. Korsos, J. Russell-  
989 Anelli, and V. Giorgio. 2008. Response of forest soil properties to urbanization  
990 gradients in three metropolitan areas. *Landscape Ecology* 23:1187-1203.

991 Pouyat, R.V., Russell-Anelli, J., Yesilonis, I.D., Groffman, P.M., 2003. Soil carbon in  
992 urban forest ecosystems. In: Kimble, J.M., Heath, L.S., Birdsey, R.A., Lal, R. (Eds.),  
993 The Potential of U.S. Forest Soils to Sequester Carbon and Mitigate the Greenhouse  
994 Effect. CRC Press, Boca Raton, pp. 347-362.

995 Qian, Y., R. F. Follett, and J. M. Kimble. 2010. Soil Organic Carbon Input from Urban  
996 Turfgrasses. *Soil Science Society of America Journal* 74:366-371.

997 Raciti, S. M., P. M. Groffman, J. C. Jenkins, R. V. Pouyat, T. J. Fahey, S. T. A. Pickett,  
998 and M. L. Cadenasso. 2011. Accumulation of Carbon and Nitrogen in Residential  
999 Soils with Different Land-Use Histories. *Ecosystems* 14:287-297.

1000 Ramalho, C. E. and R. J. Hobbs. 2012. Time for a change: dynamic urban ecology.  
1001 *Trends in Ecology & Evolution* 27:179-188.

1002 Reid, S. B., E. K. Pollard, D. C. Sullivan, and S. L. Shaw. 2010. Improvements to Lawn  
1003 and Garden Equipment Emissions Estimates for Baltimore, Maryland. *Journal of the*  
1004 *Air & Waste Management Association* 60:1452-1462.

1005 Satterthwaite, D., G. McGranahan, and C. Tacoli. 2010. Urbanization and its implications  
1006 for food and farming. *Philosophical Transactions of the Royal Society B-Biological*  
1007 *Sciences* 365:2809-2820.

1008 Schaldach, R. and J. Alcamo. 2007. Simulating the effects of urbanization, afforestation  
1009 and cropland abandonment on a regional carbon balance: a case study for Central  
1010 Germany. *Regional Environmental Change* 7:137-148.

1011 Scharenbroch, B. C. and J. E. Lloyd. 2004. A literature review of nitrogen availability  
 1012 indices for use in urban landscapes. *Journal of Arboriculture* 30:214-230.

1013 Schneider, A., M. A. Friedl, and D. Potere. 2010. Mapping global urban areas using  
 1014 MODIS 500-m data: New methods and datasets based on 'urban ecoregions'.  
 1015 *Remote Sensing of Environment* 114:1733-1746.

1016 Seto, K. C., R. Sanchez-Rodriguez, and M. Fragkias. 2010. The New Geography of  
 1017 Contemporary Urbanization and the Environment. *Annual Review of Environment*  
 1018 *and Resources*, Vol 35:167-194.

1019 Sitch, S., Smith, B., Prentice, I. C., Arneth, A., Bondeau, A., Cramer, W., Kaplan, J. O.,  
 1020 Levis, S., Lucht, W., Sykes, M. T., Thonicke, K. & Venevsky, S. (2003) Evaluation  
 1021 of ecosystem dynamics, plant geography and terrestrial carbon cycling in the LPJ  
 1022 dynamic global vegetation model. *Global Change Biology* 9 (2), 161-185.

1023 Shen, W., J. Wu, N. B. Grimm, and D. Hope. 2008. Effects of urbanization-induced  
 1024 environmental changes on ecosystem functioning in the Phoenix metropolitan region,  
 1025 USA. *Ecosystems* 11:138-155.

1026 Shinozaki, K., Yoda, K., Hozumi, K. & Kira, T. (1964) A Quantitative Analysis of Plant  
 1027 Form -- The Pipe Model Theory I. Basic Analyses. *Japanese Journal of Ecology*, 14,  
 1028 97-105.

1029 Springer, T. L. 2012. Biomass yield from an urban landscape. *Biomass & Bioenergy*  
 1030 37:82-87.

1031 Stearns, F. W.: Urban botany – an essay on survival, *Univ. Wis. Field Sta. Bull.*, 4, 1–6,  
 1032 1971.

1033 Stephenson, N.L., Das, A.J., Condit, R., Russo, S.E., Baker, P.J., Beckman, N.G.,  
 1034 Coomes, D.A., Lines, E.R., Morris, W.K., Rüger, N., Álvarez, E., Blundo, C.,  
 1035 Bunyavejchewin, S., Chuyong, G., Davies, S.J., Á, D., Ewango, C.N., Flores, O.,  
 1036 Franklin, J.F., Grau, H.R., Hao, Z., Harmon, M.E., Hubbell, S.P., Kenfack, D., Lin,  
 1037 Y., Makana, J.R., Malizia, A., Malizia, L.R., Pabst, R.J., Pongpattananurak, N., Su,  
 1038 S.H., Sun, I.F., Tan, S., Thomas, D., Mantgem, P.J.v., Wang, X., Wiser, S.K. &  
 1039 Zavala, M.A. (2014) Rate of tree carbon accumulation increases continuously with  
 1040 tree size. *Nature*, 10.1038/nature12914.

1041 Svirejeva-Hopkins, A. and H. J. Schellnhuber. 2008. Urban expansion and its  
 1042 contribution to the regional carbon emissions: Using the model based on the  
 1043 population density distribution. *Ecological Modelling* 216:208-216.

1044 Svirejeva-Hopkins, A. and Schellnhuber, H.-J.: Modelling carbon dynamics from urban  
 1045 land conversion: fundamental model of city in relation to a local carbon cycle,  
 1046 *Carbon Balance and Management*, 1, 8, doi:10.1186/1750-0680-1-8, 2006.

1047 Taha, H., S. Konopacki, and S. Gabersek. 1996. "Modeling the Meteorological and  
 1048 Energy Effects of Urban Heat Islands and their Mitigation: A 10-Region Study,"  
 1049 Lawrence Berkeley Laboratory Report LBL-38667, Berkeley, CA.

1050 Tian, H.Q., Melillo, J.M., Kicklighter, D.W., Pan, S., Liu, J., McGuire, A.D., Moore III,  
 1051 B., 2003. Regional carbon dynamics in monsoon Asia and its implications to the  
 1052 global carbon cycle. *Global and Planetary Change* 37, 201–217.

1053 Tian, H. Q., Xu, X., Lu, C., Liu, M., Ren, W., Chen, G., Melillo, J., and Liu, J.: Net  
 1054 exchanges of CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O between China's terrestrial ecosystems and the

1055 atmosphere and their contributions to global climate warming, *J. Geophys. Res.*, 116,  
 1056 G02011, doi:10.1029/2010JG001393, 2011a.

1057 Tian, H. Q., Melillo, J., Lu, C., Kicklighter, D., Liu, M., Liu, J., Ren, W., Xu, X., Chen,  
 1058 G., Zhang, C., Pan, S., and Running, S.: China's terrestrial carbon balance:  
 1059 contribution from multiple global change factors, *Global Biogeochem. Cy.*, 25,  
 1060 GB1007, doi:10.1029/2010GB003838, 2011b.

1061 Tian, H., Chen, G., Zhang, C., Liu, M., Sun, G., Chappelka, A., Ren, W., Xu, X., Lu, C.,  
 1062 Pan, S., Chen, C., Hui, D., McNulty, S., Lockaby, G., and Vance, E., 2012. Century-  
 1063 Scale Responses of Ecosystem Carbon Storage and Flux to Multiple Environmental  
 1064 Changes in the Southern United States, *Ecosystems*, 15, 674–694.

1065 Townsend-Small, A. and Czimczik, C. I.: Carbon sequestration and greenhouse gas  
 1066 emissions in urban turf, *Geophys. Res. Lett.*, 37, L02707,  
 1067 doi:10.1029/2009GL041675, 2010.

1068 Tratalos, J., R. A. Fuller, P. H. Warren, R. G. Davies, and K. J. Gaston. 2007. Urban form,  
 1069 biodiversity potential and ecosystem services. *Landscape and Urban Planning*  
 1070 83:308-317.

1071 Trusilova, K. and G. Churkina. 2008. The response of the terrestrial biosphere to  
 1072 urbanization: land cover conversion, climate, and urban pollution. *Biogeosciences*  
 1073 Discussions 5:2445-2470.

1074 Van Aardenne, J.A., Dentener, F.J., Olivier, J.G.J., Klein Goldewijk, C.G.M. and  
 1075 Lelieveld, J. 2001. A 1 x 1 degree resolution dataset of historical anthropogenic  
 1076 trace gas emissions for the period 1890-1990. *Global Biogeochemical Cycles* 15(4):  
 1077 909-928.

1078 Waisanen, P.J., Bliss, N.B. 2002. Changes in population and agricultural land in  
1079 conterminous United States counties, 1790 to 1997. *Global Biogeochemical Cycles*,  
1080 16: 1137.

1081 Wear DN. 2002. Land use. In: Wear DN, Greis JG, Eds. Southern forest resource  
1082 assessment final report. Available online at:  
1083 <http://www.srs.fs.usda.gov/sustain/report/>.

1084 Wu, J. 1999. Hierarchy and scaling: extrapolating information along a scaling ladder.  
1085 *Canadian Journal of Remote Sensing* 25:367-380.

1086 Wu, J. and M. E. Bauer. 2012. Estimating Net Primary Production of Turfgrass in an  
1087 Urban-Suburban Landscape with QuickBird Imagery. *Remote Sensing* 4:849-866.

1088 Young, R. F. 2010. Managing municipal green space for ecosystem services. *Urban*  
1089 *Forestry & Urban Greening* 9:313-321.

1090 Zhang, C., H. Q. Tian, G. S. Chen, A. Chappelka, X. F. Xu, W. Ren, D. F. Hui, M. L. Liu,  
1091 C. Q. Lu, S. F. Pan, and G. Lockaby. 2012. Impacts of urbanization on carbon  
1092 balance in terrestrial ecosystems of the Southern United States. *Environmental*  
1093 *Pollution* 164:89-101.

1094 Zhang, C., 2008. Terrestrial Carbon Dynamics of Southern United States in Responseto  
1095 Changes in Climate, Atmosphere, and Land-use/Land-cover from 1895 to 2005.  
1096 PhD dissertation, Auburn University, United States, 600pp. [Online]: etd.  
1097 [auburn.edu/etd/handle/10415/1098](http://auburn.edu/etd/handle/10415/1098) (last accessed in Jan. 2014).

1098 Zhang, C., H. Tian, S. Pan, M. Liu, G. Lockaby, E. B. Schilling, and J. Stanturf. 2008.  
1099 Effects of Forest Regrowth and Urbanization on Ecosystem Carbon Storage in a  
1100 Rural-Urban Gradient in the Southeastern United States. *Ecosystems* 11:1211-1222.

1101 Zhang, C., Wu, J., Grimm, N.B., McHale, M., Buyantuyev, A. 2013. A hierarchical patch  
 1102 mosaic ecosystem model for urban landscapes: Model development and evaluation.  
 1103 Ecological Modelling 250: 81-100.  
 1104 Zipperer, W.C. 2011. The process of natural succession in urban areas. In: Douglas,  
 1105 Ian; Goode, David; Houck, Mike; Wang, Rusong, eds. The Routledge Handbook of  
 1106 Urban Ecology. London: Routledge Press. p.187-197.  
 1107 Zipperer, W.C. and S.T.A. Pickett. 2012. Urban Ecology: Patterns of Population Growth  
 1108 and Ecological Effects. eLS. John Wiley & Sons, Ltd. DOI:  
 1109 10.1002/9780470015902.a0003246.pub2  
 1110 Zirkle, G., R. Lal, and B. Augustin. 2011. Modeling Carbon Sequestration in Home  
 1111 Lawns. Hortscience 46:808-814.  
 1112 Ziska, L. H., J. A. Bunce, and E. W. Goins. 2004. Characterization of an urban-rural  
 1113 CO<sub>2</sub>/temperature gradient and associated changes in initial plant productivity during  
 1114 secondary succession. Oecologia 139:454-458.  
 1115

Table 1 Scenario design for numeric experiments and factorial analysis

| Description                                | Scenarios                  | Factors               |                              |                 |                     |                |               |  |                                |                             |                        |   |                         |
|--|----------------------------|-----------------------|------------------------------|-----------------|---------------------|----------------|---------------|--|--------------------------------|-----------------------------|------------------------|---|-------------------------|
|  |                            | Urban land conversion | Global environmental changes |                 |                     |                |               | Urbanization induced environmental changes |                                |                             |                        | Urban managements & disturbance regimes |                         |
|  |                            |                       | Climate                      | CO <sub>2</sub> | Nitrogen deposition | Ozone exposure | Pre-urban LUC | Urban heat island                          | Elevated urban CO <sub>2</sub> | Elevated urban N deposition | Aerosol dimming effect | Urban lawn management                   | Urban forest management |
| All combined                               | S <sub>UBNZ</sub>          | ✓                     | ✓                            | ✓               | ✓                   | ✓              | ✓             | ✓  | ✓                              | ✓                           | ✓                      | ✓                                       | ✓                       |
|  | S <sub>UBNZ_Cmin</sub> *   | ✓                     | ✓                            | ✓               | ✓                   | ✓              | ✓             | ✓  | ✓                              | ✓                           | ✓                      | ✓                                       | ✓                       |
|  | S <sub>UBNZ_Cmax</sub> *   | ✓                     | ✓                            | ✓               | ✓                   | ✓              | ✓             | ✓  | ✓                              | ✓                           | ✓                      | ✓                                       | ✓                       |
| Control (global changes only)              | S <sub>GLBC</sub>          | -                     | ✓                            | ✓               | ✓                   | ✓              | ✓             | -  | -                              | -                           | -                      | -                                       | -                       |
|  | S <sub>CLM</sub>           | -                     | ✓                            | -               | -                   | -              | -             | -  | -                              | -                           | -                      | -                                       | -                       |
|  | S <sub>CO2</sub>           | -                     | -                            | ✓               | -                   | -              | -             | -  | -                              | -                           | -                      | -                                       | -                       |
|  | S <sub>NDP</sub>           | -                     | -                            | -               | ✓                   | -              | -             | -  | -                              | -                           | -                      | -                                       | -                       |
|  | S <sub>O3</sub>            | -                     | -                            | -               | -                   | ✓              | -             | -  | -                              | -                           | -                      | -                                       | -                       |
|  | S <sub>LUC</sub>           | -                     | -                            | -               | -                   | -              | ✓             | -  | -                              | -                           | -                      | -                                       | -                       |
| Urban land conversion only                 | S <sub>UBNC</sub>          | ✓                     | -                            | -               | -                   | -              | -             | -  | -                              | -                           | -                      | -                                       | -                       |
| Global changes with urban land conversion  | S <sub>GLBC&amp;UBNC</sub> | ✓                     | ✓                            | ✓               | ✓                   | ✓              | ✓             | -  | -                              | -                           | -                      | -                                       | -                       |
|  | S <sub>CLM&amp;UBNC</sub>  | ✓                     | ✓                            | -               | -                   | -              | -             | -  | -                              | -                           | -                      | -                                       | -                       |
|  | S <sub>CO2&amp;UBNC</sub>  | ✓                     | -                            | ✓               | -                   | -              | -             | -  | -                              | -                           | -                      | -                                       | -                       |
|  | S <sub>NDP&amp;UBNC</sub>  | ✓                     | -                            | -               | ✓                   | -              | -             | -  | -                              | -                           | -                      | -                                       | -                       |
|  | S <sub>O3&amp;UBNC</sub>   | ✓                     | -                            | -               | -                   | ✓              | -             | -  | -                              | -                           | -                      | -                                       | -                       |
|  | S <sub>LUC&amp;UBNC</sub>  | ✓                     | -                            | -               | -                   | -              | ✓             | -  | -                              | -                           | -                      | -                                       | -                       |
| Urbanization induced environmental changes | S <sub>UIEC&amp;UBNC</sub> | ✓                     | -                            | -               | -                   | -              | -             | ✓  | ✓                              | ✓                           | ✓                      | -                                       | -                       |
|  | S <sub>UHI&amp;UBNC</sub>  | ✓                     | -                            | -               | -                   | -              | -             | ✓  | -                              | -                           | -                      | -                                       | -                       |
|  | S <sub>UCO2&amp;UBNC</sub> | ✓                     | -                            | -               | -                   | -              | -             | -  | ✓                              | -                           | -                      | -                                       | -                       |
|  | S <sub>UNDP&amp;UBNC</sub> | ✓                     | -                            | -               | -                   | -              | -             | -  | -                              | ✓                           | -                      | -                                       | -                       |
|  | S <sub>UDIM&amp;UBNC</sub> | ✓                     | -                            | -               | -                   | -              | -             | -  | -                              | -                           | ✓                      | -                                       | -                       |
|  | S <sub>UBMG&amp;UBNC</sub> | ✓                     | -                            | -               | -                   | -              | -             | -  | -                              | -                           | -                      | ✓                                       | ✓                       |
| Urban managements                          | S <sub>LWN&amp;UBNC</sub>  | ✓                     | -                            | -               | -                   | -              | -             | -  | -                              | -                           | -                      | ✓                                       | -                       |
|  | S <sub>UFM&amp;UBNC</sub>  | ✓                     | -                            | -               | -                   | -              | -             | -  | -                              | -                           | -                      | -                                       | ✓                       |

1117 Note: “√” means changes in the environmental factor were considered, while “-” means the factor was unchanged in the simulation.  
1118 \* Following Zhang et al. (2012), UBNZ\_Cmin and UBNZ\_Cmax were designed to examine the effect of uncertainties in model parameter on the  
1119 estimated urbanization effects. Parameters of UBNZ\_Cmin were set so that carbon sequestration were minimized while carbon loss was  
1120 maximized during urbanization; UBNZ\_Cmax was the contrary (Table 2).  
1121



1122 Table 2 The parameters of urban managements and urban-induced environmental changes

| Scenarios <sup>*</sup> | Lawn Managements |                     |                          |  | Urban Forest Managements |                                  | Urbanization Induced Environmental Changes |  |                                 |
|------------------------|------------------|---------------------|--------------------------|--|--------------------------|----------------------------------|--|--|---------------------------------|
|                        | Irrigation (Y/N) | Litter remove (Y/N) | Clipping interval (days) | Nitrogen fertilization (gN m <sup>-2</sup> a <sup>-1</sup> ) | % of biomass pruned      | % of litter removed to land fill | Elevated CO <sub>2</sub> (ppmv)            | Reduced solar radiation due to aerosol (%) | Elevated NO <sub>2</sub> (ppmv) |
| S <sub>UBNZ_Cmin</sub> | Y                | Y                   | 5                        | 6.4  | 10%                      | 25%                              | 5  | -10%                                       | 0.03                            |
| S <sub>UBNZ</sub>      | Y                | Y                   | 10                       | 6.8  | 5%                       | 10%                              | 10   | -5%  | 0.045                           |
| S <sub>UBNZ_Cmax</sub> | Y                | N                   | 15                       | 7.3  | 0%                       | 0%                               | 20   | 0%   | 0.06                            |

1123 \* S<sub>UBNZ</sub> represented the normal condition (i.e. “business as usual” scenario). S<sub>UBNZ\_Cmin</sub> scenario provided a conservative estimation of the urban  
 1124 carbon storage, while the S<sub>UBNZ\_Cmax</sub> scenario simulated the maximum carbon storage of urban/developed area.

1125 Table 3, Contributions of multiple environmental controls to urbanization effect (UBNZ)  
 1126 on carbon (C) dynamic of the Forest (including needleleaf, broadleaf, mixture, and  
 1127 wetland forests), Grass (including C3 and C4 grasslands, and grassy wetland), and arid  
 1128 shrubs (including shrubland and desert) ecosystems in the Southern United States (SUS)  
 1129 from 1945-2007. Unit: g C m<sup>-2</sup>

|                            |  | Forested<br>area | Grassland<br>area | Shrubland<br>area | southern<br>US |
|----------------------------|--|------------------|-------------------|-------------------|----------------|
| Urbanization effect (UBNZ) | Urban land conversion (UBNC)   | -2655            | -1078             | -303              | -2845          |
|                            | Interactive effects between urban land conversion and global changes (GLBC-UBNC) |                  |                   |                   |                |
|                            | UBNC interact with climate change  | 43               | 33                | 21                | 39             |
|                            | UBNC interact with global CO <sub>2</sub> change                                 | -89              | -25               | -28               | -73            |
|                            | UBNC interact with N deposition change   | -86              | -19               | -10               | -69            |
|                            | UBNC interact with global O <sub>3</sub> change                                  | 70               | 28                | 27                | 58             |
|                            | UBNC interact with pre-urban land-use change                                     | -96              | 12                | 41                | -68            |
|                            | Interactive effects among the global change factors                              | -117             | 141               | -68               | -71            |
|                            | Overall effect of GLBC-UBNC  | -276             | 168               | -18               | -183           |
|                            | Urbanization-induced environmental changes (UIEC)                                |                  |                   |                   |                |
|                            | Urban heat island effect   | -136             | -98               | -33               | -120           |
|                            | Urban CO <sub>2</sub> dome effect  | 252              | 155               | 88                | 221            |
|                            | Effect from elevated N deposition in urban                                       | 270              | 245               | 92                | 249            |
|                            | Interactive effects among the UIEC   | -130             | -131              | -64               | -123           |
|                            | Overall effect of UIEC   | 256              | 171               | 82                | 226            |
|                            | Urban managements (UBMG)   |                  |                   |                   |                |
|                            | Urban lawn management  | 455              | 639               | 694               | 490            |
|                            | Urban forest management  | 525              |                   |                   | 396            |
|                            | Overall effect of UBMG   | 980              | 639               | 694               | 886            |
|                            | Overall interactive effects among UBNC, UIEC, UBMG, and GLBC-UBNC (IT_Other)     | 411              | 4                 | -6.7              | 307            |
|                            | Overall effect of UBNZ   | -2084            | -95               | 390               | -1609          |

1130

Table 4 Comparison of model predictions against observed carbon pools and fluxes of urban ecosystems

| City                              | Lon/Lat (dd) | PFT* | Age  | <u>NPP (g C/m<sup>2</sup>/yr) + SE<sup>@</sup></u> |            | <u>VEGC (kg C/m<sup>2</sup>) + SE<sup>@</sup></u> |            | <u>SOC1m (kg C/m<sup>2</sup>) + SE<sup>@</sup></u> |            | Sources of Observations                     |
|-----------------------------------|--------------|------|------|--|------------|---|------------|--|------------|---|
|                                   |              |      |      | Observation  | Prediction | Observation                                       | Prediction | Observation  | Prediction |   |
| Atlanta, GA                       | -84.4/33.65  | CF   | NA   |  |            | 9.7±0.7   | 9.3        |  |            | Nowak & Crane, 2002                         |
| Baltimore, MD                     | -76.6/39.28  | BF   | NA   |  |            | 10.0±1.3  | 11.2       | 12.1±1.8   | 11.0       | Nowak & Crane, 2002;<br>Pouyat et al., 2008 |
| Baltimore, MD                     | -76.6/39.28  | Lawn | > 50 |  |            |   |            | 12.2±1.1   | 11.9       | Pouyat et al., 2008                         |
| Baltimore, MD                     | -76.6/39.28  | Lawn | 30   |  |            |   |            | 10.7   | 9.6        | Pouyat et al., 2008                         |
| Baltimore, MD                     | -76.6/39.28  | Lawn | 20   |  |            |   |            | 8.1  | 8.9        | Pouyat et al., 2008                         |
| Boston, MA                        | -71.03/42.37 | BF   | NA   |  |            | 9.1±1.1   | 9.9        |  |            | Nowak & Crane, 2002                         |
| Fort Collins, CO <sup>&amp;</sup> | -105.1/40.6  | Lawn | > 60 |  |            |   |            | 13.1   | 13.5       | Kaye et al., 2005                           |
| Front Range, CO <sup>#</sup>      | -105/40      | Lawn | NA   | 762±92   | 731        | 1.55 ±0.16  | 1.59       | 11.6   | 12.0       | Golubiewski, 2006                           |
| Miami-Dade, FL                    | -80.20/25.77 | BF   | NA   |  |            | 7.47  | 6.83       |  |            | Escobedo et al., 2010                       |
| Philadelphia, PA <sup>\$</sup>    | -75.17/39.95 | BF   | NA   |  |            | 9.0±0.9   | 10.3       |  |            | Nowak & Crane, 2002                         |
| Syracuse, NY <sup>\$</sup>        | -76.12/43.12 | BF   | NA   |  |            | 9.4±1.0   | 9.7        |  |            | Nowak & Crane, 2002                         |
| Washington DC                     | -76.5/30.88  | Lawn | NA   | 737  | 715        | 1.49  | 1.54       |  |            | Falk, 1976                                  |

1131 \*Plant functional type (PFT): CF denotes Coniferous forest; BF denotes deciduous forests

1132 @ NPP: net primary productivity; VEGC: vegetation carbon; SOC1m: soil organic carbon (0-1 m). Table 1 values (root NPP:NPP, root C:VEGC365, C:Biomass)  
 1133 were used to convert the above-ground NPP and biomass to total NPP and VEGC. By assuming that 53% of SOC of grassland in the upper 30 cm (Jobbagy  
 1134 and Jackson, 2000), we calculated the SOC of lawns to 1 m for those studies only measuring top 30 cm.

1135 #Intensively managed sites. 23 samples. Two tailed Student's t-test indicated differences between model predictions and observations were not significant (90%  
 1136 level).

1137 & Intensively managed lawn: annual N fertilization rate = 11 gN/m<sup>2</sup>/yr; clipped litter left on site.

1138 \$ The species composition of the northern forest is not same to that of the southern forest. Before conducting simulations in NY and PA, we estimated/calibrated  
 1139 the physiological parameters of the BF functional type against the intensively studied Harvard Forest LTER site. For other cities and the regional simulations in  
 1140 the SUS, we used the parameters which were developed based on the studies in the Duke Forest.

1141

## 1142 **Figure captions**

1143 Fig. 1, The urbanization effects (UBNZ) on regional carbon dynamics are controlled by four major types of environmental changes,  
1144 including (1) urban land conversion (UBNC) during which rural land-use type is converted to urban and developed land-use composed  
1145 by impervious surfaces, managed urban lawn, and other urban vegetation (e.g., urban forest); (2) urban management (UBMG)  
1146 including lawn management (LWN) such as irrigation and fertilization, and urban forest management such as protection from logging  
1147 and fire disturbances; (3) urbanization induced environmental changes (UIEC) including effects of urban heat island (UHI), elevated  
1148 CO<sub>2</sub> (UCO<sub>2</sub>) and N deposition (UNDP), reduced solar radiation due to air pollution (UDIM), and interactions among these UIEC  
1149 factors (IT\_UIEC), and (4) the interactive effects between UBNC and multiple global environmental changes (GLBC-UBNC)  
1150 including changes in climate (CLM-UBNC), CO<sub>2</sub> (CO<sub>2</sub>-UBNC), N deposition (NDP-UBNC), ozone exposure (O<sub>3</sub>-UBNC), pre-urban  
1151 land-use change history (LUC-UBNC) such as cropland conversion and abandonment, and the interactions among all GLBC-UBNC  
1152 factors. IT\_OTHER represents the overall interactive effects among the four major controls (i.e. UBNC, UBMG, UIEC, and GLBC-  
1153 UBNC). The numbers in the figure show the carbon flux in response to each factor from 1945-2007 in the southern United States. Unit:  
1154 TgC.

1155 Fig. 2. Temporal patterns of major global change factors in the study region from 1945-2007. (a) annual precipitation; (b) temperature;  
1156 (c) relative humidity; (d) ambient ozone exposure; AOT40 is the accumulated dose over a threshold of 40 ppb during daylight hours in  
1157 a month (Felzer et al., 2004); (e) annual nitrogen deposition rate; (f) Landuse changes. The dynamic of Pre-urban cropland showed the  
1158 net balance between cropland conversion and cropland abandonment in the control scenario (i.e., S<sub>GLBC</sub>, assuming no urbanization has  
1159 taken place since 1945). In reality (i.e. in the S<sub>UBNZ</sub>) all lands have been converted into urban.

1160 Fig. 3, Cumulative effects of urbanization (UBNZ) to the carbon dynamic of the Southern United States from 1945-2007 and the  
1161 contributions of multiple environmental drivers. UBNC is the effect of urban land conversion; GLBC-UBNC is the interactive effect  
1162 between global environmental changes (GLBC, including changes in climate, CO<sub>2</sub>, N deposition, ozone, and landuse) and UBNC;  
1163 UBMG is the management effect on the carbon dynamic of urban vegetation (such as lawn and urban forest); UIEC is the effects due  
1164 to urbanization induced environmental changes (such as urban heat island, CO<sub>2</sub> dome effect, and elevated N deposition in urban); the  
1165 overall interactive effects among UBNC, GLBC-UBNC, UBMG, and UIEC is represented by IT\_OTHER. Following, Zhang et al.  
1166 (2012), UBNZ<sub>Cmin</sub> and UBNZ<sub>Cmax</sub> represent the minimum and maximum urbanization effects, respectively, as influenced by  
1167 uncertainties in model parameterization (Table 2).

1168