

1 We thank both reviewers and Yan Li for their time and effort and their helpful and constructive
2 comments. The original comments of the reviewers and Yan Li are in color. Our reply is in black.

3
4 **Comments Yan Li**

5
6 **I just happened to read this paper and I would like to pose my comments. It is a interesting
7 paper that calculate RF at very fine scale. Here are my comments.**

8 **P10127 L5: RF affects global mean temperature. Technically, estimating RF can be done at very
9 fine scale. But I am thinking RF at very fine scale, even it is positive, may have negligible impact
10 on regional/global climate. However, the local impact of forest expansion is much larger than RF
11 change. This is not for your paper, it is just my consideration on this question.**

12
13 **L6: Why forest cover increased in temperate mountains region? Is that because tree line moves
14 up to higher altitude due to global warming?**

15
16 In our discussion paper we write: "Forest cover has increased in many temperate mountainous
17 regions". We do not suggest that it increased in all temperate mountainous regions. E.g.
18 Ramankutty et al. (2010) show regional differences in the United States. The references we
19 listed in line 6-8 (P 10127) discuss both aspects, forest cover increase due to land-use change
20 and due to climatic changes (Alewell and Bebi, 2011, MacDonald et al., 2000, Ramankutty et al.,
21 2010, Kozak, 2003, Hagedorn et al., 2014).

22
23 **L20: What drives forest expansion in Switzerland? Natural causes or forestry? Or due to reasons
24 listed in P10128 L16-17?**

25
26 Forest expansion in Switzerland is mainly related to the reasons listed in P10128 L16-17 (the
27 widespread abandonment of marginal agricultural land and the subsequent expansion of forest
28 cover). However, a small amount of forest cover increase is related to climatic changes.

29 We added the following sentence to the revised discussion paper: "Land abandonment was the
30 most dominant driver for the establishment of new forest areas, however, a small fraction of
31 forest expansion at the tree line can be attributed to the recent climate warming (Gehrig-Fasel
32 et al., 2007)."

33 **For methodology part: How reliable are the climate data (global radiation) and carbon stock (soil
34 carbon...) at such high spatial scale? It seems to me that spatial data of carbon stock are derived
35 from assigning averaged values of each land class to an existing land cover map? Accurately
36 mapping carbon stock is still a challenge.**

37
38 We think that the global radiation data is of high quality (P 10133 L15-L19): "The spatial
39 resolution of the global radiation datasets is 2.2km. The derivation of the global radiation data
40 was based on the Heliosat method (Cano et al., 1986; Beyer et al., 1996; Hammer et al., 2003),

41 applied to Meteosat SEVIRI data. It was verified using high-quality surface measurements and
42 sensitivity runs for key input parameters (Durr et al., 2010).”

43 Yes, to derive carbon stocks at high spatial resolution is indeed a challenge and we thus assigned
44 average values to the land-cover classes based on a biogeographical and altitudinal
45 stratification. We mainly follow the methods in Switzerland’s Greenhouse Gas Inventory. Please
46 refer to the methods section P 10129 L 26 – P10130 L 3 (P 10129 L 24 – P10131 L 4).

47

48 Please check the label of each sub-figure and its captions of figure 3.

49 We changed the labels.

50

51 Figure 3: Does Albedo difference have seasonal variations due to phenology during snow free
52 season?
53

54 Please refer to the reply to the comments of referee 1 for a more detailed discussion on this
55 point.

56

57 P10131 L10: How do you estimate delta mc (carbon sequestrated)? Do you mean carbon stock
58 here?
59

60 We changed the description of delta mc. It is now: “... where ΔCA is the change in atmospheric
61 CO2 concentration, ΔmC the difference between carbon stocks of two LULC classes, ...”

62

63 P10139 L28-29 You said “The carbon sequestration potential of forests decreased with altitude”.
64 But why CO2 - forcing in figure 5 becomes more negative as altitude increases in the three
65 region on the righthand?
66

67 Figure 5 shows the CO2-forcing of forest expansion between 1985 and 1997. The CO2 forcing
68 becomes more negative as altitude increases because most transitions from open land to forest
69 occurred above 1000 m. Forests in high altitude will have lower carbon stocks than forests in
70 low altitudes. However, there are much more transitions from open land to forest in high
71 altitude and thus the CO2 forcing becomes more negative.

72 It seems to be a general issue, that our discussion paper does not yet clearly show, where we
73 included real forest transitions between 1985 and 1997 and where we just showed spatial
74 differences in RF of “potential” forest expansion. Please refer to the reply to the review of
75 referee 2 and the revised version of the discussion paper, where we dealt with this issue in
76 more detail.

77

78 P10140 L5: The word “carbon sequestration” sounds to me is a time dependent rate that forest
79 remove carbon from atmosphere, e.g., NEP/NEE, kgc/year. Carbon stock refers to the current
80 state about the existing mass of carbon in forest biomass.

81

82 We estimated carbon sequestration as the difference in carbon stocks between two LULC
83 classes. For a better understanding we replaced: “Transitions from Open Forest to Closed Forest
84 were generally associated with relatively high amounts of carbon sequestration...” by
85 “Transitions from Open Forest to Closed Forest were generally associated with relatively high
86 change in carbon stocks...”.

87

88 I think some contents in discussion are more suitable to appear in Results (e.g., second
89 paragraph of discussion). There are too many things in current discussion which is a bit too long
90 and lacks of focus that I get lost. It can be improved by better organize key points and
91 condensation in language.

92

93 We agree that the discussion is long. However, we considered all points in the discussion to be
94 very important and decided not to shorten it.

95

96 RFs of albedo change and CO2 have different climate sensitivities, if you want to use RF to
97 consider their contribution to temperature change, you should keep in mind about this. (see
98 Zhao, KG, 2014, Biophysical forcings of land-use changes from potential forestry activities in
99 North America; Hansen, J., et al. 2005. Efficacy of climate forcings. Journal of Geophysical
100 Research Atmospheres 110:D18104.)

101

102 We agree that climate sensitivities are a very important point. Please refer to the discussion
103 paper P 10144 L 24 – P 10145 L 2: “However, the interpretation of RF values has to be done
104 carefully. First, the concept of Radiative Forcing has been developed to compare the impact of
105 different forcing agents on the global mean temperature (Hansen et al., 2005). When applied at
106 the regional and local scales one should keep in mind that the comparison of different forcing
107 agents is far from being straightforward. For instance, the impact of albedo will remain mostly
108 local while those from CO2 will be globally distributed and therefore diluted. Furthermore, the
109 Climate sensitivities of CO2 RF and Albedo RF may differ (Davin et al., 2007).”

110 Since Zhao and Jackson (2014) refer to Davin et al. (2007) when discussing the differences in
111 climate sensitivities of CO2 and albedo, we chose to directly refer to this reference.

112

113 We thank both reviewers and Yan Li for their time and effort and their helpful and constructive
114 comments. The original comments of the reviewers and Yan Li are in color. Our reply is in black.

115

116 Comments Referee 1

117

118 The study's national level investigation into net radiative forcing of forest change is a
119 great contribution to the field. The synthesis of different data sources is well thought out
120 and well presented (especially the many assumptions required by such a synthesis). I
121 greatly appreciated the inclusion of the sensitivity analysis.

122

123 My only requested revision of any weight is at page 10133 line 26 through page 10134
124 line 2. The authors state that "seasonal variation of the albedos of different land-use
125 classes is very similar". Since the statement is in support of a central assumption of
126 the methods, some values or a citation would be helpful.

127

128 This is indeed an important assumption. We added boxplots showing the seasonal trend of the four land
129 use/land cover (LULC) classes "Forest", "Open Forest", "Intensively Used Open Land" and "Extensively
130 Used Open Land". The trends are very similar. However, especially for snow-covered albedos there are
131 also differences. For example the albedo of forest in April and May are increasing (in comparison to
132 previous values), while the albedos of the three classes "Open Forest", "Intensively Used Open Land"
133 and "Extensively Used Open Land" decrease in April and May.

134 There are mainly 2 reasons, why we decided to use average values and not differences for each month.
135 First, the strongest seasonal trend is related to snow-cover, which we explicitly included (Zhou et al.,
136 2003). Second, the use of seasonally varying albedo differences in snow-free and snow-covered albedo
137 requires albedo data for every month. Since we calculated albedo values for small biogeographical
138 regions and 4 specific LULC classes, there are sometimes only few or even missing albedo values for a
139 certain month/LULC class/biogeographical region (e.g. for snow-covered albedos in September/October
140 and May/June). Using seasonally varying albedo differences it is necessary to interpolate and extrapolate
141 albedo values for some months and accept bias when only few values are available (e.g. again for
142 September/October and May/June). Inter- and extrapolating albedo values, we calculated the spatially
143 explicit pattern of albedo RF again. Root mean square error was 4.3% and the pattern we found was
144 mostly identical. Averaging the albedo values does not account for the seasonal trends in the albedo
145 differences, however, it was a stable estimate reducing the effect of outliers and assumptions needed to
146 inter- and extrapolate albedo values.

147 We adapted the paragraph in the discussion paper:

148 "The albedo was estimated using the following equation (modified from Barnes and Roy, 2010, Roesch et
149 al., 2002):

150

$$\Delta\alpha(t) = f(t)\Delta\alpha_s + (1 - f(t))\Delta\alpha_v \quad (1)$$

151 , where $\Delta\alpha(t)$ is the monthly albedo-difference between two LULC classes, $\Delta\alpha_s$ the average albedo
152 difference between two LULC classes when snow-covered, $\Delta\alpha_v$ the average albedo difference between
153 two LULC classes when snow-free and $f(t)$ the fraction of snow-cover per month. We used average
154 albedo differences of snow-free and snow-covered albedo differences and not monthly differences
155 because of two reasons. First, the strongest seasonal trend is related to the presence of snow, which we
156 explicitly included (Zhou et al., 2003). Second, in some months, reliable albedo data was missing and we
157 considered the average to be a robust estimate. Since we found that the seasonal variation of the
158 albedos of different LULC classes is similar, the averaging of snow-covered and snow-free albedo
159 differences results in a fairly good approximation (Appendix Figure 2, Appendix Figure 3)."

160 A few minor corrections: Page 10126 line 11 - The text reads "BIOGEOPHYSICAL processes tend
161 to counter the BIOGEOPHYSICAL effect". Should one of the "biogeophysical"s be
162 "biogeochemical"?

163 Page 10126 line 21 - as above "between BIOGEOPHYSICAL (mainly albedo) and
164 BIOGEOPHYSICAL effects"
165


166 We changed the second biogeophysical to biogeochemical in both cases.
167

168 Page 10130 line 14 - Clarification needed, "and that of needles/leaves on (Perruchoud
169 et al., 1999)." Were the authors intending the this to read: "needles/leaves is based on
170 Perruchoud et al. (1999)."
171

172 Yes, it should be "needles/leaves is based on Perruchoud et al. (1999)."
173


174

175 Figures added to the appendix:



176

177 Figure 2: Seasonal variation of albedo values of the four snow-free LULC classes Closed Forest,
 178 Intensively Used Open Land, Extensively Used Open Land and Open Forest (only full BRDF inversion
 179 albedo values).



180

181 Figure 3: Seasonal variation of the albedo values of the four snow-covered LULC classes Closed Forest,
 182 Intensively Used Open Land, Extensively Used Open Land and Open Forest (full BRDF and magnitude
 183 inversion albedo values).

184

185

186

187

188 We thank both reviewers and Yan Li for their time and effort and their helpful and constructive
189 comments. The original comments of the reviewers and Yan Li are in color. Our reply is in black.
190

191 1. First, let me say that, in general, I like this analysis and think that it's important. The
192 authors have done a good job incorporating many quality dataset to address a complex
193 problem, and the spatial nature of the analysis is a major strength. However, I have a
194 major problem in that the presentation of the study design and methods are incomplete,
195 such that I cannot determine whether the study design is sound. I am left fully confused
196 by what was actually done. I therefore cannot determine whether the analysis is actually
197 fine, but the methodology simply needs to be explained better, or rather the study design
198 is flawed or could be improved. I will be more specific below. The fundamental problem is
199 that, despite an emphasis on the spatial nature of this analysis, it is not at all indicated
200 where the land use transitions you include actually happen, or even how much land area
201 is converted. Figures 3 and 4 indicate results are "wall to wall", where every pixel has
202 experienced a radiative forcing. This implies that all pixels were assigned a land use
203 transition, which seems very unreasonable. You considered five land use transitions;
204 which pixels received each transition? Figures 3 and 4 (and 5??) are presented for a
205 particular transition (Intensively Used Open Land (<1000 m) and Extensively Used Open
206 Land (>1000) to Closed Forest), so was the analysis done five times where all pixels
207 received the same transition? Where are the figures for the other transitions? Is this
208 supposed to represent a maximum afforestation case, where all open land is converted to
209 forest? Is that climatologically reasonable? Could forests grow in all of these pixels? Right
210 now your relevant study design text spans about five lines (P10129, lines 6-11). Please
211 expand this and include new figures and tables that illustrate the location and amount of
212 area where particular land use transitions occurred, and text that addresses whether
213 these transitions are supposed to represent reality (between 1985 and 1997) or a
214 hypothetical case? I cannot imagine it is the former, since every pixel seems to have been
215 altered (and experienced a radiative forcing).

216
217
218 Since most of the questions and concerns of reviewer 2 are related to the methodology
219 concerning the transitions between land use/land cover classes and the study design, we will
220 first address these issues in general and then answer the individual questions more specifically.

221
222 The methodology of the assessment of land use/land cover transitions are based on aerial
223 photograph based surveys of land use/land cover for Switzerland in the years 1985 and 1997
224 and has been described in more detail in Rutherford et al. (2008). We realize and acknowledge
225 that the description of this method has been kept very short in this manuscript and that it can
226 be difficult to understand the methodology without additional information about these data.
227 Also we acknowledge that it may have been confusing that we used these data slightly
228 differently for the analysis of (1) spatial pattern in Radiative Forcing in a temperate
229 mountainous region and (2) what the inclusion of albedo change implies for the greenhouse gas
230 inventory in Switzerland between 1985 and 1997 (calculating the total amount of albedo forcing
231 and CO2 forcing of the forest expansion between 1985 and 1997).

232
233
234

235 In the revised version of the discussion paper we will explain the methodology of land use/land
236 cover assessment more clearly and we will in particular separate more clearly between the two
237 different analysis/results for (1) spatial pattern in Radiative Forcing and (2) and Radiative
238 Forcing of forest expansion in Switzerland between 1985 and 1997.

239 We added the following paragraph to the introduction:

240 “Our study design is twofold: First, we use the spatially explicit datasets to show the pattern of
241 RF assuming that each location in Switzerland is facing a transition from agriculturally used open
242 land to forest. This is not related to any particular or realistic scenario, however, the spatial
243 pattern of RF can be of high interest for any land-use policies steering forest cover change towards
244 desired futures. In Switzerland agricultural subsidies directly influence farmers decisions on
245 whether to keep managing or abandon their land. The latter will usually result in forest expansion.
246 Second, we include the type and location of 5 different land use transitions to calculate RF in
247 Switzerland between 1985 and 1997. In summary we estimate i) to which extent albedo RF
248 offsets CO2 RF in different parts of temperate mountainous regions, ii) how each input parameter
249 influences RF, and iii) what the inclusion of albedo change implies for the greenhouse gas
250 inventory in Switzerland.”

251

252 For further changes see point 2 and 3.

253

254 **2. The fundamental problem is that, despite an emphasis on the spatial nature of this**
255 **analysis, it is not at all indicated where the land use transitions you include actually**
256 **happen, or even how much land area is converted.**

257 Unfortunately, the figure A1 and the table A4 are at the very end of this discussion paper and
258 thus not very visible. Figure A1 shows where most transitions occur. We had to use a kernel
259 density function (showing densities of land-use transitions) because it is difficult to visualize
260 single pixels in a 3000 x 2000 grid. Table A4 shows forest expansion for every biogeographical
261 region. We will refer to this figure and this table more clearly in the text. In addition we added
262 more information to Table 4.

263

264 We added:

265 “... At lower elevations, transitions from Intensively Used Open Land to Forest are frequent,
266 while in higher elevations transitions from Extensively Used Open Land to Forest are most likely
267 (Appendix, Table 4). ...”

268 Table 4 (modified version): Area affected by each type of transition between 1985 and 1997.
 269 Numbers behind each biogeographical region 1-3 indicate the elevation (1 = below 600m, 2 =
 270 600 – 1200m, 3 = above 1200m).

271

Biogeographical region	Intensively Used Open Land to Closed Forest [ha]	Extensively Used Open Land to Closed Forest [ha]	Intensively Used Open Land to Open Forest [ha]	Extensively Used Open Land to Open Forest [ha]	Open Forest to Closed Forest [ha]	Forest expansion (sum of all transitions) [ha]
Jura 1	116	31	98	35	106	386
Jura 2	113	238	73	330	522	1276
Jura 3	1	46	1	155	490	693
Plateau 1	613	87	379	52	264	1395
Plateau 2	232	60	110	44	85	531
Plateau 3	NA	NA	NA	NA	1	1
Northern Prealps 1	109	21	78	13	53	274
Northern Prealps 2	321	497	295	401	959	2473
Northern Prealps 3	34	955	77	1180	2476	4722
Alps 1	4	4	6	11	29	54
Alps 2	93	101	267	154	679	1294
Alps 3	102	739	291	1700	3687	6519


Southern Prealps 1	61	25	145	22	196	449
Southern Prealps 2	76	77	170	135	731	1189
Southern Prealps 3	23	274	71	604	1541	2513

272

273

274 We added:

275 “... Forest expansion mainly took place in elevations above 1200 m in the Prealps and the
 276 Central Alps (Figure 1). ...”



277

278 Figure 1: Spatial pattern of forest expansion. The pattern illustrates the density of forest
 279 expansion in Switzerland. The density was calculated including the area of all five transitions we
 280 used for calculating RF (see chapter “Swiss forest expansion between 1985 and 1997”) and a
 281 kernel-density function in ArcGis 10.1 (ESRI).

282

283

284 3. Right now your relevant study design text spans about five lines (P10129, lines 6-11).
285 Please expand this and include new figures and tables that illustrate the location and
286 amount of area where particular land use transitions occurred, and text that addresses
287 whether these transitions are supposed to represent reality (between 1985 and 1997) or
288 a hypothetical case?

289 We put chapter 2.6. (Spatial variability of RF and RF of Swiss forest expansion) right after the
290 description of the study area. Now chapter 2.6. and 2.2. (we renamed chapter 2.2 from “Land
291 use/Land cover (LULC)” to “Swiss forest expansion between 1985 and 1997”) are close together.
292 Both together are a description of the study design. We included some changes. The two chapters
293 are now:

294 **“Spatial variability of RF and RF of Swiss forest expansion**

295 We calculated the net RF and the offset of CO2 RF through albedo RF ($\Delta R_{FCO_2}/\Delta R_{albedo}$) to
296 show the pattern of RF in Switzerland and to calculate RF of Swiss forest-cover expansion between
297 1985 and 1997. To illustrate the pattern of RF in Switzerland we calculated a value of RF for every
298 location in Switzerland, excluding non-vegetated land, water, settlement and areas that lie above
299 the tree line (Figure 3). These are hypothetical values, because we calculated RF for the change
300 from open land to forest for all vegetated areas, and not only for the ones where forest expansion
301 was actually observed. At lower elevations, transitions from Intensively Used Open Land to forest
302 are frequent, while in higher elevations transitions are usually from Extensively Used Open Land
303 to Forest (Appendix, Table 4). We considered this by separating our estimation of the hypothetical
304 RF in transitions from Intensively Used Open Land to forest below 1000 m and transitions from
305 Extensively Used Open Land to Forest above 1000 m. The results of the spatial pattern of RF are
306 shown in maps of the study area (Figure 4 c,d).

307 To obtain results for RF of forest-cover expansion between 1985 and 1997 in Switzerland we
308 calculated net Radiative Forcing as the sum of RF for all pixels where forests expanded. This meant
309 including information on the type of forest expansion and on the location of forest expansion:

310

$$\Delta RF_{ges} = \frac{\sum_{x=1}^n RF_{x,T}}{A_E} \quad (2)$$

311

312 , where ΔRF_{ges} is the net Radiative Forcing (net RF), n the number of pixels where forests
313 expanded and RF the Radiative Forcing, which depends on the location x and the type of transition
314 T. The sum over RF is divided by the the earth’s surface A_E to convert local RF into a global
315 average RF.

316

317 **Swiss forest expansion between 1985 and 1997**

318 We use aerial photographs processed by Swiss Statistics at a spatial resolution of 100 m to derive
319 changes in land use/land cover (LULC). These aerial photographs are from the Swiss Federal
320 Office of Topography and are fully available for the two inventory periods 1979-85 and 1992-97
321 (Humbel et al., 2010). We reclassified the data of the different inventory periods into five
322 aggregated classes (Rutherford et al., 2008). While 18 classes were aggregated into four classes:
323 Closed Forest, Open Forest, Extensively and Intensively Used Open Land (Figure 2), the remaining
324 56 were classified as Other, and consisted mainly of settlements, water and non-vegetated land
325 (Appendix, Table 5). The aggregation of the original land-use classes is a simplification. It was not
326 possible to derive reliable data on albedo and carbon stocks for each LULC class. The aggregation
327 of the original land-use classes results in a sufficiently large sample of reliable albedo values and
328 carbon stocks in each of the five biogeographical regions and three elevational strata for five
329 relevant and well established land-use classes.

330 To calculate RF of land use change between 1985 and 1997, we included five transitions: 1.
331 Intensively Used Open Land -> Closed Forest, 2. Extensively Used Open Land -> Closed Forest, 3.
332 Intensively Used Open Land -> Open Forest, 4. Extensively Used Open Land -> Open Forest and 5.
333 Open Forest -> Closed Forest. Forest expansion mainly took place in elevations above 1200 m in
334 the Prealps and the Central Alps (Appendix, Figure 1). We focused on transitions where forest
335 cover and carbon stocks increase, because these transitions highly exceeded transitions with
336 forest decrease in Switzerland. In fact, the Swiss law strongly protects forests so that there have
337 been only few changes from forest to agriculturally used land during the last 30 years (Bloetzer,
338 2004, Rutherford et al., 2008).”

339

340

341 **4. Also, crucially, what is the impact of aggregating 19 land classes into five? The authors**
342 **need to include figures to clarify the impact of these simplifications in their analysis. Again,**
343 **how much area is affected?**

344 We added a table to the discussion paper that shows how we aggregated the LULC classes and
345 how much area is affected.

346 It was not possible to derive reliable data on albedo and carbon stocks for each LULC class (18
347 classes are related to forest cover change). Aggregating the land-use classes meant that it was
348 not possible anymore to differentiate between Radiative Forcings of very particular transitions
349 (e.g. from “Stony Alpine Pasture” to “Slender Forest”). However, it was possible to define average
350 albedos and carbon stocks of the aggregated land-use classes in every biogeographical region and
351 three elevational strata. The broad definitions of closed forest, open forest, intensively and
352 extensively used open land are given in Table 5. These definitions refer to well established land-
353 use classes and are very useful to reflect the most relevant categories in terms of land-use change
354 and radiative forcing.

355

356

357

358 Table 5: Aggregation of land use classes from Swiss Arealstatistik (ASCH85, ASCH97 and
 359 ASCH04) adapted from (Rutherford et al., 2008).

Aggregated class	Area [ha]	Classes from Swiss land use statistics	Area [ha]	Broad definition
Closed Forest	1121544	Afforestation*, 52	3349	Vegetation height >3m, cover density >60%, composed of tree species
		Forest dieback*, 54	14851	
		Normal forest, 50	962312	
		Slender Forest, 51	44711	
		Bushes, 57	60514	
		Groves and hedges, 58	35807	
Open Forest	150101	On non-agriculturally used land, 56	52825	Vegetation height >3m, cover density 20-60%, composed of tree species
		On agriculturally used land, 55	24108	
		Groups of trees on agriculturally used land, 59	38157	
		Groups of trees on non-agriculturally used land, 60	35011	
Extensively Use Open Land	767842	Pasture in the vicinity of settlements, 43	87303	Used for grazing, use not year-round, not machine-accessible
		Alpine meadows, 45	32316	
		Sheep alps, 49	51124	
		Favourable to pasturing, 46	368691	
		Stony alpine pasture, 48	46024	
		Grass and herb vegetation, 65	182384	
Intensively Used Open Land	837128	Arable land, 41	547754	Year-round use, in the vicinity of settlements, Mown
		Natural meadows, 42	289374	
Other		1-40, 44, 47, 61-64, 66-72		

Numbers in column 2 represent the official ASCH classes of the nomenclature 2004 (Humbel et al., 2010). The aggregation in (Rutherford et al., 2008) was adapted to the new nomenclature.

360

361 *Afforestation and *Forest dieback are LULC classes and not transitions or processes.

362

363 5. The abstract will also need to be revised so that it is very clear how the land use transitions
364 were assigned. I think the paper would be greatly improved if it was organized to address
365 a very clear and specific statement of the research objective.

366

367 We changed the abstract:

368 “In this study, we assess the climate mitigation potential from afforestation in a mountainous snow-rich
369 region (Switzerland) with strongly varying environmental conditions. Using radiative forcing calculations,
370 we quantify both the carbon sequestration potential and the effect of albedo change at high resolution.
371 We calculate the albedo radiative forcing based on remotely sensed datasets of albedo, global radiation
372 and snow cover. Carbon sequestration is estimated from changes in carbon stocks based on National
373 Inventories. We first estimate the spatial pattern of RF across Switzerland assuming homogeneous
374 transitions from open land to forest. This highlights where forest expansion still exhibits climatic benefits
375 when including the radiative forcing of albedo change. Second, given that forest expansion is currently the
376 dominant land-use change process in the swiss Alps, we calculate the radiative forcing that occurred
377 between 1985 and 1997. Our results show that the net RF of forest expansion ranges from -24 W/m² at
378 low elevations of the northern Prealps to 2 W/m² at high elevations of the Central Alps. The albedo RF
379 increases with increasing altitude, which offsets the CO₂ RF at high elevations with long snow-covered
380 periods, high global radiation and low carbon sequestration. Albedo RF is particularly relevant during
381 transitions from open land to open forest and not in later stages of forest development. Between 1985
382 and 1997, when overall forest expansion in Switzerland was approximately 4%, the albedo RF offset the
383 CO₂ RF by an average of 40%. We conclude that the albedo RF should be considered at an appropriately
384 high resolution when estimating the climatic effect of forestation in temperate mountainous regions.”

385

386

387

388

389 6. Major comment:

390 Assuming constant upward transmissivity in the radiative forcing calculation is a major
391 simplification. I would think the upward transmissivity would vary a lot over the elevation
392 gradient in this region. I appreciate that you have quantified the error associated with 30%
393 variance in this variable, but why not make an effort to include some real spatial information
394 here? I suspect this will exacerbate the elevational effects you are seeing. I think you could

395 use some archived high-resolution climate model data to develop a climatology of upward
396 transmissivity in the region and use that.

397
398 This is a very important point. The absorption will indeed vary if there is an elevation gradient (differences
399 in cloud cover etc.). We agree that a spatially explicit quantification would most likely increase the
400 elevational effect (as we have also stated in the discussion paper P 10143 L 6 – L 9). In high elevations the
401 “upward transmissivity” will usually be higher and the “upward absorption” will be lower. That means that
402 the value of 0.23 will be lower in high elevations. A lower value for the atmospheric absorption (see
403 equation (6)) will cause a higher Radiative Forcing in high elevations.

404 The simple approach of using a constant value for the absorption (or the upward transmissivity) has been
405 compared to a more complex radiative transfer model by Bright and Kvalevag (2013). They have shown
406 that the simple model performed well in comparison to the more complex model. Their result shows a
407 root mean square error of 7.2 % and a correlation of 0.93 between the forcings calculated by the simple
408 and the complex model. This is in line with our sensitivity analysis, where we assumed a high variation
409 (30%) in atmospheric absorption. However, in the sensitivity analysis we simplified matters, because we
410 assumed an independently varying atmospheric absorption. As we discussed, it will probably be linked to
411 the variation in global radiation.

412 Assuming constant albedo differences and constant global radiation, equation (6) simplifies to $I * \Delta\alpha * (1$
413 $- a)$. We use the following values:

- 414 1. $\Delta\alpha = 0.15$, $I = 176 \text{ W/m}^2$ and absorption $a=0.23$ (high elevation scenario); $\text{RF} = 20.3 \text{ W/m}^2$
- 415 2. $\Delta\alpha = 0.15$, $I = 176 \text{ W/m}^2$ and absorption $a=0.16$ (high elevation scenario); $\text{RF} = 22.2 \text{ W/m}^2$
- 416 3. $\Delta\alpha = 0.05$, $I = 120 \text{ W/m}^2$ and absorption $a=0.23$ (low elevation scenario); $\text{RF} = 4.6 \text{ W/m}^2$
- 417 4. $\Delta\alpha = 0.05$, $I = 120 \text{ W/m}^2$ and absorption $a=0.3$ (low elevation scenario); $\text{RF} = 4.2 \text{ W/m}^2$

418 This is a very simplified example. However, it illustrates that Radiative Forcing changes are small, even
419 though we used rather unrealistic high and low values for the atmospheric absorption (+/- 30%). If we use
420 the Fu/Liou online Radiative Transfer Model and calculate values of atmospheric absorption in low
421 elevation (high cloud cover fraction) and high elevation (low cloud cover fraction), we get values of
422 absorption of 0.25 and 0.19. Although the way we derive the absorption values from the Fu/Liou model is
423 a simplification, these values probably give a more realistic range of the absorption than the values 0.3
424 and 0.16. (Simplifications in the way we calculated the absorption: We used the output of the Fu/Liou
425 model together with a radiation model (Donohoe and Battisti, 2011) (assuming isotropic solar fluxes) to
426 calculate atmospheric absorption. The “isotropic assumption” is probably not exactly true and we calculate
427 the absorption without including multiple reflections.)

428 We thank you for your suggestion “to use some archived high-resolution climate model data to develop a
429 climatology of upward transmissivity in the region and use that”. This could be an option. However, using
430 data of a regional climate model for developing a climatology of upward transmissivity will also include
431 assumptions and simplifications. To our knowledge this has not been done yet (for regional models on a
432 higher resolution).

433 We acknowledge that a spatially explicit estimation would reduce uncertainty in our study. We extended
434 our discussion on this point in the manuscript. However, a spatially explicit estimation will not affect our
435 major results and findings.

436 Adapted paragraph in the discussion:

437 “The average parameter for atmospheric absorption “0.23” could be replaced by a spatially explicit
438 estimate. Including a spatially explicit parameter for atmospheric absorption would probably increase the
439 elevation gradient of RF, because atmospheric absorption should be higher in low elevations than in high
440 elevations. According to our sensitivity analysis and Bright and Kvalevag (2013) improving data on
441 atmospheric absorption will have a relatively small influence on the results.”

442

443 7. Minor comments:

444 p. 10126, line 21: change “biogeophysical (mainly albedo) and biogeophysical” to “biogeophysical
445 (mainly albedo) and biogeochemical”?

446 We changed it to biogeochemical.

447

448 8. Figure 3 captions are scrambled

449 We corrected the captions.

450

451 9. Equation 6, need to clarify whether the RF is at top of atmosphere (TOA) or at the surface. It should
452 be at the top of atmosphere.

453 We clarified that the RF is top of the atmosphere.

454

455 10. Also need to clarify whether the incoming global radiation data are for the surface, or TOA. It needs
456 to be at the surface, so that the incoming beam is already attenuated by clouds, aerosols etc. This
457 helps reduce the error associated with assuming a constant upwelling transmissivity over the
458 whole domain (although I hope you will address that problem separately).

459

460 We clarified that global radiation is the surface shortwave irradiance.

461

462

- 463 ALEWELL, C. & BEBI, P. 2011. Forest Development in the European Alps and Potential Consequences on
464 Hydrological Regime. In: BREDEMEIER, M., COHEN, S., GODBOLD, D. L., LODE, E., PICHLER, V. &
465 SCHLEPPI, P. (eds.) *Forest Management and the Water Cycle*. Springer Netherlands.
- 466 BARNES, C. A. & ROY, D. P. 2010. Radiative forcing over the conterminous United States due to
467 contemporary land cover land use change and sensitivity to snow and interannual albedo
468 variability. *Journal of Geophysical Research-Biogeosciences*, 115.
- 469 BLOETZER, G. 2004. Walderhaltungspolitik - Entwicklung und Urteil der Fachleute. In: BUNDESAMT FÜR
470 UMWELT, W. U. L. B. (ed.).
- 471 BRIGHT, R. M. & KVALEVAG, M. M. 2013. Technical Note: Evaluating a simple parameterization of
472 radiative shortwave forcing from surface albedo change. *Atmospheric Chemistry and Physics*, 13,
473 11169-11174.
- 474 DAVIN, E. L., DE NOBLET-DUCOUDRE, N. & FRIEDLINGSTEIN, P. 2007. Impact of land cover change on
475 surface climate: Relevance of the radiative forcing concept. *Geophysical Research Letters*, 34.
- 476 DONOHOE, A. & BATTISTI, D. S. 2011. Atmospheric and Surface Contributions to Planetary Albedo.
477 *Journal of Climate*, 24, 4402-4418.
- 478 GEHRIG-FASEL, J., GUISAN, A. & ZIMMERMANN, N. E. 2007. Tree line shifts in the Swiss Alps: Climate
479 change or land abandonment? *Journal of Vegetation Science*, 18, 571-582.
- 480 HAGEDORN, F., SHIYATOV, S. G., MAZEPA, V. S., DEVI, N. M., GRIGOR'EV, A. A., BARTYSH, A. A., FOMIN, V.
481 V., KAPRALOV, D. S., TERENT'EV, M., BUGMAN, H., RIGLING, A. & MOISEEV, P. A. 2014. Treeline
482 advances along the Urals mountain range – driven by improved winter conditions? *Global*
483 *Change Biology*, n/a-n/a.
- 484 HUMBEL, R., BEYELER, A., BURKHALTER, J., SAGER, J. & H.-U.ZAUGG. 2010. Arealstatistik nach
485 Nomenklatur 2004 - Standard NOAS04. Available:
486 [http://www.bfs.admin.ch/bfs/portal/de/index/dienstleistungen/geostat/datenbeschreibung/are](http://www.bfs.admin.ch/bfs/portal/de/index/dienstleistungen/geostat/datenbeschreibung/arealstatistik_noas04.html)
487 [alstatistik_noas04.html](http://www.bfs.admin.ch/bfs/portal/de/index/dienstleistungen/geostat/datenbeschreibung/arealstatistik_noas04.html) [Accessed 17.04.2010].
- 488 KOZAK, J. 2003. Forest cover change in the Western Carpathians in the past 180 years - A case study in
489 the Orawa Region in Poland. *Mountain Research and Development*, 23, 369-375.
- 490 MACDONALD, D., CRABTREE, J. R., WIESINGER, G., DAX, T., STAMOU, N., FLEURY, P., LAZPITA, J. G. &
491 GIBON, A. 2000. Agricultural abandonment in mountain areas of Europe: Environmental
492 consequences and policy response. *Journal of Environmental Management*, 59, 47-69.
- 493 RAMANKUTTY, N., HELLER, E. & RHEMTULLA, J. 2010. Prevailing Myths About Agricultural Abandonment
494 and Forest Regrowth in the United States. *Annals of the Association of American Geographers*,
495 100, 502-512.
- 496 ROESCH, A., WILD, M., PINKER, R. & OHMURA, A. 2002. Comparison of spectral surface albedos and their
497 impact on the general circulation model simulated surface climate. *Journal of Geophysical*
498 *Research*, 107, ACL13-1-ACL13-ACL13-18ACL13-ACL13-18.
- 499 RUTHERFORD, G. N., BEBI, P., EDWARDS, P. J. & ZIMMERMANN, N. E. 2008. Assessing land-use statistics
500 to model land cover change in a mountainous landscape in the European Alps. *Ecological*
501 *Modelling*, 212, 460-471.
- 502 ZHAO, K. G. & JACKSON, R. B. 2014. Biophysical forcings of land-use changes from potential forestry
503 activities in North America. *Ecological Monographs*, 84, 329-353.
- 504 ZHOU, L., DICKINSON, R. E., TIAN, Y., ZENG, X., DAI, Y., YANG, Z. L., SCHAAF, C. B., GAO, F., JIN, Y.,
505 STRAHLER, A., MYNENI, R. B., YU, H., WU, W. & SHAIKH, M. 2003. Comparison of seasonal and
506 spatial variations of albedos from Moderate-Resolution Imaging Spectroradiometer (MODIS) and
507 Common Land Model. *Journal of Geophysical Research-Atmospheres*, 108.

508

509

Carbon Storage Versus Albedo change: Radiative Forcing of Forest Expansion in Temperate Mountainous Regions of Switzerland

Jonas Schwaab^{1,5}, Mathias Bavay¹, Edouard Davin⁴, Frank Hagedorn², Fabia Huesler³, Michael Lehning^{1,6}, Martin Schneebeli¹, Esther Thürig², Peter Bebi¹

¹ WSL – Institute for Snow and Avalanche Research SLF, 7260 Davos, Switzerland

² Swiss Federal Institute for Forest, Snow and Landscape Research WSL, 8903 Birmensdorf, Switzerland

³ Institute of Geography and Oeschger Center for Climate Change Research, University of Bern, Bern, Switzerland

⁴ Institute for Atmospheric and Climate Sciences, ETH Zürich, 8092 Zürich, Switzerland

⁵ Institute for Spatial and Landscape Planning, ETH Zürich, 8093 Zürich, Switzerland

⁶ CRYOS, School of Architecture, Civil and Environmental Engineering, EPFL, 1015 Lausanne, Switzerland

Abstract

In this study, we assess the climate mitigation potential from afforestation in a mountainous snow-rich region (Switzerland) with strongly varying environmental conditions. Using radiative forcing calculations, we quantify both the carbon sequestration potential and the effect of albedo change at high resolution. We calculate the albedo radiative forcing based on remotely sensed datasets of albedo, global radiation and snow cover. Carbon sequestration is estimated from changes in carbon stocks based on National Inventories. We first estimate the spatial pattern of RF across Switzerland assuming homogeneous transitions from open land to forest. This highlights where forest expansion still exhibits climatic benefits when including the radiative forcing of albedo change. Second, given that forest expansion is currently the dominant land-use change process in the swiss Alps, we calculate the radiative forcing that occurred between 1985 and 1997. Forestation is seen as a possible option to counter climate change by sequestering carbon in forests and thus reducing the atmospheric concentration of carbon dioxide. However, previous studies suggest that the Radiative Forcing (RF) caused by forestation induced albedo change in snow-rich boreal regions may offset the carbon sequestration effect. The Swiss mountains are characterized by snow-rich areas with strongly varying environmental conditions and forest expansion is currently the dominant land-use change process. Thus, quantifying both carbon sequestration and albedo change on appropriately high resolution in this region will improve our understanding of the forests potential for climate mitigation. We calculated the albedo RF based on remotely sensed datasets of albedo, global radiation and snow cover. Carbon sequestration was estimated from changes in carbon stocks based on National Inventories.

Our results show that the net RF of forest expansion ranges from -24 W/m^2 at low elevations of the northern Prealps to 2 W/m^2 at high elevations of the Central Alps. The albedo RF increases with increasing altitude, which offsets the CO_2 RF at high elevations with long snow-covered periods, high global radiation and low

carbon sequestration. ~~Results indicate that the albedo RF is particularly relevant during transitions from open land to open forest and not in later stages of forest development. Albedo RF is particularly relevant during transitions from open land to open forest and not in later stages of forest development. Between 1985 and 1997, when overall forest expansion in Switzerland was approximately 4%,~~ the albedo RF offset the CO₂ RF by an average of 40%.

We conclude that the albedo RF should be considered at an appropriately high resolution when estimating the climatic effect of forestation in temperate mountainous regions.

Introduction

The United Nations Framework Convention on Climate Change (UNFCCC) declared in the Kyoto Protocol (Decision 11/CP.7) that changes in the carbon stocks of ecosystems, induced by LULUCF (Land Use, Land Use Change and Forestry) activities can be included in the greenhouse gas emission budget of the signatory nations (UNFCCC, 2001). Beside biogeochemical processes, LULUCF also influences biogeophysical processes (Betts, 2011, Bonan, 2008), but these effects are not yet considered in current climate policies.

Global climate models suggest that biogeochemical and biogeophysical effects vary greatly with latitude (Schaeffer et al., 2006, Bala et al., 2007, Bathiany et al., 2010, Davin and de Noblet-Ducoudre, 2010, Arora and Montenegro, 2011). In the tropics, biogeochemical and biogeophysical effects tend to act in the same direction, since tropical forests cool climate through both evaporative cooling and carbon sequestration (Costa and Foley, 2000, Gibbard et al., 2005). However, at middle and high latitudes, biogeophysical processes tend to counter the ~~biogeophysical-biogeochemical~~ effect (Gibbard et al., 2005, Betts, 2000, Govindasamy et al., 2001), thus making the net LULUCF effect more challenging to assess. Indeed, forestation in boreal regions lowers the albedo and thus counterbalances the cooling effect of carbon storage. Global climate models are important for understanding the climatic processes related to LULUCF and for quantifying the impacts on climate of forestation or deforestation over large areas. However, in highly heterogeneous landscapes such as mid-latitude mountain ranges, global climate models are limited by their relatively coarse resolution, and the concept of Radiative Forcing (RF) (Myhre et al., 2013) can provide a useful alternative. The RF concept has already been employed to investigate the balance between biogeophysical (mainly albedo) and ~~biogeophysical-biogeochemical~~ effects following forestation or deforestation. While some studies suggest that albedo RF can completely offset CO₂ RF (Betts, 2000, Bernier et al., 2011, de Wit et al., 2013), others have found that the offset is rather small (e.g. (Montenegro et al.,

2009, Kirschbaum et al., 2011). The offset seems to vary widely depending on the regional characteristics of the determining variables: global radiation, snow-cover, albedo change and carbon sequestration. These factors vary greatly, but little research has focused on how each of them influences RF (Kirschbaum et al., 2011), and only few attempts have been made to quantify RF in a spatially explicit way (e.g. Betts, 2000, Montenegro et al., 2009). Betts (2000) estimated spatially explicit RF data on a resolution of 3.75° longitude by 2.5° latitude. Montenegro et al (2009) performed their analysis on a resolution of 5 to 25 km. However, RF varies on much smaller scales. Moreover, decisions in regional planning are usually based on very local and regional information. Consequently, it is crucial to quantify RF at finer resolutions.

Forest cover has increased in many temperate mountainous regions (Alewell and Bebi, 2011, MacDonald et al., 2000, Ramankutty et al., 2010, Kozak, 2003, Hagedorn et al., 2014) and analysis of LULUCF change in the Alps suggest that changes in forest cover near the tree-line will further increase (Gehrig-Fasel et al., 2007). Some of the effects of this increase on various ecosystem services are already relatively well known (MacDonald et al., 2000, Laiolo et al., 2004, Bolliger et al., 2008) and have increasingly been considered in management strategies and subsidizing systems for agriculture and forestry (e.g. Gret-Regamey et al., 2013). However, few attempts to quantify RF in such regions have been made, even though snow-cover and hence albedo in temperate mountains varies greatly. To optimize the effects of future land-use decisions, further research on the climatic impacts of forests is essential and should be included in spatially explicit valuation methods (Bebi et al., 2012).

Switzerland is a particular suitable study area for researching the effects of changes in forest cover, since ~~forest expansion is an ongoing dominant process of land-use change and~~ many spatially explicit high quality datasets are available and forest expansion is an ongoing dominant process of land-use change. ~~Forest cover is expanding by 4% per decade at the~~

~~country scale and by 8% per decade in alpine areas (NFI).~~ The spatially explicit data series available are on land use/land cover (1 ha raster), snow cover (1 km), global radiation (2.2 km), albedo change and carbon sequestration (both explicit for biogeographical regions) for the whole area of Switzerland. Forest cover is expanding by 4% per decade at the country scale and by 8% per decade in alpine areas (NFI).⁷

Our study design is twofold: First, we use the spatially explicit datasets to show the pattern of RF assuming that each location in Switzerland is facing a transition from agriculturally used open land to forest. This is not related to any particular or realistic scenario, however, the spatial pattern of RF can be of high interest for any land-use policies steering forest cover change towards desired futures. In Switzerland agricultural subsidies directly influence farmers decisions on whether to keep managing or abandon their land. The latter will usually result in forest expansion. Second, we include the type and location of 5 different land use transitions to calculate RF in Switzerland between 1985 and 1997. In summary we estimate i) to which extent albedo RF offsets CO₂ RF in different parts of temperate mountainous regions, ii) how each input parameter influences RF, and iii) what the inclusion of albedo change implies for the greenhouse gas inventory in Switzerland. ~~to estimate i) to which extent albedo RF offsets CO₂ RF in different parts of temperate mountainous regions, ii) how each input parameter influences RF, and iii) what the inclusion of albedo change implies for the greenhouse gas inventory in Switzerland.~~

Data and Methods

Study Area

Switzerland covers an area of 41284 km² and can be divided into five biogeographical regions (Figure 1). Each region has biogeographical features that can be found globally at temperate latitudes:

1. Jura, oceanic low mountain range, elevations averaging 800 m a.s.l.; 2. Plateau, oceanic lowlands, elevations averaging 550 m a.s.l.; 3. Northern Prealps, oceanic subalpine mountain range, elevations averaging 1400 m a.s.l.; 4. Central Alps, continental alpine mountain range with elevations averaging 2150 m a.s.l.; 5. Southern Prealps, mediterranean/insubric subalpine mountain range, elevations averaging 1500 m a.s.l.. In each region, deciduous forests and mixed deciduous forest dominate at low elevations (mostly *Fagus sylvatica*), while coniferous forests are dominant at higher elevations (mostly *Picea abies*).

The Swiss landscape was strongly affected by several centuries of intensive human land-use (Bürgi and Schuler, 2003, Schneeberger et al., 2007, Gimmi et al., 2009) followed by the widespread abandonment of marginal agricultural land and the subsequent expansion of forest cover since the end of the 19th century (Baur, 2006). Due to unfavorable pedologic and climatic conditions and high slope angles, marginal land and forest expansion are mainly found at higher elevations (Baur, 2006). Land abandonment was the most dominant driver for the establishment of new forest areas, however, a small fraction of forest expansion at the tree line can be attributed to the recent climate warming (Gehrig-Fasel et al., 2007). Tree line is often the result of former land use, but also depends on various climatic factors and is generally higher in the Central Swiss Alps (approx. 2100-2300 m) than in the Northern and Southern Prealps (approx. 1800-2000 m) (Figure 3f and Körner, 2012).

Land use/Land Cover (LULC) change

~~We use aerial photographs processed by Swiss Statistics at a spatial resolution of 100 m to derive changes in land use. These aerial photographs are from the Swiss Federal Office of Topography and are fully available for the two inventory periods 1979-85 and 1992-97 (Humbel et al., 2010). We reclassified the data of the different inventory periods into five aggregated classes (Rutherford et al., 2008). While 18 classes were aggregated into four classes: Closed Forest, Open Forest, Extensively and Intensively Used Open Land (Figure 2), the remaining 56 were classified as Other, and consisted mainly of settlements, water and non-vegetated land.~~

~~To calculate RF of land use change between 1985 and 1997, we included five transitions: 1. Intensively Used Open Land \rightarrow Closed Forest, 2. Extensively Used Open Land \rightarrow Closed Forest, 3. Intensively Used Open Land \rightarrow Open Forest, 4. Extensively Used Open Land \rightarrow Open Forest and 5. Open Forest \rightarrow Closed Forest. We focused on transitions where forest cover and carbon stocks increase, because these transitions highly exceeded transitions with forest decrease in Switzerland. In fact, the Swiss law strongly protects forests so that there have been only few changes from forest to agriculturally used land during the last 30 years (Bloetzer, 2004, Rutherford et al., 2008).~~

Spatial variability of RF and RF of Swiss forest expansion

We calculated the net RF and the offset of CO₂ RF through albedo RF ($\Delta RF_{CO_2}/\Delta RF_{albedo}$) to show the spatial variability of RF in Switzerland and to calculate RF of Swiss forest-cover expansion between 1985 and 1997. To illustrate the **pattern** spatial variability of RF in Switzerland we calculated a value of RF for every location in Switzerland, excluding non-vegetated land, water, settlement and areas that lie above the tree line (Figure 3). These are hypothetical values, because we calculated RF for the change from open land to forest for all

vegetated areas, and not only for the ones where forest expansion was actually observed. At lower elevations, transitions from Intensively Used Open Land to forest are frequent, while in higher elevations transitions are almost exclusively from Extensively Used Open Land to Forest are most likely (Appendix, Table 4). We considered this by separating our estimation of the hypothetical RF in transitions from Intensively Used Open Land to forest below 1000 m and transitions from Extensively Used Open Land to Forest above 1000 m. The results of the spatial variability of RF are shown in maps of the study area (Figure 4 c,d).

To obtain results for RF of forest-cover expansion between 1985 and 1997 in Switzerland we calculated net Radiative Forcing as the sum of RF for all pixels where forests expanded. This meant including information on the type of forest expansion and on the location of forest expansion:

$$\underline{\underline{\Delta RF_{ges} = \frac{\sum_{x=1}^n RF_{x,T}}{A_E}}} \quad (1)$$

, where ΔRF_{ges} is the net Radiative Forcing (net RF), n the number of pixels where forests expanded and RF the Radiative Forcing, which depends on the location x and the type of transition T . The sum over RF is divided by the the earth's surface A_E to convert local RF into a global average RF.

Swiss forest expansion between 1985 and 1997

We use aerial photographs processed by Swiss Statistics at a spatial resolution of 100 m to derive changes in land use/land cover (LULC). These aerial photographs are from the Swiss Federal Office of Topography and are fully available for the two inventory periods 1979-85 and 1992-97 (Humbel et al., 2010). We reclassified the data of the different inventory periods into five aggregated classes (Rutherford et al., 2008). While 18 classes were aggregated into four classes: Closed Forest, Open Forest, Extensively and Intensively Used Open Land (Figure 2), the remaining 56 were classified as Other, and consisted mainly of settlements, water and non-vegetated land (Appendix, Table 5). The aggregation of the original land-use classes results in a sufficiently large sample of reliable albedo values and carbon stocks in each of the five biogeographical regions and three elevational strata for five relevant and well established land-use classes.

To calculate RF of land use change between 1985 and 1997, we included five transitions: 1. Intensively Used Open Land -> Closed Forest, 2. Extensively Used Open Land -> Closed Forest, 3. Intensively Used Open Land -> Open Forest, 4. Extensively Used Open Land -> Open Forest and 5. Open Forest -> Closed Forest. Forest expansion mainly took place in elevations above 1200 m in the Prealps and the Central Alps (Appendix, Figure 1). We focused on transitions where forest cover and carbon stocks increase, because these transitions highly exceeded transitions with forest decrease in Switzerland. In fact, the Swiss law strongly protects forests so that there have been only few changes from forest to agriculturally used land during the last 30 years (Bloetzer, 2004, Rutherford et al., 2008).

CO₂ RF

An increase in carbon stocks in terrestrial ecosystems is related to a sink of atmospheric CO₂, followed by a change in the earth's radiation balance. Myhre et al (1998) developed a parameterization to derive RF, related to a change in the CO₂ concentration in the atmosphere, based on radiative transfer schemes:

$$\Delta RF(t) = 5.35 \ln \frac{C(t)}{C_0} \quad (2)$$

, where $\Delta RF(t)$ is the Radiative Forcing, $C(t)$ is the atmospheric CO₂ concentration after perturbation and C_0 is the unperturbed atmospheric CO₂-concentration. Equation (2) can be solved if the CO₂ concentration in the atmosphere after perturbation is known.

Following Switzerland's Greenhouse Gas Inventory 1990-2010 (Heldstab et al., 2012), which is based on the Good Practice Guidance for Land Use, Land Use Change and Forestry (IPCC, 2003), we calculated the carbon stock changes resulting from land-use changes as the differences between the carbon stocks of the land-use categories before and after a transition. This takes into account changes in living plant biomass, dead wood and soil carbon stocks.

Data on carbon stocks in the living biomass and dead wood of Closed Forests and Open Forests was derived from the third National Forest Inventory (NFI) (Brändli, 2010). The data is based on 6608 field plots measured on a regular sampling 1.4 x 1.4 km grid from 2004 to 2006. Two concentric circles with 200 m² in size are used for trees with 12 cm ≤ diameter at breast height (DBH) < 36 cm, and 500 m² in size for trees with DBH ≥ 36 cm. This results in DBH measurements of approximately 11 trees per plot. On a sub-sample of approximately two trees per plot, the diameter at 7 m tree height and the tree height are measured. The biomass of all single trees is estimated according to allometric functions. The assessment of stem-wood

over bark, including stock, coarse branches (≥ 7 cm) and small branches (< 7 cm) is based on Kaufmann (2001), and that of needles/leaves on [Perruchoud et al. \(1999\)](#). Roots are estimated with equations from Wirth et al. (2004) for coniferous trees and Wutzler et al. (2008) for deciduous trees. Estimates for branches, foliage and roots were derived from the DBH only, while for stem-wood over bark including stock the diameter at 7 m tree height and the total tree height were also required (Kaufmann, 2001). For this study, only living trees were considered. The NFI data was stratified for each biogeographical region of Switzerland and three elevation strata ([Appendix, Table 2](#)). Open Forest is represented by forest plots permanently reduced in stocking, and Closed Forest by all forest plots minus plots permanently and temporarily reduced in stocking.

The estimates of soil carbon stocks were taken from previous assessments in approximately 1000 soil profiles in forests (Hagedorn et al., 2010, Nussbaum et al., 2012) and 500 soil profiles in open land across Switzerland (Leifeld et al., 2005, Bolliger et al., 2008), where each soil profile in the forest had been sampled according to horizons analyzed for their C content using a C/N analyzer. The bulk density and volumetric stone content measured, were used to calibrate pedotransfer functions. The carbon stock changes in soils were calculated as the differences between the stocks of different LULC classes. Since soil carbon stocks in agriculturally used open land and forest correlate fairly well across different altitudes (Sjörgersten-Turner et al., 2011), we decided to use the difference between mean soil carbon stocks in open land and forest for the whole Switzerland.

Changes in carbon stocks in ecosystems can be related to changes in atmospheric CO₂ concentrations by including the molecular masses of carbon, dry air and the mass of the atmosphere (Schwaiger and Bird, 2010, O'Halloran et al., 2012):

$$\Delta C_A = \frac{\Delta m_C M_a}{M_C m_a} \quad (3)$$

, where ΔC_A is the change in atmospheric CO₂ concentration, Δm_C ~~the mass of carbon sequestered in ecosystems (assuming an equilibrium before and after a change in LULC) the difference between carbon stocks of two LULC classes~~, M_C is the molecular mass of carbon, m_a is the mass of the atmosphere and M_a is the molecular mass of dry air. Before using this CO₂ value (ΔC_A) to calculate ΔRF (equation (2)), we took into consideration the fact that each CO₂ pulse emitted to the atmosphere disappears partly in sinks of the global carbon cycle (e.g. ocean). A carbon pulse response function can be used to describe such fluxes (Forster et al., 2007):

$$fr(t) = a_0 + \sum_{j=1}^3 a_j e^{-\frac{t}{\tau_j}} \quad (4)$$

, where $fr(t)$ is the fraction of a CO₂ pulse, which can still be found in the atmosphere after time t . Generally the coefficients a_0 to a_3 and τ_1 to τ_3 have no direct process-based meaning, but are fitting parameters chosen to represent a given model-based carbon-pulse response function (Joos et al., 2013). CO₂ sequestration equals negative CO₂ pulses (CO₂ removal from atmosphere). The carbon-pulse response function can be applied to negative pulses because a reduced CO₂ concentration in the atmosphere will reduce the amount of CO₂ sequestered by the terrestrial biosphere and oceans. To estimate how much carbon dioxide from continuous CO₂ pulses after time t stays airborne, a widely used convolution function can be applied (Siegenthaler and Oeschger, 1978, Cherubini et al., 2011):

$$C(t) = \int_0^t g(t') y(t-t') dt' \quad (5)$$

, where $C(t)$ stands for the yearly change in the carbon dioxide in the atmosphere that can be exclusively related to carbon sequestration during forest expansion, $g(t')$ characterizes the carbon sequestration due to forest expansion, depending on the gradient of CO₂ uptake during succession and $y(t-t')$ accounts for the reduced uptake of carbon dioxide in the carbon cycle. We estimate the integral using a simple numerical approximation and time intervals of 1 year:

$$C_t = \sum_{i=1}^t g_i y_{t-i} \quad (6)$$

We kept the background CO₂-concentration fixed to solve the radiative transfer parameterization (equation (2)) and accordingly used parameters that describe carbon sequestration in the carbon cycle for a fixed CO₂ background concentration. The CO₂ concentration and parameters for the carbon cycle will, however, change (IPCC, 2001) and it is not clear whether it is necessary to take these changes into account. Joos et al. (2013) showed that Radiative Forcing was more or less constant when CO₂ pulses were emitted to atmospheres with different CO₂ background concentrations. They suggest that not only does the carbon uptake per unit atmospheric CO₂ decrease with a high background concentration of CO₂, but so too does the RF per unit change in atmospheric CO₂. They both decrease in such a way that RF of a CO₂-emission is almost identical at preindustrial and present day conditions. Keeping background CO₂ concentrations and carbon cycle parameters constant thus seems a reasonable approximation.

Albedo RF

The yearly Radiative Forcing $\Delta RF(t)$ at the top of atmosphere of an albedo change is calculated (modified from Montenegro et al., 2009) as the seasonal average:

$$\Delta RF(t) = \frac{1}{12} \sum_{i=1}^{12} I_i \Delta \alpha_i - 0.23 I_i \Delta \alpha_i \quad (7)$$

, where I_i is the monthly global radiation at the earth surface, $\Delta \alpha_i$ is the albedo difference between two LULC classes depending on the monthly fraction of snow-cover and 0.23 is a factor to account for the absorption of the reflected radiation in the atmosphere. The first part of equation (7) describes which part of the global radiation is absorbed at the surface. The second part describes which part of the reflected shortwave radiation is absorbed in the atmosphere. The factor of 0.23 is that used by Montenegro et al. (2009) who modified the value 0.3 given by Weaver et al. (2001).

We used gridded global radiation (i.e. surface shortwave irradiance) data from MeteoSwiss (MeteoSwiss, 2012) in monthly datasets averaged over the period from 2004 to 2010 to eliminate inter-annual variability. The spatial resolution of the global radiation datasets is 2.2 km. The derivation of the global radiation data was based on the Heliosat method (Cano et al., 1986, Beyer et al., 1996, Hammer et al., 2003), applied to Meteosat SEVIRI data. It was verified using high-quality surface measurements and sensitivity runs for key input parameters (Durr et al., 2010). The albedo was estimated using the following equation (modified from Barnes and Roy, 2010, Roesch et al., 2002):

$$\Delta \alpha(t) = f(t) \Delta \alpha_s + (1 - f(t)) \Delta \alpha_v \quad (8)$$

, where $\Delta\alpha(t)$ is the monthly albedo-difference between two LULC classes, $\Delta\alpha_s$ the average albedo difference between two LULC classes when snow-covered, $\Delta\alpha_v$ the average albedo difference between two LULC classes when snow-free and $f(t)$ the fraction of snow-cover per month. We used average albedo differences of snow-free and snow-covered albedo differences and not monthly differences because of two reasons.. First, by far the strongest seasonal trend is related to the presence of snow, which we explicitly included (Zhou et al., 2003). Second, in some months reliable albedo data was missing and we considered the average to be a robust estimate. Since we found that the seasonal variation of the albedos of different LULC classes is similar, the averaging of snow-covered and snow-free albedo differences results in a fairly good approximation (Appendix Figure 2, Appendix Figure 3).~~Thus, seasonally averaging snow-covered and snow-free albedo differences results in a fairly good approximation.~~

Monthly datasets on snow-cover were provided by the Remote Sensing Research Group at the University of Bern. Raw 1-km data from AVHRR (Advanced Very High Resolution Radiometer) was processed using an algorithm to estimate the snow cover (SPARC - Separation of Pixels using Aggregated Rating Over Canada). The algorithm was adapted to the mountainous region of the European Alps and verified (Huesler et al., 2012). Again we calculated an average value for each month using data of the years 2002 – 2009 to eliminate inter-annual variability.

For assigning albedos to different LULC classes, we overlaid NASA's MODerate-resolution Imaging Spectroradiometer (MODIS) 0.5 km BRDF/Albedo Product - MCD43A and gridded 0.1 km LULC data from the Swiss Area Statistics aggregated into five categories (see [Swiss forest expansion between 1985 and 1997](#)). We used MODIS data between 2004 and 2009 and the Area Statistics from the third inventory period 2004-2009 to ensure a temporal overlap between the albedo and the LULC records. Since not all the LULC data for south-eastern

Switzerland is available yet, we complemented the Area Statistics (2004-2009) with data from the second inventory period (1992-1997) and accepted a temporal displacement for this region.

We applied several different methods for calculating land use specific albedos. First, we retained albedo pixels of the highest quality (full BRDF inversion) for our analysis. To reduce the error caused by assigning albedo values to a mixed pixel (several 0.1 km LULC classes in one 0.5 km albedo pixel), we only assigned albedo values to a specific LULC class if at least 92% of the albedo pixel were covered by just one LULC class (similar to Kvalevag et al. (2010)). The threshold of 92% (23 out of 25 pixels) is a trade-off between using as many albedo pixels in the study area as possible and at the same time reducing the error due to 8% random land cover (2 out of 25 pixels). If not enough pixels covered by at least 92% of one LULC class were available, we applied additional methods to calculate specific albedos, accepting the trade-off of using more pixels, but with less quality. For the first step, we included albedo values of lower quality produced with magnitude inversion (Liu et al., 2009). If still not enough pixels were available, we changed the cell size of the Area Statistics to 0.5 km (according to the most frequent LULC class within a pixel) and assigned albedo values to this new LULC dataset. To estimate Closed Forest albedos, we only used pixels of at least 92% land cover and best quality (full BRDF inversion). Open Forest values were mainly calculated using lower quality values (magnitude inversion) and the resampled pixels ([Appendix](#), Table 1). For Extensively and Intensively Used Open Land, we used the average albedo values for the whole study area because it was not possible to derive specific values for each region (For example, Extensively Used Open Land hardly occurs below 600 m a.s.l. and Intensively Used Open Land hardly occurs above 1200 m a.s.l.). We accepted a bias of open land albedos in these regions since they were usually not important for LULC change. For instance, LULC change below 600 m involves almost exclusively Intensively Used Open Land.

The MCD43A product is online available for free. It provides atmospherically corrected gridded albedo data for a variety of spectrums. We used broadband white-sky albedo (0.3-5.0 μm). In order to distinguish between snow-covered and snow-free areas, we applied the quality flags of the MODIS product MCD43A2. The albedo product MCD43A3 has been produced applying the MODIS BRDF/Albedo algorithm (Strahler et al., 1999). This algorithm makes use of 16 days worth of multi-date data from both the Terra and Aqua platforms and a semiempirical kernel-driven bidirectional reflectance model to determine a global set of parameters describing the BRDF (Bidirectional Reflectance Distribution Function) of the land surface (Schaaf, 2010).

Temporal signature of RF

Both, albedo RF and CO₂ RF are a function of time. The annual variation of CO₂ RF depends on the carbon pulse response function and the yearly carbon sequestration in biomass and soil during succession. The annual variation of albedo RF depends on the albedo change during succession. For both, carbon sequestration and albedo change we had to rely on the static difference between land-use classes e.g. open land (as starting point) and closed forests (end of succession). We assumed that carbon sequestration as well as albedo change follow linear trends until they reach an approximately steady state. Since a detailed description of the temporal evolution of albedo-change and carbon sequestration is complex and varies with location, we used a simplified scheme in which we assumed that albedo change completes after 30 years, carbon sequestration in biomass after 50 years and carbon sequestration in soils after 100 years. This seems to be a reasonable approximation since albedo change is likely to end before carbon accumulation in biomass does (Kirschbaum et al., 2011, de Wit et al., 2013) and carbon sequestration in soils will most likely not end before 100 years (Poeplau et al., 2011). We assumed that albedo-change and carbon sequestration stop after a certain time. However,

interactions with the global carbon cycle will still cause changes in the atmospheric CO₂-concentration. Thus, we compared the temporal mean of Radiative Forcing for two different time horizons, 100 and 1000 years. For the representation of our results, we chose the mean RF in 100 years. The temporal average of RF is not only useful when representing spatial variability, but also for a comparison with earlier studies on spatial variability of RF, which did not explicitly include temporal variation (Betts, 2000, Montenegro et al., 2009).

Spatial variability of RF and RF of Swiss forest expansion

~~We calculated the net RF and the offset of CO₂ RF through albedo RF (ARF_{CO_2}/ARF_{albedo}) to show the spatial variability of RF in Switzerland and to calculate RF of Swiss forest cover expansion between 1985 and 1997. To illustrate the spatial variability of RF in Switzerland we calculated a value of RF for every location in Switzerland, excluding non-vegetated land, water, settlement and areas that lie above the tree line (Figure 3). These are hypothetical values, because we calculated RF for the change from open land to forest for all vegetated areas, and not only for the ones where forest expansion was actually observed. At lower elevations, transitions from Intensively Used Open Land to forest are frequent, while in higher elevations transitions are almost exclusively from Extensively Used Open Land to Forest. We considered this by separating our estimation of the hypothetical RF in transitions from Intensively Used Open Land to forest below 1000 m and transitions from Extensively Used Open Land to Forest above 1000 m. The results of the spatial variability of RF are shown in maps of the study area (Figure 4 e,d).~~

~~To obtain results for RF of forest cover expansion between 1985 and 1997 in Switzerland we calculated net Radiative Forcing as the sum of RF for all pixels where forests expanded.~~

~~This meant including information on the type of forest expansion and on the location of forest expansion:~~

$$\Delta RF_{ges} = \frac{\sum_{x=1}^n RF_{x,T}}{A_E} \quad (8)$$

~~, where ΔRF_{ges} is the net Radiative Forcing (net RF), n the number of pixels where forests expanded and RF the Radiative Forcing, which depends on the location x and the type of transition T . The sum over RF is divided by the the earth's surface A_E to convert local RF into a global average RF.~~

Sensitivity Analysis

In a sensitivity analysis we tested how the spatial variability and uncertainty of each input factor influence our results. We based the sensitivity analysis on FAST (Fourier amplitude sensitivity test), developed by Saltelli et al. (1999), and used an implementation (fast99) provided in the R package “sensitivity” (Pujol et al., 2012). Applying a sensitivity analysis with FAST allowed us to show how varying input factors influenced the variance of the output including first order effects and interactions for each parameter. We approximated input as either uniform or normal distributed, according to the distribution of each input factor in the study area. We assumed input factors were uncorrelated, which only holds to a certain degree since e.g. all factors either increase or decrease with elevation. We separated our sensitivity analysis into two parts. First, we applied data on spatial variability. Each of the factors, carbon sequestration, snow-cover, global radiation, albedo difference (snow-covered) and albedo difference (snow-free) vary spatially and temporally. We averaged every input factor temporally and applied the spatial minimum and maximum of each factor to the function fast99 ([Appendix, Table 3](#)). The minimum and maximum of snow-covered albedo differences are e.g. 0.208 (Alps 600-1200) and 0.375 (Jura >1200). This analysis showed which factor had the greatest influence on the variation in RF caused by change in forest cover at a specific location. In a second sensitivity analysis, we applied the uncertainty values of each factor (~~for uncertainties, see~~ [Appendix Table 3](#)). The sensitivity analysis of uncertainties, represented by random sampling and measurement errors, indicates which parameter causes high or low uncertainty in the output.

Results

In the forest, average C stocks in biomass ranged from 95 to 170 tC/ha (NFI). In the soil (mineral soil 0-100 cm + organic layer) the average across Switzerland was 143 tC/ha (Nussbaum et al., 2012). In the Intensively Used Open Land, the average value for biomass was 4.34 tC/ha (Heldstab et al., 2012) and 91 tC/ha in the soils. In comparison, the C stocks of the Extensively Used Open Land amounted to 7 tC/ha in the biomass and 63 tC/ha in the soil. Minimum and maximum carbon sequestration in Switzerland of transitions from Intensively or Extensively Used Open Land to Closed Forest thus ranged from 143 to 241 tC/ha (Figure 3 e). This corresponded to a CO₂ RF of -16 to -27 W/m² (Figure 4 b).

The albedos of snow-covered Closed Forest ranged from 0.168 to 0.267, while those of snow-free Closed Forests ranged from 0.101 to 0.139. Similarly, the albedos of snow-covered Open Forest ranged from 0.217 to 0.307 and those of snow-free Open Forests from 0.117 to 0.141 (Appendix, Table 1). The average albedo of Intensively Used Open Land was 0.170/0.475 (snow-free/snow-covered) while that of Extensively Used Open Land was 0.154/0.549 (snow-free/snow-covered). The albedo differences between transitions from Intensively or Extensively Used Open Land to Closed Forest thus ranged from 0.208 to 0.375 (snow-covered) and from 0.023 to 0.066 (snow-free) (Figure 3c,d). Albedo change in Switzerland caused albedo RF ranging from 2 to 21 W/m² (Figure 4a).

The net RF at different locations in the study area ranged from -24 W/m² to 2 W/m² and the offset of CO₂ RF caused by albedo RF differed between 11 and 109%. The differences were particularly marked on an elevational gradient and between the 5 bioregions. Below an elevation of 600 m a.s.l., the albedo RF offset CO₂ RF by 15 % on average. Between 600 and 1200 m, the offset was 22 % and above 1200 m 54 %. The highest RF was observed in high snow-rich alpine regions. In the Central Alps, 13% of all possible LULC areas had an offset of more than 80%. In the Southern Prealps 7% of all possible LULC areas had an offset higher

than 80%. In the Central Alps as well as in the Southern Prealps we found areas where net RF was positive. However, these areas amounted to less than 0.2% of both regions. We only found positive RF at elevations above 1850 m. The albedo RF was lowest in the Plateau region and at low elevations of the Jura. Albedo RF accounted, on average, for less than 14% of the CO₂ RF in these regions. The lowest net RF (average -20 W/m²) was found in the Northern Prealps below 1200 m. Above 1200 m the net RF in the Northern Prealps was lower than in Southern Prealps and Central Alps, although the snow-cover in the Northern Prealps was persistent. However, the effect of a persistent snow-cover in the Northern Prealps was outweighed by the low global radiation, the low tree line and the high carbon sequestration in this alpine region (see Figure 3).

The net RF forcing of forests across the mountainous terrain in Switzerland strongly depended on the elevation (Figure 5). The albedo RF increased with altitude, with several factors contributing to this increase (Figure 3). First, global radiation increased with altitude, reaching maxima in continental parts of the Alps. Second, albedo (snow-free, snow-covered) of forests was in general lower above 1200m a.s.l. (where coniferous species dominate). Third, both snow cover and snow cover period strongly increased with altitude. The carbon sequestration potential of forests decreased with altitude (due to unfavorable climatic conditions towards tree line).

Radiative Forcing depended not only on location, but also on the type of LULC transition. LULC transitions from Open Land to Open Forest had a higher offset (approx. 80%) than transitions from already established Open Forest to Closed Forest (approx. 40 %.). ~~Transitions from Open Forest to Closed Forest were generally associated with relatively high amounts of carbon sequestration.~~ Transitions from Open Forest to Closed Forest were generally associated with relatively high change in carbon stocks (around 70 tC/ha), and relatively small albedo change (for snow-covered albedo < 0.06 and for snow-free albedo < 0.01).

Between 1985 and 1997, all five types of LULC transitions (Figure 2) took place on an area of 24000 ha. More than 70% of them were situated above 1000 m. Above 1000 m, the most frequent transition was the one from Open to Closed Forest (50%), followed by transitions from Extensively Used Open Land to Open Forest (20%) and from Extensively Used Open Land to Closed Forest (13%). Land use change occurred especially in high altitude regions where albedo RF strongly offsets CO₂ RF (Figure 5). At the same time, the most frequent transition was the one from Open Forest to Closed Forest, where albedo RF had the least influence on net RF. In summary, the CO₂ RF for all land use transitions that were part of the forest expansion and succession between 1985 and 1997 in Switzerland was reduced by approximately 40%, if the albedo RF is taken into account.

The net RF varied greatly on small spatial scales. In our study area, the parameter that had the most significant influence on the spatial variability of RF was snow-cover, followed by carbon sequestration, difference in snow-covered albedo, global radiation and difference in snow-free albedo (Figure 6a). Thus, if the average snow-cover (in days/year) and carbon sequestration at a certain location were known, the net RF and the offset of CO₂ RF through albedo RF could be estimated well without including all factors in an explicit calculation. We found that the offset of CO₂ RF can only be higher than 50% if the snow-cover lasted over 120 days/year. An increase in snow-cover caused an exponential increase in albedo RF. Because, first, persistent snow-cover was found in regions with high global radiation and second because the longer snow covered the surface, the longer it would be present during the days with high global radiation in March, April and May.

The uncertainty of net Radiative Forcing was mainly attributed to uncertainties in carbon sequestration, followed by snow-covered albedo difference, global radiation and snow-free albedo-difference. The main effects of uncertainty related to each input factor were much more relevant than the contribution of interactions between all input factors (Figure 6).

Discussion

Our detailed assessment of Radiative Forcing across Switzerland shows that the albedo RF increases with increasing altitude, which offsets the CO₂ RF at high elevations with long snow-covered periods, high global radiation and low carbon sequestration. The altitudinal RF gradient in Switzerland is very strong in comparison to the latitudinal gradient in boreal regions (Betts, 2000, Montenegro et al., 2009). The persistence of snow-cover increases with increasing elevation and increasing latitude. However, while persistence of snow-cover and global radiation are usually positively correlated in mountainous regions, causing high RF, they are negatively correlated in boreal regions. The strong altitudinal RF gradient found in this study is likely to be even more pronounced if altitudinal changes in forest structure and its influence on albedo and carbon sequestration are included on higher resolution because forests with very low carbon stocks can also have a low albedo (de Wit et al., 2013).

However, despite the general increase of RF with elevation, each biogeographical region has its specific characteristics. While the Jura and Plateau (under 600 m a.s.l.) are characterized by albedos of 0.136/0.139 (snow-free) and 0.272/0.276 (snow-covered), the albedo in the Southern Prealps under 600 m is respectively comparably low (0.112 and 0.185). This regional difference may be related to the different forest types and soil characteristics. Whereas beech dominates in the Jura and the Plateau, many stands in the Southern Prealps are dominated by chestnut. Moreover, forest soils in the lowlands of the Southern Prealps contain particularly high fractions of “black” fire derived carbon (Eckmeier et al., 2010). The darker soil colour may lower the albedo in addition to differences in the canopy.

Our spatially highly resolved estimates of RF are in agreement with the results of Betts (2000) and Montenegro et al. (2009) which are based on much coarser resolutions. In Betts (2000), the two pixels encompassing Switzerland show a net carbon sequestration of 100-150

and 150-200 tC/ha associated with reforestation of pasture, while in Montenegro et al. (2009), the pixels indicate a drawdown ranging between 100-150 tC/ha (maximum scenario) and 0-20 tC/ha (minimum scenario). However, many pixels are missing in alpine regions. In our study, we found similar values for the drawdown with values ranging from -10 to 160 tC/ha (minus 10 indicates emission instead of drawdown).

Small-scale variability, especially in topographically complex areas, was not captured in these former studies. Here, we used data on global radiation and snow-cover on a resolution of 2.2 km (MeteoSwiss) and 1 km (Huesler et al., 2012). To calculate differences in carbon sequestration and albedo, we relied on a biogeographical categorization and altitudinal stratification, which are based on major differences in vegetation and other ecological factors (Gutersohn, 1973, Wohlgemuth, 1996). Although the spatial resolution was high in our study, it should be refined further to allow for instance a comparison of RF on northern and southern slopes and a better capture of forest types and structures near the tree line.

The global sensitivity analysis with FAST showed that interactions between the input parameters were small in comparison to the main effects of each parameter. Thus, we verified the results of our global sensitivity analysis in a local analysis by estimating the partial derivatives for each factor. Both analyses showed good agreement since not only the interactions, but also non-linear effects had little influences. The amount of RF at a specific location is essentially influenced by carbon sequestration and snow-cover (Figure 6a). These two factors are good indicators for estimating the amount of RF. Global radiation and snow-covered albedo are also important, but they influence RF five times less than carbon sequestration and even seven times less than snow-cover. The factors with the most potential for improving our results are better estimates of carbon sequestration (Figure 6b), followed by reducing albedo uncertainty and the global radiation. The average parameter for atmospheric absorption (0.23) could be replaced by a spatially explicit specific parameter. ~~for each region or the specific top of the atmosphere~~

~~albedo could be used. If the specific absorption of the atmosphere is included, Including a spatially explicit parameter for atmospheric absorption would probably increase the elevation gradient of RF, the effect of global radiation on RF could be more pronounced since absorption of the atmosphere is higher in regions with low global radiation and lower in regions with high global radiation. because atmospheric absorption should be higher in low elevations than in high elevations. However, iAccording to our sensitivity analysis and Bright and Kvalevag (2013) improving data on atmospheric absorption will have only a relatively small influence on the results. (Figure 6b).~~

The sensitivity analysis with FAST is based on uncertainty estimates because exact values are not available. For example, MODIS values can be assigned to an average uncertainty of 10% (O'Halloran et al., 2012, Strahler et al., 1999). This uncertainty is likely to be much higher in topographically complex areas than in even terrain as the algorithm used to produce albedo values only indirectly accounts for topography. Moreover, the uncertainties associated with the different input parameters do not always refer to the same statistical measures and are thus not completely consistent (references of uncertainties are listed in [the Appendix](#), Table 3).

Regarding the different time horizons, RF increased by 17% for a time horizon of 1000 years compared with a horizon of 100 years. Larger time horizons increased RF since CO₂ RF constantly decreased due to interactions of atmospheric CO₂ with the carbon cycle, while albedo RF became constant after forests reached a steady state. Since we used a time horizon of 100 years, we rather underestimated albedo RF. This apparently goes against the findings of Schaeffer et al. (2006) and Kirschbaum et al. (2011) who both argue that CO₂ RF becomes more dominant for larger time horizons. However, they consider relatively short time periods (including rotations) where carbon sequestration does not end before the forests are removed. We think it is also necessary to include large time horizons when estimating RF, but it is of course an oversimplification to assume a fixed state after transition, since forests are frequently

disturbed (e.g. O'Halloran et al., 2012). Over longer periods snow-cover will be less persistent in the Alps, which will potentially decrease the albedo RF (de Wit et al., 2013, Pitman et al., 2011).

The maps of the possible carbon sequestration and albedo of forests in each biogeographical region (Figure 3) reflect mainly ecosystem characteristics. However, forests in Switzerland have long been under permanent anthropogenic influence and are thus not in a naturally balanced equilibrium, i.e. at the end of succession. For example, forests at high elevations in the Jura are often used for pasturing. Hence, they are less dense and sequester less carbon than they would if left to develop as undisturbed forests. The relationship between albedo RF, CO₂ RF and net RF will thus always depend on the actual and previous forest and land management. Our analysis of transitions from Extensively Used Open Land to Closed Forest and Open Forest corroborates this conclusion. The results indicated that the climatic benefit will be smaller if forests are kept in an open structure (e.g. due to pasturing) than when the canopy closes during succession. These findings are in line with former studies that estimate the effects of succession and forest structure on RF and show that changes in carbon stocks and changes in albedo are not linearly related (Kirschbaum et al., 2011, Bernier et al., 2011).

Our results should be valuable for future studies on the climatic impacts of LULUCF, especially for comparing and evaluating the results from climate models since our results are mainly based on satellite and field data. They should also be valuable for assessing the ecosystem (climate) service of forests in temperate mountains, i.e. for addressing the question of whether forest expansion in these regions is beneficial for climate or not. However, the interpretation of RF values has to be done carefully. First, the concept of Radiative Forcing has been developed to compare the impact of different forcing agents on the global mean temperature (Hansen et al., 2005). When applied at the regional and local scales one should keep in mind that the comparison of different forcing agents is far from being straightforward.

For instance, the impact of albedo will remain mostly local while those from CO₂ will be globally distributed and therefore diluted. Furthermore, the Climate sensitivities of CO₂ RF and Albedo RF may differ (Davin et al., 2007).

Finally, it is important to note that our study does not account for all possible effects of forests on climate, such as changes in evapotranspiration, surface roughness, and impacts on aerosols and other gases than CO₂. The uptake of CH₄ during forest expansion in the Alps increased according to Hiltbrunner et al. (2012), which adds to a negative CO₂ RF. This effect is, however, one magnitude smaller than the effect of CO₂-sequestration. The effect of a changing aerosol concentration, providing cooling (through cloud formation), may be very important (Spracklen et al., 2008). Changes in evapotranspiration and surface roughness, due to forest expansion, have a cooling impact in many geographical contexts (Bonan, 2008, Luyssaert et al., 2014), but we did not include the impacts of these changes in our study. In general, the influence of evapotranspiration and surface roughness will be low in those areas where snow-cover plays an important role (Bonan, 2008, Bathiany et al., 2010, Gibbard et al., 2005, Lee et al., 2011). Hence, these effects will be more important at low elevations (e.g. on the Swiss Plateau) than at high elevations in the Alps. The gap between the benefits of forest expansion at low and high elevations may thus become even wider if evapotranspiration and surface roughness are also integrated.

Conclusions

In the temperate mountainous regions of Switzerland, the net RF associated with changes in forest cover varies greatly on small spatial scales. At low elevations, with low to moderate snow-cover, RF is strongly negative due to a dominance of CO₂ RF. At high elevations in continental regions with persistent snow-cover, a very high global radiation, low carbon sequestration and low albedos of mostly evergreen tree species, RF can be positive. As a consequence, both clearly negative and positive values of RF can be found within a horizontal distance of 5 km in alpine valleys. Therefore, the climatic benefits of changes in forest cover can only be properly assessed using data at a high spatial resolution.

Our results indicate that it is very important to include albedo RF when estimating the impact on climate of changes in forest cover. Maps of RF, such as those produced in this study, indicate where climatic benefits from changes in forest cover can be expected and where not. In the Swiss Alps, the relevance of albedo RF is especially high because most transitions from open land to forest occur in regions where albedo RF causes a strong offset of CO₂ RF. Practitioners and politicians who need information about ecosystem services on local and regional scales should take into account that RF in the Swiss Alps mainly depends on the persistence of snow-cover and the potential for carbon sequestration. Moreover, late successional forest cover changes from Open Forests to Closed Forests are more beneficial for climate than early successional changes.

Our results could be improved if changes in evapotranspiration, surface roughness, aerosols and other gases than CO₂ were included. To determine the impacts of RFs better, however, further advances in climate modeling are necessary. A promising approach could thus be the coupling of regional climate models with global climate models. Regional models are able to simulate all the effects of changes in land use on climate (including evapotranspiration, surface roughness and so on) on a small scale. Coupling regional models with global models allows the

integration of feedbacks with the global circulation. This could help to close the gap between RF and temperature changes, and thus answer the question about where temperature changes caused by RF can be expected and how much change is likely.

Acknowledgments

This work was part of the Swiss research program “Forests and Climate change” and supported by the Swiss Federal Office for Environment (BAFU) and the Federal Institute for Forest, Snow and Landscape Research (WSL). We are grateful to Fortunat Joos, Stefan Wunderle, Janine Bolliger, Tobias Jonas, Henning Löwe and Michaela Teich for helpful discussions and Silvia Dingwall for language revision.

References

- ALEWELL, C. & BEBI, P. 2011. Forest Development in the European Alps and Potential Consequences on Hydrological Regime. *Forest Management and the Water Cycle: An Ecosystem-Based Approach*.
- ARORA, V. K. & MONTENEGRO, A. 2011. Small temperature benefits provided by realistic afforestation efforts. *Nature Geoscience*, 4, 514-518.
- BALA, G., CALDEIRA, K., WICKETT, M., PHILLIPS, T. J., LOBELL, D. B., DELIRE, C. & MIRIN, A. 2007. Combined climate and carbon-cycle effects of large-scale deforestation (vol 104, pg 6550, 2007). *Proceedings of the National Academy of Sciences of the United States of America*, 104, 9911-9911.
- BARNES, C. A. & ROY, D. P. 2010. Radiative forcing over the conterminous United States due to contemporary land cover land use change and sensitivity to snow and interannual albedo variability. *Journal of Geophysical Research-Biogeosciences*, 115.
- BATHIANY, S., CLAUSSEN, M., BROVKIN, V., RADDATZ, T. & GAYLER, V. 2010. Combined biogeophysical and biogeochemical effects of large-scale forest cover changes in the MPI earth system model. *Biogeosciences*, 7, 1383-1399.
- BAUR, P. 2006. Forest Expansion in the Swiss Alps: A quantitative analysis of bio-physical and socio-economic causes with an emphasis on structural changes in agriculture. Birmensdorf/Switzerland.
- BEBI, P., TEICH, M., HAGEDORN, F., ZURBRIGGEN, N., BRUNNER, S. H. & GRËT-REGAMEY, A. 2012. Veränderung von Wald und Waldleistungen in der Landschaft Davos im Zuge des Klimawandels. *Schweizerische Zeitschrift für Forstwesen*, 493-501.
- BERNIER, P. Y., DESJARDINS, R. L., KARIMI-ZINDASHTY, Y., WORTH, D., BEAUDOIN, A., LUO, Y. & WANG, S. 2011. Boreal lichen woodlands: A possible negative feedback to climate change in eastern North America. *Agricultural and Forest Meteorology*, 151, 521-528.
- BETTS, R. A. 2000. Offset of the potential carbon sink from boreal forestation by decreases in surface albedo. *Nature*, 408, 187-190.
- BETTS, R. A. 2011. CLIMATE SCIENCE Afforestation cools more or less. *Nature Geoscience*, 4, 504-505.
- BEYER, H. G., COSTANZO, C. & HEINEMANN, D. 1996. Modifications of the Heliosat procedure for irradiance estimates from satellite images. *Solar Energy*, 56, 207-212.
- BLOETZER, G. 2004. Walderhaltungspolitik - Entwicklung und Urteil der Fachleute. In: BUNDESAMT FÜR UMWELT, W. U. L. B. (ed.).
- BOLLIGER, J., HAGEDORN, F., LEIFELD, J., BOHL, J., ZIMMERMANN, S., SOLIVA, R. & KIENAST, F. 2008. Effects of land-use change on carbon stocks in Switzerland. *Ecosystems*, 11, 895-907.
- BONAN, G. B. 2008. Forests and climate change: Forcings, feedbacks, and the climate benefits of forests. *Science*, 320, 1444-1449.
- BRÄNDLI, U.-B. 2010. Schweizerisches Landesforstinventar. Ergebnisse der dritten Erhebung 2004–2006. [Results of the third Swiss national forest inventory 2004-2006]. Birmensdorf, Eidgenössische Forschungsanstalt für Wald, Schnee und Landschaft WSL. Bern, Bundesamt für Umwelt, BAFU. 312 S.
- BRIGHT, R. M. & KVALEVAG, M. M. 2013. Technical Note: Evaluating a simple parameterization of radiative shortwave forcing from surface albedo change. *Atmospheric Chemistry and Physics*, 13, 11169-11174.
- BÜRGI, M. & SCHULER, A. 2003. Driving forces of forest management—an analysis of regeneration practices in the forests of the Swiss Central Plateau during the 19th and 20th century. *Forest Ecology and Management*, 176, 173-183.
- CANO, D., MONGET, J. M., ALBUISSON, M., GUILLARD, H., REGAS, N. & WALD, L. 1986. A METHOD FOR THE DETERMINATION OF THE GLOBAL SOLAR-RADIATION FROM METEOROLOGICAL SATELLITE DATA. *Solar Energy*, 37, 31-39.

- CHERUBINI, F., PETERS, G. P., BERNTSEN, T., STROMMAN, A. H. & HERTWICH, E. 2011. CO₂ emissions from biomass combustion for bioenergy: atmospheric decay and contribution to global warming. *Global Change Biology Bioenergy*, 3, 413-426.
- COSTA, M. H. & FOLEY, J. A. 2000. Combined effects of deforestation and doubled atmospheric CO₂ concentrations on the climate of Amazonia. *Journal of Climate*, 13, 18-34.
- DAVIN, E. L. & DE NOBLET-DUCOUDRE, N. 2010. Climatic Impact of Global-Scale Deforestation: Radiative versus Nonradiative Processes. *Journal of Climate*, 23, 97-112.
- DAVIN, E. L., DE NOBLET-DUCOUDRE, N. & FRIEDLINGSTEIN, P. 2007. Impact of land cover change on surface climate: Relevance of the radiative forcing concept. *Geophysical Research Letters*, 34.
- DE WIT, H. A., BRYN, A., HOFGAARD, A., KARSTENSEN, J., KVALEVÅG, M. M. & PETERS, G. P. 2013. Climate warming feedback from mountain birch forest expansion: reduced albedo dominates carbon uptake. *Global Change Biology*, n/a-n/a.
- DURR, B., ZELENKA, A., MULLER, R. & PHILIPONA, R. 2010. Verification of CM-SAF and MeteoSwiss satellite based retrievals of surface shortwave irradiance over the Alpine region. *International Journal of Remote Sensing*, 31, 4179-4198.
- ECKMEIER, E., EGLI, M., SCHMIDT, M. W. I., SCHLUMPF, N., NOTZLI, M., MINIKUS-STARY, N. & HAGEDORN, F. 2010. Preservation of fire-derived carbon compounds and sorptive stabilisation promote the accumulation of organic matter in black soils of the Southern Alps. *Geoderma*, 159, 147-155.
- FORSTER, P., RAMASWAMY, V., ARTAXO, P., BERNTSEN, T., BETTS, R., FAHEY, D. W., HAYWOOD, J., LEAN, J., LOWE, D. C., MYHRE, G., NGANGA, J., PRINN, R., RAGA, G., SCHULZ, M. & ., R. V. D. E. 2007. Changes in Atmospheric Constituents and in Radiative Forcing. In: SOLOMON, S., D. QIN, M. MANNING, Z. CHEN, M. MARQUIS, K.B. AVERYT, M. TIGNOR AND H.L. MILLER (ed.) *In: Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge United Kingdom and New York, NY, USA: Cambridge University Press.
- FU, Q. & LIOU, K. N. 2005. *Fu-Liou Cloud/Aerosol Forcing Page* [Online]. NASA. Available: <http://snowdog.larc.nasa.gov/cgi-bin/rose/flp200503/flp200503.cgi>.
- GEHRIG-FASEL, J., GUISAN, A. & ZIMMERMANN, N. E. 2007. Tree line shifts in the Swiss Alps: Climate change or land abandonment? *Journal of Vegetation Science*, 18, 571-582.
- GIBBARD, S., CALDEIRA, K., BALA, G., PHILLIPS, T. J. & WICKETT, M. 2005. Climate effects of global land cover change. *Geophysical Research Letters*, 32.
- GIMMI, U., WOLF, A., BÜRGI, M., SCHERSTJANOI, M. & BUGMANN, H. 2009. Quantifying disturbance effects on vegetation carbon pools in mountain forests based on historical data. *Regional Environmental Change*, 9, 121-130.
- GOVINDASAMY, B., DUFFY, P. B. & CALDEIRA, K. 2001. Land use Changes and Northern Hemisphere Cooling. *Geophysical Research Letters*, 28, 291-294.
- GRET-REGAMEY, A., BRUNNER, S. H., ALTWEGG, J., CHRISTEN, M. & BEBI, P. 2013. Integrating Expert Knowledge into Mapping Ecosystem Services Trade-offs for Sustainable Forest Management. *Ecology and Society*, 18.
- GUTERSOHN, H. 1973. Naturräumliche Gliederung. *Atlas der Schweiz*.
- HAGEDORN, F., MOERL, A., WALTHERT, L. & ZIMMERMANN, S. 2010. Kohlenstoff in schweizer waldböden – bei Klima erwärmung eine potenzielle CO₂-Quelle. *Schweizerische Zeitschrift für Forstwesen*, 161, 530-535.
- HAGEDORN, F., SHIYATOV, S. G., MAZEPA, V. S., DEVI, N. M., GRIGOR'EV, A. A., BARTISH, A. A., FOMIN, V. V., KAPRALOV, D. S., TEREENT'EV, M., BUGMAN, H., RIGLING, A. & MOISEEV, P. A. 2014. Treeline advances along the Urals mountain range – driven by improved winter conditions? *Global Change Biology*, n/a-n/a.

- HAMMER, A., HEINEMANN, D., HOYER, C., KUHLEMANN, R., LORENZ, E., MULLER, R. & BEYER, H. G. 2003. Solar energy assessment using remote sensing technologies. *Remote Sensing of Environment*, 86, 423-432.
- HELDSTAB, J., SOMMERHALDER, M. & RHIM, B. 2012. Switzerland's Greenhouse Gas Inventory 1990-2010. Bern: Federal Office for the Environment FOEN.
- HILTBRUNNER, D., ZIMMERMANN, S., KARBIN, S., HAGEDORN, F. & NIKLAUS, P. A. 2012. Increasing soil methane sink along a 120-year afforestation chronosequence is driven by soil moisture. *Global Change Biology*, 18, 3664-3671.
- HUESLER, F., JONAS, T., WUNDERLE, S. & ALBRECHT, S. 2012. Validation of a modified snow cover retrieval from historical 1-km AVHRR data over the European Alps. *Journal of Geophysical Research*.
- HUMBEL, R., BEYELER, A., BURKHALTER, J., SAGER, J. & H.-U.ZAUGG. 2010. Arealstatistik nach Nomenklatur 2004 - Standard NOAS04. Available: http://www.bfs.admin.ch/bfs/portal/de/index/dienstleistungen/geostat/datenbeschreibung/arealstatistik_noas04.html [Accessed 17.04.2010].
- HUSLER, F., JONAS, T., WUNDERLE, S. & ALBRECHT, S. 2012. Validation of a modified snow cover retrieval algorithm from historical 1-km AVHRR data over the European Alps. *Remote Sensing of Environment*, 121, 497-515.
- IPCC 2001. Third Assessment Report - Appendix II, SRES Tables.
- IPCC 2003. Good Practice Guidance for Land Use, Land-Use Change and Forestry (IPCC GPG LULUCF).
- JOOS, F., ROTH, R., FUGLESTVEDT, J. S., PETERS, G. P., ENTING, I. G., VON BLOH, W., BROVKIN, V., BURKE, E. J., EBY, M., EDWARDS, N. R., FRIEDRICH, T., FROLICHER, T. L., HALLORAN, P. R., HOLDEN, P. B., JONES, C., KLEINEN, T., MACKENZIE, F. T., MATSUMOTO, K., MEINSHAUSEN, M., PLATTNER, G. K., REISINGER, A., SEGSCHNEIDER, J., SHAFFER, G., STEINACHER, M., STRASSMANN, K., TANAKA, K., TIMMERMANN, A. & WEAVER, A. J. 2013. Carbon dioxide and climate impulse response functions for the computation of greenhouse gas metrics: a multi-model analysis. *Atmospheric Chemistry and Physics*, 13, 2793-2825.
- KAUFMANN, E. 2001. Estimation of standing timber, growth and cut. In: Brassel, P., Lischke, H. (eds.): Swiss National Forest Inventory: Methods and Models of the Second Assessment. Swiss Federal Research Institute WSL, Birmensdorf: 162-196. .
- KIRSCHBAUM, M. U. F., WHITEHEAD, D., DEAN, S. M., BEETS, P. N., SHEPHERD, J. D. & AUSSEIL, A. G. E. 2011. Implications of albedo changes following afforestation on the benefits of forests as carbon sinks. *Biogeosciences*, 8, 3687-3696.
- KÖRNER, C. 2012. *Alpine Treelines, Functional Ecology of the Global High Elevation Tree Limits*, Basel, Springer.
- KOZAK, J. 2003. Forest cover change in the Western Carpathians in the past 180 years - A case study in the Orawa Region in Poland. *Mountain Research and Development*, 23, 369-375.
- KVALEVAG, M. M., MYHRE, G., BONAN, G. & LEVIS, S. 2010. Anthropogenic land cover changes in a GCM with surface albedo changes based on MODIS data. *International Journal of Climatology*, 30, 2105-2117.
- LAIOLO, P., DONDERO, F., CILIENTO, E. & ROLANDO, A. 2004. Consequences of pastoral abandonment for the structure and diversity of the alpine avifauna. *Journal of Applied Ecology*, 41, 294-304.
- LEE, X., GOULDEN, M. L., HOLLINGER, D. Y., BARR, A., BLACK, T. A., BOHRER, G., BRACHO, R., DRAKE, B., GOLDSTEIN, A., GU, L. H., KATUL, G., KOLB, T., LAW, B. E., MARGOLIS, H., MEYERS, T., MONSON, R., MUNGER, W., OREN, R., KYAW, T. P. U., RICHARDSON, A. D., SCHMID, H. P., STAEBLER, R., WOFSY, S. & ZHAO, L. 2011. Observed increase in local cooling effect of deforestation at higher latitudes. *Nature*, 479, 384-387.
- LEIFELD, J., BASSIN, S. & FUHRER, J. 2005. Carbon stocks in Swiss agricultural soils predicted by land-use, soil characteristics, and altitude. *Agriculture Ecosystems & Environment*, 105, 255-266.

- LIU, J. C., SCHAAF, C., STRAHLER, A., JIAO, Z. T., SHUAI, Y. M., ZHANG, Q. L., ROMAN, M., AUGUSTINE, J. A. & DUTTON, E. G. 2009. Validation of Moderate Resolution Imaging Spectroradiometer (MODIS) albedo retrieval algorithm: Dependence of albedo on solar zenith angle. *Journal of Geophysical Research-Atmospheres*, 114.
- LUYSSAERT, S., JAMMET, M., STOY, P. C., ESTEL, S., PONGRATZ, J., CESCHIA, E., CHURKINA, G., DON, A., ERB, K., FERLICOQ, M., GIELEN, B., GRUNWALD, T., HOUGHTON, R. A., KLUMPP, K., KNOHL, A., KOLB, T., KUEMMERLE, T., LAURILA, T., LOHILA, A., LOUSTAU, D., MCGRATH, M. J., MEYFROIDT, P., MOORS, E. J., NAUDTS, K., NOVICK, K., OTTO, J., PILEGAARD, K., PIO, C. A., RAMBAL, S., REBMANN, C., RYDER, J., SUYKER, A. E., VARLAGIN, A., WATTENBACH, M. & DOLMAN, A. J. 2014. Land management and land-cover change have impacts of similar magnitude on surface temperature. *Nature Clim. Change*, 4, 389-393.
- MACDONALD, D., CRABTREE, J. R., WIESINGER, G., DAX, T., STAMOU, N., FLEURY, P., LAZPITA, J. G. & GIBON, A. 2000. Agricultural abandonment in mountain areas of Europe: Environmental consequences and policy response. *Journal of Environmental Management*, 59, 47-69.
- METEOSWISS, F. O. O. M. A. C. 2012. Available: http://www.meteoschweiz.admin.ch/web/en/services/data_portal/grided_datasets/radiation.html [Accessed 28.02.2012 2012].
- MONTENEGRO, A., EBY, M., MU, Q. Z., MULLIGAN, M., WEAVER, A. J., WIEBE, E. C. & ZHAO, M. S. 2009. The net carbon drawdown of small scale afforestation from satellite observations. *Global and Planetary Change*, 69, 195-204.
- MYHRE, G., HIGHWOOD, E. J., SHINE, K. P. & STORDAL, F. 1998. New estimates of radiative forcing due to well mixed greenhouse gases. *Geophysical Research Letters*, 25, 2715-2718.
- MYHRE, G., SHINDELL, D., BRÉON, F.-M., COLLINS, W., FUGLESTVEDT, J., HUANG, J., KOCH, D., LAMARQUE, J.-F., LEE, D., MENDOZA, B., NAKAJIMA, T., ROBOCK, A., STEPHENS, G., TAKEMURA, T. & ZHANG, H. 2013. Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. *Anthropogenic and Natural Radiative Forcing*.
- NUSSBAUM, M., PAPRITZ, A., BALTENSWEILER, A. & WALTHERT, L. 2012. Organic Carbon Stocks of Swiss Forest Soils. Final Report.: Institute of Terrestrial Ecosystems, ETH Zürich and Swiss Federal Institute for Forest, Snow and Landscape Research (WSL), Zürich and Birmensdorf.
- O'HALLORAN, T. L., LAW, B. E., GOULDEN, M. L., WANG, Z. S., BARR, J. G., SCHAAF, C., BROWN, M., FUENTES, J. D., GOCKEDE, M., BLACK, A. & ENGEL, V. 2012. Radiative forcing of natural forest disturbances. *Global Change Biology*, 18, 555-565.
- PERRUCHOUD, D., KIENAST, F., KAUFMANN, E. & BRAKER, O. U. 1999. 20th century carbon budget of forest soils in the Alps. *Ecosystems*, 2, 320-337.
- PITMAN, A. J., AVILA, F. B., ABRAMOWITZ, G., WANG, Y. P., PHIPPS, S. J. & DE NOBLET-DUCOUDRE, N. 2011. Importance of background climate in determining impact of land-cover change on regional climate. *Nature Climate Change*, 1, 472-475.
- POEPLAU, C., DON, A., VESTERDAL, L., LEIFELD, J., VAN WESEMAEL, B., SCHUMACHER, J. & GENSJØR, A. 2011. Temporal dynamics of soil organic carbon after land-use change in the temperate zone - carbon response functions as a model approach. *Global Change Biology*, 17, 2415-2427.
- PUJOL, G., IOOSS, B. & JANON, A. 2012. Sensitivity Analysis. Version 1.6-1.
- RAMANKUTTY, N., HELLER, E. & RHEMTULLA, J. 2010. Prevailing Myths About Agricultural Abandonment and Forest Regrowth in the United States. *Annals of the Association of American Geographers*, 100, 502-512.
- ROESCH, A., WILD, M., PINKER, R. & OHMURA, A. 2002. Comparison of spectral surface albedos and their impact on the general circulation model simulated surface climate. *Journal of Geophysical Research*, 107, ACL13-1-ACL13-ACL13-18ACL13-ACL13-18.

- RUTHERFORD, G. N., BEBI, P., EDWARDS, P. J. & ZIMMERMANN, N. E. 2008. Assessing land-use statistics to model land cover change in a mountainous landscape in the European Alps. *Ecological Modelling*, 212, 460-471.
- SALTELLI, A., TARANTOLA, S. & CHAN, K. P. S. 1999. A quantitative model-independent method for global sensitivity analysis of model output. *Technometrics*, 41, 39-56.
- SCHAAF, C. 2010. *MODIS BRDF/Albedo Product (MCD43) User's Guide* [Online]. Available: <http://www-modis.bu.edu/brdf/userguide/intro.html> [Accessed 30.03.2012 2012].
- SCHAEFFER, M., EICKHOUT, B., HOOGWJK, M., STRENGERS, B., VAN VUUREN, D., LEEMANS, R. & OPSTEEGH, T. 2006. CO₂ and albedo climate impacts of extratropical carbon and biomass plantations. *Global Biogeochemical Cycles*, 20.
- SCHNEEBERGER, N., BÜRGI, M. & KIENAST, P. D. F. 2007. Rates of landscape change at the northern fringe of the Swiss Alps: Historical and recent tendencies. *Landscape and Urban Planning*, 80, 127-136.
- SCHWAIGER, H. P. & BIRD, D. N. 2010. Integration of albedo effects caused by land use change into the climate balance: Should we still account in greenhouse gas units? *Forest Ecology and Management*, 260, 278-286.
- SIEGENTHALER, U. & OESCHGER, H. 1978. PREDICTING FUTURE ATMOSPHERIC CARBON-DIOXIDE LEVELS. *Science*, 199, 388-395.
- SJÖRGERSTEN-TURNER, S., ALEWELL, C., CÉCILLON, L., HAGEDORN, F., JANDL, R., LEIFELD, J., VEGARD, M., SCHINDLBACHER, A., SEBASTIÁ, M. T. & VAN MIEGROET, H. 2011. Mountain soils in a changing climate - vulnerability of carbon stocks and ecosystem feedbacks. In: JANDL, R., MIRCO, R. & MATS, O. (eds.) *Soil Carbon in Sensitive European Ecosystems*. Wiley-Blackwell.
- SPRACKLEN, D. V., BONN, B. & CARSLAW, K. S. 2008. Boreal forests, aerosols and the impacts on clouds and climate. *Philosophical Transactions of the Royal Society a-Mathematical Physical and Engineering Sciences*, 366, 4613-4626.
- STRAHLER, A. H., LUCHT, W., SCHAAF, C. B., TSANG, T., GAO, F., LI, X., MULLER, J.-P., LEWIS, P. & BARNESLEY, M. J. 1999. MODIS BRDF/Albedo Product: Algorithm Theoretical Basis Document Version 5.0.
- UNFCCC 2001. Decision 11/CP.7: Land Use, Land-use Change and Forestry. In: CHANGE, U. N. F. C. O. C. (ed.).
- WIRTH, C., SCHUMACHER, J. & SCHULZE, E. D. 2004. Generic biomass functions for Norway spruce in Central Europe - a meta-analysis approach toward prediction and uncertainty estimation. *Tree Physiology*, 24, 121-139.
- WOHLGEMUTH, T. 1996. Biogeographical Regionalization of Switzerland Based on Floristic Data: How Many Species are Needed? *Biodiversity Letters*.
- WUTZLER, T., WIRTH, C. & SCHUMACHER, J. 2008. Generic biomass functions for Common beech (*Fagus sylvatica*) in Central Europe: predictions and components of uncertainty. *Canadian Journal of Forest Research-Revue Canadienne De Recherche Forestiere*, 38, 1661-1675.
- ZHOU, L., DICKINSON, R. E., TIAN, Y., ZENG, X., DAI, Y., YANG, Z. L., SCHAAF, C. B., GAO, F., JIN, Y., STRAHLER, A., MYNENI, R. B., YU, H., WU, W. & SHAIKH, M. 2003. Comparison of seasonal and spatial variations of albedos from Moderate-Resolution Imaging Spectroradiometer (MODIS) and Common Land Model. *Journal of Geophysical Research-Atmospheres*, 108.

Figure legends

Figure 1: Digital elevation model and biogeographical regions of Switzerland.

Figure 2: Examples of a) Intensively Used Open Land b) Extensively Used Open Land c) Open Forest d) Closed Forest

Figure 3 a) – f): a) Mean yearly global radiation. b) Days with snow-cover per year. c) Albedo difference without snow (difference between Intensively/Extensively Used Open Land and Closed Forest) d) Albedo difference with snow (difference between Intensively/Extensively Used Open Land and Closed Forest) e) Carbon sequestration (difference between intensively/Extensively and Closed Forest) f) Tree-line elevation calculated using the Swiss Area Statistics of 1997 and applying the method of Paulsen and Körner (2001).

~~a) Albedo difference without snow (difference between Intensively/Extensively Used Open Land and Closed Forest) b) Albedo difference with snow (difference between Intensively/Extensively Used Open Land and Closed Forest) c) Carbon sequestration (difference between intensively/Extensively and Closed Forest) d) Days with snow cover per year. e) Mean yearly global radiation. f) Tree line elevation calculated using the Swiss Area Statistics of 1997 and applying the method of Paulsen and Körner (2001).~~

Figure 4 a)-d): a) albedo RF b) CO₂ RF c) Offset: albedo RF/CO₂ RF. d) Net RF: albedo RF plus CO₂ RF. All datasets were derived for transitions from Intensively Used Open Land (<1000 m) and Extensively Used Open Land (>1000) to Closed Forest.

Figure 5: Radiative Forcings of the forest expansion between 1985 and 1997 for three elevations in the biogeographical regions Jura, Plateau, Northern Prealps, Central Alps, Southern Prealps.

Figure 6a,b: Global sensitivity analysis FAST a) sensitivity analysis of spatial variability of c_seq (carbon sequestration), sc (snow-cover), glob (global radiation), a_ns (albedo difference no snow-cover) and a_s (albedo difference snow-covered) b) sensitivity analysis of uncertainty for each pixel: c_seq, sc, glob, a_ns, a_s, c_cycle (carbon cycle), rt (radiative transfer) and atm_absorb (atmospheric absorption).

Figures

Figure 1




Figure 2




Figure 3 a) – f)




Figure 4 a)-d)




Figure 5






Figure 6 a) and b)



Appendix, Tables

Table 1: Albedo of forest (snow, no snow) and Open Forest (snow, no snow) and corresponding standard errors of each biogeographical region. Values marked with * were derived using majority pixels and/or magnitude inversion (see chapter albedo RF), all other values have been derived using 92% pixel cover and full inversion. Numbers 1-3 indicate the elevation level of each biogeographical region: Jura 1 = Jura below 600 m, Jura 2 = Jura between 600 and 1200 m and Jura 3 = Jura above 1200 m.

Region	Forest				Open Forest			
	Snow	Ste(%)	Not Snow	Ste(%)	Snow	Ste(%)	Not Snow	Ste(%)
Jura 1	0.263	1.30	0.139	0.08	NA	NA	NA	NA
Jura 2	0.221	0.50	0.133	0.04	0.303*	1.80*	0.141*	0.23*
Jura 3	0.175	0.64	0.109	0.12	0.232*	0.67*	0.117*	0.13*
Plateau 1	0.266	0.61	0.136	0.03	NA	NA	NA	NA
Plateau 2	0.251	0.53	0.134	0.05	NA	NA	NA	NA
Plateau 3	NA	NA	0	NA	NA	NA	NA	NA
Northern Prealps 1	0.240	3.16	0.127	0.45	NA	NA	NA	NA
Northern Prealps 2	0.194	0.38	0.118	0.06	0.217*	2.36*	0.114*	0.49*

Northern Prealps 3	0.203	0.40	0.112	0.11	0.275*	0.42*	0.121	0.095
Central Alps 1	0.168*	2.1*	0.107	1.31	0.353*	15.7*	0.141*	0.58*
Central Alps 2	0.174	0.61	0.104	0.15	0.285*	3.09*	0.134*	0.18*
Central Alps 3	0.192	0.23	0.101	0.08	0.274*	0.33*	0.118*	0.068*
Southern Prealps 1	0.185*	4.9*	0.111	0.23	NA	NA	0.127*	0.78*
Southern Prealps 2	0.190	3.13	0.121	0.07	0.258*	3.28*	0.126*	0.27*
Southern Prealps 3	0.194	0.78	0.107	0.12	0.307*	0.50*	0.119	0.089

Table 2: Carbon stocks of Closed Forests and Open Forests in biomass and soils. The standard errors (Stes) refer to the deviation from the sample mean in every biogeographical region. Not included are errors of the model parameters which have been used to derive carbon stocks from tree measurements.

	Forest		Open Forest	
	biomass (tC/ha)	Ste (%)	biomass (tC/ha)	Ste (%)
Jura 1	134.1	4.6	NA	NA
Jura 2	144.3	2.3	101.27	27.2
Jura 3	103.7	5.5	68.44	13.4
Plateau 1	140.4	2.7	21.08	51.0
Plateau 2	159.7	3.2	92.80	46.2
Plateau 3	137.58	14.1	71.26	33.6
Northern Prealps 1	169.8	9.5	NA	NA
Northern Prealps 2	166.8	2.7	85.60	22.7
Northern Prealps 3	152.5	4.5	72.21	11.0
Central Alps 1	122	11.2	NA	NA

Central Alps 2	124.7	4.0	70.41	29.8
Central Alps 3	115.5	2.3	57.90	7.9
Southern Prealps 1	95.2	6.8	NA	NA
Southern Prealps 2	94.7	4.7	37.40	39.3
Southern Prealps 3	100.6	4.5	41.60	11.6

Table 3: Spatial variability and uncertainties applied in the sensitivity analysis.

	Spatial variability	Uncertainty (standard deviation)
Carbon sequestration [tC/ha]	143-241	+/- 35% (Hagedorn et al., 2010, Heldstab et al., 2012)
Snow-cover [days/year]	10-240	+/- 10% (Husler et al., 2012)
Global radiation [W/m ²]	117-180	+/- 28% (Durr et al., 2010)
Albedo snow-covered	0.208-0.375	+/- 36% (Combined error of MODIS data and methods for assigning LULC specific albedos)
Albedo snow-free	0.025 - 0.066	+/- 28% (Combined error of MODIS data and methods for assigning LULC specific albedos)
Carbon cycle	-	+/- 15% (Joos et al., 2013)
Radiative transfer	-	+/- 10% (Myhre et al., 1998)

Atmospheric absorption	-	+/- 30% radiative transfer tests*
------------------------	---	--------------------------------------

*For the uncertainty in atmospheric absorption, we relied on experiments with Fu and Liou online model (Fu and Liou, 2005). We tested different scenarios of cloudiness, aerosol concentration and elevation to determine how much atmospheric absorption could vary over Switzerland.

Table 4: Area affected by each type of transition between 1985 and 1997. Numbers 1-3 behind the biogeographical regions indicate the elevation (1 = below 600m, 2 = 600 – 1200m, 3 = above 1200m).

	<u>Area affected by each transition between 1985 and 1997 [in ha]</u>					
<u>Biogeographical region</u>	<u>Intensively Used Open Land to Closed Forest</u>	<u>Extensively Used Open Land to Closed Forest</u>	<u>Intensively Used Open Land to Open Forest</u>	<u>Extensively Used Open Land to Open Forest</u>	<u>Open Forest to Closed Forest</u>	<u>Forest expansion (sum of all transitions)</u>
<u>Jura 1</u>	<u>116</u>	<u>31</u>	<u>98</u>	<u>35</u>	<u>106</u>	<u>386</u>
<u>Jura 2</u>	<u>113</u>	<u>238</u>	<u>73</u>	<u>330</u>	<u>522</u>	<u>1276</u>
<u>Jura 3</u>	<u>1</u>	<u>46</u>	<u>1</u>	<u>155</u>	<u>490</u>	<u>693</u>
<u>Plateau 1</u>	<u>613</u>	<u>87</u>	<u>379</u>	<u>52</u>	<u>264</u>	<u>1395</u>
<u>Plateau 2</u>	<u>232</u>	<u>60</u>	<u>110</u>	<u>44</u>	<u>85</u>	<u>531</u>
<u>Plateau 3</u>	<u>NA</u>	<u>NA</u>	<u>NA</u>	<u>NA</u>	<u>1</u>	<u>1</u>
<u>Northern Prealps 1</u>	<u>109</u>	<u>21</u>	<u>78</u>	<u>13</u>	<u>53</u>	<u>274</u>
<u>Northern Prealps 2</u>	<u>321</u>	<u>497</u>	<u>295</u>	<u>401</u>	<u>959</u>	<u>2473</u>
<u>Northern Prealps 3</u>	<u>34</u>	<u>955</u>	<u>77</u>	<u>1180</u>	<u>2476</u>	<u>4722</u>
<u>Central Alps 1</u>	<u>4</u>	<u>4</u>	<u>6</u>	<u>11</u>	<u>29</u>	<u>54</u>
<u>Central Alps 2</u>	<u>93</u>	<u>101</u>	<u>267</u>	<u>154</u>	<u>679</u>	<u>1294</u>

<u>Central</u>	<u>102</u>	<u>739</u>	<u>291</u>	<u>1700</u>	<u>3687</u>	<u>6519</u>
<u>Alps 3</u>						
<u>Southern</u> <u>Prealps 1</u>	<u>61</u>	<u>25</u>	<u>145</u>	<u>22</u>	<u>196</u>	<u>449</u>
<u>Southern</u> <u>Prealps 2</u>	<u>76</u>	<u>77</u>	<u>170</u>	<u>135</u>	<u>731</u>	<u>1189</u>
<u>Southern</u> <u>Prealps 3</u>	<u>23</u>	<u>274</u>	<u>71</u>	<u>604</u>	<u>1541</u>	<u>2513</u>
<u>Biogeogra</u>	<u>Forest expansion (ha)</u>					
<u>Jura 1</u>	<u>386</u>					
<u>Jura 2</u>	<u>1276</u>					
<u>Jura 3</u>	<u>693</u>					
<u>Plateau 1</u>	<u>1395</u>					
<u>Plateau 2</u>	<u>531</u>					
<u>Plateau 3</u>	<u>1</u>					
<u>Pre-Alps 1</u>	<u>274</u>					
<u>Pre-Alps 2</u>	<u>2473</u>					
<u>Pre-Alps 3</u>	<u>4722</u>					
<u>Alps 1</u>	<u>54</u>					
<u>Alps 2</u>	<u>1294</u>					
<u>Alps 3</u>	<u>6519</u>					
<u>Southern</u>	<u>449</u>					
<u>Southern</u>	<u>1189</u>					
<u>Southern</u>	<u>2513</u>					

Table 5 Aggregation of land use classes from Swiss Arealstatistik (ASCH85, ASCH97 and ASCH04) adapted from Rutherford et al. (2008)

<u>Aggregated class</u>	<u>Area [ha]</u>	<u>Classes from Swiss land use statistics</u>	<u>Area [ha]</u>	<u>Broad definition</u>
<u>Closed Forest</u>	<u>1121544</u>	<u>Afforestation, 52</u> <u>Forest dieback, 54</u> <u>Normal forest, 50</u> <u>Slender Forest, 14</u> <u>Bushes, 57</u> <u>Groves and hedges, 58</u>	<u>3349</u> <u>14851</u> <u>962312</u> <u>44711</u> <u>60514</u> <u>35807</u>	<u>Vegetation height >3m, cover density >60%, composed of tree species</u>
<u>Open Forest</u>	<u>150101</u>	<u>On non-agriculturally used land, 56</u> <u>On agriculturally used land, 55</u> <u>Groups of trees on agriculturally used land, 59</u> <u>Groups of trees on non-agriculturally used land, 60</u>	<u>52825</u> <u>24108</u> <u>38157</u> <u>35011</u>	<u>Vegetation height >3m, cover density 20-60%, composed of tree species</u>
<u>Extensively Used Open Land</u>	<u>767842</u>	<u>Pasture in the vicinity of settlements, 43</u> <u>Alpine meadows, 45</u> <u>Sheep alps, 49</u> <u>Favourable to pasturing, 46</u> <u>Stony alpine pasture, 48</u> <u>Grass and herb vegetation, 65</u>	<u>87303</u> <u>32316</u> <u>51124</u> <u>368691</u> <u>46024</u> <u>182384</u>	<u>Used for grazing, use not year-round, not machine-accessible</u>
<u>Intensively Used Open Land</u>	<u>837128</u>	<u>Arable land, 41</u> <u>Natural meadows, 42</u>	<u>547754</u> <u>289374</u>	<u>Year-round use, in the vicinity of settlements, Mown</u>
<u>Other</u>	<u>125186</u>	<u>1-40, 44, 47, 61-64, 66-72</u>	<u>125186</u>	

Numbers in column 3 represent the official ASCH classes of the nomenclature 2004 (Humbel et al., 2010). The aggregation in (Rutherford et al., 2008) was adapted to the new nomenclature.

|

Appendix, Figures




Figure 1: Spatial pattern of forest expansion. The pattern illustrates the density of forest expansion in Switzerland. The density was calculated including the area of all five transitions we used for calculating RF ([see chapter “Swiss forest expansion between 1985 and 1997”](#)[see Land use/Land Cover \(LULC\) change](#)) and a kernel-density function in ArcGis 10.1 (ESRI).




Figure 2: Seasonal variation of albedo values of the four snow-free LULC classes Closed Forest, Intensively Used Open Land, Extensively Used Open Land and Open Forest (only full BRDF inversion albedo values).




Figure 3: Seasonal variation of the albedo values of the four snow-covered LULC classes Closed Forest, Intensively Used Open Land, Extensively Used Open Land and Open Forest (full BRDF albedo values and magnitude inversion albedo values for open forest).