1 Identifying environmental controls on vegetation

2 greenness phenology through model-data integration

3

M. Forkel¹, N. Carvalhais^{1,2}, S. Schaphoff³, W. v. Bloh³, M. Migliavacca¹, M. Thurner^{1,4} and K. Thonicke³

- 6 [1]{Max-Planck-Institute for Biogeochemistry, Department for Biogeochemical Integration,
- 7 Hans-Knöll-Str. 10, 07745 Jena, Germany }
- 8 [2] {Universidade Nova de Lisboa, Faculdade de Ciências e Tecnologia, 2829-516, Caparica,
- 9 Portugal}
- 10 [3]{Potsdam Institute for Climate Impact Research, Earth System Analysis, Telegraphenberg
- 11 A31, 14473 Potsdam, Germany}
- 12 [4]{Stockholm University, Department of Applied Environmental Science and Bolin Centre
- 13 for Climate Research, Svante Arrhenius väg 8, 10691 Stockholm, Sweden}

14 Correspondence to: M. Forkel (mforkel@bgc-jena.mpg.de)

15

16 Abstract

17 Existing dynamic global vegetation models (DGVMs) have a limited ability in reproducing 18 phenology and decadal dynamics of vegetation greenness as observed by satellites. These 19 limitations in reproducing observations reflect a poor understanding and description of the 20 environmental controls on phenology, which strongly influence the ability to simulate longer 21 term vegetation dynamics, e.g. carbon allocation. Combining DGVMs with observational data 22 sets can potentially help to revise current modelling approaches and thus to enhance the 23 understanding of processes that control seasonal to long-term vegetation greenness dynamics. 24 Here we implemented a new phenology model within the LPJmL (Lund Potsdam Jena 25 managed lands) DGVM and integrated several observational data sets to improve the ability 26 of the model in reproducing satellite-derived time series of vegetation greenness. Specifically, we optimized LPJmL parameters against observational time series of the fraction of absorbed 27 28 photosynthetic active radiation (FAPAR), albedo and gross primary production to identify the

1 main environmental controls for seasonal vegetation greenness dynamics. We demonstrated 2 that LPJmL with new phenology and optimized parameters better reproduces seasonality, 3 inter-annual variability and trends of vegetation greenness. Our results indicate that soil water 4 availability is an important control on vegetation phenology not only in water-limited biomes 5 but also in boreal forests and the arctic tundra. Whereas water availability controls phenology 6 in water-limited ecosystems during the entire growing season, water availability co-modulates 7 jointly with temperature the beginning of the growing season in boreal and arctic regions. 8 Additionally, water availability contributes to better explain decadal greening trends in the 9 Sahel and browning trends in boreal forests. These results emphasize the importance of 10 considering water availability in a new generation of phenology modules in DGVMs in order 11 to correctly reproduce observed seasonal to decadal dynamics of vegetation greenness.

12

13 **1** Introduction

The greenness of the terrestrial vegetation is directly linked to plant productivity, surface 14 15 roughness and albedo and thus affects the climate system (Richardson et al., 2013). Vegetation greenness can be quantified from satellite observations for example as Normalized 16 17 Difference Vegetation Index (NDVI) (Tucker, 1979). NDVI is a remotely sensed proxy for structural plant properties like leaf area index (LAI) (Turner et al., 1999), green leaf biomass 18 19 (Gamon et al., 1995) and plant productivity. Especially, NDVI of green vegetation has a 20 linear relationship with the fraction of absorbed photosynthetic active radiation (FAPAR) 21 (Fensholt et al., 2004; Gamon et al., 1995; Myneni et al., 1995, 1997b; Myneni and Williams, 22 1994). Satellite-derived FAPAR estimates are often used to estimate terrestrial photosynthesis (Beer et al., 2010; Jung et al., 2008, 2011; Potter et al., 1999). Decadal 23 24 satellite observations of NDVI demonstrate widespread positive trends ("greening") especially in the high latitude regions (Lucht et al., 2002; Myneni et al., 1997a; Xu et al., 25 26 2013) but also in the Sahel, southern Africa and southern Australia (Fensholt and Proud, 27 2012; de Jong et al., 2011, 2013b). Surprisingly, these trends are accompanied by negative 28 trends ("browning") which were observed regionally in parts of the boreal forests of North 29 America and Eurasia, and in parts of eastern Africa and South America (Baird and Verbyla, 30 2012; Bi et al., 2013; de Jong et al., 2013b). Regionally different causes have been identified 31 for the observed greening and browning trends. The greening of the high latitudes is supposed

to be mainly induced by rising air temperatures (Lucht et al., 2002; Myneni et al., 1997a; Xu 1 2 et al., 2013). Browning trends in subtropical regions were related to changing drought 3 conditions and land use change (Cook and Pau, 2013; van Leeuwen et al., 2013). On the other 4 hand, the environmental controls on the browning of boreal forests have been intensively 5 investigated but no concluding or general explanation has been found so far (Barichivich et al., 2014; Beck et al., 2011; Beck and Goetz, 2011; Bunn et al., 2007; Goetz et al., 2005; Piao 6 7 et al., 2011; Wang et al., 2011). Trends in vegetation greenness are often related to changes 8 in vegetation phenology like an earlier onset and an associated lengthening of the growing 9 season in mid- and high-latitude regions (Atzberger et al., 2013; Høgda et al., 2001, 2013; 10 Tucker et al., 2001; Zeng et al., 2011). Changes in vegetation greenness are linked to changes 11 in primary production and thus affect atmospheric CO₂ concentrations and the terrestrial 12 carbon cycle (Barichivich et al., 2013; Keeling et al., 1996; Myneni et al., 1997a). 13 Additionally, vegetation greenness affects the climate system by influencing surface albedo. 14 For example, greening trends in high-latitudes are associated with decreasing surface albedo 15 (Urban et al., 2013) which alters the surface radiation budget (Loranty et al., 2011). This can 16 potentially further contribute to a warming of arctic regions (Chapin et al., 2005). Thus, 17 satellite observations of vegetation greenness demonstrate the recent interactions and changes 18 between terrestrial vegetation dynamics and the climate system.

19 Dynamic global vegetation models (DGVM) or generally climate/carbon cycle models are 20 used to analyze and project the response of the terrestrial vegetation to the past, recent and 21 future climate variability (Prentice et al., 2007). DGVMs can be used to explain observed trends in vegetation greenness (Lucht et al., 2002) or to quantify the related terrestrial CO₂ 22 uptake. While most global models simulate an increasing uptake of CO₂ by the terrestrial 23 vegetation under future climate change scenarios, the magnitude of future changes in land 24 25 carbon uptake largely differs among models (Friedlingstein et al., 2006; Sitch et al., 2008). 26 The spread of land carbon uptake estimates among DGVMs might be partly related to 27 insufficient representations of vegetation phenology and greenness (Richardson et al., 2012). Coupled climate-carbon cycle models and uncoupled DGVMs have been compared against 30 28 29 year satellite-derived time series of LAI (Anav et al., 2013; Murray-Tortarolo et al., 2013; 30 Zhu et al., 2013). Models usually overestimate mean annual LAI in all biomes and have a too 31 long growing season because of a delayed season end (Anav et al., 2013; Murray-Tortarolo et 32 al., 2013; Zhu et al., 2013). Additionally, most DGVMs have more positive LAI trends than

the satellite-derived LAI product, i.e. they underestimate browning trends in boreal forests 1 2 while a few DGVMs do not reproduce the general greening of the high latitudes (Murray-3 Tortarolo et al., 2013). The limitations of DGVMs in reproducing observed LAI or FAPAR 4 time series is mostly related to limited phenology routines that often miss environmental 5 controls on seasonal leaf development (Kelley et al., 2013; Murray-Tortarolo et al., 2013; 6 Richardson et al., 2012). In conclusion, with improved modelling approaches for vegetation 7 phenology and greenness, DGVMs can potentially more accurately reproduce the recent, and 8 project the future response of the terrestrial vegetation to climate variability.

9 Past studies successfully demonstrated the use of vegetation greenness observations to 10 improve stand-alone phenology models or to optimize phenology and productivity-related 11 parameters in DGVMs. The growing season index (GSI) is an empirical phenology model that is used to estimate seasonal leaf developments (Jolly et al., 2005). Empirical parameters of 12 13 GSI have been optimized against globally distributed 10 year FAPAR and LAI time series 14 from MODIS to reanalyze climatic drivers for vegetation phenology (Stöckli et al., 2008, 15 2011). This optimization resulted in a good representation of temporal FAPAR and LAI 16 dynamics in all major biomes except evergreen tropical forests (Stöckli et al., 2011). Model 17 parameters of the Biome-BGC model were optimized against eddy covariance flux observations and NDVI time series from MODIS for poplar plantations in Northern Italy 18 19 which resulted in a more accurate representation of carbon fluxes and NDVI (Migliavacca et 20 al., 2009). The BETHY-CCDAS model was optimized against FAPAR time series from 21 MERIS for seven eddy covariance sites (Knorr et al., 2010) and later for 170 land grid cells using coarse 8 by 10° spatial resolution (Kaminski et al., 2012). These studies demonstrated 22 23 the improvements in simulated vegetation phenology by optimizing model parameters against 24 observations of vegetation greenness.

25 Nevertheless, spatial patterns and temporal dynamics of vegetation greenness were not yet 26 optimized in a DGVM globally at a higher spatial resolution (0.5°) and by using long-term 27 (30 year) satellite-derived time series of vegetation greenness. Newly developed 30 year time series of LAI or FAPAR from the GIMMS3g dataset (Global Inventory Modeling and 28 Mapping Studies, 3rd generation of datasets) (Zhu et al., 2013) allow improving DGVMs not 29 only based on seasonal cycles of single years (i.e. phenology) but additionally against decadal 30 31 time series properties including inter-variability and trends. By integrating the GIMMS3g 32 FAPAR data set in a DGVM, we can potentially improve spatial patterns and seasonal to

long-term temporal dynamics of vegetation greenness. We are using the LPJmL DGVM
 (Lund-Potsdam-Jena managed lands). Similar to other DGVMs, LPJmL does not accurately
 reproduce the growing season onset and seasonal amplitude of observed LAI and FAPAR
 time series presumably because of a limited phenology model (Kelley et al., 2013; Murray Tortarolo et al., 2013). Thus integrating long-term observations of FAPAR in the LPJmL
 DGVM potentially requires the development of an improved phenology scheme.

7 We are aiming to improve environmental controls on vegetation phenology and greenness in 8 LPJmL by 1) developing a new phenology module for LPJmL, by 2) optimizing FAPAR, 9 productivity and phenology-related parameters of LPJmL against 30 year satellite-derived time series of FAPAR, against 10 year satellite-derived time series of vegetation albedo and 10 11 against spatial patterns of mean annual gross primary production (GPP) from a data-oriented estimate and by 3) integrating further data streams in LPJmL to constrain land cover 12 13 dynamics and disturbance effects on vegetation greenness in diagnostic model simulations. This model-data integration approach for LPJmL (LPJmL-MDI) will be applied to identify 14 15 the environmental controls on vegetation greenness phenology.

16

17 2 Model, data sets and model-data integration

18 **2.1 Overview**

19 LPJmL is a dynamic global vegetation model that simulates ecosystem processes as carbon 20 and water fluxes, carbon allocation in plants and soils, permafrost dynamics, fire spread and 21 behaviour and vegetation establishment and mortality. We were using LPJmL version 3.5. 22 This version is based on the original LPJ model (Sitch et al., 2003). The model has been 23 extended for human land use (Bondeau et al., 2007), and agricultural water use (Rost et al., 2008). It includes a process-oriented fire model (Thonicke et al., 2010), an improved 24 25 representation of surface albedo and snow coverage (Strengers et al., 2010) and a newly 26 implemented soil hydrology scheme and permafrost module (Schaphoff et al., 2013). This 27 study focusses on the natural vegetation plant functional types (PFTs) (Sitch et al., 2003), i.e. 28 our model developments and optimizations were not applied for crop functional types (CFTs) 29 (Bondeau et al., 2007) because crop phenology is highly driven by human practices.

1 We developed a model-data integration approach for the LPJmL DGVM (LPJmL-MDI, 2 Figure 1). LPJmL-MDI allows to 1) directly insert land cover, tree cover and burnt area data 3 sets in LPJmL for diagnostic model applications (section 2.4.1); 2) to optimize LPJmL model 4 parameters against datasets (here FAPAR, GPP, albedo; section 2.4.2); and 3) to evaluate and 5 benchmark LPJmL simulations against observations or observation-based data sets (section 6 2.4.3). Like in a prognostic mode, LPJmL was driven by climate forcing data. Additionally, 7 observed burnt areas were directly inserted in LPJmL to consider observed fire dynamics in 8 diagnostic model applications. For this, we directly replaced the simulated burnt area in the 9 LPJmL-SPITFIRE fire module (Thonicke et al., 2010) by observed burnt areas using the 10 approach of Lehsten et al. (2008). Thus, the timing and location of fire spread is constrained 11 by observations whereas fire effects on vegetation are still simulated by LPJmL-SPITFIRE. 12 We further prescribed observed land cover and tree cover fractions to control for vegetation dynamics in parameter optimization experiments. Observed FAPAR and albedo time series as 13 14 well as observation-based mean annual spatial patterns of GPP were used in a joint cost 15 function to optimize productivity, phenology, radiation, and albedo-related model parameters 16 using a genetic optimization algorithm.

17 LPJmL was previously evaluated against site measurements of net carbon ecosystem 18 exchange (Schaphoff et al., 2013; Sitch et al., 2003), atmospheric CO₂ fractions (Sitch et al., 19 2003), soil moisture (Wagner et al., 2003), evapotranspiration and runoff (Gerten et al., 2004; 20 Schaphoff et al., 2013), fire regimes (Thonicke et al., 2010), and permafrost distribution 21 (Schaphoff et al., 2013). Here we evaluate LPJmL against additional and partly new available 22 global data sets of FAPAR (Baret et al., 2013; Zhu et al., 2013), GPP and evapotranspiration (ET) (Jung et al., 2011), tree cover (Townshend et al., 2011) and biomass (Carvalhais et al., 23 24 2014; Saatchi et al., 2011; Thurner et al., 2014).

25 2.2 FAPAR and phenology in the LPJmL DGVM

26 **2.2.1 FAPAR**

FAPAR is defined as the ratio between the photosynthetic active radiation absorbed by the green canopy (APAR) and the total incident photosynthetic active radiation (PAR). Thus, the total FAPAR of a grid cell is the sum of FAPAR that is distributed among the individual PFTs:

$$1 \qquad FAPAR_{PFT} = \frac{APAR_{PFT}}{PAR} \tag{1}$$

2
$$FAPAR_{gridcell} = \sum_{PFT=1}^{PFT=n} FAPAR_{PFT}$$
 (2)

3 where *n* is the number of established PFTs in a grid cell. The FAPAR of a PFT depends on 4 the annual maximum foliar projective cover (FPC), on the daily snow coverage in the green 5 canopy ($F_{snow,gv}$), green-leaf albedo (β_{leaf}) and the daily phenology status (*Phen*):

6
$$FAPAR_{PFT} = FPC_{PFT} \times (Phen_{PFT} - (Phen_{PFT} \times F_{snow,gv,PFT})) \times (1 - \beta_{leaf,PFT}) (3)$$

7 Thus, the temporal dynamic of FAPAR in LPJmL is affected on an annual time step by 8 changes in foliar projective cover (FPC_{PFT}) and on daily time steps by changes in phenology 9 (*Phen*_{PFT}) and snow coverage in the green canopy ($F_{snow,gv,PFT}$) (Figure S1). This approach 10 extends the previous implementation of Sitch et al. (2003) where FAPAR depends only on 11 FPC and phenology but leaf albedo and snow effects on FAPAR were missing.

12 FPC_{PFT} expresses the land cover fraction of a PFT. It is the annual maximum fractional green 13 canopy coverage of a PFT and is annually calculated from crown area (*CA*), population 14 density (*P*) and LAI (Sitch et al., 2003):

15
$$FPC_{PFT} = CA_{PFT} \times P_{PFT} \times (1 - e^{-k_{PFT} \times LAI_{PFT}})$$
 (4)

16 The last term expresses the light extinction in the canopy which depends exponentially on LAI and the light extinction coefficient k of the Lambert-Beer law (Monsi and Saeki, 1953). 17 18 The parameter k had a constant value of 0.5 for all PFTs in the original LPJmL formulation 19 (Sitch et al., 2003). We changed k to a PFT-dependent parameter because it varies for 20 different plant species as seen from field observations (Bolstad and Gower, 1990; Kira et al., 21 1969; Monsi and Saeki, 1953). Crown area and leaf area index are calculated based on 22 allocation rules and are depending on the annual biomass increment (Sitch et al., 2003). 23 Population density depends on establishment and mortality processes in LPJmL (Sitch et al., 24 2003).

1 2.2.2 Phenology

The daily phenology and green leaf status of a PFT (*Phen_{PFT}*) in LPJmL expresses the fractional cover of green leafs (from 0 = no leafs to 1 = full leaf cover). Thus, it represents the temporal dynamic of the canopy greenness. We explored two phenology models in this study: First, we were trying to optimize model parameters of the original phenology module in LPJmL (LPJmL-OP, Sitch et al., 2003). Secondly, we implemented a new phenology module based on the growing season index (GSI) concept (Jolly et al., 2005), hereinafter called LPJmL-GSI.

9 LPJmL-OP has three different routines for summergreen (i.e. temperature-driven deciduous),
10 evergreen (no seasonal variation) and rain-green (i.e. water-driven deciduous) PFTs (details in
11 Supplement 1.1). Obviously, LPJmL-OP misses important controls on phenology like effects
12 of light in all PFTs or effects of water in summergreen and herbaceous PFTs. Additionally, in
13 herbaceous PFTs the end of the growing season is not controlled by environmental conditions
14 but is defined based on fixed calendar dates.
15 Because of the obvious limitations of LPJmL-OP, we developed the alternative LPJmL-GSI

16 phenology module. The growing season index (GSI) is an empirical phenology model that 17 multiplies limiting effects of temperature, day length and vapour pressure deficit (VPD) to a common phenology status (Jolly et al., 2005). We modified the GSI concept for the specific 18 19 use in LPJmL (LPJmL-GSI). We defined the phenology status as a function of cold 20 temperature, short-wave radiation and water availability. Additionally to the original GSI 21 model, we added a heat stress limiting function because it has been suggested that vegetation 22 greenness is limited by temperature-induced heat stress in several ecosystems (Bunn et al., 23 2007; Verstraeten et al., 2006) and has been demonstrated that heat stress reduces plant productivity also without additional water stress (Jiang and Huang, 2001; Van Peer et al., 24 25 2004; Poirier et al., 2012). Thus, the daily phenology status of a PFT is the product of the 26 daily cold temperature ($f_{cold,PFT}$), light ($f_{light,PFT}$), water ($f_{water,PFT}$) and heat stress ($f_{heat,PFT}$) 27 limiting functions:

28
$$Phen_{PFT} = f_{cold,PFT} \times f_{light,PFT} \times f_{water,PFT} \times f_{heat,PFT}$$
 (5)

- 29 Examples for the four functions are shown in Figure 2.
- 30 The cold temperature limiting function at a daily time step *t* is defined as:

$$1 \qquad f_{cold,PFT}^{t} = f_{cold,PFT}^{t-1} + \left(\frac{1}{1 + e^{-sl_{cold,PFT} \times (T - base_{cold,PFT})}} - f_{cold,PFT}^{t-1}\right) \times \tau_{cold,PFT}$$
(6)

where $sl_{cold,PFT}$ and $base_{cold,PFT}$ are PFT-dependent slope and inflection point parameters of a logistic function based on mean daily air temperature T (°C). The parameter $\tau_{cold,PFT}$ is the change rate parameter based on the difference between the actual predicted limiting function value and the previous-day cold temperature limiting function value. This parameter introduces a temporal autocorrelation in the phenology status and avoids abrupt phenological changes because of changing weather conditions.

8 The light-limiting function was implemented accordingly:

9
$$f_{light,PFT}^{t} = f_{light,PFT}^{t-1} + \left(\frac{1}{1 + e^{-sl_{light,PFT} \times (SW - base_{light,PFT})}} - f_{light,PFT}^{t-1}\right) \times \tau_{light,PFT}$$
(7)

10 where $sl_{light,PFT}$ and $base_{light,PFT}$ are the PFT-dependent slope and inflection point parameters of 11 a logistic function based on daily shortwave downward radiation SW (W m⁻²). The parameter 12 $\tau_{light,PFT}$ is the temporal change rate for the light-limiting function.

13 The water-limiting function $f_{water,PFT}$ depends on the daily water availability W(%) in LPJmL:

14
$$f_{water,PFT}^{t} = f_{water,PFT}^{t-1} + \left(\frac{1}{1 + e^{-sl_{water,PFT} \times (W - base_{water,PFT})}} - f_{water,PFT}^{t-1}\right) \times \tau_{water,PFT}$$
15 (8)

16 where $sl_{water,PFT}$ and $base_{water,PFT}$ are the PFT-dependent slope and inflection point parameters 17 of a logistic function based on daily water availability. *W* is a ratio between water supply from 18 soil moisture and atmospheric water demand (Supplement 1.2) (Gerten et al., 2004). The 19 parameter $\tau_{water,PFT}$ is the temporal change rate for the water-limiting function.

The heat-stress limiting function is defined as the cold-temperature limiting function based ondaily air temperature but with a negative slope parameter:

22
$$f_{heat,PFT}^{t} = f_{heat,PFT}^{t-1} + \left(\frac{1}{1 + e^{sl_{heat,PFT} \times (T - base_{heat,PFT})}} - f_{heat,PFT}^{t-1}\right) \times \tau_{heat,PFT}$$
(9)

where $sl_{heat,PFT}$ and $base_{heat,PFT}$ are the PFT-dependent slope and inflection point parameters of a logistic function based on *T*. The parameter $\tau_{heat,PFT}$ is the temporal change rate for the heat limiting function.

1 Besides the additional use of the heat stress limiting function, LPJmL-GSI has important 2 differences to the original GSI phenology model (Jolly et al., 2005). We made the water limiting function dependent on water availability. VPD has been used instead in the original 3 4 GSI phenology model. Nevertheless, it has been shown that phenology is more driven by soil 5 moisture and plant available water than by atmospheric water demand especially in 6 Mediterranean and grassland ecosystems (Archibald and Scholes, 2007; Kramer et al., 2000; 7 Liu et al., 2013; Yuan et al., 2007) and that GSI performed better when using a soil moisture 8 limiting function instead of the VPD limiting function (Migliavacca et al., 2011). With the 9 implementation of the water limiting function in LPJmL-GSI, phenology depends not only on 10 atmospheric water demand as in the original GSI model but also on water supply from soil 11 moisture. Additionally, the soil moisture can be modulated through seasonal freezing and 12 thawing in permafrost soils according to the permafrost routines in LPJmL (Schaphoff et al., 2013). Another important difference to the original GSI phenology model is the use of logistic 13 14 functions instead of stepwise linear functions with fixed thresholds because smooth functions 15 are generally easier to optimize than functions with abrupt thresholds and potentially better 16 represent biological processes. A moving average of 21 days has been used in the original 17 GSI model to create smooth phenological cycles and to avoid abrupt phenology changes 18 because of daily weather variability (Jolly et al., 2005). It has been shown that PFT- and 19 limiting function-dependent time averaging parameters are needed instead of one single time 20 averaging parameter (Stöckli et al., 2011). We implemented change rate parameters τ_{cold} , τ_{light} , 21 τ_{water} and τ_{heat} that are PFT- and limiting function-dependent instead of moving average 22 window lengths because LPJmL cannot use the same running window time averaging 23 approach as a prognostic model.

24

25 2.3 Data sets

26 **2.3.1** Data sets for parameter optimization: FAPAR, albedo and GPP

We used FAPAR, albedo and GPP data sets to optimize phenology, FAPAR, productivity and vegetation albedo-related parameters in LPJmL (Figure 2). We require long-term FAPAR datasets to improve vegetation greenness in LPJmL on seasonal to decadal time scales. Two recently developed datasets provide 30-year time series of FAPAR. The Geoland2 BioPar (GEOV1) FAPAR dataset (Baret et al., 2013) (hereinafter called GL2 FAPAR) and the
 GIMMS3g FAPAR (Zhu et al., 2013) datasets were used in this study.

3 GL2 FAPAR is defined as the black-sky green canopy FAPAR at 10:15 solar time and has been produced based on SPOT VGT (1999-2012) and AVHRR (1981-2000) observations 4 5 (Baret et al., 2013). The GL2 FAPAR dataset has a temporal resolution of 10 days and a spatial resolution of 0.05° for the AVHRR-period and of 1/112° for the SPOT VGT period. 6 GIMMS3g FAPAR corresponds to black-sky FAPAR at 10:35 solar time and has been 7 8 produced based on the GIMMS3g NDVI dataset (Pinzon and Tucker, 2014; Zhu et al., 2013). 9 GIMMS3g FAPAR has a 15-day temporal resolution and a 1/12° spatial resolution and covers July 1981 to December 2011. We excluded in both FAPAR datasets observations that were 10 11 flagged as contaminated by snow, aerosols or clouds. Additionally, we excluded FAPAR observations for months with temperatures $< 0^{\circ}$ C to exclude potential remaining distortions of 12 13 snow cover. Both datasets were aggregated to a 0.5° spatial and monthly temporal resolution 14 to be comparable with LPJmL simulations. We found that the GL2 AVHRR and GL2 VGT 15 FAPAR datasets have not been well harmonized (Supplement 2.1). Thus, we did not use the 16 combined GL2 VGT and AVHRR FAPAR dataset for parameter optimization and for 17 analyses of inter-annual variability and trends but only for analyses and evaluations of mean seasonal cycles and spatial patterns of FAPAR. The GIMMS3g FAPAR dataset has no 18 19 uncertainty estimates. Uncertainty estimates are necessary in multiple data stream parameter 20 optimization to weight single data streams in the total cost function. As a workaround we 21 estimated the uncertainty based on monthly-varying quantile regressions to the 0.95 quantile 22 between FAPAR and the FAPAR uncertainty in the GL2 VGT dataset. We applied the fitted 23 regressions to the GIMMS3g dataset to estimate FAPAR uncertainties (Supplement 2.2). The 24 fit to the upper quantile provides conservative uncertainty estimates for the GIMMS3g 25 FAPAR dataset.

We used monthly shortwave white-sky albedo time series ranging from 2000 to 2010 from the MODIS C5 dataset (Lucht et al., 2000; Schaaf et al., 2002) to constrain vegetation albedo parameters. Albedo observations in months with < 5°C air temperature and above an albedo of 0.3 were excluded from optimization because we are optimizing only vegetation-related albedo parameters. High albedo values at low temperatures are probably affected by changing snow regimes which is not within our focus of model development and optimization. Thus we are only optimizing growing season albedo.

We used mean annual total GPP patterns from the data-oriented MTE (model tree ensemble) 1 2 GPP estimate (Jung et al., 2011). This GPP estimate uses FLUXNET eddy covariance 3 observations together with satellite observations and climate data to upscale GPP using a 4 machine learning approach (Jung et al., 2011). This dataset is not an observation but a result 5 of an empirical model. Nevertheless, evaluation and cross-validation analyses have shown 6 that this dataset well represents the mean annual spatial patterns and mean seasonal cycles of 7 GPP whereas it has a poor performance in representing temporal GPP anomalies (trends and 8 extremes) (Jung et al., 2011). Thus, we are only using the mean annual total GPP from this 9 dataset for parameter optimization to constrain LPJmL within small biases of mean annual 10 GPP. We used the mean seasonal cycle from the MTE GPP product as an independent 11 benchmark for model evaluation.

12

2.3.2 Data sets for the prescription of land cover, tree cover and burnt area

The FAPAR, albedo and GPP data sets do not presumably contain enough information to constrain all processes that control FAPAR dynamics. Especially, processes like establishment, mortality, competition between PFTs, allocation and disturbances control FPC and thus FAPAR. The optimization of parameters of these processes against appropriate data streams is not feasible within this study. Thus, we directly prescribed land and tree cover fractions as well as burnt areas from observed data to control for some of these processes.

21 To prescribe land and tree cover in LPJmL, we combined several datasets to create 22 observation-based maps of FPC (Supplement 3.1). Land cover maps from remote sensing 23 products are not directly comparable with PFTs in global vegetation models due to 24 differences in classification systems (Jung et al., 2006; Poulter et al., 2011a). PFTs in LPJmL 25 are defined according to biome (tropical, temperate or boreal), leaf type (needle-leaved, broadleaved) and phenology type (summergreen, evergreen, rain green). We extracted the 26 27 biome information from the Köppen-Geiger climate classification (Kottek et al., 2006) 28 whereas leaf type and phenology were extracted from the SYNMAP land cover map (Jung et 29 al., 2006). FPC was derived from MODIS tree cover (Townshend et al., 2011). Because 30 LPJmL so far classified herbaceous vegetation according to their photosynthetic pathway (i.e. 31 C_3 , temperate herbaceous and C_4 , tropical herbaceous), we further sub-divided herbaceous PFTs according to biome and introduced a polar herbaceous PFT (PoH) based on the existing
 temperate herbaceous PFT (TeH) to differentiate tundra from temperate grasslands.

3 Burnt area data was prescribed directly in LPJmL by combining three data sets, the Global Fire Emissions Database (GFED) burnt area dataset (Giglio et al., 2010), the Alaska Large 4 5 Fire Database (ALFDB) (Frames, 2012; Kasischke et al., 2002) and the Canadian National Fire Database (CNFDB) (CFS, 2010; Stocks et al., 2002). GFED provides monthly burnt area 6 estimates in 0.5° resolution from 1996 to 2011. Burnt areas from the Alaska (ALFDB) and 7 8 Canada (CNFDB) fire databases were used to extent burnt area time series before 1996 for 9 boreal North America. Fire perimeter observations from 1979 to 1996 from ALFDB and CNFDB were aggregated to $0.5^{\circ} \times 0.5^{\circ}$ gridded monthly burnt area time series. Observations 10 11 before 1979 were excluded because fires were not reported for all provinces in Canada. Although the CNFDB contains only fire perimeters > 200ha, in both databases some fires are 12 13 missing due to different mapping techniques, and fire perimeters do not agree with burned 14 area, the integration of these datasets provides unique information about spatial-temporal 15 patterns of disturbances especially in boreal ecosystems. It is necessary to simulate fire 16 activity also during the model spin-up as fire influences the equilibrium between vegetation, 17 soil and climate as well. Otherwise biomass would be overestimated at the beginning of the 18 transient model run. For this purpose, we created artificial burnt area time series for the 19 periods 1901-1978 (North America) and 1901-1995 (rest of the world). For this observed 20 annual total burnt areas from the periods 1979-2011 (North America) and 1996-2011 (rest of 21 the world) were resampled according to temperature and precipitation conditions and assigned 22 to the pre-data period in order to include fire regimes that agree with observed fire regimes in 23 the spin-up of LPJmL. This approach assumes that fire regimes in the pre-data period were 24 not different than in the observation period.

25

26 **2.3.3 Data sets for model evaluation**

LPJmL was evaluated against data sets that are independent from the optimization and prescription data sets and against independent temporal or spatial scales of the optimization and prescription data sets. We compared LPJmL against mean annual patterns and mean seasonal cycles of ET from the MTE estimate (Jung et al., 2011). Further, we evaluated model results against spatial patterns of biomass. Ecosystem biomass estimates were taken from satellite-derived forest biomass maps for the tropics (Saatchi et al., 2011) and for the temperate and boreal forests (Thurner et al., 2014) including an estimation of herbaceous biomass (Carvalhais et al., 2014). Additionally, we evaluated LPJmL against independent temporal and spatial scales of the integration data (mean seasonal cycle of GPP, tree cover, inter-annual variability and trends of FAPAR). We were using tree cover from MODIS to evaluate LPJmL model runs with dynamic vegetation.

7

8 2.3.4 Climate forcing data and model spin-up

9 LPJmL was driven by observed monthly temperature and precipitation data from the CRU
10 TS3.1 dataset ranging from 1901 to 2011 (Harris et al., 2013) as well as by monthly
11 shortwave downward radiation and long wave net radiation re-analysis data from ERA12 Interim (Dee et al., 2011).

LPJmL needs a model spin-up to establish PFTs and to bring vegetation and soil carbon pools into equilibrium. The spin-up was performed according to the standard LPJmL modelling protocol (Schaphoff et al., 2013; Thonicke et al., 2010): LPJmL was run for 5000 years by repeating the climate data from 1900-1930. After the spin-up model run, the transient model run was restarted from the spin-up conditions in 1901 and LPJmL was run for the period 1901-2011. Model results were analyzed for the observation period (1982-2011).

19 For model optimization experiments we used a different spin-up scheme because the spin-up 20 is computational time demanding and many model runs are needed during optimization 21 experiments. As in the standard modelling protocol, we firstly spin-up the model for 5000 22 years by repeating the climate from 1901-1930. Secondly, a transient model run was restarted 23 from the spin-up conditions in 1901 and was performed for the period 1901-1979. Thirdly, 24 each optimization experiment was restarted from the conditions in 1979 and a second spin-up 25 for 100 years by recycling the climate from 1979 to 1988 was performed. The transient model 26 run was restarted from the conditions of the second spin-up and simulated for the period 27 1979-2011. This second spin-up is needed to bring the vegetation into a new equilibrium 28 which can be caused by a new parameter combination during optimization. From visual 29 analyses of model results, we found that a spin-up time of 100 years for the second spin-up 30 was enough to eliminate trends in FAPAR and GPP that resulted from other equilibrium 31 conditions.

2 2.4 Model-data integration

3 **2.4.1** Prescription of land and tree cover

4 Land cover is expressed as FPC in LPJmL. We used the observation-based FPC dataset to prescribe land and tree cover in LPJmL (section 2.3.2, Supplement 3.1). The presence of a 5 6 PFT in a grid cell depends on establishment and mortality in LPJmL (Sitch et al., 2003). A 7 PFT establishes in a grid cell if the climate is within the bioclimatic limits of the PFT for 8 establishment and survival. On the other hand, a PFT dies in a grid cell if the climate is no 9 longer suitable for the PFT. Additionally, mortality occurs because of heat stress, low 10 productivity, competition among PFTs for light, and because of fire disturbance (Sitch et al., 2003; Thonicke et al., 2010). 11

12 FPC is the major variable that contributes to inter-annual variability of FAPAR in LPJmL 13 despite the daily phenological status. Thus fixing FPC to the observed value is not a desired solution to prescribe land cover in LPJmL. Fixing FPC would neglect mortality effects on 14 15 land cover but would also permit the simulation of post-fire succession trajectories. Consequently, we prescribed land cover in LPJmL using a hybrid diagnostic-dynamic 16 17 approach. In this approach we prescribed the annual maximum FPC in LPJmL similar to 18 previous approaches (Poulter et al., 2011b). Firstly, we switched off the effects of bioclimatic 19 limits on establishment and mortality. Only these PFTs were allowed to establish in a grid cell 20 that occurred in the observed land-cover data set. Vegetation growth depends on the annual 21 biomass increment and allocation rules in LPJmL. This leads to an extension of FPC of each 22 PFT. We limited a further expansion of FPC if the simulated FPC exceeded the observed FPC 23 by replacing the simulated FPC with the observed FPC (prescribed maximum FPC). 24 Consequently, the simulated FPC can be lower than the observed FPC because the PFT is still 25 growing or because the FPC was reduced due to fire, heat stress or low productivity. For 26 herbaceous PFTs we only reduced the FPC if the observed total fractional vegetation cover in 27 a grid cell was exceeded. This allowed herbaceous PFTs to replace tree PFTs if the FPC of 28 trees was reduced due to fire or other mortality effects in the model. With this approach a 29 prescription of land cover can be achieved in LPJmL which well represents observed PFT 30 distributions (Supplement 3.2) but still allows for main processes of dynamic vegetation.

2 **2.4.2 Parameter optimization**

3 Photosynthesis, albedo, FAPAR and phenology-related model parameters of LPJmL were 4 optimized against observed FAPAR and albedo satellite observations and data-oriented 5 estimates of GPP. A description of all parameters including parameter values is given in Supplement 4.1. The parameter α_a is the most important parameter in LPJmL for 6 7 photosynthesis (Zaehle et al., 2005). This parameter accounts for the amount of radiation that 8 is absorbed at leaf level in comparison to the total canopy. Thus, this parameter is a 9 replacement for a more enhanced model formulation for canopy structure and leaf clumping. 10 We used this parameter to adjust biases in GPP. The PFT-dependent leaf, stem and litter albedo parameters (β_{leaf} , β_{stem} and β_{litter}) are mostly sensitive for model simulations of albedo. 11 12 The parameter β_{leaf} affects additionally the maximum FAPAR of a PFT. The light extinction coefficient k controls the FPC of a PFT and thus affects mainly land cover, maximum FAPAR 13 and the available radiation for photosynthesis. All other parameters that were considered in 14 optimization experiments are the parameters of the LPJmL-OP and LPJmL-GSI phenology 15 16 modules. These parameters contribute mainly to seasonal variations in FAPAR. Some 17 parameters were excluded from optimization experiments that were identified as insensitive to 18 GPP and FAPAR simulations in PFTs. The temporal change rate parameters τ_{tmin} , τ_{light} , τ_{heat} 19 and τ_{water} are insensitive in most PFTs because of the monthly temporal resolution of the used 20 climate forcing data.

The optimization of model parameters was performed by minimizing a cost function between model simulations and observations using a combined genetic and gradient-based optimization algorithm (GENOUD, genetic optimization using derivatives, Mebane and Sekhon, 2011, see Supplement 4.2 for details). The cost function *J* of LPJmL for a single model grid cell (*gc*) depends on the scaled model parameter vector *d* (*d* = proposed parameter value / prior parameter value) and is the sum of square error (SSE) between model simulation and observation weighted by the number of observations (*nobs*) for each data stream (DS):

$$28 J(d)_{gc} = \sum_{DS=1}^{DS=n} \frac{SSE_{DS}(d)}{nobs_{DS}} (11)$$

The SSE for a single data stream is calculated from the LPJmL simulation of this data stream
 (x_{LPJmL}) and the corresponding observed values (x_{obs}) weighted by the uncertainty of the
 observations (x_{obsunc}) for each time step *t*:

4
$$SSE(d) = \sum_{t=1}^{t=n} \frac{(x_{LPJmL,t}(d \times p_0) - x_{obs,t})^2}{x_{obsunc,t}^2}$$
 (12)

5 where p_0 are LPJmL prior parameters. That means the minimization of the cost function *J* is 6 based on scalars of LPJmL parameters relative to the prior parameter values.

7 Different model optimization experiments were performed for individual grid cell and for 8 multiple grid cells of the same PFT for LPJmL-OP as well as for LPJmL-GSI (Table 1). In the 9 grid cell-based optimization experiments model parameters of the established target tree PFT 10 and the established herbaceous PFT were optimized at the same time. The purpose of grid 11 cell-level optimization experiments was to explore the variability of parameters within 12 different regions and PFTs. In the PFT-level optimization experiments the cost of LPJmL was 13 calculated as the sum of the cost for each grid cell weighted by the grid cell area A:

14
$$J(d)_{PFT} = \frac{\sum_{gc=1}^{gc=n} J(d)_{gc} \times A_{gc}}{\sum_{gc=1}^{n} A_{gc}}$$
 (13)

For PFT-level optimizations parameters of herbaceous PFTs were first optimized for grid cells where only the herbaceous PFT was dominant. In a second step, the optimized parameters of the herbaceous PFTs were used in the optimization of the target tree PFT (Figure S9). The purpose of PFT-level optimization experiments is to derive optimized parameter sets that can be used for one PFT in global model runs.

For grid cell as well PFT-level optimization experiments, we only used grid cells that are vegetated, dominated by one PFT and that are only marginally affected from agricultural use or fire disturbances. These grid cells are called candidate grid cells in the following. We randomly selected grid cells from the set of candidate grid cells to perform grid cell- or PFTlevel optimization experiments. Table 1 gives an overview of all optimization experiments for LPJmL-OP and LPJmL-GSI with the number of used grid cells. Grid cells that were selected for optimization experiments are also shown in Figure 3. The PFT-level optimization of LPJmL-OP (OP.pft) did not result in plausible posterior parameter sets because of structural
 limitations of the LPJmL-OP phenology model for herbaceous PFTs (i.e. no water effects,
 calendar day as end of growing season), raingreen PFT (i.e. binary phenology) and evergreen
 PFTs (i.e. constant phenology) and was therefore excluded from further analysis.

Posterior parameter sensitivities, uncertainties and correlations were explored by analyzing
the maximum likelihood and the posterior range of each parameter as derived from all
parameter sets from the genetic optimization algorithm (Supplement 4.3).

8

9

2.4.3 Model evaluation and time series analysis

10 Global model runs of LPJmL were performed in order to evaluate model results against the 11 integration data, against independent metrics of the integration data and against independent 12 data streams. We evaluated results from LPJmL-OP with standard parameters (LPJmL-OP-13 prior), from LPJmL-OP with optimized productivity, albedo and FAPAR parameters from 14 grid-cell level optimization experiments (LPJmL-OP-gc) and from LPJmL-GSI with 15 optimized parameters from PFT-level optimization experiments (Table 2). We did not use optimized phenology parameters in the LPJmL-OP-gc model run because we were not able to 16 17 derive plausible phenology parameters in optimization experiments of LPJmL-OP. All model 18 runs were performed with dynamic vegetation and prescribed burnt areas.

19 We aggregated monthly FAPAR time series to mean annual FAPAR to evaluate inter-annual 20 variability and trends. Mean annual FAPAR time series were averaged from all monthly 21 values with mean monthly air temperatures $> 0^{\circ}$ C to exclude potential remaining effects of 22 snow in the observed FAPAR time series. Trends in mean annual FAPAR time series and 23 trend breakpoints were computed using the "greenbrown" package for the R software (Forkel 24 et al., 2013). In this implementation, trends are computed by fitting piece-wise linear trends to 25 the annual FAPAR time series using ordinary least squares regression. The significance of 26 trends was computed using the Mann-Kendall trend test (Kendall, 1975; Mann, 1945).

1 **3** Results and Discussion

2 **3.1 Parameter optimization**

3 **3.1.1 Performance of phenology models**

The newly developed LPJmL-GSI phenology model resulted in significantly higher 4 5 correlations with monthly GIMMS3g FAPAR than LPJmL-OP in all PFTs except in the tropical broadleaved evergreen (TrBE) and boreal broadleaved summergreen (BoBS) PFTs 6 7 (Figure 4). LPJmL-OP with prior parameters had high correlations with monthly GIMMS3g 8 FAPAR in broad-leaved summergreen PFTs (TeBS median r = 0.87, BoBS median r = 0.92) 9 PFTs and medium correlations in boreal needle-leaved PFTs (BoNE median r = 0.53, BoNS 10 median r = 0.6). In all other PFTs, LPJmL-OP had low correlations with monthly GIMMS3g 11 FAPAR. The correlation against monthly GIMMS3g FAPAR did not significantly improve in 12 all PFTs after grid cell-level optimization experiments of LPJmL-OP (Figure 4). The use of the newly developed LPJmL-GSI phenology model already significantly improved the 13 14 correlation with monthly GIMMS3g FAPAR in all PFTs except in the temperate herbaceous 15 (TeH) and BoBS PFTs. LPJmL-GSI had significantly higher correlations with monthly 16 GIMMS3g FAPAR after grid cell-level optimization experiments in the TrBR, TeNE, TeBS, 17 TeH, BoBS and BoNS PFTs. After PFT-level optimization experiments, LPJmL-GSI had 18 median correlation coefficients > 0.5 in all PFTs except in broadleaved evergreen PFTs 19 (TrBE, TeBE). These results prove that the rain-green, evergreen and herbaceous phenology 20 schemes of LPJmL-OP were not able to reproduce temporal FAPAR dynamics despite the 21 attempt of parameter optimization and that LPJmL-GSI can reproduce seasonal FAPAR 22 dynamics in most PFTs.

23 The low correlations coefficients between LPJmL-GSI and GIMMS3g FAPAR after optimization experiments in broadleaved evergreen PFTs (TrBE, TeBE) might be caused by 24 25 the specific properties of the FAPAR dataset in these PFTs. GIMMS3g FAPAR does not have a clear seasonal cycle but a high short-term variability in broadleaved evergreen forests. 26 27 These regions are often covered by clouds that inhibit continuous optical satellite 28 observations. The high short-term variability results ultimately in low correlation coefficients 29 between both LPJmL versions (LPJmL-OP and LPJmL-GSI) and GIMMS3g FAPAR time 30 series. In temperate broadleaved evergreen forests, the GIMMS3g FAPAR dataset might have a wrong seasonality. In these regions, the mean seasonal FAPAR cycles from the GIMMS3g
and GL2 VGT FAPAR datasets are anti-correlated and FAPAR from LPJmL-GSI agrees
better with the GL2 VGT dataset. Because of these reasons, we did not expect to improve
seasonal FAPAR dynamics in broadleaved evergreen forests with the current model-data
integration setup.

6 All optimization experiments of LPJmL-OP and LPJmL-GSI resulted in a significant 7 reduction of the cost in comparison to the respective prior models (Supplement 4.4, Figure 8 S10). Nevertheless, the prior parameter set of LPJmL-GSI resulted already in a significant 9 lower cost than the grid cell-level optimized parameter sets of LPJmL-OP in tropical and 10 polar herbaceous PFTs, and in temperate broad-leaved summergreen and boreal needle-leaved 11 summergreen PFTs. The reduction of the overall cost was in all model optimization experiments usually associated with a significant reduction of the annual GPP bias (Figure 12 13 S11). LPJmL-OP with prior parameters underestimated mean annual GPP in the tropical 14 broad-leaved evergreen PFT and overestimated mean annual GPP in all other PFTs. Grid cell-15 level optimization experiments of LPJmL-OP resulted in a significant reduction of the GPP 16 bias in all PFTs except in the polar herbaceous PFT (PoH). We were not able to remove the 17 GPP bias and to reduce the cost of LPJmL-OP and of LPJmL-GSI in the PoH PFT in 18 optimization experiments because of inconsistencies between the FAPAR and GPP datasets 19 or in the LPJmL formulation. LPJmL was not able to sustain the relatively high peak FAPAR 20 in Tundra regions as seen in the GIMMS3g dataset given the low mean annual GPP of the 21 MTE dataset (Supplement 4.4). These inconsistencies might be related to higher uncertainties 22 of the GPP and FAPAR datasets in tundra regions where the MTE GPP dataset is not covered by many eddy covariance measurement sites, and where satellite-based FAPAR observations 23 24 are affected from high sun zenith angles (Tao et al., 2009; Walter-Shea et al., 1998). On the 25 other hand, dominant tundra plant communities like mosses and lichen are not represented in 26 LPJmL (Supplement 4.4). All model optimizations experiments kept growing season albedo 27 within reasonable ranges in comparison to MODIS albedo (Figure S12). These results 28 demonstrate an improved performance of optimized model parameter sets over prior model 29 parameter sets and of LPJmL-GSI over LPJmL-OP regarding a cost that is defined based on 30 30 years of monthly FAPAR, mean annual GPP and 10 years of monthly vegetation albedo.

3.1.2 Parameter sensitivities and uncertainties

The uncertainty of productivity and albedo-related parameters was reduced after optimization of LPJmL-GSI in most PFTs while the reduction of the uncertainty of phenology-related parameters depended often on plant functional type (Figure 5). Prior and posterior parameter values from each optimization experiment are listed in the Supplement (Tables S2 to S5).

6 The parameter α_a (absorption of light at leaf level in relation to canopy level) was sensitive 7 within a narrow parameter range for all PFTs. The posterior α_a parameter range was smaller 8 than the uniform prior range in all PFTs. In all optimization experiments we found for the 9 parameter α_a a gradient from high values in tropical to low values in boreal PFTs (Figure 10 S13). This pattern reflects the initial overestimation of mean annual GPP in temperate and boreal PFTs and underestimation of GPP in tropical regions with the prior parameter set of 11 12 LPJmL-OP. Thus, the low α_a parameter values probably accounts for nitrogen limitation 13 effects on productivity in boreal forests (Vitousek and Howarth, 1991) that are currently not 14 considered in LPJmL. A future implementation of nitrogen limitation processes in LPJmL 15 requires a re-optimization of the α_a parameter.

16 The leaf albedo parameter β_{leaf} was sensitive in all PFTs and the posterior β_{leaf} parameter 17 range was smaller than the prior parameter range in evergreen PFTs. In these evergreen PFTs 18 the β_{leaf} parameter was well constrained because albedo satellite observations are less affected 19 by variations in background albedo (soil, snow) than in deciduous PFTs. In all other PFTs the 20 β_{leaf} posterior parameter range was equal the prior parameter range or the optimized parameter 21 value was close to a boundary of the prior parameter range. This result indicates that the 22 albedo routines in LPJmL should consider variations in background albedo caused by changes 23 in soil properties, soil moisture, or snow conditions in order to accurately reproduce satellite-24 observed albedo time series (see supplementary discussion in Supplement 4.5). Nevertheless, 25 the optimization of the leaf albedo parameter β_{leaf} resulted in values that differed especially 26 between broadleaved and needle-leaved evergreen PFTs as well as herbaceous PFTs (Figure 27 5, Figure S14). Low leaf albedo parameters in needle-leaved evergreen PFTs (TeNE and 28 BoNE) and high leaf albedo parameters in broad-leaved summergreen and herbaceous PFTs 29 agree well with the patterns reported by Cescatti et al. (2012).

The light extinction coefficient k was sensitive for all PFTs but the posterior parameter range was only in herbaceous PFTs and in the BoBS PFT smaller than the prior parameter range

1 (Figure 5). In all PFTs this parameter had a large spatial variability (Figure S15). The 2 parameter k affects mostly the FPC and thus the maximum FAPAR. Thus, this parameter 3 cannot be well constrained for tree PFTs in the current optimization setup because the 4 maximum FPC of trees was prescribed from the land and tree cover dataset. On the other 5 hand, the maximum FPC of herbaceous PFTs was not prescribed from observations which 6 resulted in narrow k posterior parameter ranges for herbaceous PFTs. The parameter k was 7 optimized towards a very high value in the BoNS PFT (k = 0.7) due to high tree mortality 8 rates after low-productivity years (Supplement 4.5). This parameter would result in an 9 overestimated PFT coverage in model runs with dynamic vegetation. Thus, we performed a 10 second optimization experiment for this PFT (blue in Figure 5) where k_{BoNS} was limited to 0.65. This optimization experiment resulted in similar posterior values for the other 11 12 parameters. Although the k parameter was well constrained for the TrH, TeH and PoH PFTs, these parameters cannot be used in the final parameter set of LPJmL-GSI. In dynamic 13 14 vegetation model runs, the relatively low k parameter values for the TrH and TeH PFTs and 15 relatively high values for the PoH PFT would result in an underestimation of herbaceous 16 coverage in temperate and tropical climates and an overestimation of herbaceous coverage in 17 boreal and polar climates, respectively. Therefore, we performed three more optimization 18 experiments for herbaceous PFTs where we fixed k at 0.5 (blue in Figure 5). These 19 optimization experiments resulted in similar α_a parameters but different albedo parameters 20 and phenology parameters in order to compensate for biases in FAPAR and albedo that were 21 introduced by the fixed k parameter. Thus, the high spatial variability and the large 22 uncertainty of the light extinction coefficient k require re-addressing this parameter in a model 23 optimization setup with dynamic vegetation using tree and vegetation cover data or perhaps a replacement by a better representation of canopy architecture and radiative transfer. 24

25 The sensitivity and posterior uncertainty of phenology-related model parameters depended 26 often on plant functional type. The parameter base_{tmin} which controls the effect of cold 27 temperature on phenology was sensitive in all PFTs except the TrBE and TrH PFTs. The 28 posterior parameter range was smaller than the prior parameter range in temperate PFTs 29 (TeNE, TeBS and TeH). The parameter base_{heat} which controls the effect of heat stress on 30 phenology was sensitive in TrBR, TrH, TeH, BoNE and BoNS PFTs while in other PFTs this 31 parameter was only sensitive towards the boundaries of the prior parameter range. 32 Nevertheless, the posterior parameter range was only smaller than the prior parameter range

in TrBR and TrH PFTs. The parameter baselight was sensitive in temperate and boreal PFTs. In 1 tropical PFTs this parameter is only sensitive above a certain threshold (i.e. 60 W m⁻² for 2 TrBE and 100 W m⁻² for TrBR). The parameter base_{water} was sensitive in all PFTs. The 3 posterior parameter range of this parameter was smaller in all PFTs except in TeBS, BoNE, 4 BoBS and BoNS PFTs. Although, the parameter basewater had a large variability among PFTs, 5 6 it was generally optimized towards higher values in PFTs that are presumably water-7 controlled (TrBR, TeBS, TrH, TeH) and optimized towards lower values in PFTs that are 8 presumably less water controlled (TrBE, TeNE, BoNE, BoNS, PoH). This result indicates that FAPAR of water-controlled PFTs reacts already to small decreases in water availability 9 10 whereas other PFTs react only to strong decreases in water availability. We found no strong 11 correlations between posterior values of the phenology-related model parameters (Figure S16) 12 which indicates the ability to disentangle the relative effects of temperature, light and water on phenology. As the base_{water} parameter was the only phenology parameter which was 13 14 sensitive in all PFTs, this indicates that water availability is the only phenological control that acts in all PFTs. 15

16

17 **3.2** Effects of an improved phenology module in LPJmL

18 **3.2.1** Effects on seasonal and inter-annual greenness dynamics

19 LPJmL-GSI represents better the observed spatial patterns and seasonal to decadal temporal 20 dynamics of vegetation greenness (FAPAR) than LPJmL-OP (Figure 6, Supplement 5.3). 21 Whereas LPJmL-OP overestimated mean annual FAPAR in many high-latitude and semi-arid 22 regions, LPJmL-GSI was closer to both datasets and within the uncertainty of the GL2 VGT 23 FAPAR dataset in most regions and under most climate conditions (Figure S22). LPJmL-GSI 24 still overestimated mean annual FAPAR in temperate dry regions, but this overestimation was 25 reduced in comparison to LPJmL-OP.

We further observe a substantial improvement in LPJmL-GSI regarding the seasonal cycles, monthly and annual dynamics of FAPAR as retrieved from the GIMMS3g and GL2 VGT FAPAR datasets (Figure 6, Figures S23-S25). Monthly FAPAR time series from LPJmL-GSI were significantly ($p \le 0.05$) higher correlated with GIMMS3g than from LPJmL-OP in boreal forests of eastern Siberia, in the North American tundra, in temperate and tropical

1 grasslands of central Asia, North America, Australia and especially, in the Sahel (Figure 6a). 2 This is because of an improved representation of spring onset and the end of the growing 3 season in temperate and boreal forests and in herbaceous PFTs (Figure S24). The highest 4 differences between simulated and observed mean seasonal FAPAR cycles where observed in 5 the temperate broad-leaved evergreen PFT, where both model versions had opposite, although 6 insignificant, relationships to the GIMMS3g datasets. For this PFT, the observational 7 constraints are particularly problematic, where a weak agreement and opposite relationship is 8 observed between the two datasets (r = -0.48).

9 Globally, LPJmL-GSI describes better the inter-annual dynamics of GIMMS3g FAPAR when 10 compared to the previous model versions (Figure 6 b). In 20% of the land the difference to 11 other model versions is statistically significant, and in 40% does not detract from the previous model versions. This improvement in inter-annual variability is especially seen in temperate 12 13 and tropical dry regions, with sparse tree cover and grassland dominated ecosystems (western 14 United States, central Asia, the Sahel, southern Africa, and Australia) (Figure S25). In the 15 arctic, boreal and temperate climates LPJmL usually shows a higher correlation with the 16 GIMMS3g dataset than the correlation observed between both datasets (GIMMS3g and GL2 17 VGT). These results demonstrate that LPJmL-GSI can explain the inter-annual variability of 18 the GIMMS3g FAPAR dataset especially in temperate and boreal forests and temperate and 19 tropical grasslands.

Overall, the global spatial representation of phenological dynamics in LPJmL-GSI improves significantly over the previous model versions from seasonal to inter-annual time scales. Given the inclusion of water controls on phenological development, these results emphasize the importance of water availability in explaining the mean spatial patterns of vegetation greenness, but also the seasonal phenology as well as inter-annual dynamics in vegetation development.

26

3.2.2 Effects on trends in vegetation greenness

The role of different climate drivers underlying the greening and browning trends in vegetation activity is still highly debated and the dominant factors show a strong spatial variability (de Jong et al., 2013a). The consideration of different environmental controls on the phenological development in LPJmL shows a significant improvement in representing
 such dynamics when compared to the previous model formulations (Figure 7).

3 Both LPJmL-OP and LPJmL-GSI reproduced the observed greening trends in tundra regions and in boreal forests of Siberia. In both model versions this greening is mostly driven by 4 5 annual changes in foliar projective cover and effects of temperature on spring phenology. This agrees with observational studies that identified temperature increases as drivers for an 6 increasing shrub cover in tundra ecosystems (Blok et al., 2011; Forbes et al., 2010; Myers-7 Smith et al., 2011; Raynolds et al., 2013; Sturm et al., 2001) and that found a positive 8 9 association between warming, increasing tree ring widths and NDVI greening in boreal 10 forests of eastern Siberia (Berner et al., 2011, 2013).

11 Parts of the boreal forests in North America showed significant browning trends in the 12 GIMMS3g dataset but a tendency to positive trends in the GL2 dataset. The simulation results 13 from LPJmL-GSI are in agreement with the GIMMS3g-based browning trends, rather than 14 greening trends. However, these model-based browning trends were not as strong as in the GIMMS3g dataset. In LPJmL-GSI these browning trends are caused by the effects of seasonal 15 16 light and water effects on phenology, and by fire activity. In the GIMMS3g dataset these browning trends were related to several environmental factors like fire activity (Goetz et al., 17 18 2005), temperature-induced drought stress (Beck et al., 2011; Bunn and Goetz, 2006) and to 19 snow-regulated changes in soil water availability (Barichivich et al., 2014).

The Sahel had widespread greening trends in the GIMMS3g FAPAR dataset. Whereas LPJmL-OP simulated browning trends, the implementation of water availability effects on phenology enabled LPJmL-GSI to reproduce the observed greening trends. Increases in precipitation and rain-use efficiency were also identified in observational studies as the main drivers of positive trends in vegetation greenness in the Sahel (Fensholt et al., 2013).

Overall, we observed that both LPJmL-OP and LPJmL-GSI reproduced the greening trends in tundra, boreal and temperate forests, although LPJmL-GSI showed a wider agreement in the extent of browning trends in the boreal forests of North America. Further, in the Sahel region, the greening trends can only be reproduced through the inclusion of water availability controls on the phenology development. These results demonstrate that environmental controls like light, heat stress and water availability contribute to a better description of regional greening and browning trends in very different bioclimatic regions of the globe. Hence, a comprehensive characterization of the different environmental controls on
 phenological development is essential in performing model-based analysis of long term trends
 in vegetation activity.

4

5 **3.2.3 Effects on carbon fluxes and stocks**

6 LPJmL-GSI and LPJmL-OP-gc with optimized parameters represented better the global 7 patterns and mean seasonal cycles of gross primary production and biomass than LPJmL with 8 original phenology and prior parameters (LPJmL-OP-prior) (Figure 8). LPJmL-OP-prior 9 overestimated mean annual GPP and biomass in most polar, boreal and temperate regions. LPJmL-OP-prior underestimated mean annual GPP but overestimated mean annual biomass 10 in tropical regions around the Equator. These biases were reduced in LPJmL-OP-gc and 11 12 LPJmL-GSI. LPJmL generally overestimated GPP also in arid regions but these biases were 13 reduced after optimization in LPJmL-OP-gc and LPJmL-GSI (Figure S18). We also found 14 that the mean seasonal cycle of GPP from LPJmL-GSI agreed better with the mean seasonal 15 GPP cycle from the MTE estimate especially in temperate forests and in tropical, temperate 16 and polar grasslands (Supplement 5.1, Figure S17) although no information about the 17 seasonality of GPP was included in optimization experiments. LPJmL-GSI still overestimated biomass in some tropical regions (African Savannas, south-east Brazil, south and south-east 18 19 Asia) (Figure S19). These regions were mainly simulated as agricultural lands in LPJmL, i.e. 20 as different crop functional types (CFTs). The LPJmL-GSI phenology module was not 21 applied or optimized for agricultural regions, where the seasonal phenological development is 22 prescribed according to the CFTs parameterizations from Bondeau et al. (2007). Generally, LPJmL-GSI performed substantially better than LPJmL-OP-prior and LPJmL-OP-gc when 23 24 comparing the global total carbon fluxes and stocks to the data-oriented estimates 25 (Supplement 5.1, Table S6). These results demonstrate that in addition to the optimization of 26 productivity parameters in LPJmL, the implementation of the new GSI-based phenology 27 improved estimates of spatial patterns, seasonal dynamics, and global totals of gross primary 28 production and biomass.

3.2.4 Effects on forest distribution

2 LPJmL-GSI with dynamic vegetation better represented the observed tree cover in high 3 latitude regions than LPJmL-OP-prior and LPJmL-OP-gc (Figure 8 d). LPJmL-OP-prior 4 highly overestimated tree cover in boreal and arctic regions and simulated a too northern arctic tree line in comparison with tree cover from MODIS observations. Although this 5 6 overestimation was reduced after optimization, LPJmL-OP-gc still highly overestimated tree 7 cover in boreal and temperate regions. The occurrence of trees was shifted southwards in 8 LPJmL-GSI. Although LPJmL-GSI still overestimated tree cover in boreal regions, this 9 overestimation was much lower than in LPJmL-OP-gc. LPJmL-OP-prior and LPJmL-OP-gc 10 slightly underestimated tree cover in temperate regions around 45°N but this was well reproduced by LPJmL-GSI. We found no differences in tree cover between LPJmL-OP and 11 12 LPJmL-GSI in other parts of the world where tree cover is highly affected from agricultural 13 land use and thus implicitly prescribed to LPJmL. These results demonstrate that additional to the optimization of productivity parameters in LPJmL-OP-gc, the newly developed GSI-based 14 15 phenology model and the optimized model parameters contribute to a better representation of 16 tree cover in high-latitude regions.

17

3.2.5 Effects on evapotranspiration processes

19 Evapotranspiration from LPJmL agreed well with the data-oriented MTE estimates (Figure 8 20 b). The implementation and optimization of the new GSI-based phenology did not affect ET 21 much. ET increased only in tropical rainforests around the Equator in LPJmL-GSI and 22 LPJmL-OP-gc in comparison to LPJmL-OP-prior because of the increased GPP in these 23 regions. In other regions ET remained almost unchanged. But this does not imply that the 24 structural improvements in LPJmL-GSI did not affect the transpiration processes (Figures 25 S20, S21). Indeed, LPJmL-GSI had lower interception losses than LPJmL-OP in boreal 26 forests because of the reduced tree cover. On the other hand this implies that simulated soil 27 evaporation was increased. Furthermore, interception and soil evaporation had slightly shifted 28 seasonal cycles in LPJmL-GSI compared to LPJmL-OP due to the seasonal differences in 29 timing of leaf development and senescence stages (Figure S21). Consequently, small 30 differences in total evapotranspiration result from opposite and compensatory changes in 31 interception and soil evaporation and slight changes in transpiration fluxes in LPJmL-GSI.

2 3.3 Applicability and challenges of the LPJmL-GSI phenology module

3 The LPJmL-GSI phenology module is part of a DGVM that is applied for climate impact 4 studies. In order to assess how well the model performs under different climate conditions, we 5 additionally tested how the model performance changes in grid cells that were not used during 6 optimization (Figure S26). We found no general decrease in model performance with distance 7 to the nearest grid cell used for optimization, or under different temperature conditions. 8 Especially, no significantly lower correlations ($p \le 0.05$, Wilcoxon rank-sum test, Figure S26) 9 were found between simulated and observed FAPAR time series in grid cells that were 3° to 10 5°C warmer than the closest optimization grid cell. From a typical perspective of space for 11 time substitution, this could indicate that the confidence in the simulation of FAPAR 12 dynamics should not detract under climate warming scenarios of 0.3° to 4.8°C (IPCC, 2014).

13 Nevertheless, model optimization experiments and model evaluation indicated further needs 14 for improvement in future studies, for example, simulations of surface albedo could improve 15 through the implementation of time-varying effects of snow conditions and surface moisture 16 on albedo. Also, an enhanced representation of canopy architecture and canopy radiative 17 transfer could reduce the large spatial variability and parameter uncertainty found for the light 18 extinction coefficient and hence improve the simulation of tree coverage and peak FAPAR. In 19 addition to temperature, light and water availability, phenology also depends on other factors 20 that are not considered in LPJmL-GSI. Phenology is also driven by leaf age (Caldararu et al., 21 2012, 2014) and nutrient availability (Wright, 1996). These effects are neither considered in 22 the original GSI phenology model (Jolly et al., 2005; Stöckli et al., 2011) nor in the LPJmL-23 GSI or other traditional formulations. Here, the lower posterior values found for the parameter 24 α_a may be compensating for missing nitrogen limitation effects on productivity in boreal 25 forests (Vitousek and Howarth, 1991). Thus a future implementation of nitrogen limitation 26 processes in LPJmL requires a re-optimization of the α_a parameter. Additionally, the current 27 implementation of phenology in LPJmL affects photosynthesis only through changes in 28 APAR. In future model developments a stronger coupling between phenology and ecosystem 29 carbon cycle dynamics could be explored. For example, the LPJmL-GSI phenology module 30 could demand carbon for leaf development from photosynthesis or additional storage pools on 31 the one hand and could trigger carbon turnover through litterfall on the other hand. In this

case a phenology module could partly regulate an optimal carbon gain for a canopy similar to 1 2 the approach of Caldararu et al. (2014). Nevertheless, such an analysis needs to go beyond the 3 approach of Caldararu et al. (2014) and demands for additional observational constraints on 4 ecosystem carbon fluxes, leaf area, biomass and litterfall. In order to better understand 5 couplings between leaf phenology, changes in carbon allocation and photosynthesis it will be 6 of benefit to use site-level eddy covariance measurements from the FLUXNET network 7 (Baldocchi et al., 2001) together with ancillary data in ecosystem-scale model optimization 8 experiments (Carvalhais et al., 2010; Kuppel et al., 2012; Williams et al., 2009). Thus the 9 LPJmL-GSI phenology module and the LPJmL model-data integration approach can serve as 10 a framework to further explore hypotheses of ecosystem processes and vegetation dynamics.

11 We demonstrated the improved performance of LPJmL-GSI over LPJmL-OP in representing observed carbon fluxes and stocks, forest cover and seasonal to decadal dynamics of 12 13 vegetation greenness. Thus, similar approaches to the LPJmL-GSI phenology module can be 14 applied in other DGVMs to improve model simulations in comparison with observations. 15 However, the adaptation of current results to other models should be cautionary because the 16 phenology scheme of LPJmL-GSI is an empirical approach with PFT-dependent parameters 17 that need to be estimated. This estimation is model-specific because (1) different models do 18 not necessarily use the same definition and set of PFTs; (2) our parameterizations depend on 19 model structure, e.g. different models often use different hydrology routines; and (3) our 20 posterior parameters for phenology were also constrained by using albedo and GPP data. Thus 21 LPJmL-GSI model parameters cannot be easily transferred to other models. It might be 22 possible to use the parameters of the temperature and light limiting functions in other models 23 because these functions depend uniquely on the forcing data. On the other hand, the 24 parameters for the water availability limiting function might need to be re-optimized because 25 of differences in soil moisture computations. However, depending on the co-variability 26 between forcing variables and simulated water availability by other models, the best 27 parameterizations may differ from the ones presented here. Consequently, we acknowledge 28 the potential need to optimize parameters of the LPJmL-GSI phenology model in order to 29 obtain plausible results in other modelling structures. However, it is likely that the LPJmL-30 GSI phenology model can be easily applied to other models of the LPJ group of models 31 (Prentice et al., 2011; Smith et al., 2001) that are using the hydrology routines of Gerten et al.

(2004) while probably additional parameter optimization exercise are required to adapt the
 model to other types of DGVMs or ecosystem models.

3

4 **3.4** Environmental controls on vegetation greenness phenology

5 As the newly developed GSI-based phenology model of LPJmL can reproduce the seasonality and monthly dynamics of observed FAPAR in most biomes, it can be used to identify 6 7 environmental controls on vegetation greenness phenology. The importance of phenological 8 controls differed by climate regions, ecosystems and season (Figure 9). We identified 9 environmental controls on seasonal FAPAR dynamics by analyzing the mean seasonal cycles 10 of FAPAR, of the cold temperature, light, water availability and heat stress limiting functions 11 for phenology from the LPJmL-GSI model run. This analysis is comparable to previous 12 investigations of limiting factors for vegetation phenology (e.g. Jolly et al., 2005; Caldararu et 13 al., 2014). FAPAR seasonality in high-latitude regions (tundra, boreal forests) was mainly 14 controlled by cold temperature (entire year) and light (October to February). We also found an important control by water availability in February to April in the tundra and in boreal forests 15 16 of North America and eastern Siberia. This water limitation in early spring was due to the 17 seasonal freezing of the upper permafrost layer in LPJmL. FAPAR seasonality in temperate 18 grasslands in western North America and central Asia was controlled from a mixture of cold 19 temperature (January to April), of water availability (May to November) and light (November 20 to January). FAPAR seasonality in temperate forests in Europe was mainly limited by cold 21 temperature in spring and by a combination of cold temperature and light in autumn. 22 Additionally, heat stress and water availability contributed to a small reduction in summer 23 FAPAR in temperate and boreal forests. The FAPAR seasonality in savannas (Sahel) was 24 limited by water availability in the entire year and additionally by heat stress before the 25 beginning of the rain season. The FAPAR seasonality of temperate regions in South America 26 was limited by water availability in the entire year. Cold temperature was additionally 27 limiting between May and September. Thus, water availability was the only environmental 28 factor in LPJmL-GSI that controlled phenology globally from tropical to arctic biomes.

The implementation of the water limiting function on phenology in LPJmL-GSI resulted in unique patterns of phenological controls that were different from results reported in similar analyses (Caldararu et al., 2014; Jolly et al., 2005). LPJmL-GSI showed water limitation on

1 phenology in many sub-tropical and dry temperate regions (especially Mediterranean, Pampas 2 and Patagonia in South America, Mongolia, and northern Great Plains). The original GSI 3 model showed mainly temperature and light limitation in these regions. In contrast to the 4 original GSI, our implementation considers water limitations on phenology based on plant 5 available water and not on VPD (Jolly et al., 2005). As considered by Caldararu et al. (2014), 6 soil water availability exerts a more direct control on phenology development, which has been 7 demonstrated for Mediterranean ecosystems (Kramer et al., 2000; Richardson et al., 2013) 8 and in dry temperate grasslands (Liu et al., 2013; Yuan et al., 2007).

9 Additionally, we identify water availability as an important limiting function for spring 10 phenology in boreal and arctic regions in LPJmL-GSI because of the seasonal freezing of the 11 upper active layer in permafrost soils. Although no relationships between active layer depth and vegetation greenness were found so far (Mcmichael et al., 1997), frozen grounds limit the 12 13 seasonal tree growth in boreal forests because of limited water supply and nutrient uptake 14 (Benninghoff, 1952; Jarvis and Linder, 2000). As the seasonal freezing and thawing of 15 permafrost soils is to a large extent driven by changes in air temperature, one might argue that 16 air temperature is enough to explain phenology dynamics in boreal and arctic regions. 17 Nevertheless, we found weak correlations between posterior model parameters for the cold 18 temperature and water limiting function for phenology in PFTs that experience strong 19 permafrost dynamics (BoNS r = 0.2, PoH r = -0.28) (Figure S16). This indicates that the 20 water and cold temperature limiting functions in boreal and arctic regions are only weakly 21 correlated. Indeed, we did not find a completely synchronized temporal dynamic of the cold 22 temperature and water limiting functions for phenology (Figure 9). These results emphasize 23 the ability to disentangle effects of seasonal air temperature and soil moisture on phenology in 24 boreal and arctic regions. Air temperature and soil thawing are not completely synchronized 25 because soil temperature depends also on topography, substrate, and the insulating effects of 26 the snow, litter and vegetation cover (Jorgenson et al., 2010; Shur and Jorgenson, 2007; 27 Zhang, 2005). Soils might be still frozen if air temperature is already positive or vice versa. 28 Also experimental studies highlighted the role of permafrost-regulated soil moisture on 29 phenology and productivity in boreal and arctic ecosystem (Natali et al., 2012; Schuur et al., 30 2007). It also has been observed that the seasonal freezing and thawing in permafrost regions 31 regulates ecosystem evapotranspiration (Ohta et al., 2008) and fire activity (Forkel et al., 32 2012) especially during extreme dry years. Thus, although temperature might be enough to

explain average spatial patterns of phenology in boreal and arctic regions we acknowledge
 that variations in snow or vegetation cover that affects soil temperature and thus moisture
 might be important factors in explaining inter-annual variations of land surface phenology.

The heat stress limiting function was additionally introduced in LPJmL-GSI. Heat stress had 4 5 no importance for seasonal FAPAR dynamics in most regions except in temperate and 6 tropical grasslands. The heat stress function was highly correlated with the water availability 7 function in temperate grasslands. This suggests that summer FAPAR is both regulated by 8 water-induced and temperature-induced drought conditions in temperate grasslands. In 9 tropical grasslands, heat stress and water availability were driving the temporal dynamics of 10 seasonal FAPAR but asynchronously (in the Sahel). These results suggest that soil moisture 11 needs to be considered in observational data analyses and in other ecosystem models as 12 controlling factor for vegetation phenology in all biomes.

13 Interestingly, Caldararu *et al.* (2014) identify leaf age as the dominant factor for phenology 14 development in many permanent moist subtropical and tropical forests, but also in several water limited regions which were here identified as seasonally controlled by water 15 16 availability. We cannot identify a dominant control on seasonal FAPAR dynamics in evergreen forests, as leaf age is not explicitly simulated in LPJmL-GSI. We acknowledge that 17 18 the consideration of leaf age effects on phenology could further enhance the representation of ecosystem processes. However, the seasonal co-variation between LAI or FAPAR and 19 20 environmental controls on phenology complicates the ability to disentangle the leaf aging 21 signal from a temperature, light or water availability-driven signal, especially in seasonally 22 deciduous vegetation types, where climate-driven models explain a significant fraction of 23 seasonal variability and the realized age of leafs is shorter than a year. In addition, cloud 24 cover contamination over moist tropical or subtropical forests pertain usually a weak seasonal 25 signal and a high short-term variability, hinging on the reliability of the seasonal signal. 26 Especially, Morton et al. (2014) show that seasonal changes in MODIS LAI in the Amazon 27 forests are linked to insufficient corrections of the sun-sensor geometry, which challenge the 28 representation of vegetation phenology. However, in these tropical moist regions, where we 29 find no environmental seasonal controls, and the realized age of oldest leafs are higher than a 30 year, leaf age may be an important contributor for further consideration regarding the above-31 seasonal frequency of phenology. Hence, grasping the relevance of leaf longevity, especially 32 in tropical perennial systems, would necessarily require ground observations of leaf development and litter fall to constrain leaf age parameters, as well as measurements of soil
 water content to address the appropriateness of soil moisture effects.

3

4 4 Conclusions

5 We have demonstrated a major improvement of the LPJmL dynamic global vegetation model 6 by implementing a new set of phenological controls on vegetation greenness and by 7 integrating multiple decadal satellite observations. We have proven that the original 8 phenology model in LPJmL is unable to explain temporal dynamics of FAPAR. As an 9 alternative we implemented a new phenology model (LPJmL-GSI) which considers effects of 10 cold temperature, heat stress, light, and water availability on vegetation phenology. We 11 developed a model-data integration approach for LPJmL (LPJmL-MDI) to 1) constrain model 12 parameters against observations, 2) to directly integrate observed land cover fractions and 13 burnt area time series and 3) to evaluate LPJmL against independent data streams. Specifically, phenology, productivity, and albedo-related model parameters of LPJmL-GSI 14 15 were optimized jointly against 30 year time series of satellite observations of FAPAR, against 10 year time series of vegetation albedo and mean annual patterns of gross primary 16 17 production using a genetic optimization algorithm.

18 The new phenology model and the parameter optimization clearly improved LPJmL model 19 simulations. LPJmL-GSI better reproduces observed spatial patterns of gross primary 20 production, tree cover, biomass and FAPAR than the original model. LPJmL-GSI simulates 21 global total carbon stocks and fluxes that are closer to independent estimates than from the 22 original model. LPJmL-GSI better represents observed seasonal, monthly, inter-annual and 23 decadal FAPAR dynamics than the original model. The improvements of LPJmL in 24 representing observed patterns and temporal dynamics of vegetation greenness allows 25 assessing environmental controls on vegetation phenology and greenness. Contrasting to 26 previous studies (Jolly et al., 2005; Stöckli et al., 2011), our results indicate that soil water 27 availability is a major control of seasonal FAPAR dynamics not only in water-limited biomes 28 but also in boreal forests and the arctic tundra where water availability is regulated through 29 seasonal thawing and freezing of the active permafrost layer. Until now phenology of these 30 ecosystems was mostly considered as temperature-limited. The consideration of the effect of 31 soil water availability on phenology in LPJmL improved model simulations of greening

trends in the Sahel and of browning trends in parts of the boreal forests of North America.
Our results demonstrate that improved phenology models that consider seasonal effects of
water availability on a continuous canopy development are needed in order to correctly
explain seasonal to long-term dynamics in vegetation greenness.

5

6 Acknowledgements

7 We thank Maarten Braakhekke and Enrico Tomelleri for testing model optimization 8 algorithms. We thank Ulrich Weber for his assistance in data processing. We thank Ranga 9 Myneni and Frédéric Baret for their comments on FAPAR datasets. We thank the IT 10 departments at MPI-BGC Jena and PIK Potsdam for providing and maintaining the respective 11 high performance computing infrastructures.

We gratefully thank the following researchers, groups and institutes for producing, providingand hosting their datasets:

- Compton Tucker, Jorge Pinzon, Ranga Myneni and the GIMMS group for the
 GIMMS FPAR3g dataset
- 16 INRA, CNES and VITO for the Geoland2 FAPAR dataset
- 17 Martin Jung for the MTE GPP and ET data, and the SYNMAP land cover map
- 18 Sassan Saatchi for the tropical forest biomass map
- Markus Kottek, Franz Rubel and the University of Veterinary Medicine Vienna for
 providing the Koeppen-Geiger climate map
- 21 Louis Giglio and Guido Van der Werf for providing the GFED database
- Jennifer L. Northway and Gary Schmunk for compiling the Alaska Large Fire
 Database
- Natural Resources Canada for providing the Canada National Fire Database
- Phil Jones, Ian Harris and the CRU research unit for providing the CRU climate
 dataset
- 27 ECMWF for ERA-Interim climate reananlysis data.

- Matthias Forkel received funding from the Max Planck Institute for Biogeochemistry and
 from the European Commission's 7th Framework Programme project CARBONES (grant
 agreement 242316). Matthias Forkel conducted this work under the International Max Planck
- 4 Research School for Global Biogeochemical Cycles.

1 References

Anav, A., Murray-Tortarolo, G., Friedlingstein, P., Sitch, S., Piao, S. and Zhu, Z.: Evaluation
of Land Surface Models in Reproducing Satellite Derived Leaf Area Index over the High-

4 Latitude Northern Hemisphere. Part II: Earth System Models, Remote Sens., 5(8), 3637–

5 3661, doi:10.3390/rs5083637, 2013.

- 6 Archibald, S. and Scholes, R. J.: Leaf green-up in a semi-arid African savanna separating 7 tree and grass responses to environmental cues, J. Veg. Sci., 18(4), 583–594, 2007.
- Atzberger, C., Klisch, A., Mattiuzzi, M. and Vuolo, F.: Phenological Metrics Derived over the
 European Continent from NDVI3g Data and MODIS Time Series, Remote Sens., 6(1), 257–
 284, doi:10.3390/rs6010257, 2013.
- Baird, R. A. and Verbyla, D.: Browning of the landscape of interior Alaska based on 1986-2009 Landsat sensor NDVI, Can. J. Of, 1382, 1371–1382, doi:10.1139/x2012-088, 2012.
- 13 Baldocchi, D., Falge, E., Lianhong Gu, Olson, R., Hollinger, D., Running, S., Anthoni, P.,
- 14 Bernhofer, C., Davis, K., Evans, R., Fuentes, J., Goldstein, A., Katul, G., Law, B., Xuhui Lee,
- 15 Malhi, Y., Meyers, T., Munger, W., Oechel, W. and Paw U, K. T.: FLUXNET: A New Tool
- 16 to Study the Temporal and Spatial Variability of Ecosystem-Scale Carbon Dioxide, Water
- 17 Vapor, and Energy Flux Densities, Bull. Am. Meteorol. Soc., 82(11), 2415, 2001.
- 18 Baret, F., Weiss, M., Lacaze, R., Camacho, F., Makhmara, H., Pacholcyzk, P. and Smets, B.:
- 19 GEOV1: LAI and FAPAR essential climate variables and FCOVER global time series
- 20 capitalizing over existing products. Part1: Principles of development and production, Remote
- 21 Sens. Environ., 137, 299–309, doi:10.1016/j.rse.2012.12.027, 2013.
- 22 Barichivich, J., Briffa, K. R., Myneni, R. B., Osborn, T. J., Melvin, T. M., Ciais, P., Piao, S.
- and Tucker, C.: Large-scale variations in the vegetation growing season and annual cycle of
 atmospheric CO2 at high northern latitudes from 1950 to 2011, Glob. Change Biol., 19(10),
- 25 3167–3183, doi:10.1111/gcb.12283, 2013.
- 26 Barichivich, J., Briffa, K. R., Myneni, R., Schrier, G. van der, Dorigo, W., Tucker, C. J.,
- Osborn, T. J. and Melvin, T. M.: Temperature and Snow-Mediated Moisture Controls of Summer Photosynthetic Activity in Northern Terrestrial Ecosystems between 1982 and 2011,
- 29 Remote Sens., 6(2), 1390–1431, doi:10.3390/rs6021390, 2014.
- 30 Beck, P. S. A. and Goetz, S. J.: Satellite observations of high northern latitude vegetation
- 31 productivity changes between 1982 and 2008: ecological variability and regional differences,
- 32 Environ. Res. Lett., 6(4), 045501, doi:10.1088/1748-9326/6/4/045501, 2011.
- Beck, P. S. A., Juday, G. P., Alix, C., Barber, V. A., Winslow, S. E., Sousa, E. E., Heiser, P.,
 Herriges, J. D. and Goetz, S. J.: Changes in forest productivity across Alaska consistent with
 biome shift, Ecol. Lett., 14(4), 373–9, doi:10.1111/j.1461-0248.2011.01598.x, 2011.
- Beer, C., Reichstein, M., Tomelleri, E., Ciais, P., Jung, M., Carvalhais, N., Rödenbeck, C.,
 Arain, M. A., Baldocchi, D., Bonan, G. B., Bondeau, A., Cescatti, A., Lasslop, G., Lindroth,
- 38 A., Lomas, M., Luyssaert, S., Margolis, H., Oleson, K. W., Roupsard, O., Veenendaal, E.,

- 1 Viovy, N., Williams, C., Woodward, F. I. and Papale, D.: Terrestrial Gross Carbon Dioxide
- 2 Uptake: Global Distribution and Covariation with Climate, Science, 329(5993), 834–838,
- 3 doi:10.1126/science.1184984, 2010.
- Benninghoff, W. S.: Interaction of Vegetation and Soil Frost Phenomena, Arctic, 5(1), 34–44,
 1952.
- Berner, L. T., Beck, P. S. A., Bunn, A. G. and Goetz, S. J.: Plant response to climate change
 along the forest-tundra ecotone in northeastern Siberia, Glob. Change Biol., 19(11), 3449–
 3462, doi:10.1111/gcb.12304, 2013.
- 9 Berner, L. T., Beck, P. S. A., Bunn, A. G., Lloyd, A. H. and Goetz, S. J.: High-latitude tree 10 growth and satellite vegetation indices: Correlations and trends in Russia and Canada (1982-
- 11 2008), J. Geophys. Res.-Biogeosciences, 116, doi:10.1029/2010jg001475, 2011.
- Bi, J., Xu, L., Samanta, A., Zhu, Z. and Myneni, R.: Divergent Arctic-Boreal Vegetation
 Changes between North America and Eurasia over the Past 30 Years, Remote Sens., 5(5),
 2093–2112, doi:10.3390/rs5052093, 2013.
- 15 Blok, D., Sass-Klaassen, U., Schaepman-Strub, G., Heijmans, M. M. P. D., Sauren, P. and
- Berendse, F.: What are the main climate drivers for shrub growth in Northeastern Siberian tundra?, Biogeosciences, 8(5), 1169–1179, doi:10.5194/bg-8-1169-2011, 2011.
- 17 tunuta:, Diogeosciences, o(5), 1109-1179, u01.10.5194/0g-6-1109-2011, 2011.
- Bolstad, P. V. and Gower, S. T.: Estimation of leaf area index in fourteen southern Wisconsin
 forest stands using a portable radiometer, Tree Physiol., 7(1_2_3_4), 115–124, 1990.
- Bondeau, A., Smith, P. C., Zaehle, S., Schaphoff, S., Lucht, W., Cramer, W., Gerten, D.,
 Lotze-Campen, H., Müller, C., Reichstein, M. and Smith, B.: Modelling the role of
 agriculture for the 20th century global terrestrial carbon balance, Glob. Change Biol., (13),
 679–706, 2007.
- Bunn, A. G. and Goetz, S. J.: Trends in Satellite-Observed Circumpolar Photosynthetic
 Activity from 1982 to 2003: The Influence of Seasonality, Cover Type, and Vegetation
 Density, Earth Interact., 10(12), 1–19, doi:10.1175/ei190.1, 2006.
- Bunn, A. G., Goetz, S. J., Kimball, J. S. and Zhang, K.: Northern High-Latitude Ecosystems
 Respond to Climate Change, Eos Trans. Am. Geophys. Union, 88(34), 333–333,
 doi:10.1029/2007eo340001, 2007.
- Caldararu, S., Palmer, P. I. and Purves, D. W.: Inferring Amazon leaf demography from satellite observations of leaf area index, Biogeosciences, 9(4), 1389–1404, doi:10.5194/bg-9-1389-2012, 2012.
- Caldararu, S., Purves, D. W. and Palmer, P. I.: Phenology as a strategy for carbon optimality:
 a global model, Biogeosciences, 11(3), 763–778, doi:10.5194/bg-11-763-2014, 2014.
- Carvalhais, N., Forkel, M., Khomik, M., Bellarby, J., Jung, M., Migliavacca, M., Mu, M.,
 Saatchi, S., Santoro, M., Thurner, M., Weber, U., Ahrens, B., Beer, C., Cescatti, A.,
 Randerson, J. T. and Reichstein, M.: Global covariation of carbon turnover times with climate
- in terrestrial ecosystems, Nature, advance online publication, doi:10.1038/nature13731, 2014.

- 1 Carvalhais, N., Reichstein, M., Ciais, P., Collatz, G. J., Mahecha, M. D., Montagnani, L.,
- 2 Papale, D., Rambal, S. and Seixas, J.: Identification of vegetation and soil carbon pools out of
- 3 equilibrium in a process model via eddy covariance and biometric constraints, Glob. Change
- 4 Biol., 16(10), 2813–2829, doi:10.1111/j.1365-2486.2010.02173.x, 2010.
- Cescatti, A., Marcolla, B., Santhana Vannan, S. K., Pan, J. Y., Román, M. O., Yang, X.,
 Ciais, P., Cook, R. B., Law, B. E., Matteucci, G., Migliavacca, M., Moors, E., Richardson, A.
 D., Seufert, G. and Schaaf, C. B.: Intercomparison of MODIS albedo retrievals and in situ
 measurements across the global FLUXNET network, Remote Sens. Environ., 121, 323–334,
 doi:10.1016/j.rse.2012.02.019, 2012.
- 10 CFS: Canadian Large Fire Database. Natural Resources Canada, Canadian Forest Service, 11 edited by C. F. Service, [online] Available from: http://cwfis.cfs.nrcan.gc.ca/en CA/nfdbID -
- 12 427, 2010.
- 13 Chapin, F. S., Sturm, M., Serreze, M. C., McFadden, J. P., Key, J. R., Lloyd, A. H., McGuire,
- 14 A. D., Rupp, T. S., Lynch, A. H., Schimel, J. P., Beringer, J., Chapman, W. L., Epstein, H. E.,
- 15 Euskirchen, E. S., Hinzman, L. D., Jia, G., Ping, C.-L., Tape, K. D., Thompson, C. D. C.,
- 16 Walker, D. A. and Welker, J. M.: Role of Land-Surface Changes in Arctic Summer Warming,
- 17 Science, 310(5748), 657–660, doi:10.1126/science.1117368, 2005.
- Cook, B. I. and Pau, S.: A Global Assessment of Long-Term Greening and Browning Trends
 in Pasture Lands Using the GIMMS LAI3g Dataset, Remote Sens., 5(5), 2492–2512,
 doi:10.3390/rs5052492, 2013.
- 21 Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., 22 23 Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., 24 Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., K\rallberg, P., Köhler, M., Matricardi, 25 M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J. J., Park, B. K., Peubey, C., Rosnay, P. de, Tavolato, C., Thépaut, J. N. and Vitart, F.: The ERA-Interim reanalysis: configuration and 26 27 performance of the data assimilation system, Q. J. R. Meteorol. Soc., 137(656), 553-597, 28 doi:10.1002/qj.828, 2011.
- 29 Fensholt, R. and Proud, S. R.: Evaluation of Earth Observation based global long term
- 30 vegetation trends Comparing GIMMS and MODIS global NDVI time series, Remote Sens.
- 31 Environ., 119, 131–147, doi:10.1016/j.rse.2011.12.015, 2012.
- Fensholt, R., Rasmussen, K., Kaspersen, P., Huber, S., Horion, S. and Swinnen, E.: Assessing
 Land Degradation/Recovery in the African Sahel from Long-Term Earth Observation Based
 Primary Productivity and Precipitation Relationships, Remote Sens., 5(2), 664–686,
 doi:10.3390/rs5020664, 2013.
- Fensholt, R., Sandholt, I. and Rasmussen, M. S.: Evaluation of MODIS LAI, fAPAR and the
 relation between fAPAR and NDVI in a semi-arid environment using in situ measurements,
 Remote Sens. Environ., 91(3-4), 490–507, doi:10.1016/j.rse.2004.04.009, 2004.
- Forbes, B. C., Macias Fauria, M. and Zetterberg, P.: Russian Arctic warming and `greening' are closely tracked by tundra shrub willows, Glob. Change Biol., 16(5), 1542–1554, 2010.

- Forkel, M., Carvalhais, N., Verbesselt, J., Mahecha, M., Neigh, C. and Reichstein, M.: Trend
 Change Detection in NDVI Time Series: Effects of Inter-Annual Variability and
- 3 Methodology, Remote Sens., 5(5), 2113–2144, doi:10.3390/rs5052113, 2013.
- Forkel, M., Thonicke, K., Beer, C., Cramer, W., Bartalev, S. and Schmullius, C.: Extreme fire
 events are related to previous-year surface moisture conditions in permafrost-underlain larch
 forests of Siberia, Environ. Res. Lett., 7(4), 044021, doi:10.1088/1748-9326/7/4/044021,
 2012.
- 8 Frames: Alaska Large Fire Database, [online] Available from: 9 http://www.frames.gov/rcs/10000/10465.html, 2012.
- 10 Friedlingstein, P., Cox, P., Betts, R., Bopp, L., von Bloh, W., Brovkin, V., Cadule, P., Doney,
- 11 S., Eby, M., Fung, I., Bala, G., John, J., Jones, C., Joos, F., Kato, T., Kawamiya, M., Knorr,
- 12 W., Lindsay, K., Matthews, H. D., Raddatz, T., Rayner, P., Reick, C., Roeckner, E.,
- Schnitzler, K.-G., Schnur, R., Strassmann, K., Weaver, A. J., Yoshikawa, C. and Zeng, N.:
 Climate–Carbon Cycle Feedback Analysis: Results from the C⁴ MIP Model Intercomparison,
- 14 Chinale–Carbon Cycle Feedback Analysis. Results from the C MIP Model intercomparis
- 15 J. Clim., 19(14), 3337–3353, doi:10.1175/JCLI3800.1, 2006.
- 16 Gamon, J. A., Field, C. B., Goulden, M. L., Griffin, K. L., Hartley, E., Joel, G., Peñuelas, J.
- 17 and Valentini, R.: Relationships Between NDVI, Canopy Structure, and Photosynthesis in
- 18 Three Californian Vegetation Types, Ecol. Appl., 5(1), 28–41, 1995.
- 19 Gerten, D., Schaphoff, S., Haberlandt, U., Lucht, W. and Sitch, S.: Terrestrial vegetation and
- water balance hydrological evaluation of a dynamic global vegetation model, J. Hydrol.,
 (286), 249–270, 2004.
 - 22 Giglio, L., Randerson, J. T., van der Werf, G. R., Kasibhatla, P. S., Collatz, G. J., Morton, D.
 - 23 C. and DeFries, R. S.: Assessing variability and long-term trends in burned area by merging
 - 24 multiple satellite fire products, Biogeosciences, 7(3), 1171–1186, 2010.
 - Goetz, S. J., Bunn, A. G., Fiske, G. J. and Houghton, R. A.: Satellite-observed photosynthetic
 trends across boreal North America associated with climate and fire disturbance, Proc. Natl.
 Acad. Sci. U. S. A., 102(38), 13521–13525, doi:10.1073/pnas.0506179102, 2005.
 - Harris, I., Jones, P. D., Osborn, T. J. and Lister, D. H.: Updated high-resolution grids of
 monthly climatic observations the CRU TS3.10 Dataset., Int. J. Climatol., n/a–n/a,
 doi:10.1002/joc.3711, 2013.
 - Høgda, K. A., Karlsen, S. R. and Solheim, I.: Climatic change impact on growing season in
 Fennoscandia studied by a time series of NOAA AVHRR NDVI data, in Geoscience and
 Remote Sensing Symposium, 2001. IGARSS '01. IEEE 2001 International, vol. 3, pp. 1338–
 1340 vol.3., 2001.
 - Høgda, K. A., Tømmervik, H. and Karlsen, S. R.: Trends in the Start of the Growing Season
 in Fennoscandia 1982–2011, Remote Sens., 5(9), 4304–4318, doi:10.3390/rs5094304, 2013.
 - 37 IPCC, I. P. on C. C.: Climate Change 2013 The Physical Science Basis Working Group I
 - 38 Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate
 - 39 Change, Cambridge University Press, Cambridge., 2014.

Jarvis, P. and Linder, S.: Botany: Constraints to growth of boreal forests, Nature, 405(6789),
 904–905, doi:10.1038/35016154, 2000.

Jiang, Y. and Huang, B.: Physiological Responses to Heat Stress Alone or in Combination with Drought: A Comparison between Tall Fescue and Perennial Ryegrass, HortScience, 5 36(4), 682–686, 2001.

- Jolly, W. M., Nemani, R. and Running, S. W.: A generalized, bioclimatic index to predict
 foliar phenology in response to climate, Glob. Change Biol., 11(4), 619–632,
 doi:10.1111/j.1365-2486.2005.00930.x, 2005.
- 9 De Jong, R., Schaepman, M. E., Furrer, R., de Bruin, S. and Verburg, P. H.: Spatial 10 relationship between climatologies and changes in global vegetation activity, Glob. Change 11 Biol., 19(6), 1953–1964, doi:10.1111/gcb.12193, 2013a.
- De Jong, R., Verbesselt, J., Schaepman, M. E. and Bruin, S.: Trend changes in global
 greening and browning: contribution of short-term trends to longer-term change, Glob.
 Change Biol., 18(2), 642–655, doi:10.1111/j.1365-2486.2011.02578.x, 2011.
- 15 De Jong, R., Verbesselt, J., Zeileis, A. and Schaepman, M.: Shifts in Global Vegetation 16 Activity Trends, Remote Sens., 5(3), 1117–1133, doi:10.3390/rs5031117, 2013b.
- 17 Jorgenson, M. T., Romanovsky, V. E., Harden, J. W., Shur, Y. L., O'Donnell, J., Schuur, E.
- 18 A. G., Kanevskiy, M. and Marchenko, S.: Resilience and vulnerability of permafrost to 19 climate change, Can. J. For. Res. - Rev. Can. Rech. For., 40(7), 1219–1236, 2010.
- Jung, M., Henkel, K., Herold, M. and Churkina, G.: Exploiting synergies of global land cover
 products for carbon cycle modeling, Remote Sens. Environ., 101, 534–553,
 doi:10.1016/j.rse.2006.01.020, 2006.
- 23 Jung, M., Reichstein, M., Margolis, H. A., Cescatti, A., Richardson, A. D., Arain, M. A., Arneth, A., Bernhofer, C., Bonal, D., Chen, J., Gianelle, D., Gobron, N., Kiely, G., Kutsch, 24 25 W., Lasslop, G., Law, B. E., Lindroth, A., Merbold, L., Montagnani, L., Moors, E. J., Papale, D., Sottocornola, M., Vaccari, F. and Williams, C.: Global patterns of land-atmosphere fluxes 26 27 of carbon dioxide, latent heat, and sensible heat derived from eddy covariance, satellite, and 28 meteorological observations. J. Geophys. Res.. 116. G00J07-G00J07, 29 doi:10.1029/2010jg001566, 2011.
- Jung, M., Verstraete, M., Gobron, N., Reichstein, M., Papale, D., Bondeau, A., RobusTelli, M.
 and Pinty, B.: Diagnostic assessment of European gross primary production, Glob. Change
 Biol., 14(10), 2349–2364, doi:10.1111/j.1365-2486.2008.01647.x, 2008.
- Kaminski, T., Knorr, W., Scholze, M., Gobron, N., Pinty, B., Giering, R. and Mathieu, P. P.:
 Consistent assimilation of MERIS FAPAR and atmospheric CO2 into a terrestrial vegetation
 model and interactive mission benefit analysis, Biogeosciences, 9(8), 3173–3184,
 doi:10.5194/bg-9-3173-2012, 2012.
- Kasischke, E. S., Williams, D. and Barry, D.: Analysis of the patterns of large fires in the
 boreal forest region of Alaska, Int. J. Wildland Fire, 11(2), 131–144, 2002.

- Keeling, C. D., Chin, J. F. S. and Whorf, T. P.: Increased activity of northern vegetation
 inferred from atmospheric CO2 measurements, Nature, 382(6587), 146–149,
 doi:10.1038/382146a0, 1996.
- Kelley, D. I., Prentice, I. C., Harrison, S. P., Wang, H., Simard, M., Fisher, J. B. and Willis,
 K. O.: A comprehensive benchmarking system for evaluating global vegetation models,
 Biogeosciences, 10(5), 3313–3340, doi:10.5194/bg-10-3313-2013, 2013.
- 7 Kendall, M. G.: Rank Correlation Methods, Griffin, London., 1975.
- 8 Kira, T., Shinokazi, K. and Hozumi, K.: Structure of forest canopies as related to their 9 primary productivity, Plant Cell Physiol., 10(1), 129–142, 1969.
- Knorr, W., Kaminski, T., Scholze, M., Gobron, N., Pinty, B., Giering, R. and Mathieu, P.-P.:
 Carbon cycle data assimilation with a generic phenology model, J. Geophys. Res.
 Biogeosciences, 115(G4), G04017, doi:10.1029/2009JG001119, 2010.
- Kottek, M., Grieser, J., Beck, C., Rudolf, B. and Rubel, F.: World Map of the Köppen-Geiger
 climate classification updated, Meteorol. Z., 15(3), 259–263, doi:10.1127/09412948/2006/0130, 2006.
- Kramer, K., Leinonen, I. and Loustau, D.: The importance of phenology for the evaluation of
 impact of climate change on growth of boreal, temperate and Mediterranean forests
 ecosystems: an overview, Int. J. Biometeorol., 44(2), 67–75, 2000.
- Kuppel, S., Peylin, P., Chevallier, F., Bacour, C., Maignan, F. and Richardson, A. D.:
 Constraining a global ecosystem model with multi-site eddy-covariance data, Biogeosciences,
 9(10), 3757–3776, doi:10.5194/bg-9-3757-2012, 2012.
- Van Leeuwen, W. J. D., Hartfield, K., Miranda, M. and Meza, F. J.: Trends and ENSO/AAO
 Driven Variability in NDVI Derived Productivity and Phenology alongside the Andes
 Mountains, Remote Sens., 5(3), 1177–1203, doi:10.3390/rs5031177, 2013.
- Lehsten, V., Tansey, K. J., Balzter, H., Thonicke, K., Spessa, A., Weber, U., Smith, B. and
 Arneth, A.: Estimating carbon emissions from African wildfires, Biogeosciences Discuss,
 5(4), 3091–3122, doi:10.5194/bgd-5-3091-2008, 2008.
- Liu, H., Tian, F., Hu, H. C., Hu, H. P. and Sivapalan, M.: Soil moisture controls on patterns of
 grass green-up in Inner Mongolia: an index based approach, Hydrol. Earth Syst. Sci.,
 17(2004), 805–815, doi:10.5194/hess-17-805-2013, 2013.
- Loranty, M. M., Goetz, S. J. and Beck, P. S. A.: Tundra vegetation effects on pan-Arctic albedo, Environ. Res. Lett., 6(2), 024014, doi:10.1088/1748-9326/6/2/024014, 2011.
- 33 Lucht, W., Prentice, I. C., Myneni, R. B., Sitch, S., Friedlingstein, P., Cramer, W., Bousquet,
- 34 P., Buermann, W. and Smith, B.: Climatic Control of the High-Latitude Vegetation Greening
- 35 Trend and Pinatubo Effect, Science, 296(5573), 1687–1689, doi:10.1126/science.1071828,
 36 2002.

- 1 Lucht, W., Schaaf, C. B. and Strahler, A. H.: An algorithm for the retrieval of albedo from
- 2 space using semiempirical BRDF models, Geosci. Remote Sens. IEEE Trans. On, 38(2), 977–
- 3 998, doi:10.1109/36.841980, 2000.
- 4 Mann, H. B.: Nonparametric Tests Against Trend, Econometrica, 13(3), 245–259, 1945.
- 5 Mcmichael, C. E., Hope, A. S., Stow, D. A. and Fleming, J. B.: The relation between active
- 6 layer depth and a spectral vegetation index in arctic tundra landscapes of the North Slope of
- 7 Alaska, Int. J. Remote Sens., 18(11), 2371–2382, doi:10.1080/014311697217666, 1997.
- Mebane, W. R. and Sekhon, J. S.: Genetic Optimization Using Derivatives : The rgenoud
 Package for R, J. Stat. Softw., 42(11), 2011.
- 10 Migliavacca, M., Galvagno, M., Cremonese, E., Rossini, M., Meroni, M., Sonnentag, O.,
- 11 Cogliati, S., Manca, G., Diotri, F., Busetto, L., Cescatti, A., Colombo, R., Fava, F., Morra di
- 12 Cella, U., Pari, E., Siniscalco, C. and Richardson, A. D.: Using digital repeat photography and
- 13 eddy covariance data to model grassland phenology and photosynthetic CO2 uptake, Agric.
- 14 For. Meteorol., 151(10), 1325–1337, doi:10.1016/j.agrformet.2011.05.012, 2011.
- 15 Migliavacca, M., Meroni, M., Busetto, L., Colombo, R., Zenone, T., Matteucci, G., Manca, G.
- and Seufert, G.: Modeling Gross Primary Production of Agro-Forestry Ecosystems by Assimilation of Satellite-Derived Information in a Process-Based Model, Sensors, 9(2), 922–
- 18 942, doi:10.3390/s90200922, 2009.
- Monsi, M. and Saeki, T.: Über den Lichtfaktor in den Pflanzengesellschaften und seine
 Bedeutung für die Stoffproduktion, Jpn. J. Bot., 14, 22–52, 1953.
- 21 Morton, D. C., Nagol, J., Carabajal, C. C., Rosette, J., Palace, M., Cook, B. D., Vermote, E.
- 22 F., Harding, D. J. and North, P. R. J.: Amazon forests maintain consistent canopy structure
- and greenness during the dry season, Nature, 506, 221–224, doi:10.1038/nature13006, 2014.
- 24 Murray-Tortarolo, G., Anav, A., Friedlingstein, P., Sitch, S., Piao, S., Zhu, Z., Poulter, B., 25 Zaehle, S., Ahlström, A., Lomas, M., Levis, S., Viovy, N. and Zeng, N.: Evaluation of Land 26 Surface Models in Reproducing Satellite-Derived LAI over the High-Latitude Northern 27 Hemisphere. Part I: Uncoupled DGVMs. Remote Sens., 5(10). 4819-4838. 28 doi:10.3390/rs5104819, 2013.
- Myers-Smith, I. H., Forbes, B. C., Wilmking, M., Hallinger, M., Lantz, T., Blok, D., Tape, K.
 D., Macias-Fauria, M., Sass-Klaassen, U., Lévesque, E., Boudreau, S., Ropars, P.,
 Hermanutz, L., Trant, A., Collier, L. S., Weijers, S., Rozema, J., Rayback, S. a, Schmidt, N.
 M., Schaepman-Strub, G., Wipf, S., Rixen, C., Ménard, C. B., Venn, S., Goetz, S., AndreuHayles, L., Elmendorf, S., Ravolainen, V., Welker, J., Grogan, P., Epstein, H. E. and Hik, D.
 S.: Shrub expansion in tundra ecosystems: dynamics, impacts and research priorities, Environ.
- 35 Res. Lett., 6(4), 045509–045509, doi:10.1088/1748-9326/6/4/045509, 2011.
- Myneni, R. B., Hall, F. G., Sellers, P. J. and Marshak, A. L.: The interpretation of spectral vegetation indexes, Geosci. Remote Sens. IEEE Trans. On, 33(2), 481–486, doi:10.1109/36.377948, 1995.

1 Myneni, R. B., Keeling, C. D., Tucker, C. J., Asrar, G. and Nemani, R. R.: Increased plant 2 growth in the northern high latitudes from 1981 to 1991, Nature, 386(6626), 698–702, 3 doi:10.1038/386698a0, 1997a.

Myneni, R. B., Ramakrishna, R., Nemani, R. and Running, S. W.: Estimation of global leaf
area index and absorbed par using radiative transfer models, IEEE Trans. Geosci. Remote
Sens., 35(6), 1380–1393, doi:10.1109/36.649788, 1997b.

- Myneni, R. B. and Williams, D. L.: On the relationship between FAPAR and NDVI, Remote
 Sens. Environ., 49(3), 200–211, doi:10.1016/0034-4257(94)90016-7, 1994.
- 9 Natali, S. M., Schuur, E. A. G. and Rubin, R. L.: Increased plant productivity in Alaskan
 10 tundra as a result of experimental warming of soil and permafrost, J. Ecol., 100(2), 488–498,
 11 doi:10.1111/j.1365-2745.2011.01925.x, 2012.
- 12 Ohta, T., Maximov, T. C., Dolman, A. J., Nakai, T., van der Mo
- 12 Ohta, T., Maximov, T. C., Dolman, A. J., Nakai, T., van der Molen, M. K., Kononov, A. V., 13 Maximov, A. P., Hiyama, T., Iijima, Y., Moors, E. J., Tanaka, H., Toba, T. and Yabuki, H.:
- 14 Interannual variation of water balance and summer evapotranspiration in an eastern Siberian
- 15 larch forest over a 7-year period (1998-2006), Agric. For. Meteorol., 148(12), 1941–1953,
- 16 2008.
- Van Peer, L., Nijs, I., Reheul, D. and De Cauwer, B.: Species richness and susceptibility to
 heat and drought extremes in synthesized grassland ecosystems: compositional vs
 physiological effects, Funct. Ecol., 18(6), 769–778, doi:10.1111/j.0269-8463.2004.00901.x,
 2004.
- Piao, S., Wang, X., Ciais, P., Zhu, B., Wang, T. and Liu, J.: Changes in satellite-derived
 vegetation growth trend in temperate and boreal Eurasia from 1982 to 2006, Glob. Change
 Biol., 17(10), 3228–3239, doi:10.1111/j.1365-2486.2011.02419.x, 2011.
- Pinzon, J. E. and Tucker, C. J.: A Non-Stationary 1981–2012 AVHRR NDVI3g Time Series,
 Remote Sens., 6(8), 6929–6960, doi:10.3390/rs6086929, 2014.
- Poirier, M., Durand, J.-L. and Volaire, F.: Persistence and production of perennial grasses
 under water deficits and extreme temperatures: importance of intraspecific vs. interspecific
 variability, Glob. Change Biol., 18(12), 3632–3646, doi:10.1111/j.1365-2486.2012.02800.x,
 2012.
- Potter, C. S., Klooster, S. and Brooks, V.: Interannual Variability in Terrestrial Net Primary
 Production: Exploration of Trends and Controls on Regional to Global Scales, Ecosystems,
 2(1), 36–48, doi:10.1007/s100219900056, 1999.
- Poulter, B., Ciais, P., Hodson, E., Lischke, H., Maignan, F., Plummer, S. and Zimmermann,
 N. E.: Plant functional type mapping for earth system models, Geosci Model Dev, 4(4), 993–
 1010, doi:10.5194/gmd-4-993-2011, 2011a.
- Poulter, B., Frank, D. C., Hodson, E. L. and Zimmermann, N. E.: Impacts of land cover and
 climate data selection on understanding terrestrial carbon dynamics and the CO2 airborne
 fraction, Biogeosciences, 8(8), 2027–2036, doi:10.5194/bg-8-2027-2011, 2011b.

1 Prentice, I. C., Bondeau, A., Cramer, W., Harrison, S. P., Hickler, T., Lucht, W., Sitch, S., 2 Smith, B. and Sykes, M. T.: Dynamic Global Vegetation Modeling: Quantifying Terrestrial 3 Ecosystem Responses to Large-Scale Environmental Change, in Terrestrial Ecosystems in a 4 Changing World, edited by J. G. Canadell, D. E. Pataki, and L. F. Pitelka, pp. 175-192, 5 Heidelberg. Springer Berlin [online] Available from: http://link.springer.com/chapter/10.1007/978-3-540-32730-1 15 (Accessed 6 January 2014), 6 7 2007.

- Prentice, I. C., Kelley, D. I., Foster, P. N., Friedlingstein, P., Harrison, S. P. and Bartlein, P.
 J.: Modeling fire and the terrestrial carbon balance, Glob. Biogeochem. Cycles, 25(3),
 GP3005 doi:10.1020/2010GP003206.2011
- 10 GB3005, doi:10.1029/2010GB003906, 2011.
- 11 Raynolds, M. K., Walker, D. A., Verbyla, D. and Munger, C. A.: Patterns of Change within a
- 12 Tundra Landscape: 22-year Landsat NDVI Trends in an Area of the Northern Foothills of the
- 13 Brooks Range, Alaska, Arct. Antarct. Alp. Res., 45(2), 249–260, doi:10.1657/1938-4246-
- 14 45.2.249, 2013.
- 15 Richardson, A. D., Anderson, R. S., Arain, M. A., Barr, A. G., Bohrer, G., Chen, G., Chen, J.
- 16 M., Ciais, P., Davis, K. J., Desai, A. R., Dietze, M. C., Dragoni, D., Garrity, S. R., Gough, C.
- 17 M., Grant, R., Hollinger, D. Y., Margolis, H. A., McCaughey, H., Migliavacca, M., Monson,
- R. K., Munger, J. W., Poulter, B., Raczka, B. M., Ricciuto, D. M., Sahoo, A. K., Schaefer, K.,
 Tian, H., Vargas, R., Verbeeck, H., Xiao, J. and Xue, Y.: Terrestrial biosphere models need
- better representation of vegetation phenology: results from the North American Carbon
- 21 Program Site Synthesis, Glob. Change Biol., 18(2), 566–584, doi:10.1111/j.1365-
- 22 2486.2011.02562.x, 2012.
- Richardson, A. D., Keenan, T. F., Migliavacca, M., Ryu, Y., Sonnentag, O. and Toomey, M.:
 Climate change, phenology, and phenological control of vegetation feedbacks to the climate
 system, Agric. For. Meteorol., 169, 156–173, doi:10.1016/j.agrformet.2012.09.012, 2013.
- Rost, S., Gerten, D., Bondeau, A., Lucht, W., Rohwer, J. and Schaphoff, S.: Agricultural
 green and blue water consumption and its influence on the global water system, Water
 Resour. Res., 44(9), W09405, doi:10.1029/2007wr006331, 2008.
- Saatchi, S. S., Harris, N. L., Brown, S., Lefsky, M., Mitchard, E. T. A., Salas, W., Zutta, B.
 R., Buermann, W., Lewis, S. L., Hagen, S., Petrova, S., White, L., Silman, M. and Morel, A.:
 Benchmark map of forest carbon stocks in tropical regions across three continents, Proc. Natl.
- 32 Acad. Sci., 108(24), 9899–9904, doi:10.1073/pnas.1019576108, 2011.
- Schaaf, C. B., Gao, F., Strahler, A. H., Lucht, W., Li, X., Tsang, T., Strugnell, N. C., Zhang,
 X., Jin, Y., Muller, J.-P., Lewis, P., Barnsley, M., Hobson, P., Disney, M., Roberts, G.,
 Dunderdale, M., Doll, C., d' Entremont, R. P., Hu, B., Liang, S., Privette, J. L. and Roy, D.:
 First operational BRDF, albedo nadir reflectance products from MODIS, Remote Sens.
 Environ., 83(1–2), 135–148, doi:10.1016/S0034-4257(02)00091-3, 2002.
- Schaphoff, S., Heyder, U., Ostberg, S., Gerten, D., Heinke, J. and Lucht, W.: Contribution of
 permafrost soils to the global carbon budget, Environ. Res. Lett., 8(1), 014026, 2013.

1 Schuur, E. A. G., Crummer, K. G., Vogel, J. G. and Mack, M. C.: Plant Species Composition and Productivity following Permafrost Thaw and Thermokarst in Alaskan Tundra, 2 3 Ecosystems, 10(2), 280-292, doi:10.1007/s10021-007-9024-0, 2007.

4 Shur, Y. L. and Jorgenson, M. T.: Patterns of permafrost formation and degradation in 5 relation to climate and ecosystems, Permafr. Periglac. Process., 18(1), 7–19, 2007.

- 6 Sitch, S., Huntingford, C., Gedney, N., Levy, P. E., Lomas, M., Piao, S. L., Betts, R., Ciais, 7 P., Cox, P., Friedlingstein, P., Jones, C. D., Prentice, I. C. and Woodward, F. I.: Evaluation of
- 8 the terrestrial carbon cycle, future plant geography and climate-carbon cycle feedbacks using
- 9 five Dynamic Global Vegetation Models (DGVMs), Glob. Change Biol., 14(9), 2015–2039,
- 10 doi:10.1111/j.1365-2486.2008.01626.x, 2008.
- Sitch, S., Smith, B., Prentice, I. C., Arneth, A., Bondeau, A., Cramer, W., Kaplan, J. O., 11 Levis, S., Lucht, W., Sykes, M. T., Thonicke, K. and Venevsky, S.: Evaluation of ecosystem 12 13 dynamics, plant geography and terrestrial carbon cycling in the LPJ dynamic global 14 vegetation model, Glob. Change Biol., (9), 161–185, 2003.
- Smith, B., Prentice, I. C. and Sykes, M. T.: Representation of vegetation dynamics in the 15 16 modelling of terrestrial ecosystems: comparing two contrasting approaches within European 17 climate space, Glob. Ecol. Biogeogr., 10(6), 621-637, doi:10.1046/j.1466-822X.2001.t01-1-18 00256.x, 2001.
- 19 Stöckli, R., Rutishauser, T., Baker, I., Liniger, M. a and Denning, a S.: A global reanalysis of 20 vegetation phenology, J. Geophys. Res., 116(G3), G03020-G03020, 21 doi:10.1029/2010jg001545, 2011.
- 22 Stöckli, R., Rutishauser, T., Dragoni, D., O'Keefe, J., Thornton, P. E., Jolly, M., Lu, L. and
- 23 Denning, A. S.: Remote sensing data assimilation for a prognostic phenology model, J. 24 Geophys. Res., 113(G4), G04021–G04021, 2008.
- 25 Stocks, B. J., Mason, J. A., Todd, J. B., Bosch, E. M., Wotton, M., Amiro, B. D., Flannigan, M. D., Hirsch, K. G., Logan, K. A., Martell, D. L. and Skinner, W. R.: Large forest fires in 26 27 Canada, 1959-1997, J. Geophys. Res., 107(D1), 8149-8149, doi:10.1029/2001jd000484, 28 2002.
- 29 Strengers, B. J., Müller, C., Schaeffer, M., Haarsma, R. J., Severijns, C., Gerten, D., Schaphoff, S., van den Houdt, R. and Oostenrijk, R.: Assessing 20th century climate-30 vegetation feedbacks of land-use change and natural vegetation dynamics in a fully coupled 31 32 vegetation-climate model, Int. J. Climatol., 30(13), 2055-2065, doi:10.1002/joc.2132, 2010.
- 33 Sturm, M., Racine, C. H. and Tape, K.: Climate change: Increasing shrub abundance in the 34 Arctic, Nature, 411(6837), 546–547, doi:10.1038/35079180, 2001.
- 35 Tao, X., Wang, D., Wu, D., Yan, B., Fan, W., Xu, X. and Yao, Y.: A model for instantaneous 36 FAPAR retrieval: Theory and validation, in Geoscience and Remote Sensing
- 37 Symposium, 2009 IEEE International, IGARSS 2009, vol. 1, pp. I-144-I-147., 2009.

- 1 Thonicke, K., Spessa, A., Prentice, I. C., Harrison, S. P., Dong, L. and Carmona-Moreno, C.:
- 2 The influence of vegetation, fire spread and fire behaviour on biomass burning and trace gas
- 3 emissions: results from a process-based model, Biogeosciences, 7(6), 1991–2011, 2010.

Thurner, M., Beer, C., Santoro, M., Carvalhais, N., Wutzler, T., Schepaschenko, D.,
Shvidenko, A., Kompter, E., Ahrens, B., Levick, S. R. and Schmullius, C.: Carbon stock and
density of northern boreal and temperate forests, Glob. Ecol. Biogeogr., 23(3), 297–310,
doi:10.1111/geb.12125, 2014.

- 8 Townshend, J., Carroll, M., Dimiceli, C., Sohlberg, R. a, Hansen, M. C. and DeFries, R. S.:
- 9 Vegetation Continuous Fields MOD44B. 2000 Percent Tree Cover, Collection 5, [online]
- 10 Available from: http://glcf.umd.edu/data/vcf/ (Accessed 15 January 2013), 2011.
- Tucker, C. J.: Red and Photographic Infrared Linear Combinations for Monitoring
 Vegetation, Remote Sens. Environ., 150(8), 127–150, 1979.
- 13 Tucker, C. J., Slayback, D. A., Pinzon, J. E., Los, S. O., Myneni, R. B. and Taylor, M. G.:
- Higher northern latitude normalized difference vegetation index and growing season trends
 from 1982 to 1999, Int. J. Biometeorol., 45(4), 184–190, doi:10.1007/s00484-001-0109-8,
 2001.
- Turner, D. P., Cohen, W. B., Kennedy, R. E., Fassnacht, K. S. and Briggs, J. M.:
 Relationships between Leaf Area Index and Landsat TM Spectral Vegetation Indices across
- 19 Three Temperate Zone Sites, Remote Sens. Environ., 70(April 1998), 52–68, 1999.
- 20 Urban, M., Forkel, M., Schmullius, C., Hese, S., Hüttich, C. and Herold, M.: Identification of
- 21 land surface temperature and albedo trends in AVHRR Pathfinder data from 1982 to 2005 for
- 22 northern Siberia, Int. J. Remote Sens., 34(12), 4491–4507,
- 23 doi:10.1080/01431161.2013.779760, 2013.
- Verstraeten, W. W., Veroustraete, F. and Feyen, J.: On temperature and water limitation of
 net ecosystem productivity: Implementation in the C-Fix model, Ecol. Model., 199(1), 4–22,
 doi:10.1016/j.ecolmodel.2006.06.008, 2006.
- Vitousek, P. M. and Howarth, R. W.: Nitrogen limitation on land and in the sea: How can it
 occur?, Biogeochemistry, 13(2), 87–115, doi:10.1007/BF00002772, 1991.
- Wagner, W., Scipal, K., Pathe, C., Gerten, D., Lucht, W. and Rudolf, B.: Evaluation of the
 agreement between the first global remotely sensed soil moisture data with model and
 precipitation data, J. Geophys. Res. Atmospheres, 108(D19), 4611,
 doi:10.1029/2003JD003663, 2003.
- Walter Shea, E. A., Blad, B. L., Mesarch, M. A., Hays, C. J., Deering, D. W. and Eck, T. F.:
 Absorbed photosynthetically active radiation and sun view geometry effects on remote
 sensing relationships, Remote Sens. Rev., 17(1-4), 89–102,
 doi:10.1080/02757259809532365, 1998.
- Wang, X., Piao, S., Ciais, P., Li, J., Friedlingstein, P., Koven, C. and Chen, A.: Spring
 temperature change and its implication in the change of vegetation growth in North America

1 from 1982 to 2006, Proc. Natl. Acad. Sci. U. S. A., 108(4), 1240–1245, 2 doi:10.1073/pnas.1014425108, 2011.

Williams, M., Richardson, A. D., Reichstein, M., Stoy, P. C., Peylin, P., Verbeeck, H.,
Carvalhais, N., Jung, M., Hollinger, D. Y., Kattge, J., Leuning, R., Luo, Y., Tomelleri, E.,
Trudinger, C. M. and Wang, Y. P.: Improving land surface models with FLUXNET data,
Biogeosciences, 6(7), 1341–1359, doi:10.5194/bg-6-1341-2009, 2009.

Wright, S. J.: Phenological Responses to Seasonality in Tropical Forest Plants, in Tropical 7 Forest Plant Ecophysiology, edited by S. S. Mulkey, R. L. Chazdon, and A. P. Smith, pp. 8 9 440-460, Springer US. [online] Available from: 10 http://link.springer.com/chapter/10.1007/978-1-4613-1163-8 15 (Accessed 26 February 11 2014), 1996.

- Xu, L., Myneni, R. B., Chapin Iii, F. S., Callaghan, T. V., Pinzon, J. E., Tucker, C. J., Zhu, Z.,
 Bi, J., Ciais, P., Tømmervik, H., Euskirchen, E. S., Forbes, B. C., Piao, S. L., Anderson, B. T.,
 Ganguly, S., Nemani, R. R., Goetz, S. J., Beck, P. S. A., Bunn, A. G., Cao, C. and Stroeve, J.
 C.: Temperature and vegetation seasonality diminishment over northern lands, Nat. Clim.
 Change, 3(6), 581–586, doi:10.1038/nclimate1836, 2013.
- Yuan, W., Zhou, G., Wang, Y., Han, X. and Wang, Y.: Simulating phenological
 characteristics of two dominant grass species in a semi-arid steppe ecosystem, Ecol. Res.,
 22(5), 784–791, doi:10.1007/s11284-006-0318-z, 2007.
- Zaehle, S., Sitch, S., Smith, B. and Hatterman, F.: Effects of parameter uncertainties on the
 modeling of terrestrial biosphere dynamics, Glob. Biogeochem. Cycles, 19(3), n/a–n/a,
 doi:10.1029/2004gb002395, 2005.

Zeng, H., Jia, G. and Epstein, H.: Recent changes in phenology over the northern high
latitudes detected from multi-satellite data, Environ. Res. Lett., 6(4), 045508–045508,
doi:10.1088/1748-9326/6/4/045508, 2011.

- Zhang, T.: Influence of the seasonal snow cover on the ground thermal regime: An overview,
 Rev. Geophys., 43(4), 2005.
- Zhu, Z., Bi, J., Pan, Y., Ganguly, S., Anav, A., Xu, L., Samanta, A., Piao, S., Nemani, R. and
 Myneni, R.: Global Data Sets of Vegetation Leaf Area Index (LAI)3g and Fraction of
 Photosynthetically Active Radiation (FPAR)3g Derived from Global Inventory Modeling and
 Mapping Studies (GIMMS) Normalized Difference Vegetation Index (NDVI3g) for the
- 32 Period 1981 to 2011, Remote Sens., 5(2), 927–948, doi:10.3390/rs5020927, 2013.

Table 1: Overview of optimization experiments with information sources for prior and posterior parameter sets. Parameter values and prior parameter ranges for each parameter set are listed in Supplement 4.1. 3

Experiment	Description	Number of randomly selected grid cells	Prior parameter set and sources	Posterior parameter set
OP.prior	Parameters or model results of LPJmL-OP with standard parameters		Table S2	
			Sitch et al. (2003): α_a , k, ramp, aphen _{min} , aphen _{max} , Wscal _{min}	
			Strengers et al. (2010): sfc and albedo parameters (partly estimated from MODIS albedo)	
OP.gc	Optimization of	530 in total	Table D2	One optimized
	single grid cells of LPJmL-OP.	TrBE 66, TrBR 51, TeNE 46, TeBE 32, TeBS 32, BoNE 68, BoBS 40, BoNS 49, TeH 66, TrH 80	Parameters as in OP.prior parameter set per grid cell. Median- averaged values for PFTs (Table S3)	
OP.pft	Optimization of	673 in total	Median-averaged values for PFTs from posterior values of OP.gc (Table S3)	(No useful posterior parameter sets were found)
(results not shown)	multiple grid of LPJmL-OP. Multiple grid cells of the same dominant PFT were optimized at the same time.	TrBE 50, TrBR 80, TeNE 50, TeBE 50, TeBS 80, BoNE 50, BoBS 80, BoNS 158, TeH 50, TrH 25		
GSI.prior	Parameters or model results of LPJmL-GSI with standard		Table S4	
			$\begin{array}{l} OP.gc: \alpha_{a}, k, sfc, \beta_{leaf}, \beta_{litter}, \\ and \beta_{stem} \end{array}$	
	parameters.		Stöckli et al. (2011): parameters for cold and light limiting functions derived from fitting logistic functions to stepwise functions as reported in Stöckli et al. (2011)	
GSI.gc	Optimization of single grid cells of LPJmL-GSI.	348 in total	Parameters as in GSI.prior (Table S4)	One optimized parameter set per grid cell.
		TrBE 33, TrBR 33, TeNE 32, TeBE 22, TeBS 43, BoNE 30, BoBS 41, BoNS 30, TeH 46, TrH 38		
GSI.pft	Optimization of multiple grid of LPJmL-GSI. Multiple grid cells of the same	500 in total TrBE 30, TrBR 30, TeNE 30, TeBE 30, TeBS 30, BoNE 50,	Parameters as in GSI.prior (Table S4)	Table S5 (one optimized parameter set per PFT)

dominant PFT	BoBS 30, BoNS 60,
were optimized at	TeH 70, TrH 70, PoH
the same time.	70

_

Table 2: Overview of global model runs that were used in this study for model evaluation.

Model run	Phenology model	Parameter set	Further settings
LPJmL- OP-prior	original phenology	LPJmL standard parameters as in the OP.prior experiment (Table S2)	dynamic vegetation/no prescribed land cover, prescribed agricultural land use, prescribed observed burnt area
LPJmL- OP-gc	original phenology	Optimized productivity, FAPAR and albedo parameters from the OP.gc optimization experiment, but original phenology parameters as in the OP.prior experiment (Table S3)	
LPJmL- GSI	GSI-based phenology	Parameters from the GSI.pft optimization experiment (Table S5)	



- 2 Figure 1. Structure of the model-data integration approach for LPJmL (LPJmL-MDI). The
- 3 LPJmL model structure is highly simplified.



Figure 2: Examples of the cold temperature, heat stress, light and water limiting functions for
 phenology in LPJmL-GSI. Depending on the chosen parameters the functions have different

4 shapes for each PFT.



Figure 3: Map of the dominant PFT in each grid cell as derived from SYNMAP, Köppen-Geiger climate zones and MODIS VCF. Grid cells that were used in any of the optimization experiments are shown as black crosses. Some grid cells were used in multiple optimization experiments. Grid cells that are dominated by agriculture were not used for optimization (TrML, tropical managed lands and TeML, temperate managed lands).

7



2 Figure 4: Distribution of the correlation coefficient between monthly LPJmL and GIMMS3g 3 FAPAR (1982-2011) for several grid cells in prior model runs and optimization experiments 4 grouped by plant functional types and biomes. (a) Correlation coeffcient for LPJmL-OP with 5 default parameters (a, OP.prior), after grid cell-level optimizations (b, OP.gc), cost for 6 LPJmL-GSI with prior parameters (c, GSI.prior), after grid cell-level optimizations (d, 7 GSI.gc) and after PFT-level optimizations (e, GSI.pft). Biomes are Tr (tropical), Te 8 (temperate) and Bo (boreal/polar). (b) Legend for the plot. Each distribution is plotted 9 according to usual boxplot statistics. The point symbols indicate the plant functional type. The significance flag on top of each distribution shows if a distribution is significant different ($p \le p$ 10 11 0.01) to the corresponding distribution of the same PFT in another optimization experiment. The significance is based on the Wilcoxon rank-sum test. For example "acd" indicates a 12 13 significant difference to the main categories a (OP.prior), c (GSI.prior) and d (GSI.gc) but no 14 significant difference to b (OP.gc) and e (GSI.pft).

15



1

2 Figure 5: Uncertainty and sensitivity of LPJmL-GSI parameters derived from all individuals 3 of genetic optimizations at PFT level. Shown is the relationship between parameter values and 4 the likelihood of the corresponding parameter vector. The likelihood is normalized with the 5 likelihood of the optimum parameter set. Only individuals with dAIC < 2 are shown. Grey 6 areas indicate the uniform prior parameter range. Red crosses indicate the optimum parameter 7 value. The optimum parameter value is indicated as text in a plot if it is outside of the plotting 8 range. Results from two independent optimization experiments are shown for the BoNS, TrH, 9 TeH and PoH PFTs (black and blue colours, respectively) but not all parameters were 10 included in both experiments. The parameter ALBEDO LITTER in the TrBE and TeBE 11 PFTs was not considered in optimization experiments.





2 Figure 6: Best LPJmL model runs for (a) monthly FAPAR dynamics (1982-2011, n = 3603 months) and (b) time series of mean annual FAPAR (1982-2011, n = 30 years). The best 4 LPJmL model run has the highest correlation coefficient between simulated LPJmL FAPAR and GIMMS3g FAPAR. If one model run is shown the correlation coefficient of this best 5 model is significant higher than of the second best model run ($p \le 0.05$, Fisher z-6 7 transformation on difference in correlation). If two model runs are shown the correlation 8 coefficients of the first and second best model runs are not significantly different from each 9 other (p > 0.05).



1

2 Figure 7: Comparison of trends in mean annual FAPAR from LPJmL and from satellite 3 datasets. Trends were computed between 1982 and 2011 as linear trends. The significance of 4 a trend was determined using the Mann-Kendall trend test. Only significant trends slopes ($p \le p$ 5 0.05) are displayed in each map. Spatial correlations of trend slopes (Spearman coefficient) 6 between LPJmL and the GIMMS3g dataset are given in the map titles. Time series are 7 showing mean annual FAPAR time series and trends spatially averaged for the regions as 8 indicated in the first map. The blue area in time series represents the uncertainty of the GL2 9 VGT FAPAR dataset. Numbers in the time series plot are correlation coefficients between 10 mean annual FAPAR time series from GIMMS3g and from GL2 or LPJmL model runs, respectively. The significance of a trend and of the correlation is indicated as point symbol: 11 *** $(p \le 0.001)$, ** $(p \le 0.01)$, * $(p \le 0.05)$, . $(p \le 0.1)$. 12



Figure 8: Latitudinal gradients of (a) gross primary production (GPP), (b) evapotranspiration,
(c) biomass and (d) tree cover from data-oriented estimates and from LPJmL model
simulations. Gradients were spatially averaged (median) from all 0.5° grid cells for latitudinal
bands of 1° width. Grey areas represent uncertainty estimates for the data-oriented estimates.





2 Figure 9: Phenological controls on seasonal FAPAR dynamics. The maps are red-green-blue 3 composites of the mean monthly values for the water (red), light (green) and cold temperature 4 (blue) phenology limiting function values from the LPJmL-GSI model run. White regions in 5 the maps are without vegetation or dominated by croplands for which the LPJmL-GSI 6 phenology model was not applied. Time series represent the mean seasonal cycles (January to 7 December) (averaged over 1982-2011) of simulated and observed FAPAR and phenology 8 limiting function values averaged for different regions as indicated in the first map. 9 Phenology limiting function values close to 0 indicate a strong control by phenology limiting 10 functions whereas values close to 1 indicate no phenological control. The correlation coefficients of each time series with the simulated FAPAR time series are shown in each time 11 12 series plot. The significance of the correlation is indicated as point symbol: *** ($p \le 0.001$), ** $(p \le 0.01)$, * $(p \le 0.05)$, . $(p \le 0.1)$. 13