

Protist community composition during early phytoplankton blooms in the naturally iron-fertilized Kerguelen area (Southern Ocean)

Clément Georges¹, Sébastien Monchy¹, Savvas Genitsaris¹, Urania Christaki^{1*}

5 [1]{INSU-CNRS, UMR 8187 LOG, Laboratoire d’Océanologie et de Géosciences, Université du Littoral Côte d’Opale, ULCO, 32 avenue Foch, F-62930 Wimereux, France}

* Corresponding author: Urania.Christaki@univ-littoral.fr

10

15

Abstract

Microbial eukaryotic community composition was examined by 18S rRNA gene tag pyrosequencing, during the early phase of spring phytoplankton blooms induced by natural iron fertilization, off Kerguelen Island in the Southern Ocean (KEOPS2 cruise). A total of 999
 25 operational taxonomical units (OTUs), affiliated to 30 known high-level taxonomic groups, were retrieved from 16 samples collected in the upper 300 m water column. The alveolata group was the most abundant in terms of sequence number and diversity (696 OTUs). The majority of alveolata sequences were affiliated to Dinophyceae and to two major groups of marine alveolates (MALV-I and MALV-II). In the upper 180 m, only 13% of the OTUs were
 30 shared between of the fertilized stations and the reference site characterized by high nutrient low chlorophyll (HNLC) waters. Fungi and Cercozoa were present in iron-fertilized waters, but almost absent in the HNLC samples, while Haptophyta and Chlorophyta characterized the HNLC sample. Finally, the 300 m depth samples of all stations were differentiated by the presence of MALV-II and Radiolaria. Multivariate analysis, examining the level of similarity
 35 between different samples, showed that protistan assemblages differed significantly between the HNLC and iron-fertilized stations, but also between the diverse iron-fertilized blooms.

Keywords: Planktonic protists, natural iron fertilization, 18S rRNA, tag pyrosequencing, Southern Ocean

1. Introduction

45 Molecular investigations into the planktonic protists of natural microbial communities have revealed an astonishing diversity (e.g. Caron et al., 2012 and references therein) and a variety of novel and/or previously unobserved groups of saprophytes, parasites, and intracellular symbionts (e.g Guillou et al., 2008; Massana and Pedrós-Alió, 2008; Bråte et al., 2012). The wide ecological roles of protists include: phototrophic and mixotrophic species, 50 belonging to the primary producers; heterotrophic species, acting as a 'link' between the microbial food web and the higher trophic levels; as well as decomposers and parasitic taxa (Caron et al., 2009 and references therein). A series of molecular studies have examined spatial or temporal patterns in protistan community structure and diversity. These have indicated that the microbial community structure is generally highly responsive to 55 environmental forcing, and that dominant protistan taxa can differ markedly over temporal and spatial scales associated with common oceanographic features (e.g Countway et al., 2007; 2010; Nolte et al., 2010; Gilbert et al., 2012; Mangot et al., 2013; Lie et al., 2013; Wolf et al., 2014; Christaki et al., 2014).

The Southern Ocean has a unique geography with several large-scale water masses 60 separated by oceanic fronts, and has major implications for the global ocean circulation and climate system. It is also the largest high nutrient-low chlorophyll (HNLC) ocean, where iron limits phytoplankton production, resulting in a large stock of major inorganic nutrients (Martin and Fitzwater, 1990). A pronounced shift to larger phytoplankton cells, in particular diatoms, has been generally observed resulting upon natural (Blain et al., 2007; Pollard et al., 65 2009) or artificial (Boyd et al., 2007; Smetacek et al., 2012) iron additions. While evidence of iron limitation of phytoplankton growth is unequivocal, the subsequent direct or indirect

impact of iron on heterotrophic eukaryotes of the microbial food web is less clear. For example, a moderate increase in microzooplankton biomass was observed during the iron-fertilization experiment IronEx-2 in the Equatorial Pacific sector and the SOIREE in the Southern Ocean (Landry et al., 2000; Hall and Safi, 2001). In contrast, the microzooplankton grazing pressure on the total phytoplankton community decreased during the iron-fertilization experiment SERIES in the Gulf of Alaska and the SEEDS1 in the western subarctic Pacific (Boyd et al., 2004; Saito et al., 2005). In the Kerguelen region, the iron limitation of the Southern Ocean is relieved by natural iron-fertilization (Blain et al., 2007). **Natural iron-fertilization is an uncommon process in which iron supply of the surface waters from iron-rich deep water is observed. Only two studies referred to natural iron fertilization in the vicinity of Crozet (Pollard et al., 2009) and Kerguelen Islands (Blain et al., 2007). The KEOPS 1 cruise demonstrated that the phytoplankton bloom was sustained by iron supply from iron-rich deep water below, representing natural iron fertilization (Blain et al., 2007). This study also showed that microzooplankton grazing was an important factor for phytoplankton biomass decrease in the bloom area (Brussaard et al., 2008) mainly affecting the small sized phytoplankton population (Brussaard et al., 2008; Christaki et al., 2008).**

The KEOPS2 cruise sampling strategy covered spatially diverse iron-fertilized stations at early bloom stages in the Kerguelen plateau and ocean region (Oct-Nov 2011). This data showed that natural iron-fertilization of the Southern Ocean on the scale of hundreds of thousands of square kilometers produced a mosaic of blooms, and that the biological and biogeochemical response to fertilization was diverse (Blain et al., this volume, a).

The objective of this study was to explore the microbial eukaryotic community structure using 18S rRNA gene tag pyrosequencing during the onset of spring phytoplankton blooms in the context of natural iron-fertilization of the Southern Ocean. The hypothesis tested was that the protistan communities would differ between the blooms, and between the iron-fertilized

blooms and the HNLC waters. The use of tag pyrosequencing provided a unifying approach for assessing the breadth of protistan communities, including the groups that are quasi impossible to characterize using traditional approaches of microscopy and culture (e.g. MAST, MALV, Fungi, and others).

2. Materials and Methods

2.1 Sample collection and DNA extraction

The present study was carried out during the KEOPS2 cruise from October 15th to November 20th 2011. Water samples were collected from four stations above and off the Kerguelen plateau (Fig. 1a, b). Stations A3-2, E-4W, and F-L were located in the blooms, while the reference station R-2 was located in the HNLC region (Fig. 1a, b). All water samples were collected with 12 L Niskin bottles mounted on a rosette equipped with a CTDO Seabird SBE911-plus. According to CTD profiles, four sampling depths were chosen at each station in order to represent the mixed layer (ML), the bottom of the ML, and the deeper waters (Table 1). Five to 7.5 liters of each depth were subsequently filtered on 10, 3, and 0.6 μ m, 47 mm nucleopore filters (Whatman, USA) using a serial filtration system at very low pressure (15 rpm). The serial filtration was performed in order to avoid filter clumping and to minimize disruption of fragile protists. The filters were immediately frozen in liquid nitrogen and then stored at -80 °C until analysis. After pooling together and cutting into small pieces the 10, 3, and 0.6 μ m filters, DNA extractions were carried out using the MO BIO PowerWater DNA Isolation Kit (MO BIO Laboratories, Inc, Carlsbad, CA), following the manufacturer's protocol instructions.

2.2 PCR and tag pyrosequencing

The DNA samples were amplified using the two universal eukaryote primers 18S-82F (5'-GAAACTGCGAATGGCTC-3', López-García et al., 2003) and Euk-516r (5'-ACCAGACTTGCCCTCC-3', Amann et al., 1990). These primers have been designed to amplify the variable V2 and V3 eukaryote 18S rRNA gene regions. A 10 bp tag sequence specific to each sample, a 4 bp TCAG key, and a 26 bp adapter for the GS FLX technology, were added to the primers. Polymerase chain reactions were carried out according to standard conditions for Platinum Tag High-Fidelity DNA polymerase (Invitrogen) with 10 ng of environmental DNA as a template. After the denaturation step at 94 °C for 2 min, 30 cycles of amplification were performed with a GeneAmp PCR System Apparatus (Applied Biosystems) as follows: 15 s at 94 °C, 30 s at 50 °C, 1 min at 72 °C, and 7 mins at 72 °C. Tag pyrosequencing was carried out by the company GenoScreen (Lille, France). The library was prepared following the procedures described by Roche (Basel, Switzerland) and used in a ¼ plate run on a 454 GS FLX Titanium sequencer. Pyrosequences were submitted on GenBank-SRA under the accession number SRP041236.

2.3 Quality filtering and taxonomic affiliations of the sequences

The sequences were processed using the MOTHUR 1.28.0 software (Schloss, 2009) following the standard operating procedure (http://www.mothur.org/wiki/Schloss_SOP) (Schloss et al., 2011). First, flowgrams were extracted and demultiplexed according to their tag. The resulting sixteen flowgrams were denoised using the MOTHUR 1.28.0 implementation of PyroNoise (Quince, 2009). Primer sequences, TAG, and key fragments were subsequently removed, and only sequences above 200 bp long, displaying less than eight homopolymers, were kept. The remaining sequences were dereplicated to unique sequences and aligned against the SILVA 108 database (<http://www.arb-silva.de/>) containing 62,587 eukaryotes SSU-18S rRNA sequences. Around 7 % of the sequences suspected of being chimeras were removed using the UCHIME software

(http://drive5.com/usearch/manual/uchime_algo.html) (Edgar, 2011). The remaining sequences were clustered into operational taxonomical units (OTUs) at 97 % similarity threshold. Single singletons (unique amplicons after 97 % clustering that occurred exclusively in only one sample) were removed from downstream analyses, as these are most likely erroneous sequencing products (Reeder and Knight, 2009; Kunin et al., 2010; Behnke et al., 2010). This dataset showed a representative overview of the diversity as indicated by the rarefaction curves reaching a plateau in most cases (Fig. S1; supplementary material). All OTUs were given a putative taxonomic affiliations based on BLAST (Altschul et al., 1990) identification of the closest cultured or uncultured relatives against the PR2 (Guillou et al., 2013) and the GenBank databases. The OTUs identified as metazoan, were removed from downstream analysis. However, the metazoan OTUs displayed high and heterogeneous number of sequences between samples, making subsampling of the remaining OTUs unsuitable as it resulted in a drastic loss of diversity. For this reason, the data are presented based on the relative abundance of OTUs in each sample.

2.4 Data Analysis

Rarefaction curves and alpha diversity estimators within particular samples (richness estimator S_{Chao1} ; the heterogeneity of the diversity; Simpson and Berger–Parker indices) were calculated with the PAST 2.17c software (Hammer et al., 2001). The S_{Chao1} approach uses the numbers of singletons and doubletons to estimate the number of expected species. According to S_{Chao1} , ‘missing’ species information is mostly concentrated on those of low frequency counts. The Simpson index measures the ‘evenness’ of the community and ranges from 0 (one taxon dominates the community) to 1 (all taxa are represented equally). Berger–Parker indicates the relative abundance of the dominant OTU in each sample (for more details, see

Maguran, 2004). Protistan assemblages, from the different samples, were compared using the Plymouth routines in the multivariate ecological research (PRIMER v.6) software package (Clarke and Gorley, 2006). In order to identify inter-relationships between samples, Bray–Curtis similarities were analyzed by cluster analysis and non-metric MDS on square-root sequence abundance. The similarity profile (SIMPROF) permutation test was conducted in PRIMER v.6 to establish the significance of dendrogram branches resulting from cluster analysis. Similarity percentage (SIMPER) analysis, also performed with PRIMER, was used to identify of the contribution of different OTUs to the observed similarity pattern.

3. Results

3.1 Study site

The hydrographic conditions during KEOPS2 are reported in detail in Blain et al. (this volume, a). The 'historical' A3 station situated ~ 500 m on the Kerguelen plateau (Blain et al., 2007; 2008) was characterized by a deep mixed layer (ML) (153 ± 15 m) (Table 1, Fig. 2). Stations F-L and E-4W revealed concentrations of 4.0 and $2.38 \mu\text{g L}^{-1}$ Chl *a*, respectively, constrained to shallow ML (38 ± 7 m and 61 ± 11 m, respectively; Table 1). The highest temperature was recorded in the ML of the F-L station (4.2°C , Fig. 2), indicating the influence of sub-Antarctic waters. The reference site (station R-2) in HNLC waters had low concentrations of Chl *a* ($0.25 \pm 0.08 \mu\text{g L}^{-1}$), and a temperature of 2.1°C (Fig. 2) in the ML (105 ± 15 m). The macronutrient concentrations in all 16 sampling points were high: ~ 20 – $26 \mu\text{M}$ for nitrate plus nitrite; ~ 1 – $1.8 \mu\text{M}$ for phosphate; ~ 8 – $19 \mu\text{M}$ for silicate; while dissolved iron was lower at the reference HNLC R-2 station (0.08 nM) relative to the iron-fertilized stations (0.16 – 0.22 nM ; Table 2).

3.2 Composition and distribution of protistan assemblages

After quality filtering and normalization, 999 unique OTUs, clustering 50,674 sequences
 190 (average length: 240 bp), were revealed for the 16 samples. The mean ratio of observed
 (Table 2) to expected (S_{chao1} , Table 2) OTUs was 75 ± 10 % (mean \pm sd) for all depths and
 stations. The highest number of unique OTUs, considering all depths, was observed at the F-L
 station (711 OTUs), and the lowest at the E-4W station (387 OTUs), while A3-2 and the
 HNLC R-2 stations had similar number of OTUs (550 and 496, respectively). The Simpson
 195 index, was relatively high, ranging from 0.76 (F-L station in the ML) to 0.99 (HNLC, R-2
 station at 300m). The Berger-Parker, indicating the relative abundance of the dominant OTU
 was generally low, except at the F-L station, where it reached its' highest value (0.48; Table
 2).

3.2.1 High-level taxonomic groups

200 The 999 OTUs were affiliated into 30 higher taxonomic groups distributed in all the
 samples (Table 3) and shown as pie charts for each of the four stations (Fig. 3). At all stations,
 Alveolata was the most diverse group (696 OTUs, mainly composed of MALV-II,
 Dinophyceae, MALV-I, and Ciliophora). The iron-fertilized stations accounted for the highest
 percentages of Alveolata while the lowest percentage was observed at the HNLC station R-2
 205 (Fig. 4). Stramenopiles were represented by 133 OTUs belonging to 10 higher taxonomic
 groups (Table 3). The most representative Stramenopile groups, in terms of OTUs number,
 were MAST, followed by Bacillariophyceae, and Labyrinthulomycetes (Table 3). The relative
 abundance of sequences of Stramenopiles ranged between 8 and 29 % in the mixed layer
 samples (Fig. 4). Radiolaria (belonging to Rhizaria) were present at all stations and were
 210 more abundant in the 300 m depth samples. Their relative abundance was particularly
 pronounced at station F-L, where they represented 55 % of all sequences (Fig. 4). The
 fertilized stations were characterized by lower relative abundances of Haptophyta and
 Chlorophyta compared with the HNLC R-2 station (Fig. 4). Fungi were represented by

relatively high OTU richness (28 OTUs; Table 3). They were found almost exclusively at the
fertilized stations, when only three OTUs were detected at the HNLC R-2 station (Fig. 3).

Regarding lineages distribution according to depth, the proportions of phototrophic protists
(e.g. Bacillariophyceae and Haptophyta) generally decreased below the ML. The relative
contribution of MALV-I and MALV-II increased with depth, at all stations except at station
F-L.

3.2.2 Most abundant OTUs

The most abundant 207 OTUs, representing > 1% of the sequences for each higher
taxonomic group, accounted for 95 % of the total sequences.

The heterotrophic *Gyrodinium* spp. was the dominant Dinophyceae genus in all samples,
while the small autotrophic *Gymnodinium* spp., also present in all samples, displayed higher
relative abundance in the HNLC R-2 samples (Table 4). Among Ciliophora, the genus
Strombidium was the most abundant, while different OTUs belonging to Tintinnid species
(Choreotrichia) were detected at all stations. The 17 most representative MAST-related OTUs
were distributed in eight clades, with a MAST-9 sp. prevailing at the surface F-L station
(Table 4).

At the fertilized stations, Bacillariophyceae-related OTUs were dominated by small sized
species such as *Planktoniella*, *Thalassiosira*, and *Minidiscus* spp., while *Pseudo-nitzschia* was
relatively abundant at the HNLC R-2 station (Table 4). Regarding the rest of the
Stramenopiles, the photosynthetic picoalgae of the genus *Bolidomonas* prevailed at all
stations. The non-photosynthetic Labyrinthulomycetes were more often found at the iron-
fertilized stations, with the parasitic genus *Oblongichytrium* sp. being relatively more
abundant at the E-4W and A3-2 stations (Table 4).

In all samples, the Haptophyta were dominated by *Phaeocystis antarctica*. Among
Chlorophyta, *Micromonas* were better represented at the F-L and R-2 stations, while

Pyramimonas spp. accounted for most of the Chlorophyta sequences at the A3-2 and E-4W
stations. Choanoflagellates comprised eight OTUs, all belonging to the Stephanoecidae. Fungi
were poorly represented at the HNLC R-2 station. Finally, Cercozoa were present at the iron-
fertilized stations, but almost absent at the HNLC station R-2 (Table 4).

3.3 Similarity of protistan assemblages

Altogether, the stations shared 197 OTUs, with 40 OTUs specific to the fertilized stations
(Fig. 5). The F-L station contained the highest number of exclusive OTUs (Fig. 5). The Bray-
Curtis similarity analysis of 999 OTUs indicated four major clusters (Fig. 6a). The SIMPROF
significance test indicated significant differences ($P < 0.05$) between these four groups and
showed significant differences within the groups (i) to (iv) (Fig. 6a). The two-dimensional
space nMDS visual representation, based on Bray-Curtis similarity analysis highlighted two
major clusters ('shallow' and 'deep' samples). An overall low similarity ($>15\%$) was observed
within each group (Fig. 6b). At a higher level of similarity (40-50 %), the clusters broke
roughly into individual stations: HNLC (cluster i); A3-2 (cluster ii); and E-4W (cluster iii);
while the F-L 20m and 65 m samples clustered with E-4W and the HNLC stations,
respectively (Fig. 6b). Within the 'deep' assemblage (cluster iv), the similarity between
samples was low, except for samples R 300m and F-L 180m, which displayed 40 % similarity
(Fig. 6b). The SIMPER test highlighted the most relevant OTUs forming each cluster (Table
5). In the first cluster (i), the major contributor was Haptophyta (in particular *P. antarctica*),
followed by Dinophyceae, and Chlorophyta. In the second cluster (ii), Dinophyceae
contributed to 49.2 % of the similarity, with *G. spirale*, having an important contribution
together with 10 other Dinophyceae and Bacillariophyceae-related OTUs. In the third cluster
(iii), Dinophyceae also prevailed (58.6 % of the similarity), with two OTUs affiliated to *G.*
spirale, where *G. rubrum* was the most important. Finally, the last cluster (iv), representing
the 'deep' samples, was characterized by MALV-II and Radiolaria.

4. Discussion

4.1 Overview of the commonly occurring taxa according to tag pyrosequencing

This is the first broad study of protist community composition in the natural iron-fertilized Kerguelen area of the Southern Ocean. The overall taxonomic diversity of protists recovered included 999 OTUs, belonging to 30 high level taxonomic groups. A total of 207 OTUs were classified as 'abundant' (each representing $\geq 1\%$ of sequences in their higher taxonomic group) (Table 4); the most frequent OTUs belonged to Alveolata, followed by Stramenopiles, then Hacrobia (Table 3).

4.1.1 Phytoplankton

Although the tag pyrosequencing of the 18S rRNA gene has become a routine method in marine microbial diversity studies, it is itself subjected to several limitations, including, DNA extraction and PCR-related biases, chimera formation, and primer non-universality (e.g. Prokopowich et al., 2003; Ki and Han, 2005; Zhu et al., 2005; Edgcomb et al., 2011). Although it has been established that Bacillariophyceae respond to iron-fertilization by rapidly forming extensive blooms (for a review see Quéguiner, 2013), concerning this study only 11 out of the 38 OTUs affiliated to Bacillariophyceae were found in common with the 52 diatom taxa morphologically identified in the Kerguelen area during the KEOPS 1 cruise at the end of the bloom period (Armand et al., 2008).

According to KEOPS2's microscopical observations and pigment analysis data, Bacillariophyceae dominated the phytoplankton community in the blooms (Sackett et al., this volume, Lasbleiz et al., this volume). In particular, *Fragilariopsis kerguelensis*, *Pseudonitzschia* spp., *Eucampia antarctica*, and *Chaetoceros* spp. were found to be the four dominant diatom taxa, via microscopy (Sackett et al., this volume). However, while *Pseudonitzschia*, *Eucampia*, and *Chaetoceros*-related OTUs represented 14 % of the Bacillariophyceae-related sequences, no *Fragilariopsis*-related OTUs were detected. Potential

limitation regarding pyrosequencing detection of Bacillariophyceae have been reported recently in an extensive study at the San Pedro Ocean Time-Series station (SPOT, Lie et al; 2013). They can be related to extraction efficiency from thick walled diatoms (Medinger et al., 2010) and/or amplification biases favouring species with high 18S rRNA gene copy number, such as ciliates and dinoflagellates (Potvin and Lovejoy, 2009). It is also worth noting that 28 out of the 52 taxa identified by microscopy (Armand et al., 2008) were not referenced in the GenBank. Finally, regarding the 27 diatom taxa that were 'identified' only by pyrosequencing -based on sequence similarity with the closest existing cultured relatives in GenBank-, they mainly belonged to the genera previously observed in this area (Armand et al., 2008). The accuracy of BLAST-derived taxonomy, especially at low-level taxa, depends on, sequence length, variability of the 18S region, database coverage for the specific taxonomic group, and correct identification of the reference sequence (Bik et al., 2012).

Sequences belonging to the nano- and pico-phytoplanktonic groups of Bolidophyceae, Pelagophyceae, Chrysophyceae, and Cryptophyta were found at relatively low abundances in all samples. Moreover, Haptophyta were dominated by an OTU affiliated as *Phaeocystis antarctica* (100 % sequence identity). This phylotype has been previously reported as dominant in the south of the polar front (Wolf et al., 2014), in the Ross Sea waters (DiTullio et al., 2000), and in the naturally iron-fertilized bloom around the Crozet plateau (Poulton et al., 2007).

4.1.2 Microzooplankton: Dinoflagellates, Ciliates, and Radiolaria

Although, Dinophyceae might be over-represented in the sequence data, possibly due to its high 18S gene copy number (e.g. Prokopowich et al., 2003; Zhu et al., 2005), tag pyrosequencing has allowed the highlighting of its extensive diversity (161 OTUs) in the Southern ocean; previously missed by conventional microscopy and/or pigment analysis (see also Wolf et al., 2014). For example, based on microscopy, *Gyrodinium* is the most abundant

dinoflagellate analyzed; however, no reliable distinction has been made between *G. spirale*
315 and *G. rubrum* with morphological observations (Saito et al., 2005; Georges et al., in prep.)

Ciliophora, which are ecologically important grazers of small sized phytoplankton, accounted for a relative high number of OTUs (60 OTUs). As with previous microscopic observations in the Kerguelen area (Christaki et al., 2008), the most representative ciliate sequences in this study belonged to Strombidiidae. The relatively large sized *Strombidium*
320 spp. ($\geq 50 \mu\text{m}$) can be plastidic (mixotrophic) and, along with *Tontonia* spp. and *Laboea* spp. -also present in sequences-, were found to contribute to 40-60 % of the aloricate ciliate biomass during the late bloom on the Kerguelen plateau (KEOPS1, Christaki et al., 2008). Finally, the most relatively abundant sequences of tintinnid taxa -which are also important nanophytoplankton consumers- belonged to the large *Cymatocyclis calyciformis* (Christaki et
325 al., 2008).

Radiolaria were another well-represented microzooplankton group (35 OTUs). These can act as particle feeders, by trapping their prey on the peripheral network of rhizopodia, or capture diatoms. They are also hosts of dinoflagellate symbionts and parasites, and may be important reservoirs of MALV taxa (e.g. Bråte et al., 2012). In this study, the relative increase
330 of MALV with depth was consistent with a parallel increase of Radiolaria. This observation is also supported by the hypothesis that MALV taxa are able to parasitize 'deeper' planktonic organisms such as Spumellarida (Guillou et al., 2008), which were the most common group and were always well represented in the deeper water samples in this study (Fig.4, Table 4). Radiolaria and MALV taxa characterizing deeper protistan assemblages have also been
335 reported in the North Atlantic (Countway et al., 2007; 2010; Not et al., 2007) and deep Antarctic polar front samples (López-García et al., 2001).

4.1.3 Symbionts, Parasites, and Decomposers

This assemblage included the taxonomic groups of MALV-I, MALV-II, Labyrinthulomycetes, Pirsonia, Oomyeta, Apicomplexa, Perkinsea, Fungi, and Cercozoa.

340 Many of these groups have a zooflagellate-stage in their life cycles; and are classified together in microscopical studies as ‘heterotrophic nanoflagellates’. MALV-I and MALV-II, appearing in virtually all marine surveys (López-García et al., 2001; Massana and Pedrós-Alió, 2008). Their considerable abundance and diversity suggests interactions with various hosts, and therefore, it has been proposed that the whole MALV assemblage is composed of marine
345 parasites (Skovgaard et al., 2005; Massana and Pedrós-Alió, 2008).

Fungi and Cercozoa accounted for 28 and 17 OTUs, respectively. In a recent succession study in the English Channel, it was observed that these groups mostly co-occurred with Bacillariophyceae (Christaki et al., 2014). Fungi are possibly related to the polysaccharide degradation of the freshly produced organic material by primary producers (Kimura and
350 Naganuma 2001; Raghukumar, 2004). It is known for diatoms that polysaccharides are their main exudates (Myklestad, 1995 and references therein), and these sugars could promote the growth of Fungi. Many Cercozoa are parasites of marine organisms, including large heavily silicified diatoms (e.g. Tillman et al., 1999; Schnepf and Kühn, 2000), which could explain why Fungi and Cercozoa were detected in the bloom stations and were poorly represented (2-
355 3 OTUs, Table 4) at the HNLC R-2 station. Labyrinthulomycetes were also better represented in terms of numbers of sequences in the bloom stations (Table 4). Labyrinthulomycetes (19 OTUs) are common osmo-heterotrophic marine protists (López-García et al., 2001) having parasitic, commensalistic, or mutualistic relationships with their hosts. They play an important role in decomposition processes (Collado-Mercado et al., 2010) by colonizing fecal pellets,
360 including under deep-sea conditions (Raghukumar, 2004).

4.1.4. Small heterotrophic protists

Among the small heterotrophic protists found in the samples, there were a variety of MAST (46 OTUs), Choanoflagellida (10 OTUs), and Telonemia (12 OTUs). MAST taxa are widely distributed in the world's oceans, and have been identified as free-living bacterivorous heterotrophic flagellates through a combination of FISH and other measurements (Massana et al., 2006, Jürgens and Massana, 2008 for a review). Choanoflagellida of the genus *Stephanotheca* sp. were also observed by epifluorescence microscopy in KEOPS2 samples, and were more abundant and diversified in the 0 - 200 m layer (KEOPS2 data, Georges et al., in prep.).

4.2 Variability of protistan assemblages relative to iron-fertilization

In general, the stability of OTUs richness and diversity indices between the HNLC R-2 and iron-fertilized stations indicated that the environment maintained an overall diversity across stations and depths (Table 2). These observations are in agreement with previous molecular studies based on protistan diversity (e.g. Countway et al., 2007; Monchy et al., 2012). However, community structure analysis showed clear differences inside and outside the blooms (Fig. 6a).

4.2.1 HNLC station

Based on trophic organization, HNLC areas seem conceptually similar to oligotrophic regions dominated by small producers and an active microbial food web (e.g. Hall and Safi, 2001; Oliver et al., 2004; Christaki et al., 2008; Obernosterer et al., 2008; Christaki et al., this volume). The characteristic contributors of the HNLC cluster (i) were Haptophyta, Chlorophyta, and MAST, which included mainly nanoplanktonic organisms. During KEOPS2, the relative importance of small-sized cells at the HNLC station is in accordance with the flow cytometry data ($4.8 \pm 1.9 \times 10^3 \text{ mL}^{-1}$ nano-picophytoplankton cells in comparison to $1.8 \pm 1.3 \times 10^3 \text{ mL}^{-1}$ at the bloom stations; KEOPS2 data). The factors influencing phytoplankton community composition (e.g. diatoms vs. *Phaeocystis* sp.) in the Southern

Ocean are a complex interplay between bottom up (iron-silicate-light availability; controlling growth) and top down effects (grazing; controlling mortality) (Cullen, 1991; Arrigo et al., 1999; Smetacek et al., 2004; Schoemann et al., 2005). Live plankton observations completed
390 on board (<https://www.youtube.com/watch?v=KPgoz8bWRJU>) revealed the presence of small colonies and free-living cells belonging to the Haptophyta *Phaeocystis* sp. at all stations. It seems that *Phaeocystis* species cope best with the environmental conditions in the open ocean waters south of the Polar Front, where it was found to be the most dominant phylotype (Wolf et al., 2014).

4.2.2 Iron-fertilized sites

The mechanisms that fertilize the surface water in the region around Kerguelen are complex, which results in a patchwork of blooms with diverse biological and biogeochemical response (Blain et al., this volume, a). The phytoplankton bloom at the 'historical' A3 station situated on the Kerguelen plateau is bottom-up sustained by low-level supplies of iron and
400 other nutrients (Blain et al., 2007). Drifters have revealed a north-eastward driven circulation pattern in the Kerguelen Plateau and oceanic area, while strong horizontal mixing have been found in the East Kerguelen Basin off the plateau (Zhou et al; this volume; Fig. 1b). Station E-4W is located at the shelf break in a region with very strong currents (Zhou et al., this volume), and consequently receives iron-rich waters from the Kerguelen Island and Plateau
405 (A3 station area) which mix with Polar Front waters that cross the Kerguelen plateau while traveling northeast (Fig. 1b). The depth of the ML varied considerably, from 40 m north of the Polar Front at station F-L to 170 m above the plateau at station A3. In accordance with these hydrographic characteristics, multivariate analysis of sequences showed that the ML sample of the F-L (20 m) was found in the same cluster as the E-4W samples, while the 65 m
410 F-L sample was grouped with the HNLC samples. The OTUs putatively affiliated to heterotrophic dinoflagellate taxa (Table 5) were the major contributors of clusters (ii) and (iii)

(Fig. 6a, b). Dinoflagellate increase during iron-fertilized blooms, in particular, *Gyrodinium* spp. has been observed with microscopic counts during the iron addition experiments, and has been attributed to the increase of their diatoms prey (Hall and Safi, 2001; Saito et al. 2005; Henjes et al. 2007).

Concluding, the tag pyrosequencing approach in this study has provided an overview of the protistan assemblages present in the naturally fertilized blooms and the HNLC waters in the Southern Ocean. Despite the under-representation of Bacillariophyceae diversity and the over-representation of Dinophyceae in the sequences, the community similarity analysis showed clear differences between the iron-fertilized and the HNLC waters, and among the blooms, in regards to their location and the fertilization mechanisms. The molecular approach has also highlighted a rich assemblage of potential phytoplankton parasites and organic matter decomposers mostly present in the iron-fertilized blooms.

Acknowledgements

KEOPS was financed by INSU-CNRS, IPEV and ANR and the French ministry of higher education through a PhD grant to CG. We thank our many colleagues who participated in the collection of various data sets and the crew aboard the R/V 'Marion Dufresne' for their help in the successful completion of the cruise. We also thank www.englisheditor.webs.com for the paper's English proofing.

References

- Altschul, S.F., Gish, W., Miller, W., Myers, E.W., and Lipman D.J.: Basic local alignment search tool, *J. Mol. Biol.*, 215, 403–410, 1990.
- Amann, R.I., Binder, B.J., Olson, R.J., Chisholm, S.W., Devereux, R., and Stahl, D.A.: Combination of 16S rRNA-targeted oligonucleotide probes with flow cytometry for analyzing mixed microbial populations, *Appl. Environ. Microbiol.*, 56, 1919–192, 1990.

- Armand, L.K., Crosta, X., Quéguiner, B., Mosseri, J., and Garcia, N.: Diatoms preserved in surface sediments of the northeastern Kerguelen Plateau, *Deep Sea Res. II*, 55, 677-692, 2008.
- Arrigo, K.R., Robinson, D.H., Worthen, D.L., Dunbar, R.B., DiTullio, G.R., VanWoert, M.,
440 and Lizotte, M.P.: Phytoplankton community structure and the drawdown of nutrients and CO₂ in the Southern Ocean, *Science*, 283, 365-367, 1999.
- Behnke, A., Engel, M., Christen, R., Nebel, M., Klein, R.R., Stoeck, T.: Depicting more accurate pictures of protistan community complexity using pyrosequencing of hypervariable SSU rRNA gene regions, *Environ. Microbiol.*, 13, 340-349, 2010.
- 445 Bik, H.M., Porazinska, D.L., Creer, S., Caporaso, J.G., Knight, R., and Thomas, K.W.: Sequencing our ways towards understanding global eukaryotic biodiversity, *Trends Ecol. Evol.*, 27, 233-243, 2012.
- Blain, S., Queguinblainer, B., Armand, L., Belviso, S., Bombled, B., Bopp, L., Bowie, A., Brunet, C., Brussaard, C., Carlotti, F., Christaki, U., Corbiere, A., Durand, I., Ebersbach,
450 F., Fuda, J.L., Garcia, N., Gerringa, L., Griffiths, B., Guigue, C., Guillerm, C., Jacquet, S., Jeandel, C., Laan, P., Lefevre, D., Lo Monaco, C., Malits, A., Mosseri, J., Obernosterer, I., Park, Y.H., Picheral, M., Pondaven, P., Remenyi, T., Sandroni, V., Sarthou, G., Savoye, N., Scouarnec, L., Souhaut, M., Thuiller, D., Timmermans, K., Trull, T., Uitz, J., van Beek, P., Veldhuis, M., Vincent, D., Viollier, E., Vong, L., and Wagener, T.: Effect of
455 natural iron fertilization on carbon sequestration in the southern ocean, *Nature*, 446, 1070-1074, 2007.
- Blain, S., Quéguiner, B., and Trull, T.: The natural iron fertilization experiment KEOPS (Kerguelen Ocean and Plateau compared Study): an overview, *Deep-Sea Res. II*, 55, 559-565, 2008.
- 460 Blain, S.: KEOPS2: Implementation and overview, this volume (a).

Blain, S., Oriol, L., Capparos, J., Guéneuguès, A., and Obernosterer, I.: Distributions and stoichiometry of dissolved nitrogen and phosphorus in the iron fertilized region near Kerguelen (Southern Ocean), this volume. (b)

Boyd, P.W., Law, C.S., Wong, C.S., Nojiri, Y., Tsuda, A., Levasseur, M., Takeda, S., Rivkin, R., Harrison, P.J., Strzepek, R., Gower, J., McKay, R.M., Abraham, E., Arychuk, M., Barwell-Clarke, J., Crawford, W., Crawford, D., Hale, M., Harada, K., Johnson, K., Klyosawa, H., Kudo, I., Marchetti, A., Miller, W., Needoba, J., Nishioka, J., Ogawa, H., Page, J., Robert, M., Saito, H., Sastri, A., Nelson, S., Soutar, T., Sutherland, N., Taira, Y., Whitney, F., Wong, S.-K.E., and Yoshimura, T.: The decline and fate of an iron-induced subarctic phytoplankton bloom, *Nature* 428, 549–553, 2004.

Boyd, P.W., Jickells, T., Law, C.S., Blain, S., Boyle, E.A., Buesseler, K.O., Coale, K.H., Cullen, J.J., de Baar, H.J. W., Follows, M., Harvey, M., Lancelot, C., Levasseur, M., Owens, N.P.J., Pollard, R., Rivkin, R.B., Sarmiento, J., Schoemann, V., Smetacek, V., Takeda, S., Tsuda, A., Turner, S., and Watson, A.J.: Mesoscale iron enrichment experiments 1993–2005: Synthesis and future directions, *Science*, 315, 612–617, 2007.

Bråte, J., Krabberød, A.K., Dolven, J.K., Ose, R.F., Kristensen, T., Bjørklund, K.R.: Radiolaria associated with large diversity of marine alveolates, *Protist*, 163, 767–777, 2012.

Brussaard, C.P.D., Timmermans, K.R., Uitz, J., and Veldhuis, M.J.W.: Virioplankton dynamics and virally induced phytoplankton lysis versus microzooplankton grazing southeast of the Kerguelen (Southern Ocean), *Deep-Sea Res. II*, 55, 752–765, 2008.

Caron, D.A., Countway, P.D., Savai, P., Gast, R.J., Schnetzer, A., Moorthi, S.D, Dennett M.R., Moran, D.M. and Jones, A.C.: Defining DNA-based operational taxonomic units for microbial–eukaryote ecology, *Appl. Environ. Microbiol.*, 75, 5797–5808, 2009.

- 485 Caron, D.A., Countway, P.D., Jones, A.C., Kim, D.Y., and Schnetzer, A.: Marine protistan
diversity, *Ann. Rev. Mar. Sci.*, 4, 467–493, 2012.
- Christaki, U., Obernosterer, I., Van Wambeke, F., Veldhuis, M.J.W., Garcia, N., and Catala,
P.: Microbial food web structure in a naturally iron fertilized area in the Southern Ocean
(Kerguelen Plateau), *Deep-Sea Res. II*, 55, 706-719, 2008.
- 490 Christaki, U., Kormas, K.A., Genitsaris, S., Georges, C., Sime-Ngando, T., Viscogliosi, E.,
and Monchy, S.: Winter-summer succession of unicellular eukaryotes in a meso-eutrophic
coastal system, *Microb. Ecol.*, 67, 13-23, 2014.
- Christaki, U., Lefèvre, D., Georges, C., Colombet, J., Catala, P., Courties, C., Sime-Ngando,
T., Blain, S., and Obernosterer I. (2014) Microbial food web dynamics during spring
495 phytoplankton blooms in the naturally iron-fertilized Kerguelen area (Southern Ocean),
this volume.
- Clarke K.R., and Gorley, R.N.: *Primer v6: user manual/tutorial*, PRIMER-E, Plymouth, 2006.
- Closset, I., Lasbleiz, M., Leblanc, K., Quéguiner, B., Cavagna, A.-J., Elskens, M., Navez, J.,
and Cardinal D.: Seasonal evolution of net and regenerated silica production around a
500 natural Fe-fertilized area in the Southern Ocean estimated from Si isotopic approaches, this
volume.
- Collado-Mercado E., Radway, J.C., and Collier, J.L.: Novel uncultivated labyrinthulomycetes
revealed by 18S rDNA sequences from seawater and sediment samples, *Aquat. Microb.
Ecol.*, 58, 215-228, 2010.
- 505 Countway, P.D., Gast, R.J., Dennett, M.R., Savai, P., Rose J.M., and Caron, D.A.: Distinct
protistan assemblages characterize the euphotic zone and deep sea (2500 m) of the western
North Atlantic (Sargasso Sea and Gulf Stream), *Environ. Microbiol.*, 9, 1219-1232, 2007.

Countway, P.D., Vigil, P.D., Schnetzer, A., Moorthi, S.D., and Caron, D.A.: Seasonal analysis of protistan community structure and diversity at the USC Microbial Observatory (San Pedro Channel, North Pacific Ocean), *Limnol. Oceanogr.*, 55, 2381–2396, 2010.

Cullen, J.J.: Hypotheses to explain high-nutrient conditions in the open sea, *Limnol. Oceanogr.*, 36, 1578–1599, 1991.

DiTullio, G.R., Grebmeier, J.M., Arrigo, K.R., Lizotte, M.P., Robinson, D.H., Leventer, A., Barry, J.P., VanWoert, M.L., and Dunbar, R.B.: Rapid and early export of *Phaeocystis Antarctica* blooms in the Ross Sea, Antarctica, *Nature*, 404, 595–598, 2000.

Edgar, R.C.: Search and clustering orders of magnitude faster than BLAST, *Bioinformatics*, 26, 2460–2461, 2010.

Edgcomb, V., Orsi, W., Bunge, J., Jeon, S., Christen, R., Leslin, C., Holder, M., Taylor, G.T., Suarez, P., Varela, R., and Epstein, S.: Protistan microbial observatory in the Cariago Basin, Caribbean. I. Pyrosequencing vs. Sanger insights into species richness, *ISME J.*, 5, 1344–1356, 2011.

Gilbert, J.A., Steele, J.A., Caporaso, J.G., Steinbruck, L., Reeder, J., Temperton, B., Huse, S., McHardy, A.C., Knight, R., Joint, I., Somerfield, P., Fuhrman, J.A., and Field, D.: Defining seasonal marine microbial community dynamics, *ISME J.*, 6, 298–308, 2012.

Guillou, L., Viprey, M., Chambouvet, A., Welsh, R.M., Kirkham, A.R., Massana, R., Scanlan, D.J., and Worden A.Z.: Widespread occurrence and genetic diversity of marine parasitoids belonging to Syndiniales (Alveolata), *Environ. Microbiol.*, 10, 3349–3365, 2008.

Guillou, L., Bachar, D., Audic, S., Bass, D., Berney, C., Bittner, L., Boutte, C., Burgaud, G., de Vargas, C., Decelle, J., Del Campo, J., Dolan, J.R., Dunthorn, M., Edvardsen, B., Holzmann, M., Kooistra, W.H., Lara, E., Le Bescot, N., Logares, R., Mahé, F., Massana, R., Montresor, M., Morard, R., Not, F., Pawlowski, J., Probert, I., Sauvadet, A.L., Siano,

- R., Stoeck, T., Vaulot, D., Zimmermann, P., and Christen, R.: The Protist Ribosomal Reference database (PR2): a catalog of unicellular eukaryote small sub-unit rRNA sequences with curated taxonomy, *Nucleic Acid. Res.*, 41, 597–604, 2013.
- 535
- Hall, J.A., and Safi, K.: The impact of in situ Fe fertilization on the microbial food web in the Southern Ocean, *Deep-Sea Res. II*, 48, 2591–2613, 2001.
- Hammer Ø, Harper D.A.T., and Ryan P.D.: PAST: paleontological statistics software package for education and data analysis, *Palaeontol. Electr.*, 4, 9, 2001.
- 540
- Henjes, J., Assmy, P., Klaas, C., and Smetacek, V.: Response of the larger protozooplankton to an iron-induced phytoplankton bloom in the Polar Frontal Zone of the Southern Ocean (EisenEx), *Deep Sea Res. I*, 54, 774-791.
- Jürgens, K., and Massana, R.: Protistan grazing on marine bacterioplankton, In: Kirchman D.L. (Ed): *Microbial Ecology of the Oceans*, 2nd Edition, pp. 383-441, 2008.
- 545
- Kimura H., and Naganuma T.: Thraustochytrids: a neglected agent of the marine microbial food chain, *Aquat. Ecosyst. Health Manag.*, 4, 13–18, 2001.
- Ki, J.-S., and Han, M.-S.: Efficient 5' ETS walking from conserved 18S rDNA sequences of the dinoflagellates *Alexandrium* and *Akashiwo sanguinea* (Dinophyceae), *J. Appl. Phycol.*, 17, 475-481, 2005.
- 550
- Kunin, V., Engelbrektson, A., Ochman, H., and Hugenholtz, P.: Wrinkles in the rare biosphere: pyrosequencing errors can lead to artificial inflation of diversity estimates, *Environ. Microbiol.*, 12, 118–123, 2010.
- Landry, M.R., Ondrusek, M.E., Tanner, S.J., Brown, S.L., Costantinou, J., Bidigare, R.R., Coale, K.H., and Fitzwater, S.: Biological response to iron fertilization in the eastern equatorial Pacific (IronEx II). I. Microplankton community abundances and biomass, *Mar. Ecol. Prog. Ser.*, 201, 27–42, 2000.
- 555

- Lasbleiz, M., Lefèvre, D., Leblanc, K., and Quéguiner, B.: Biological productivity regime and *in-situ* methods comparison around the Kerguelen Island in the Southern Ocean, this volume.
- 560 Lie, A.A.Y., Kim, D.Y., Schnetzer, A., and Caron, D.A.: Small-scale temporal and spatial variations in protistan community composition at the San Pedro Ocean Time-series station off the coast of southern California, *Aquat. Microb. Ecol.*, 2, 93-110, 2013.
- López-García, P., Rodríguez-Valera, F., Pedrós-Alió, C., and Moreira, D.: Unexpected diversity of small eukaryotes in deep-sea Antarctic plankton, *Nature*, 409, 603-607, 2001.
- 565 López-García P., Philippe H., Gail F., and Moreira D.: Autochthonous eukaryotic diversity in hydrothermal sediment and experimental microcolonizers at the Mid-Atlantic Ridge, *P. Natl. Acad. Sci. USA*, 100, 697–702, 2003.
- Magurran, A.E.: *Measuring biological diversity*. Blackwell, Malden, 2004
- Mangot, J.F., Domaizon, I., Taib, N., Marouni, N., Duffaud, E., Bronner, G., and Debroas, D.: Short-term dynamics of diversity patterns: evidence of continual reassembly within lacustrine small eukaryotes, *Environ. Microbiol.*, 15, 1745–1758, 2013.
- 570 Martin, J.H., and Fitzwater, S.E.: Iron deficiency limits phytoplankton growth in Antarctic waters, *Global Biogeochem. Cycles*, 4, 5–12, 1990.
- Massana, R., Terrado, R., Forn, I., Lovejoy, C. and Pedrós-Alió, C.: Distribution and abundance of uncultured heterotrophic flagellates in the world oceans, *Environ. Microbiol.*, 8, 1515–1522, 2006.
- 575 Massana, R., and Pedrós-Alió, C.: Unveiling new microbial eukaryotes in the surface ocean, *Curr. Opin. Microbiol.*, 11, 213–218, 2008.
- Medinger, R., Nolte, V., Pandey, R.V., Jost, S., Ottenwalder, B., Schlotterer, C., and Boenigk, J.: Diversity in a hidden world: potential and limitation of next-generation sequencing for
- 580

surveys of molecular diversity of eukaryotic microorganisms, *Mol. Ecol.*, 19 (Suppl 1), 32–40, 2010.

Monchy, S., Grattepanche, J.D., Breton, E., Meloni, D., Sancier, G., Chabe, M., Delhaes, L., Viscogliosi, E., Sime-Ngando, T., and Christaki U.: Microplanktonic community structure
585 in a coastal system relative to a *Phaeocystis* bloom inferred from morphological and tag
pyrosequencing methods, *PLoS ONE* 7, e39924, 2012.

Myklestad, S.M.: Release of extracellular products by phytoplankton with special emphasis
on polysaccharides, *Sci. Total Environ.*, 165, 155–164, 1995.

Nolte, V., Pandey, R.V., Jost, S., Medinger, R., Ottenwälder, B., Boenigk, J., and Schlötterer,
590 C.: Contrasting seasonal niche separation between rare and abundant taxa conceals the
extent of protist diversity, *Mol. Ecol.*, 19, 2908–2915, 2010.

Not, F., Valentin, K., Romari, K., Lovejoy, C., Massana, R., Tobe, K., Vaultot, D., and
Medlin, L.K.: Picobiliphytes: a marine picoplanktonic algal group with unknown affinities
to other eukaryotes, *Science*, 315, 253-255, 2007.

595 Obernosterer, I., Christaki, U., Lefevre, D., Catala, P., Van Wambeke, F., and LeBaron, P.:
Rapid bacterial mineralization of organic carbon produced during a phytoplankton bloom
induced by natural iron fertilization in the Southern Ocean, *Deep-Sea Res. II*, 55, 777-789,
2008.

Oliver, J.L., Barber, R.T., Smith, W.O., and Ducklow, H.W.: The heterotrophic bacterial
600 response during Southern Ocean Iron Experiment (SOFEX), *Limnol. Oceanogr.* 49, 2129–
2140, 2004.

Pollard, R., Salter, I., Sanders, R., Lucas, M., Moore, C., Mills, R., Statham, P., Allen, J.,
Baker, A. and Fones, G.: Southern Ocean deep-water carbon export enhanced by natural
iron fertilization, *Nature*, 457, 577-581, 2009.

- 605 Potvin, M., and Lovejoy, C.: PCR-based diversity estimates of artificial and environmental
18S rRNA gene libraries, *J. Eukaryot. Microbiol.*, 56, 174-181, 2009.
- Poulton, A.J., Moore, M., Seeyave, S., Lucas, M.I., Fielding, S., and Ward, P.: Phytoplankton
community composition around the Crozet Plateau, with emphasis on diatoms and
Phaeocystis, *Deep Sea Res. II*, 54, 2085-2105, 2007.
- 610 Prokopowich, C.D., Gregory, T.R., and Crease, T.J. : The correlation between rDNA copy
number and genome size in eukaryotes, *Genome*, 46, 48-50, 2003.
- Quéguiner, B.: Iron fertilization and the structure of planktonic communities in high nutrient
regions of the Southern Ocean, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 90, 43–54,
2013.
- 615 Quérroué, F., Sarthou, G.: The distribution of dissolved trace elements around the Kerguelen
plateau region, this volume
- Quince, C., Lanzen, A., Curtis, T.P., Davenport, R.J., Hall, N., Head, I.M., Read, LF, and
Sloan WT.: Accurate determination of microbial diversity from 454 pyrosequencing data,
Nat. Methods, 6, 639–641, 2009.
- 620 Raghukumar, S.: The role of fungi in marine detrital processes, In: Ramaiah, N.E. (ed) *Marine
microbiology: facets & opportunities*, National Institute of Oceanography, Goa, pp 91–
101, 2004.
- Reeder, J., and Knight, R.: The ‘rare biosphere’: a reality check, *Nat. Methods*, 6, 636–637,
2009.
- 625 Saito, H., Suzuki, K., Hinuma, A., Takahashi, O., Fukami, K., Kiyosawa, H., Saino, T., Saino,
T., and Tsuda, A.: Responses of microzooplankton to in situ iron fertilization in the
Western Subarctic Pacific (SEEDS), *Prog. Oceanogr.*, 64, 223–236, 2005.

Sackett, O., Armand, L., Beardall, J., Hill, R., Doblin, M., Connelly, C., Howes, J., Stuart, B.,
 Ralph, P., and Heraud, P.: Taxon specific responses of Southern Ocean diatoms to Fe
 630 enrichment revealed by FTIR microspectroscopy, this volume.

Schloss, P.D., Westcott, S.L., Ryabin, T., Hall, J.R., Hartmann, M., Hollister, E.B.,
 Lesniewski, R.A., Oakley, B.B., Parks, D.H., Robinson, C.J., Sahl, J.W., Stres, B.,
 Thallinger, G.G., Van Horn, D.J., and Weber C.F.: Introducing mothur: open-source,
 platform-independent, community-supported software for describing and comparing
 635 microbial communities, *Appl. Environ. Microbiol.*, 75, 7537–7541, 2009.

Schloss, P.D., Gevers, D., and Westcott, S.L.: Reducing the effects of PCR amplification and
 sequencing artifacts on 16S rRNA-based studies, *PLoS ONE*, 6, e27310, 2011.

Schnepf, E., and Kühn, S.F.: Food uptake and fine structure of *Cryothecomonas longipes* sp.
 nov., a marine nanoflagellate incertae sedis feeding phagotrophically on large diatoms,
 640 *Helgol. Mar. Res.*, 54, 18–32, 2000.

Schoemann, V., Becquevort, S., Stefels, J., Rousseau, V., and Lancelot, C.: *Phaeocystis*
 blooms in the global ocean and their controlling mechanisms: a review, *J. Sea Res.*, 53, 43-
 66, 2005.

Skovgaard, A., Massana, R., Balague, V., and Saiz E.: Phylogenetic position of the copepod-
 645 infesting parasite *Syndinium turbo* (Dinoflagellata, Syndinea), *Protist*, 156, 413–423, 2005.

Smetacek, V., Assmy, P., and Henjes, J.: The role of grazing in structuring Southern Ocean
 pelagic ecosystems and biogeochemical cycles, *Antartct. Sci.*, 16, 541-558, 2004.

Smetacek, V., Klaas, C., Strass, V.H., Assmy, P., Montresor, M., Cisewski, B., Savoye, N.,
 Webb, A., D’ Ovidio, F., Arrieta, J.M., Bathmann, U., Bellerby, R., Berg, G.M., Croot, P.,
 650 Gonzalez, S., Henjes, J., Herndl, G.J., Hoffmann, L.J., Leach, H., Losch, M., Mills, M.M.,
 Neill, C., Peeken, I., Röttgers, R., Sachs, O., Sauter, E., Schmidt, M.M., Schwarz, J.,

Terbrüggen, A., and Wolf-Gladrow, D.: Deep carbon export from a Southern Ocean iron-fertilized diatom bloom, *Nature*, 487: 313-319, 2012.

655 Tillmann, U., Hesse, K.-J., and Tillmann, A.: Large-scale parasitic infection of diatoms in the North Frisian Wadden Sea, *J. Sea Res.*, 42, 255–261, 1999.

Wolf, C., Frickenhaus, S., Kiliyas, E.S., Peeken, I., and Metfies, K.: Protist community composition in the Pacific sector of the Southern Ocean during austral summer 2010, *Polar Biol.* 37, 375-389, 2014.

660 Zhou, M., Zhu, Y., d'Ovidio, F., Park, Y.H., Durand, I., Kesternare, E., Sanial, E., Van-Beek, P., Quéguiner, B., Carlotti, F., and Blain, F.: Surface currents and upwelling in Kerguelen Plateau regions, this volume.

Zhu, F., Massana, R., Not, F., Marie, D., and Vaulot, D.: Mapping of picoeucaryotes in marine ecosystems with quantitative PCR of the 18S rRNA gene, *FEMS Microbiol. Ecol.*, 52, 79–92, 2005.

Table 1. Brief description of the stations. The depth of the mixed layer (ML) is based on a difference in sigma of 0.02 to the surface value. The mean ML (\pm SD) of all CTD casts performed during the occupation of the stations is given. Ze: the euphotic layer depth. For Chl *a* and major inorganic nutrients mean values \pm SD for the mixed layer.

| Station | Date (2011) | Latitude (°N) | Longitude (°E) | Station depth (m) | Sampling depths (m) | ML (m) | Ze (m) | Chl <i>a</i> ($\mu\text{g L}^{-1}$) ^a | NO ₃ + NO ₂ (μM) ^b | PO ₄ (μM) ^b | Si(OH) ₄ (μM) ^c | DFe (nM) ^d |
|---------|----------------|------------------|-------------------|----------------------|------------------------|--------------|-----------|---|---|---|---|--------------------------|
| R-2 | 26/10 | -50.359 | 66.717 | 2450 | 20, 60, 150, 300 | 105 \pm 15 | 92 | 0.25 \pm 0.08 | 26.0 \pm 0.2 | 1.83 \pm 0.03 | 12.3 \pm 0.3 | 0.08 \pm 0.07 |
| F-L | 07/11 | -48.505 | 74.614 | 2690 | 20, 65, 180, 300 | 38 \pm 7 | 28 | 4.00 \pm 1.58 | 20.5 \pm 1.9 | 1.06 \pm 0.21 | 7.7 \pm 0.8 | 0.22 \pm 0.06 |
| E-4W | 10/11 | -48.765 | 71.425 | 1398 | 30, 80, 150, 300 | 61 \pm 11 | 31 | 2.38 \pm 0.31 | 25.4 \pm 1.0 | 1.79 \pm 0.10 | 18.5 \pm 1.2 | 0.17 \pm 0.03 |
| A3-2 | 16/11 | -50.624 | 72.056 | 528 | 20, 80, 160, 300 | 153 \pm 15 | 38 | 2.03 \pm 0.33 | 26.2 \pm 0.4 | 1.78 \pm 0.03 | 18.9 \pm 0.5 | 0.16 \pm 0.03 |

^a Lasbleiz et al., (this volume)

670 ^b Blain et al., (this volume,b)

^c Closset et al., (this volume)

^d Qu  rou   et al., (this volume)

Table 2. Number of OTUs, the richness estimator (S_{chao1}), Simpson and Berger-Parker indices for each sample. **Nb of seqs before and after removing metazoan and single singletons sequences**

| Station | Depths (m) | Nb OTUs | Nb seqs before | Nb seqs after | S_{chao1} | Simpson (1-D) | Berger-Parker |
|---------|------------|---------|----------------|---------------|--------------------|---------------|---------------|
| R-2 | 20 | 157 | 5448 | 4714 | 198 | 0.95 | 0.18 |
| | 60 | 170 | 6346 | 1522 | 218 | 0.95 | 0.16 |
| | 150 | 233 | 4407 | 1562 | 390 | 0.97 | 0.13 |
| | 300 | 282 | 1098 | 950 | 409 | 0.99 | 0.05 |
| F-L | 20 | 186 | 5586 | 3028 | 253 | 0.76 | 0.48 |
| | 65 | 508 | 7305 | 5730 | 663 | 0.98 | 0.08 |
| | 180 | 265 | 7818 | 905 | 382 | 0.98 | 0.05 |
| | 300 | 284 | 10205 | 2026 | 383 | 0.83 | 0.40 |
| E-4W | 30 | 173 | 7151 | 6108 | 198 | 0.85 | 0.33 |
| | 80 | 209 | 10977 | 6674 | 236 | 0.92 | 0.23 |
| | 150 | 191 | 11989 | 5771 | 255 | 0.94 | 0.19 |
| | 300 | 97 | 3178 | 242 | 174 | 0.97 | 0.08 |
| A3-2 | 20 | 215 | 10666 | 1803 | 285 | 0.93 | 0.22 |
| | 80 | 200 | 3866 | 2118 | 273 | 0.98 | 0.08 |
| | 160 | 181 | 5986 | 2022 | 219 | 0.95 | 0.13 |
| | 300 | 330 | 11590 | 5662 | 385 | 0.94 | 0.23 |

Table 3: Higher-level taxonomic distribution of protistan OTUs defined at 97 % sequence similarity.

| Supergroup | Taxonomic groups | OTUs |
|-------------------|-------------------------|-------------|
| Alveolata | MALV-II | 339 |
| | Dinophyceae | 161 |
| | MALV-I | 101 |
| | Ciliophora | 60 |
| | MALV-III | 21 |
| | MALV-IV | 8 |
| | Apicomplexa | 3 |
| | MALV-V | 2 |
| | Perkinsea | 1 |
| Stramenopiles | MAST | 46 |
| | Bacillariophyceae | 37 |
| | Labyrinthulomycetes | 19 |
| | Bolidophyceae | 13 |
| | Pirsonia | 6 |
| | Dictyochophyceae | 4 |
| | Pelagophyceae | 3 |
| | Hyphochrytriaceae | 2 |
| | Oomyceta | 2 |
| | Chrysophyceae | 1 |
| | | |
| Hacrobia | Haptophyta | 20 |
| | Picobiliphyta | 15 |
| | Telonemia | 12 |
| | Centroheliozoa | 2 |
| | Cryptophyta | 1 |
| Opisthokonta | Fungi | 28 |
| | Choanoflagellida | 10 |
| Rhizaria | Radiolaria | 35 |
| | Cercozoa | 17 |
| Archaeplastida | Chlorophyta | 29 |
| Apusozoa | Hilomonadea | 1 |

710

715

720

725

730

32

| | | | | | | | | | | | | | | | | | | | |
|-----------------------|--------------------|-----------------------------|-----|------|------|-------|------|------|------|-------|------|------|------|-------|-------|-------|------|------|-------|
| Choanoflagellida (1%) | Otu1941 | StephanoecidaeGroupD sp. | 97 | 22.2 | 10.0 | 11.1 | 25.0 | 33.3 | 44.1 | 25.0 | | 83.3 | 4.0 | 21.5 | | 73.3 | 66.7 | 4.0 | 25.0 |
| | Otu1928 | Stephanoeca cauliculata | 99 | 57.5 | 72.7 | 55.6 | 25.0 | | 11.8 | 25.0 | | 16.7 | 1.0 | 1.8 | | 13.3 | 16.7 | 6.0 | 25.0 |
| | Otu1710 | StephanoecidaeGroupD sp. | 100 | 18.5 | 18.2 | | | | 23.5 | | 66.7 | | 5.0 | 43.8 | | 6.7 | 13.3 | | 25.0 |
| | Otu0960 | StephanoecidaeGroupH sp. | 90 | | | | | 66.7 | 2.9 | 25.0 | | | | 24.6 | | | | | |
| | Otu1959 | StephanoecidaeGroupD sp. | 95 | | | 11.1 | | | 11.8 | | | | | | | | 3.3 | | 25.0 |
| | Otu1706 | StephanoecidaeGroupD sp. | 93 | 1.9 | | | | | | 25.0 | 33.3 | | | | | 6.7 | | | |
| | Otu1905 | StephanoecidaeGroupH sp. | 91 | | | 22.2 | | | | | | | | | | | | | |
| | Otu1828 | StephanoecidaeGroupD sp. | 94 | | | | 5.0 | | | | | | | | | | | | |
| Radiolaria (4%) | Otu1699 | Spumellarida-Group-I sp. | 99 | | | | 67.9 | | 4.2 | 8.0 | 73.7 | | | | 18.8 | | | | 50.7 |
| | Otu1655 | Spumellarida-Group-I sp. | 100 | | | | | 16.7 | 1.1 | 4.0 | 13.9 | | | | | | | | 1.0 |
| | Otu1138 | Stylodictya sp. | 99 | | | | 1.9 | 25.0 | 42.1 | 16.0 | 0.4 | | | | 37.5 | 33.3 | | | 7.7 |
| | Otu1589 | Spumellarida-Group-I sp. | 100 | | | | | | | 4.0 | 6.7 | | | | | | | | 0.3 |
| | Otu0699 | Triastrum aurivillii | 95 | 25.0 | | | | | 1.1 | | | | 76.2 | | | | | | 1.7 |
| | Otu1856 | RAD-B-Group-IV sp. | 99 | | 50.0 | 1.9 | | 12.6 | 8.0 | 0.3 | | | 6.3 | 33.3 | | 66.7 | | 5.8 | |
| | Otu0036 | RAD-B-Group-IV sp. | 97 | | | 1.9 | | | 8.0 | | | | | | | | | | 7.5 |
| | Otu0686 | RAD-B-Group-II sp. | 100 | | | | | | | | | | | | | | | | 7.4 |
| Otu1654 | RAD-B-Group-II sp. | 99 | | | | 5.7 | | 2.1 | 4.0 | | | | | | | | | 5.8 | |
| Otu1349 | RAD-B-Group-IV sp. | 100 | | 75.0 | | | | 17.9 | 16.0 | | | 19.0 | | | | 100.0 | 33.3 | | |
| Cerczoa (1%) | Otu1449 | Protocystis iphodon | 100 | | | | | 2.6 | | | | | | 100.0 | 100.0 | | | | |
| | Otu1378 | Protaspa-lineage sp. | 98 | | | 100.0 | | 30.8 | 26.7 | | | 6.7 | | | | 33.3 | 28.6 | 33.3 | 100.0 |
| | Otu1257 | Ebria tripartita | 100 | | | | | 51.3 | 20.0 | | | | | | | | | | |
| | Otu0591 | Protaspa-lineage sp. | 99 | | | | | 2.6 | 6.7 | | | 46.7 | 13.3 | | | | 28.6 | | |
| | Otu0887 | TAGIRI1-lineage sp. | 98 | | | | | | | | | | 40.0 | | | | 28.6 | | |
| | Otu1806 | Protaspa sp. | 99 | | | 100.0 | | 10.3 | | | | | | | | 16.7 | | | |
| | Otu0881 | Cryothecomonas-lineage sp. | 99 | | | | | | | | | | 33.3 | | | | | | |
| | Otu0170 | Cryothecomonas-lineage sp. | 100 | | | | | | 6.7 | | | | | | | 33.3 | | 33.3 | |
| | Otu0742 | TAGIRI1-lineage sp. | 98 | | | | | | | | | | 13.3 | | | 16.7 | 14.3 | | |
| | Otu0201 | Cryothecomonas sp. | 100 | | | | | 20.0 | | | | | | | | | | | |
| | Otu1040 | Mataza-lineage sp. | 100 | | | | | 2.6 | | | | 13.3 | | | | | | | |
| | Otu1368 | Protaspa-lineage sp. | 100 | | | | | | 6.7 | | | 6.7 | | | | | | 33.3 | |
| | Otu0857 | Cryothecomonas sp. | 99 | | | | | | | | | 13.3 | | | | | | | |
| | Otu0941 | Marimonadida sp. | 92 | | | | | | | | | 13.3 | | | | | | | |
| Otu1624 | Endo4-lineage sp. | 99 | | | | | | | | 100.0 | | | | | | | | | |
| Chlorophyta (4%) | Otu1717 | Micromonas pusilla (RCC658) | 100 | 2.0 | 28.8 | 3.5 | | 25.4 | 38.6 | 68.4 | | 1.0 | 6.6 | 5.2 | | | 5.9 | 3.3 | |
| | Otu1962 | Pyramimonas gelidicola | 97 | 6.9 | 4.0 | 2.3 | | | 0.9 | | | 1.8 | 72.5 | 29.6 | | 17.4 | 2.9 | 53.3 | |
| | Otu1742 | Micromonas pusilla (RCC418) | 100 | 22.7 | 29.9 | 3.5 | | 13.6 | 21.6 | 5.3 | | 30.0 | 6.1 | | | | 3.3 | | |
| | Otu1918 | Bathycoccus prasinos | 100 | 12.8 | 26.6 | 34.5 | | 6.8 | 16.5 | | | 10.0 | 4.4 | 1.0 | | | 17.6 | | |
| | Otu1791 | Prasinoderma coloniale | 95 | 33.5 | 3.4 | | 5.0 | 8.5 | 2.5 | 15.8 | | | 0.9 | 3.8 | | | | 3.3 | |
| | Otu0166 | Pyramimonas disomata | 96 | 2.7 | 1.1 | | 5.0 | | | | | | | 33.3 | | | | 6.7 | |
| | Otu1766 | Pyramimonas olivacea | 99 | 2.2 | 1.7 | 1.4 | | 3.4 | 2.2 | | 1.0 | 5.5 | | 18.5 | | 47.8 | 14.8 | 6.7 | 48.3 |
| | Otu0940 | Pyramimonas sp. | 100 | | 0.6 | 0.7 | | 1.2 | 1.5 | | | 13.6 | 4.4 | 4.6 | | 13.4 | 41.2 | 1.0 | |
| | Otu1775 | Crustomastigaceae sp. | 99 | 0.2 | | | | 16.9 | 0.4 | 5.3 | | 10.0 | 1.3 | 3.6 | | | 17.6 | | 51.7 |
| | Otu0151 | Mamiella sp. | 100 | | | 0.7 | | 6.7 | 2.5 | | | 6.3 | 1.7 | | | 4.3 | | | |

Table 5: Results of SIMPER (similarity percentages) following the Bray-Curtis cluster analysis (Fig. 6a). Forty-one OTUs contributing for at least 1% of the similarity of each cluster are listed in this table. In parenthesis, the mean of Bray-Curtis similarity is given for each cluster.

| OTUs | Taxonomic Groups | Putative Taxonomic Affiliation | Cluster (i) (43.8%) | Cluster (ii) (51.8%) | Cluster (iii) (47.6%) | Cluster (iv) (20.7%) |
|--------------|-------------------|------------------------------------|------------------------|-------------------------|--------------------------|-------------------------|
| Otu1951 | Dinophyceae | <i>Gyrodinium spirale</i> | 5.1 | 17 | 33.5 | 1.3 |
| Otu1914 | | <i>Gyrodinium rubrum</i> | | 5.2 | 17.9 | |
| Otu1967 | | <i>Gymnodinium</i> sp. | 8.8 | 3.5 | 2.6 | 4.9 |
| Otu1898 | | <i>Karlodinium micrum</i> | | 5.9 | 1.9 | 1.3 |
| Otu1770 | | <i>Warnowia</i> sp. | | 3 | 1.5 | 4.1 |
| Otu1016 | | <i>Gyrodinium rubrum</i> | 1.3 | 3.3 | | 3.6 |
| Otu1763 | | <i>Dinophyceae</i> sp. | 1.5 | 3.7 | | 1.5 |
| Otu1808 | | <i>Warnowia</i> sp. | 1.8 | 3.6 | | |
| Otu1953 | | <i>Peridinium tyrrhenicum</i> | | 1.4 | | 1.8 |
| Otu1871 | | <i>Gymnodinium catenatum</i> | | | | 1.7 |
| Otu1816 | | <i>Karlodinium micrum</i> | | | | 1.3 |
| Otu1722 | | <i>Gymnodinium</i> sp. | | 1.3 | | |
| Otu1454 | | <i>Islandinium minutum</i> | | 1.3 | | |
| Otu1793 | | <i>Amphidinium semilunatum</i> | | | 1.1 | |
| Total | | | 18.4 | 49.2 | 58.6 | 21.5 |
| Otu1653 | MALV-I | Dino-Group-I-Clade-1 sp. | 1.3 | 2.2 | 2.1 | 6 |
| Otu1912 | | Dino-Group-I-Clade-1 sp. | | 1.9 | 3.2 | 1.4 |
| Otu1292 | | Dino-Group-I-Clade-5 sp. | | 2.9 | | 1.6 |
| Otu1285 | | Dino-Group-I-Clade-1 sp. | 1.3 | | 2.6 | |
| Otu1790 | | Dino-Group-I-Clade-4 sp. | 1.2 | | 1.4 | |
| Otu1393 | | Dino-Group-I-Clade-1 sp. | 1 | | | |
| Total | | | 4.7 | 6.9 | 9.2 | 9 |
| Otu1211 | MALV-2 | Dino-Group-II-Clade-10 sp. | | | | 3.7 |
| Otu1116 | | Dino-Group-II-Clade-10 sp. | | | | 3.2 |
| Otu1663 | | Dino-Group-II-Clade-7 sp. | | | | 2 |
| Otu1613 | | Dino-Group-II-Clade-7 sp. | | | | 2 |
| Otu1513 | | Dino-Group-II-Clade-6 sp. | | | | 1.1 |
| Otu1183 | | Dino-Group-II-Clade-7 sp. | | | | 1.1 |
| Total | | | | | | 13.1 |
| Otu1799 | Ciliophora | <i>Strombidium biarmatum</i> | 1.2 | | | |
| Otu1447 | Bacillariophyceae | <i>Thalassiosira tenera</i> | | 6.4 | 2 | 1.9 |
| Otu0978 | | <i>Thalassiosira delicatula</i> | | 2.7 | | |
| Total | | | | 9.1 | 2 | 1.9 |
| Otu1932 | Pelagophyceae | <i>Aureococcus anophagefferens</i> | 4.7 | | | |
| Otu1762 | MAST | MAST-1B sp. | 1.3 | | | |
| Otu1923 | | MAST-1C sp. | 1.1 | | | |
| Total | | | 2.4 | | | |
| Otu1717 | Chlorophyta | <i>Micromonas pusilla</i> | 4 | | | |
| Otu1918 | | <i>Bathycoccus prasinos</i> | 3.8 | | | |
| Otu1742 | | <i>Micromonas pusilla</i> | 3.8 | | | |
| Total | | | 11.5 | | | |
| Otu1782 | Haptophyta | <i>Phaeocystis antarctica</i> | 18.3 | 2.7 | 5.2 | 1.3 |
| Otu1884 | | <i>Chrysochromulina strobilus</i> | 4.5 | | | |
| Otu1907 | | <i>Chrysochromulina</i> sp. | 2.7 | | 1.3 | |
| Total | | | 25.5 | 2.7 | 6.4 | 1.3 |
| Otu1863 | Fungi | <i>Malassezia restricta</i> | 1.3 | | | |
| Otu1699 | Radiolaria | <i>Spumellarida</i> sp. | | | | 6.2 |
| Otu1138 | | <i>Stylodictya</i> sp. | | | | 1 |
| Total | | | | | | 7.2 |

770 Fig. 1. Bathymetry of the study area and location of the sampled stations (a), and Chl *a* (color scale), surface velocity fields (arrows), the polar front (PF, black line) (b). The chlorophyll content represented on the map corresponds to the last week of the KEOPS2 and the cross indicates the position of the North-South and East-West transects sampled to provide an overview of the blooms. Map is courtesy of Y. Park and colleagues.

775 Fig. 2. Profiles of Temperature (a), Chl *a* as derived from *in vivo* Fluorescence (b) and Salinity (c) for each of the four sampling stations.

Fig. 3. Overall diversity of major high-level taxonomic groups and number of OTUs indicated in parenthesis at each station.

780

Fig. 4. Relative abundance of major high-level taxonomic groups at each station and depth.

Fig. 5. Venn diagrams representing the number of OTUs shared between the different stations.

785 Fig. 6. Cluster diagram for the 16 samples constructed from a Bray-Curtis similarity matrix of square-root-transformed OTU abundances. Asterisks at nodes in the dendrogram indicate significant differences between bifurcations ($P < 0.05$) (a). Nonmetric multidimensional (nMDS) scaling plots in two dimensions constructed from a Bray-Curtis similarity matrix. Bray-Curtis similarity contours are 15% (solid lines) and 40% (dashed lines) (b).

790

Suppl. fig 1: Rarefaction curves representing the numbers of OTUs versus the number of reads. The OTUs were determined using the program Mothur, with a cutoff value set to 0.03 (OTUs were grouped when their level of sequence similarity was $\geq 97\%$) for the analysis

795




Figure 1






Figure 2


R-2




F-L



E-4W



A3-2






Figure 4




Figure 5





Figure 6