

This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Spring bloom community change modifies carbon pathways and C : N : P : Chl *a* stoichiometry of coastal material fluxes

K. Spilling^{1,2}, A. Kremp¹, R. Klais³, K. Olli³, and T. Tamminen¹

¹Marine Research Centre, Finnish Environment Institute, P.O. Box 140, 00251 Helsinki, Finland

²Tvärminne Zoological Station, University of Helsinki, 10900 Hanko, Finland

³Institute of Ecology and Earth Sciences, University of Tartu, Lai 40, 51005 Tartu, Estonia

Received: 3 June 2014 – Accepted: 8 June 2014 – Published: 6 August 2014

Correspondence to: K. Spilling (kristian.spilling@environment.fi)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Coastal carbon pathways and stoichiometry

K. Spilling et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Abstract

Diatoms and dinoflagellates are major bloom-forming phytoplankton groups competing for resources in the oceans and coastal seas. Recent evidence suggests that their competition is significantly affected by climatic factors under ongoing change, modifying especially the conditions for cold-water, spring bloom communities in temperate and arctic regions. We investigated the effects of phytoplankton community composition on spring bloom carbon flows and nutrient stoichiometry in multi-year mesocosm experiments. Comparison of differing communities showed that community structure significantly affected C accumulation parameters, with highest particulate organic carbon (POC) build-up and dissolved organic carbon (DOC) release in diatom-dominated communities. In terms of inorganic nutrient drawdown and bloom accumulation phase, the dominating groups behaved as functional surrogates. Dominance patterns, however, significantly affected C : N : P : Chl *a* ratios over the whole bloom event: when diatoms were dominant, these ratios increased compared to dinoflagellate dominance or mixed communities. Diatom-dominated communities sequestered carbon up to 3.6-fold higher than the expectation based on the Redfield ratio, and 2-fold higher compared to dinoflagellate dominance. To our knowledge, this is the first experimental report of consequences of climatically driven shifts in phytoplankton dominance patterns for carbon sequestration and related biogeochemical cycles in coastal seas. Our results also highlight the need for remote sensing technologies with taxonomical resolution, as the C : Chl *a* ratio was strongly dependent on community composition and bloom stage. Climate-driven changes in phytoplankton dominance patterns will have far-reaching consequences for major biogeochemical cycles and need to be considered in climate change scenarios for marine systems.

Coastal carbon pathways and stoichiometry

K. Spilling et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



1 Introduction

Coastal seas and shelf areas (<200 m deep) constitute approximately 5% of the ocean, but are among the most vital marine biotopes, both from an ecological and from a socio-economical perspective. They connect terrestrial, atmospheric, and marine biogeochemical cycles, and it has been estimated that ~ 12% of the marine primary production and ~ 86% of the total carbon burial in the ocean takes place in coastal regions (Dunne et al., 2007). Coastal seas also play a pivotal role in trophic transfer of organic carbon from primary producers through the food web, and include some of the richest fisheries in the world. At the same time, these areas are the most affected by direct and indirect anthropogenic pressures and are highly vulnerable to projected global change (Halpern et al., 2008). Multiple drivers of the marine food web, such as temperature, UV irradiation, $p\text{CO}_2$, and runoff of nutrients and freshwater, are affecting the ecosystem on different levels. One of the key issues for predicting how global change will affect coastal marine environments is to identify population dynamics and feedback loops under a changing environment (Harley et al., 2006; Eggers et al., 2014).

Temperate aquatic systems are characterized by high productivity, especially of new production, as opposed to recycled production (Dugdale and Goering, 1967). Their production is highly seasonal, and the annual spring bloom represents the most significant production phase. High initial concentrations of inorganic nutrients, increasing solar radiation, and emerging stratification of water layers trigger the onset of photosynthetic production. Several bloom-forming phytoplankton groups compete for resources in marine environments during spring, the most conspicuous being diatoms, dinoflagellates and prymnesiophytes. Recent evidence from both coastal and offshore environments shows decadal shifts in the relative proportions of diatoms and dinoflagellates at different seasons, and suggests that their competition is significantly affected by climatic factors under ongoing change (Leterme et al., 2005; Hinder et al., 2012), modifying especially the spring bloom conditions of temperate and arctic regions; mild winters and more storms have been shown to favor dinoflagellates (Klais et al., 2013), and

BGD

11, 11863–11902, 2014

Coastal carbon pathways and stoichiometry

K. Spilling et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Coastal carbon pathways and stoichiometry

K. Spilling et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



also changes in thermal stratification patterns and freshwater runoff are thought to affect phytoplankton community composition (Smayda and Reynolds, 2001). The extensive arctic shelf seas and marginal ice zones are globally among the most susceptible biotopes for climate change, and their changing production preconditions will potentially have a great impact on global carbon budgets and interconnected biogeochemical cycles. The consequences of climate-driven phytoplankton community change represent therefore urgent challenges for reliable climate change scenarios.

The physiology and morphology of different phytoplankton species and groups vary considerably, with direct impacts on ecosystem-wide nutrient cycling and cascading food web effects. Differences in species-specific traits like growth rate, nutrient affinities and biochemical composition, cell size, motility, and life cycle strategies govern the outcome of resource competition, and therefore the community composition in a set of environmental conditions. They also directly affect system-level carbon sequestration, stoichiometry of material flows, and the export of organic carbon to the sea floor. Several functional aspects of algal physiology are thus relevant for large-scale biogeochemical cycles, and their incorporation in trait-based models of phytoplankton production (Litchman and Klausmeier, 2008; Litchman et al., 2010) would significantly enhance the predictive potential of marine biogeochemical models under climate change.

Among the temperate coastal seas projected to change most rapidly is the Baltic Sea, due to its close interaction with the intensively modified catchment, the predicted changes in annual precipitation patterns over northern Europe, its reduced alkalinity, and heavy fishing pressure (Niiranen et al., 2013). In the Baltic Sea, cold-water dinoflagellates and diatoms have been considered functional surrogates during the spring bloom, as both effectively deplete the wintertime inorganic nutrient concentrations (Tamminen, 1995; Kremp et al., 2008), and the bloom terminates in most basins once nitrate has been consumed below analytical detection limits (Tamminen and Andersen, 2007). However, there are obvious differences with respect to life cycle strategies and sedimentation patterns of the competing phytoplankton groups. In general, diatoms sink quickly to the sea floor once nutrients are depleted, and it has been shown

that the fraction of the population forming resting spores is highly species-specific (Rynearson et al., 2013). Dinoflagellates, on the other hand, lyse before reaching the sediment, or alternatively go through a life cycle transformation producing decomposition-resistant resting cysts (Heiskanen, 1998).

The differences in sedimentation patterns have a large impact on decomposition of the bloom biomass in sediments, with consequences on oxygen consumption and release of phosphorus (Spilling and Lindström, 2008). This indicates strong cascading effects of bloom community composition on benthic food webs and material cycles. Although grazing pressure is relatively low during the spring bloom period in the Baltic Sea (Lignell et al., 1993), phytoplankton species composition has been shown to affect also the planktonic grazer communities because of species-specific differences in food quality for the emerging copepod populations (Vehmaa et al., 2011). Therefore, the cascading effects of bloom composition are potentially pervasive within the whole ecosystem.

In this study, we investigate the effects of phytoplankton community composition on stoichiometry of planktonic biogeochemical processes, in a coastal model system displaying ongoing, climate-driven community change (the Baltic Sea; Klais et al., 2011). We hypothesize that a change in phytoplankton community composition (here, diatom vs. dinoflagellate dominance) will significantly modify the carbon budget and stoichiometric composition of the seston. The data originate from coastal mesocosm experiments with nutrient enrichments performed in 5 consecutive years, with multi-week time series of phytoplankton species composition, primary production and nutrient fractions in a total of 40 mesocosms. The experiments displayed natural phytoplankton communities of interannually highly variable relative contributions of diatoms and cold-water dinoflagellates (< 10 % to > 90 % of either group) to the total bloom biomass.

BGD

11, 11863–11902, 2014

Coastal carbon pathways and stoichiometry

K. Spilling et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2 Materials and methods

2.1 Experimental set-up

The mesocosm experiments were conducted in spring 2004 to 2008 under laboratory conditions at the Tvärminne Zoological Station, University of Helsinki. The experimental set-up consisted of a control unit of natural sea water, and three nutrient manipulation treatments: a combined nitrate (N) and phosphate (P) addition; dissolved silicate (DSi) addition; and combined N, P, and DSi addition. On top of the nutrient manipulations, different treatments were added. In 2004, 2006, and 2008, two different light environments were used, in a full factorial 2^3 design; in 2005 and 2007, two cultured diatoms typical for the spring bloom in the area: *Thalassiosira levanderi* ($\sim 10 \mu\text{m}$ diameter) and *T. baltica* ($20\text{--}30 \mu\text{m}$ diameter), were added to the natural communities. A summary of the experimental design and initial conditions in the experimental units is given in Table 1.

For each experiment, containers were filled with natural surface water and pre-screened with a $200 \mu\text{m}$ mesh-size net to remove metazooplankton. Water was collected during ice break-up from the ice edge near the Storfjärden monitoring station at the SW coast of Finland ($59^\circ 51' \text{N}$; $23^\circ 13' \text{E}$). In 2004, white plastic (PE) 80 L barrels were used as experimental units, whereas for the years 2005 to 2008, 25 L transparent polycarbonate carboys were used.

Ice break-up typically coincides with the initiation of the annual spring bloom in the area (Niemi, 1975), and the captured phytoplankton community was assumed to represent the seed community for the spring bloom. The timing of ice break-up varied between years, and accordingly the experiments were started on different dates in subsequent years, depending on the ice situation.

After filling containers on the ice, they were immediately brought to the laboratory; the water was divided into mesocosm carboys, and placed into a walk-in incubator set to 2°C . The mesocosms were illuminated by daylight spectrum, fluorescent tubes (Philips TLD-95) at a 12 : 12 L/D cycle, corresponding to the ambient light cycle. Differ-

BGD

11, 11863–11902, 2014

Coastal carbon pathways and stoichiometry

K. Spilling et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ent irradiance was used for different treatments (Table 1). Pre-filtered ($0.2\ \mu\text{m}$) air was bubbled into the mesocosms to keep a low level of turbulence.

Natural nutrient conditions were manipulated by additions of $\text{NO}_3\text{-N}$, $\text{PO}_4\text{-P}$ (N + P treatment), and/or $\text{SiO}_2\text{-Si}$ (DSi treatment), with similar nutrient manipulations carried out each year. Nutrient additions (see Table 1 for the initial NO_3 concentrations) were targeted at approximately doubling the typical wintertime concentrations in the area, while maintaining a balanced Redfield ratio (a molar N : P ratio of 16). Irradiance was adjusted to $20\ \mu\text{mol photons m}^{-2}\ \text{s}^{-1}$ for the low light treatment (LL), which was applied in the 2004, 2006, and 2008 experiments; and to $90\ \mu\text{mol photons m}^{-2}\ \text{s}^{-1}$ for the high light treatment (HL). In 2005 and 2007, all treatments received $90\ \mu\text{mol photons m}^{-2}\ \text{s}^{-1}$.

The successive years represented different initial conditions, due to differences in ice break-up and other meteorological history of the winter-early spring season. The ambient nutrient concentrations of each experiment thus represented varying phases of the early bloom period despite similar experimental additions, between full wintertime levels and the spring depletion period. The initial, natural phytoplankton community varied from year to year.

2.2 Sampling protocol and measurements

Mesocosms were sampled for Chlorophyll *a* (Chl *a*), nutrient and phytoplankton immediately after the addition of nutrients on day 0, and subsequently every 2 to 3 days. The duration of the experiment differed between years, depending on how fast nutrients were exhausted. Prior to sampling, which took place at the beginning of the daily light period; the contents of the mesocosms were stirred with a polycarbonate rod to ensure an even distribution of phytoplankton and other particulate matter. The samples thus represent bloom development without sedimentation losses.

Samples for dissolved and particulate nutrients and Chl *a* were processed immediately. Nutrient concentrations ($\text{NO}_3\text{-N}$, $\text{PO}_4\text{-P}$ and DSi) were determined manually in duplicate according to the standard colorimetric methods (Grasshoff et al., 1983). Dis-

Coastal carbon pathways and stoichiometry

K. Spilling et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



solved organic carbon (DOC) concentrations were measured by the High Temperature Catalytic Oxidation (HTCO) method using a Shimadzu TOC-V CPH carbon and nitrogen analyzer. Sub-samples (< 0.45 μm Supor Acrodisc PES filter, Gelman Sciences) were acidified to pH 2.5 with 2 M HCl and stored in darkness at room temperature.

5 The 20 mL glass ampoules were stored for 4 to 6 months, before determining the DOC concentration according to Sharp et al. (1993).

For determining Chlorophyll *a* (Chl *a*), 50 mL duplicate samples were filtered onto glass fiber filters (Whatman GF/F) and extracted in 10 mL of 94 % ethanol for 24 h in the dark at room temperature. Chl *a* was measured on a Shimadzu RFPC-5001
10 fluorometer, calibrated with pure Chl *a* (Sigma). Duplicate filters were also prepared for determination of particular organic carbon (POC), nitrogen (PON) and phosphorus (POP). For all samples acid washed, pre-combusted GF/F filters were used. The filters were allowed to dry and stored at room temperature (20 °C) until nutrient determination. POC and PON were measured from the same filter with a mass spectrometer (Europa Scientific). POP was determined according to Solórzano and Sharp (1980).

Phytoplankton samples were preserved with acid Lugol's solution. Prior to microscopic analysis, volumes of 50 and 25 mL were set up for concentration in Utermöhl chambers and allowed to settle for at least 24 h. Diatoms and dinoflagellates, identified to species or genus level, were counted with an inverted light microscope (Leica DM
20 IRB Wetzlar, Germany). Cells were grouped into two size categories (> and < 10 μm), which were counted separately at $\times 200$ and $\times 787$ magnification. At least half of the chamber bottom was screened when cell densities were low, otherwise 400 cells were counted, if possible, for each category. Cell dimensions of diatoms and dinoflagellates were measured on 25 randomly selected cells of each species, and biovolumes were calculated using formulas given for standard geometric shapes of phytoplankton taxa (Sun and Liu, 2003). Biovolume values were converted to carbon according to the recommendations of Menden-Deuer and Lessard (2000).

25 Radiolabeled ^{14}C was used for determining the total primary production, and this was determined on all sampling days. An activity of 0.15 kBq was added to 10 mL

BGD

11, 11863–11902, 2014

Coastal carbon pathways and stoichiometry

K. Spilling et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Coastal carbon pathways and stoichiometry

K. Spilling et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



of sample and incubations carried out in the same light and temperature conditions as for the mesocosms. After an incubation period of 3 h, 4 mL sample was extracted, 150 μL of 37 % formaldehyde was added to fix the sample, and 100 μL of 1 M HCl was added in ventilation cupboard to remove unassimilated (inorganic) ^{14}C isotope. The samples were left with the lid open for 24 h before 7 mL of liquid scintillation cocktail (Hi Safe) was added. The radioactivity was measured using a liquid scintillation counter (PerkinElmer Inc., Wallac Winspectral 1414). The amount of total dissolved inorganic carbon (DIC) was measured with a high-temperature combustion IR carbon analyzer (Unicarbo, Electro Dynamo, Finland). Primary production was calculated from uptake of ^{14}C knowing the total amount of added isotope and total DIC. Due to the relatively short incubation period, measured primary production was assumed to represent gross primary production (Sakshaug et al., 1997).

2.3 Data treatment

The development of dissolved inorganic nutrients and particulate organic carbon (POC), nitrogen (PON), phosphorus (POP), Chlorophyll *a* (Chl *a*), and dissolved organic carbon (DOC) were organized as a function of time (e.g. Fig. 1). Background levels of refractory DOC are very high in the Baltic Sea (Hoikkala, 2012), so the start concentration was subtracted from all values to express DOC change during the experiment. The phytoplankton development was divided into two growth stages: exponential and stationary growth phase. The exponential growth phase was defined from the start of the experiment until the primary production peak per volume (i.e. not normalized to biomass); stationary growth phase was defined as the period after this point until the end of the experiment.

The community growth rate (d^{-1}) was determined during the exponential growth phase for the biomass-related parameters (μ_{POC} , μ_{Chla}) by linear regression of the natural log transformed data. The exponential growth of DOC (χ_{DOC}) was done in the same way. During stationary growth phase a linear regression (without log transformation) was fitted to the data parameters in order to find the rate of change.

Coastal carbon pathways and stoichiometry

K. Spilling et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Primary production was modeled from the ^{14}C incubations, assuming the measured production to represent the whole light period (12 h d^{-1}). Sampling did not take place every day, and we estimated the carbon fixation between sampling days by linear regression. A simple model was created, summing the gross carbon fixation for each day, and this was termed total gross production (TGP). This accumulated gross primary production would be the theoretical development of POC without any loss processes.

The growth rate of TGP (μ_{TGP}) was calculated similarly to the other biomass-related parameters described above, and carbon assimilation efficiency (CAE) was calculated from the ratio between the measured carbon accumulation and total gross production:

$$\text{CAE} = \mu_{\text{POC}} / \mu_{\text{TGP}} \quad (1)$$

Carbon loss rate (CLR) was calculated as fraction of TGP not entering the POC pool:

$$\text{CLR} = 1 - \text{CAE} \quad (2)$$

Respiration (RES) was calculated as the part of the loss rate not released as DOC, assuming that all carbon not adding to the POC or DOC pools was used for respiration.

$$\text{RES} = \text{CLR} - (\chi_{\text{DOC}} / \mu_{\text{TGP}}) \quad (3)$$

The CAE, CLR and RES parameters were also calculated for the stationary growth phases with the difference that the rate of change where used instead of growth rate, e.g. Δ_{POC} instead of μ_{POC} .

Phytoplankton community composition data were used for calculating the proportion of dinoflagellates and diatoms of the total community biomass. Species evenness (Shannon equitability, H_E) was calculated as follows:

$$H_E = -[p_i \cdot \ln(p_i/S)] \quad (4)$$

where p_i is the proportion of i th species biovolume from total biovolume of the sample, and S is the number of species present in the sample.

Coastal carbon pathways and stoichiometry

K. Spilling et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ANOVA on ranks were used to check for statistical significance ($\alpha = 0.05$) between different phytoplankton community composition for the different carbon budget parameters. “On ranks” were used because of a low normal distribution score for several parameters using the Anderson-Darling test ($A^2 > 1$ and $p < 0.05$). The ANOVA and regression analysis were carried out in SigmaPlot (SPSS).

3 Results

3.1 Phytoplankton community

The initial phytoplankton community composition varied from year to year, with relative proportions of the total biomass ranging from > 90 % diatoms to > 90 % dinoflagellates. In general, there were more species of diatoms present in the mesocosms compared to dinoflagellates. The most abundant diatoms were *Thalassiosira baltica*, *T. levan-deri*, *Chaetoceros wighamii*, *Skeletonema marinoi* and *Achnanthes taeniata*, and two dinoflagellates were dominating: *Biecheleria baltica* and *Peridiniella catenata*. Species evenness was highest in a mixed community, when dinoflagellates constituted 20–70 % (i.e. 30–80 % diatoms) of the total population (Fig. 2). The effect on evenness was less pronounced when diatoms dominated. During diatom dominance, several species were represented whereas during dinoflagellate dominance, only one out of the two species was accounted for most of the biomass (*B. baltica* in 2004 and *P. catenata* in 2007).

The dominance of either diatoms or dinoflagellates were almost complete being at > 90 % of the total phytoplankton biomass during the experiments with the exception of 2008 (mixed community) when Chrysophytes made up 10–20 % of the biomass.

3.2 Carbon budget

The community carbon growth rate (μ_{POC}) was clearly affected by the light conditions, but not by the community composition. The average μ_{POC} under the low and

Coastal carbon pathways and stoichiometry

K. Spilling et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



high light conditions (20 and $90 \mu\text{mol photons m}^{-2} \text{s}^{-1}$) were $0.08 (\text{d}^{-1}) \pm 0.01$ (SD) and $0.15 (\text{d}^{-1}) \pm 0.02$ (SD) respectively. There was no significant difference in μ_{POC} between different community compositions (Fig. 3a). However, growth rate calculated from Chl *a* ($\mu_{\text{Chl}a}$) was highest in diatom dominated communities and decreased linearly ($p < 0.0001$) with increasing dinoflagellate proportion (Fig. 3a). The carbon assimilation efficiency (CAE) was positively correlated ($p = 0.04$) with the growth rate (Fig. 3b), indicating lower loss rates with increasing growth rate.

During exponential growth, there was no apparent difference between phytoplankton communities in gross primary production (Fig. 4, Table 2). There was, however, an effect on the CAE and respiration (RES). The mixed community had on average a $\sim 30\%$ higher CAE and $\sim 75\%$ lower RES compared to the situations when dinoflagellates or diatoms dominated (Fig. 4, Tables 2 and 3).

In the stationary growth phase, the community composition clearly had an effect on the carbon budget (Fig. 4, Tables 2 and 4). When diatoms constituted $> 80\%$ of the population, there was on average 3–7 times higher buildup of particulate organic carbon (POC), ~ 2 fold higher total gross production (TGP), and ~ 2 -fold higher release of dissolved organic carbon (DOC) compared to the situations with mixed or dinoflagellate dominated communities (Fig. 5, Tables 2 and 4). The diatoms also had the highest CAE during stationary growth, but with no statistical difference with a mixed community (Table 4).

3.2.1 Stoichiometry

The phytoplankton community clearly affected the stoichiometry of the seston, with C : N and C : P ratio being higher during diatom dominance (Fig. 6). The drawdown of inorganic N and P was close to 100% and stayed stable after the onset of stationary growth phase (e.g. Fig. 1a). There was a significant, negative correlation between C : N and C : P ratios with increasing dinoflagellate proportion during both exponential and stationary growth phase ($p \leq 0.03$). The C : N ratio was a factor of 1.2–1.7 times higher

than the Redfield ratio during exponential growth phase, and increased to 1.7–3.6 times higher than the Redfield ratio in all communities during stationary growth phase.

The C : P ratio was up to 1.4 times higher than the Redfield ratio during exponential growth in diatom dominated communities; dinoflagellate dominated communities were approximately on par with the Redfield ratio. During stationary growth phase the C : P ratio increased to 1.4–2.8 times higher than the Redfield ratio.

The N : P ratio was for most samples fell below the Redfield ratio of 16 and did not vary between communities during exponential growth ($p = 0.23$). After nutrients had been depleted, however, there was a negative correlation of N : P ratio with increasing dinoflagellate proportion ($p < 0.001$) (Fig. 6).

The C : Chl *a* ratio was clearly affected by the phytoplankton community composition and growth phase (Fig. 7), and there was on average a trend of decreasing ratios during exponential growth followed by increasing ratios during stationary growth phase. At the start of the experiment, the average C : Chl *a* ratios (comparing only high light treatments) were 477 ± 338 (SD), 84 ± 51 (SD) and 55 ± 31 (SD) g C (g Chl *a*)⁻¹ for diatom, mixed, and dinoflagellate-dominated communities, respectively. The initial decrease in the C : Chl *a* ratio was most rapid during diatom dominance (95 ± 74 (SD) at the primary production peak), whereas there was less change for mixed communities or during dinoflagellate dominance (60 ± 60 (SD) and 45 ± 17 (SD) respectively at the primary production peak). The C : Chl *a* ratio started to increase again during the stationary growth phase and were at the end of the experiment 208 ± 62 (SD), 218 ± 155 (SD) and 387 ± 49 (SD) for diatom, mixed and dinoflagellate dominated communities respectively.

BGD

11, 11863–11902, 2014

Coastal carbon pathways and stoichiometry

K. Spilling et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4 Discussion

4.1 Natural mixed communities as an experimental system

Differences in the physiology and cellular composition of diatoms and dinoflagellates have been recurrently established in monocultures (Chan, 1980; Banse, 1982; Menden-Deuer and Lessard, 2000), with the general conclusion that diatoms show higher maximum growth rates, higher photosynthetic rates per unit carbon, and lower C:Chl *a* ratios compared to dinoflagellates. The conclusions are based on monoculture growth under saturating light and nutrient abundance, or on continuous monocultures with established light or nutrient limitation, with the implicit or explicit assumption that the empirically derived traits can be utilized in modeling the performance of natural planktonic populations or communities (Cloern et al., 1995; Sarthou et al., 2005). Other angles to the phylogenetic-metabolic differences have been to address the evolutionary inheritance of elemental stoichiometry between phyla and superfamilies (Quigg et al., 2003), or to mechanistically model the stoichiometry of the nested biochemical processes underlying primary production of unicellular algae (Geider et al., 1998; Falkowski, 2000; Pahlow, 2005).

Our experiment series departs from these approaches by empirically studying the carbon flow and community stoichiometry over the full duration of natural, mixed community spring bloom events. During a multispecies bloom, the abiotic conditions and species interactions go through a continuous transformation, inducing transitory physiological acclimation responses and changes in competitive advantage between species. This seriously complicates prediction of bloom development with species-specific properties originating from growth in controlled, artificial monoculture conditions (Sathyendranath et al., 2009; Mateus et al., 2012). We used standardized, representative environmental conditions (light, temperature, nutrient supplies) and exclusion of advective and sedimentation flows, to specifically address the net effects of variable diatom-to-dinoflagellate proportions of the bloom community on modification of coastal biogeochemistry.

BGD

11, 11863–11902, 2014

Coastal carbon pathways and stoichiometry

K. Spilling et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Coastal carbon pathways and stoichiometry

K. Spilling et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Our experimental setup eliminated sinking losses that affect the overall bloom dynamics in open natural systems. Diatoms, in particular, are known to aggregate and sink efficiently out of the photic layer after bloom culmination (Kjørboe et al., 1990; Underwood et al., 2004). Our results therefore represent an upper limit for bloom C draw-down. However, the stoichiometric differences between different communities evolved fast after the bloom peak, and the variable physical forcing in coastal seas include changes in mixing of the surface layer, resuspension, and lateral transport, which counteract permanent sedimentation of fresh biogenic material, prolonging the stationary phase of bloom communities.

4.2 Dominance patterns in experimental communities

The 5 initial communities represented the natural variability of phytoplankton in the respective years, as mesocosm communities developed from natural inocula. Inter-annual variability in community composition was considerable: years of dinoflagellate dominance alternated with years of diatom dominance or evenly mixed communities. The experimental treatments of light, nutrient supplies and community structure amplified or further diversified the dominance patterns of the natural inoculum communities. This provided a wide range of dominance conditions in the altogether 40 mesocosms over multi-week bloom events, thus representing an ideal semi-natural experimental system for community-level comparisons.

In the coastal study area, pre-bloom and bloom period weather patterns have been found to be significantly related to high dinoflagellate proportions during spring (Klais et al., 2013). Klais et al. (2013) reported that mild winters with thin ice cover more storms favored dinoflagellates, suggesting that changing climate conditions are likely to drive the increasing frequency of coastal dinoflagellate-dominated spring blooms. Recent biodiversity shifts in offshore phytoplankton communities have been repeatedly linked to changing climate conditions (Reid et al., 1998; Hinder et al., 2012), modifying hydrographic properties of the water column, and thus selecting for specifically adapted taxonomic groups, most notably dinoflagellates (Hallegraeff, 2010).

Coastal carbon pathways and stoichiometry

K. Spilling et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



While diatom dominance was in most cases caused by several co-occurring diatom species, the dinoflagellate blooms in the mesocosms consisted of a single species – *Biecheleria baltica* (formerly known as *Woloszynskia halophila*) or *Peridiniella catenata*. This seems to be a general phenomenon in a wide range of marine habitats: diatoms behave as guild members sharing the habitat, whereas dinoflagellates usually follow a taxonomical hierarchical pathway towards domination of one species (Smayda and Reynolds, 2001). Diatoms are, in general, tolerant to habitat diversity and are adapted to habitats with several ecological niches, whereas dinoflagellates often are habitat specialists where typically the best adapted species outcompete the rest (Smayda and Reynolds, 2001).

In dinoflagellate-dominated blooms, the respective species already constituted a major fraction of the initial community, by far outnumbering any other phytoplankton species, and were thus able to maintain dominance under several experimental treatments despite their relatively low species-specific growth rates (Kremp et al., 2008). Dinoflagellates have been shown to possess compensatory strategies to compete with fast growing phytoplankton groups, such as allelopathy, mixotrophy, and internal nutrient storages (Legrand and Carlsson, 1998; Collos et al., 2004; Tillmann et al., 2008). *B. baltica* has recently been confirmed to effectively suppress growth of co-occurring diatoms by excretion of allelochemicals (Suikkanen et al., 2011), and utilization of residual P has been suggested to facilitate sustained growth of *B. baltica* in the 2004 mesocosms (Kremp et al., 2008).

4.3 Carbon production and losses during developing and late bloom stages

There were no significant differences between communities of diatom or dinoflagellate dominance in carbon-based growth rates (μ_{POC} or μ_{TGP}) during the exponential bloom phase. This is somewhat counterintuitive, taking into account the general conclusions from monoculture studies, and previous evidence that Baltic Sea dinoflagellates exhibit lower growth rates in mixed communities than the competing diatoms (Kremp et al., 2008). It should be noted that the carbon budgets are cumulative for the whole ex-

ponential phase, including also the variable delay periods from the experiment onset. Also, we are dealing with natural, mixed communities even in both “dominance” categories. Despite the differences in instantaneous growth rates between individual diatom and dinoflagellate species, the varying mixed communities thus performed production-wise comparably during the bloom accumulation phase.

Growth rates based on increase in Chl *a* were higher than carbon-based growth rates when diatoms were dominating. The faster accumulation of Chl *a* than carbon, on a community scale, could be caused by rapid synthesis of Chl *a* in diatoms based on reserve storage (Ross and Geider, 2009). This was supported by the difference in C : Chl *a* ratio between diatom and dinoflagellate-dominated communities. During diatom dominance, the C : Chl *a* ratio was rapidly decreasing during exponential growth phase, reflecting the difference in μ_{POC} and $\mu_{\text{Chl}a}$. The Chl *a* based measurements overestimated the production rate under diatom dominance, and the results emphasize the importance of considering the currency of planktonic production measurements in large-scale estimates of aquatic primary production.

Our data showed that assimilation efficiency was highest in mixed communities, compared to either diatom or dinoflagellate dominance. This is in line with recent studies on the effects of biodiversity on community functioning, indicating that more diverse communities support higher resource use efficiency and productivity (Ptacnik et al., 2008; Worm et al., 2006; Stockenreiter et al., 2013; Striebel et al., 2009). Different species have different environmental requirements, occupying different niches in the ecosystem. With increased diversity, the probability of occupying more of the total niche space increases, leading to better utilization of resources.

The net metabolic differences within variable community dominance manifested themselves only after the exponential growth phase, when nutrients were effectively incorporated to biomass, and loss processes became prominent. Net growth of primary producers is regulated by the balance of production and loss processes, such as respiration, excretion, sedimentation, and grazing. Sedimentation losses were eliminated in our experimental setup. Grazing effects were assumed to be minor, because there

Coastal carbon pathways and stoichiometry

K. Spilling et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Coastal carbon pathways and stoichiometry

K. Spilling et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



are no overwintering populations of large copepods in the Baltic Sea, our experiments started with 200 μm pre-screening, and the grazing pressure by other heterotrophs on the large-celled (mostly $> 20 \mu\text{m}$) spring bloom diatoms or dinoflagellates is negligible (Lignell et al., 1993). The main loss pathways are therefore respiration and excretion of dissolved organic matter. The latter evidently possess high carbon-to-nutrient ratios, as particulate nutrient fractions remained relatively stable once inorganic nutrient pools were depleted during the exponential growth phase.

There was no statistically significant difference between respiration rates of communities dominated by diatoms or dinoflagellates. In monocultures, dinoflagellates have generally higher respiration ratio than diatoms (Spilling and Markager, 2008; Falkowski and Owens, 1978), but our experiments with natural mixed communities did not reproduce this difference reliably. During the exponential phase, respiration ratios were equal, whereas for the stationary stage, high variability in the generally high respiration ratios of dinoflagellate-dominated communities (median 78 %) failed to yield significance for the apparent difference to diatom dominance. However, significantly lower assimilation efficiencies of dinoflagellate-dominated communities (median 10 %) were clearly driven by respiration, not by DOC release (Fig. 5).

Especially in diatom-dominated communities, POC continued to increase significantly after the primary production peak, and the communities kept fixing $^{14}\text{CO}_2$, as indicated by the modeled carbon accumulation and relatively low (median 45 %) respiration ratios. Concomitant late bloom DOC release in diatom-dominated communities increased significantly over other communities, as well. The results from our natural diatom-dominated communities reproduced the early observations obtained with diatom batch monocultures by (Goldman et al., 1992), who pointed out that the conventional new production concept, based on Redfield ratios (Dugdale and Goering, 1967), neglects the “excess” carbon fixation, due to uncoupling between photosynthesis and nutrient acquisition.

Our results showed that the post-peak bloom DOC accumulation was an order of magnitude lower than the parallel POC increase in diatom-dominated communities

(note y-axis scales in Fig. 1). DOC is by definition a pragmatic concept (organic carbon passing a glass-fiber filter with a nominal 0.7 μm porosity). Diatoms are known to excrete C-rich organic compounds of variable molecular weight as means to dissipate harvested light energy, once biomass synthesis becomes limited by nutrient deficiency (Kjørboe et al., 1990; Underwood et al., 2004). It is likely that the colloidal and mucoid DOC fractions were progressively trapped on POC filters during the late bloom stages, when coagulation and aggregation of detrital matter with the continuum of “dissolved” organic carbon continued in the absence of sedimentation flows. Additionally, any labile DOC excreted was probably quickly utilized by bacteria and did not add to the measured DOC pool.

The overall conclusion of the C budgets for different communities is that during the build-up of the bloom, differences between varying community dominance were far smaller than anticipated from monoculture studies. With regard to inorganic nutrient drawdown, exponential biomass development, and assimilation efficiency, diatoms and dinoflagellates acted to a large extent as functional surrogates. Major differences evolved only after the bloom culmination (coinciding with the depletion of inorganic nutrient pools and primary productivity peak), with significant consequences for carbon sequestration, C:N:P stoichiometry of spring bloom material flows, and the carbon-to-chlorophyll ratio of the communities.

4.4 Effects of community composition on nutrient stoichiometry and C drawdown during bloom events

The stationary phase of diatom-dominated communities strongly influenced the stoichiometry of seston, by doubling (112% increase) the C content from the exponential phase compared both to N and P, and 1.6 times higher compared to stationary phase dinoflagellate dominance (Fig. 6). The net effect was therefore a 3.6-fold enhanced CO_2 sequestration to that expected from the Redfield ratio. Even dinoflagellate-dominated communities exhibited corresponding CO_2 drawdown enhancement to POC in the stationary phase, but at a significantly lower level (1.7-fold higher than the C:N, Red-

BGD

11, 11863–11902, 2014

Coastal carbon pathways and stoichiometry

K. Spilling et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



field prediction). Carbon assimilation efficiencies in late bloom stages were very low for dinoflagellate-dominated communities due to high respiration rates, therefore preventing significant accumulation of organic C despite ongoing ^{14}C fixation. Dinoflagellate communities had a lower N : P ratio than diatom communities in stationary growth phase. Dinoflagellates have a high uptake affinity for P and keep assimilating it after growth stops (Kremp et al., 2008) probably due to their large genome, of which P is an essential component.

The excess carbon fixation noted in a stationary diatom batch culture (Goldman et al., 1992) was supported in field conditions by chemical proxies and discussed within the “biological pump” framework, as a vehicle transporting more CO_2 -derived carbon from the atmosphere to the oceans than expected from nutrient availability and fixed Redfield ratios (Sambrotto et al., 1993). Engel et al. (2002) showed that a major component of the emerging high POC : PON ratios in an experimentally induced natural diatom community bloom was aggregation of “marine snow” (72 % more dissolved inorganic carbon fixation than inferred from nitrate supply and Redfield stoichiometry), following a large late-bloom flow of carbon into transparent exopolymer particles (TEP). Schartau et al. (2007) modeled this “carbon overconsumption” flux based on the experimental results, and addressed 30 % of the POC increase to TEP formation.

Our results clearly support the “overconsumption” carbon flow pattern for a natural diatom-dominated bloom presented by Engel et al. (2002) and modeled by Schartau et al. (2007), but the difference of our results to Redfield-based estimates was even higher. The diatom community of Engel et al. (2002) exhibited close-to-Redfield stoichiometry during the bloom accumulation phase, while our diatom-dominated exponential phase communities showed seston C : N and C : P ratios almost double and triple the corresponding Redfield ratios, respectively. In the stationary phase, our strongest diatom-dominated communities had up to 3.6 times higher seston C : N content (regression in Fig. 6) than anticipated from Redfield ratios, as compared to 72 % by Engel et al. (2002).

BGD

11, 11863–11902, 2014

Coastal carbon pathways and stoichiometry

K. Spilling et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Coastal carbon pathways and stoichiometry

K. Spilling et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Estimates of offshore carbon overconsumption in the field, based on integrative geochemical approaches to in situ variations of chemical species, were reported up to 300 % in the “Vanishing in Bermuda” debate and Joint Global Ocean Flux Study (Toggweiler, 1994; Michaels et al., 1994; Marchal et al., 1996), soon after the experimental observations of Goldman et al. (1992), who found C : N ratio enhancement of ca. 200 % in late phase diatom cultures. Our N-based “C overconsumption” for coastal, diatom-dominated natural community bloom events (up to 3.6 times higher) therefore support, but also expand these observations, in terms of (a) the observed ranges of “C overconsumption”, (b) direct measurements of bloom events by natural mixed phytoplankton communities, and (c) geographically covering a coastal regime. Most importantly, however, we also show that the bloom community composition significantly affects the level of “C overconsumption”. Dinoflagellate-dominated communities showed a similar pattern of increasing carbon-to-nutrient ratios of seston from exponential to stationary phases, but with clearly smaller departures from the Redfield stoichiometry than under diatom dominance (1.7 times higher than Redfield C : N).

4.5 Carbon-to-chlorophyll ratio and community composition

A stoichiometric ratio of particular interest for large-scale estimates of aquatic primary production, either for geographically defined provinces or globally, is the carbon-to-chlorophyll ratio. Implementing numerical models of primary productivity requires either direct carbon-based phytoplankton observations or incorporation of fixed (Cloern et al., 1995) or dynamic C : Chl *a* ratios (Taylor et al., 1997). Phytoplankton C is notoriously difficult to separate from seston C, and no methods for direct measurements in the field are available. Most available spatially extensive observations originate from satellite remote sensing of chlorophyll, which requires bridging to carbon-based models (Behrenfeld and Falkowski, 1997). Currently, advanced oceanic biogeochemical models include dynamic C : Chl *a* ratios with photoacclimation parameterization, the most common of which is the Geider et al. (1998) model or its derivatives (Sathyendranath et al., 2009; Baird et al., 2013).

Coastal carbon pathways and stoichiometry

K. Spilling et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Carbon-to-chlorophyll ratios are known to be highly variable both in monocultures and in nature (Taylor et al., 1997; Chan, 1980), and the photoacclimation models are generally parameterized with monoculture responses to controlled laboratory conditions, most often highly departing from any set of natural conditions. Major uncertainty is introduced when laboratory models are translated for application to field models, if the sources of C:Chl *a* variability are not sufficiently understood and accounted for (Sathyendranath et al., 2009). These sources are normally addressed as responses of cultured algae to light and nutrient availability, which certainly are the key drivers for photosynthesis and the maintenance of the photosynthetic machinery, including cellular quotas for C, N, and P. However, species- or group-specific differences in these responses have rarely been incorporated, and the ability of photoacclimation models to cope with functionally different phytoplankton groups and non-steady state natural conditions is a major current challenge for variable stoichiometry models (De La Rocha et al., 2010).

Our results showed that both the growth stage of a bloom and the species dominance patterns strongly affected the community C:Chl *a* ratios. The lowest ratios (30 to 80; g:g) were encountered during the primary productivity and Chl *a* peak phases, when the community composition had a minor effect. During senescent bloom stages, diatom-dominated communities developed 4-fold C:Chl *a* ratios (median 200), whereas dinoflagellate-dominated communities showed median values of ca. 400, in similar irradiance, temperature, and nutrient-depleted conditions. These transient, order-of-magnitude changes within a few weeks during bloom events, with a strong component of species composition, present so far overlooked challenges for models of phytoplankton acclimation and geographically extensive production estimates based on satellite remote sensing.

5 Community change in changing climate

Competition between cold-water dinoflagellates and diatoms represents an important aspect of community change especially in changing climatic conditions of coastal temperate and arctic environments. Other well-documented, increasing dinoflagellate occurrences amidst diatom dominance that suggest climatic connotations are warm-water harmful algal blooms (Hallegraeff et al., 2010), while in several marine habitats, prymnesiophytes (especially *Emiliana huxleyi* and *Phaeocystis*; (Breton et al., 2006) are the main competitors for diatoms. Our results indicate that such variation in phytoplankton community dominance patterns have potentially significant consequences for marine biogeochemical cycles that need to be addressed, as ca. half of global primary production is attributed to marine systems, with phytoplankton as the main component (Field et al., 1998).

Application of either fixed Redfield stoichiometry, or uniform “phytoplankton” stoichiometry, can lead to several-fold errors or uncertainties in estimates for CO₂ sequestration, especially during temperate and arctic spring blooms. In dynamic natural bloom conditions, distinguishing between community composition and physiological acclimation during varying bloom stages, as sources of variation in seston stoichiometry, therefore remains a major challenge for trait-based phytoplankton ecology.

Marine biota are often regarded and studied as mere passive objects of climate change, and their responses to changes in e.g. temperature, acidification, stratification, and light climate of the mixed surface layer are accordingly actively studied. It appears essential to focus equally on the active role of phytoplankton in climate change: how marine carbon sequestration and interconnected biogeochemical cycles are directly modified by community change. Incorporating functional diversity and stoichiometric flexibility of primary producers into marine biogeochemical models is therefore a pending task for climate change research. An obvious parallel challenge for geographically extensive estimates of marine primary production is enhanced taxonomical

BGD

11, 11863–11902, 2014

Coastal carbon pathways and stoichiometry

K. Spilling et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



resolution of remote sensing technologies, to cope with the ongoing large-scale community change and its biogeochemical consequences.

Acknowledgements. This study was funded by Academy of Finland (decision no 111336 & 263862), funding was also provided by Walter and Andrée de Nottbeck Foundation (K. S.), and Estonian Research Council grant PUT39 (K. O. and R. K.). We would like to thank all the staff at Tvärminne Zoological Station (Univ. Helsinki) for all technical assistance and Päivi Hakanen for her assistance in the microscopy work.

References

Baird, M. E., Ralph, P. J., Rizwi, F., Wild-Allen, K., and Steven, A. D.: A dynamic model of the cellular carbon to chlorophyll ratio applied to a batch culture and a continental shelf ecosystem, *Limnol. Oceanogr.*, 58, 1215–1226, 2013.

Banse, K.: Cell volumes, maximal growth rates of unicellular algae and ciliates, and the role of ciliates in marine pelagial, *Limnol. Oceanogr.*, 27, 1059–1071, 1982.

Behrenfeld, M. J. and Falkowski, P. G.: A consumer's guide to phytoplankton primary productivity models, *Limnol. Oceanogr.*, 42, 1479–1491, 1997.

Breton, E., Rousseau, V., Parent, J.-Y., Ozer, J., and Lancelot, C.: Hydroclimatic modulation of diatom/Phaeocystis blooms in nutrient-enriched Belgian coastal waters (North Sea), *Limnol. Oceanogr.*, 51, 1401–1409, 2006.

Chan, A. T.: Comparative physiological study of marine diatoms and dinoflagellates in relation to irradiance and cell size. II. Relationship between photosynthesis, growth, and carbon/chlorophyll *a* ratio, *J. Phycol.*, 16, 428–432, 1980.

Cloern, J. E., Grenz, C., and Vidergar-Lucas, L.: An empirical model of the phytoplankton chlorophyll: carbon ratio-the conversion factor between productivity and growth rate, *Limnol. Oceanogr.*, 40, 1313–1321, 1995.

Collos, Y., Gagne, C., Laabir, M., Vaquer, A., Cecchi, P., and Souchu, P.: Nitrogenous nutrition of *Alexandrium catenella* (Dinophyceae) in cultures and in Thau Lagoon, Southern France, *J. Phycol.*, 40, 96–103, 2004.

De La Rocha, C., Terbrueggen, A., Völker, C., and Hohn, S.: Response and recovery of *Thalassiosira weissflogii* to nitrogen and silicon starvation: growth rates, nutrient uptake, and C, Si, and N per cell, *Mar. Ecol.-Prog. Ser.*, 412, 57–68, 2010.

Coastal carbon pathways and stoichiometry

K. Spilling et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Coastal carbon pathways and stoichiometry

K. Spilling et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Dugdale, R. C. and Goering, J. J.: Uptake of new and regenerated forms of nitrogen in primary productivity, *Limnol. Oceanogr.*, 12, 196–206, 1967.
- Dunne, J. P., Sarmiento, J. L., and Gnanadesikan, A.: A synthesis of global particle export from the surface ocean and cycling through the ocean interior and on the seafloor, *Global Biogeochem. Cy.*, 21, GB4006, doi:10.1029/2006GB002907, 2007.
- Eggers, S. L., Lewandowska, A. M., Barcelos e Ramos, J., Blanco-Ameijeiras, S., Gallo, F., and Matthiessen, B.: Community composition has greater impact on the functioning of marine phytoplankton communities than ocean acidification, *Global Change Biol.*, 20, 713–723, 2014.
- Engel, A., Goldthwait, S., Passow, U., and Alldredge, A.: Temporal decoupling of carbon and nitrogen dynamics in a mesocosm diatom bloom, *Limnol. Oceanogr.*, 47, 753–761, 2002.
- Falkowski, P. G.: Rationalizing elemental ratios in unicellular algae, *J. Phycol.*, 36, 3–6, 2000.
- Falkowski, P. G. and Owens, T. G.: Effects of light intensity on photosynthesis and dark respiration in six species of marine phytoplankton, *Mar. Biol.*, 45, 289–295, 1978.
- Field, C. B., Behrenfeld, M. J., Randerson, J. T., and Falkowski, P.: Primary production of the biosphere: integrating terrestrial and oceanic components, *Science*, 281, 237–240, 1998.
- Geider, R. J., MacIntyre, H. L., and Kana, T. M.: A dynamic regulatory model of phytoplankton acclimation to light, nutrients and temperature, *Limnol. Oceanogr.*, 43, 679–694, 1998.
- Goldman, J., Hansell, D., and Dennett, M.: Chemical characterization of three large oceanic diatoms: potential impact on water column chemistry, *Mar. Ecol.-Prog. Ser.*, 88, 257–270, 1992.
- Grasshoff, K., Ehrhardt, M., and Kremling, K.: *Methods of Seawater Analysis*, Verlag Chemie, Weinheim, 419 pp., 1983.
- Hallegraeff, G. M.: Ocean climate change, phytoplankton community responses, and harmful algal blooms: a formidable predictive challenge, *J. Phycol.*, 46, 220–235, 2010.
- Halpern, B. S., Walbridge, S., Selkoe, K. A., Kappel, C. V., Micheli, F., D'Agrosa, C., Bruno, J. F., Casey, K. S., Ebert, C., and Fox, H. E.: A global map of human impact on marine ecosystems, *Science*, 319, 948–952, 2008.
- Harley, C. D., Randall Hughes, A., Hultgren, K. M., Miner, B. G., Sorte, C. J., Thornber, C. S., Rodriguez, L. F., Tomanek, L., and Williams, S. L.: The impacts of climate change in coastal marine systems, *Ecol. Lett.*, 9, 228–241, 2006.
- Heiskanen, A. S.: Factors governing sedimentation and pelagic nutrient cycles in the northern Baltic Sea, *Monograph Boreal Env. Res.*, 8, 1–80, 1998.

Coastal carbon pathways and stoichiometry

K. Spilling et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Hinder, S. L., Hays, G. C., Edwards, M., Roberts, E. C., Walne, A. W., and Gravenor, M. B.: Changes in marine dinoflagellate and diatom abundance under climate change, *Nat. Clim. Change*, 2, 271–275, 2012.

Hoikkala, L.: Dynamics of dissolved organic matter and its bioavailability to heterotrophic bacteria in the Gulf of Finland, Northern Baltic Sea, University of Helsinki, 62 pp., 2012.

Kjørboe, T., Andersen, K. P., and Dam, H. G.: Coagulation efficiency and aggregate formation in marine phytoplankton, *Mar. Biol.*, 107, 235–245, 1990.

Klais, R., Tamminen, T., Kremp, A., Spilling, K., An, B. W., Hajdu, S., and Olli, K.: Spring phytoplankton communities shaped by interannual weather variability and dispersal limitation: mechanisms of climate change effects on key coastal primary producers, *Limnol. Oceanogr.*, 58, 753–762, 2013.

Kremp, A., Tamminen, T., and Spilling, K.: Dinoflagellate bloom formation in natural assemblages with diatoms: nutrient competition and growth strategies in Baltic spring phytoplankton, *Aquat. Microb. Ecol.*, 50, 181–196, 2008.

Legrand, C. and Carlsson, P.: Uptake of high molecular weight dextran by the dinoflagellate *Alexandrium catenella*, *Aquat. Microb. Ecol.*, 16, 81–86, 1998.

Leterme, S. C., Edwards, M., Seuront, L., Attrill, M., Reid, P., and John, A.: Decadal basin-scale changes in diatoms, dinoflagellates, and phytoplankton color across the North Atlantic, *Limnol. Oceanogr.*, 50, 1244–1253, 2005.

Lignell, R., Heiskanen, A. S., Kuosa, H., Gundersen, K., Kuuppo-Leinikki, P., Pajuniemi, R., and Uitto, A.: Fate of a phytoplankton spring bloom-sedimentation and carbon flow in the planktonic food web in the northern Baltic, *Mar. Ecol.-Prog. Ser.*, 94, 239–252, 1993.

Litchman, E. and Klausmeier, C. A.: Trait-based community ecology of phytoplankton, *Annu. Rev. Ecol. Evol. S.*, 39, 615–639, 2008.

Litchman, E., de Tezanos Pinto, P., Klausmeier, C. A., Thomas, M. K., and Yoshiyama, K.: Linking traits to species diversity and community structure in phytoplankton, *Hydrobiologia*, 653, 15–28, 2010.

Marchal, O., Monfray, P., and Bates, N. R.: Spring–summer imbalance of dissolved inorganic carbon in the mixed layer of the north-western Sargasso Sea, *Tellus B*, 48, 115–134, 1996.

Mateus, M., Leitão, P., de Pablo, H., and Neves, R.: Is it relevant to explicitly parameterize chlorophyll synthesis in marine ecological models?, *J. Marine Syst.*, 94, S23–S33, 2012.

Menden-Deuer, S. and Lessard, E. J.: Carbon to volume relationships for dinoflagellates, diatoms, and other protist plankton, *Limnol. Oceanogr.*, 45, 569–579, 2000.

Coastal carbon pathways and stoichiometry

K. Spilling et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Michaels, A. F., Bates, N. R., Buesseler, K. O., Carlson, C. A., and Knap, A. H.: Carbon-cycle imbalances in the Sargasso Sea, *Nature*, 372, 537–540, 1994.
- Niemi, Å.: Ecology of phytoplankton in the Tvärminne area, SW coast of Finland. II. Primary production and environmental conditions in the archipelago and the sea zone, *Acta Botanica Fennica*, 105, 1–73, 1975.
- 5 Niiranen, S., Yletyinen, J., Tomczak, M. T., Blenckner, T., Hjerne, O., MacKenzie, B. R., Müller-Karulis, B., Neumann, T., and Meier, H.: Combined effects of global climate change and regional ecosystem drivers on an exploited marine food web, *Global Change Biol.*, 19, 3327–3342, 2013.
- 10 Pahlow, M.: Linking chlorophyll-nutrient dynamics to the Redfield N:C ratio with a model of optimal phytoplankton growth, *Mar. Ecol.-Prog. Ser.*, 287, 33–43, 2005.
- Ptacnik, R., Solimini, A. G., Andersen, T., Tamminen, T., Brettum, P., Lepistö, L., Willén, E., and Rekolainen, S.: Diversity predicts stability and resource use efficiency in natural phytoplankton communities, *P. Natl. Acad. Sci. USA*, 105, 5134–5138, 2008.
- 15 Quigg, A., Finkel, Z. V., Irwin, A. J., Rosenthal, Y., Ho, T.-Y., Reinfelder, J. R., Schofield, O., Morel, F. M., and Falkowski, P. G.: The evolutionary inheritance of elemental stoichiometry in marine phytoplankton, *Nature*, 425, 291–294, 2003.
- Reid, P. C., Edwards, M., Hunt, H. G., and Warner, A. J.: Phytoplankton change in the North Atlantic, *Nature*, 391, 546–546, 1998.
- 20 Ross, O. N. and Geider, R. J.: New cell-based model of photosynthesis and photo-acclimation: accumulation and mobilisation of energy reserves in phytoplankton, *Mar. Ecol.-Prog. Ser.*, 383, 53–71, 2009.
- Rynearson, T., Richardson, K., Lampitt, R., Sieracki, M., Poulton, A., Lyngsgaard, M., and Perry, M.: Major contribution of diatom resting spores to vertical flux in the sub-polar North Atlantic, *Deep-Sea Res. Pt. I*, 82, 60–71, 2013.
- 25 Sakshaug, E., Bricaud, A., Dandonneau, Y., Falkowski, P. G., Kiefer, D. A., Legendre, L., Morel, A., Parslow, J., and Takahashi, M.: Parameters of photosynthesis: definitions, theory and interpretation of results, *J. Plankton Res.*, 19, 1637–1670, 1997.
- Sambrotto, R. N., Savidge, G., Robinson, C., Boyd, P., Takahashi, T., Karl, D. M., Langdon, C., Chipman, D., Marra, J., and Codispoti, L.: Elevated consumption of carbon relative to nitrogen in the surface ocean, *Nature*, 363, 248–250, 1993.
- 30 Sarthou, G., Timmermans, K. R., Blain, S., and Tréguer, P.: Growth physiology and fate of diatoms in the ocean: a review, *J. Sea Res.*, 53, 25–42, 2005.

Coastal carbon pathways and stoichiometry

K. Spilling et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Sathyendranath, S., Stuart, V., Nair, A., Oka, K., Nakane, T., Bouman, H., Forget, M.-H., Maass, H., and Platt, T.: Carbon-to-chlorophyll ratio and growth rate of phytoplankton in the sea, *Mar. Ecol.-Prog. Ser.*, 383, 73–84, 2009.

Schartau, M., Engel, A., Schröter, J., Thoms, S., Völker, C., and Wolf-Gladrow, D.: Modelling carbon overconsumption and the formation of extracellular particulate organic carbon, *Biogeosciences*, 4, 433–454, doi:10.5194/bg-4-433-2007, 2007.

Sharp, J. H., Benner, R., Bennett, L., Carlson, C. A., Dow, R., and Fitzwater, S. E.: Re-evaluation of high temperature combustion and chemical oxidation measurements of dissolved organic carbon in seawater, *Limnol. Oceanogr.*, 38, 1774–1782, 1993.

Smayda, T. J. and Reynolds, C. S.: Community assembly in marine phytoplankton: application of recent models to harmful dinoflagellate blooms, *J. Plankton Res.*, 23, 447–461, 2001.

Solórzano, L. and Sharp, J. H.: Determination of total dissolved phosphorus and particulate phosphorus in natural waters, *Limnol. Oceanogr.*, 25, 754–758, 1980.

Spilling, K. and Lindström, M.: Phytoplankton life cycle transformation lead to species-specific effects on sediment processes in the Baltic Sea, *Cont. Shelf. Res.*, 28, 2488–2495, 2008.

Spilling, K. and Markager, S.: Ecophysiological growth characteristics and modeling of the onset of the spring bloom in the Baltic Sea, *J. Marine Syst.*, 73, 323–337, 2008.

Stockenreiter, M., Haupt, F., Graber, A. K., Seppälä, J., Spilling, K., Tamminen, T., and Stibor, H.: Functional group richness: implications of biodiversity for light use and lipid yield in microalgae, *J. Phycol.*, 49, 838–847, 2013.

Striebel, M., Behl, S., and Stibor, H.: The coupling of biodiversity and productivity in phytoplankton communities: consequences for biomass stoichiometry, *Ecology*, 90, 2025–2031, 2009.

Suikkanen, S., Hakanen, P., Spilling, K., and Kremp, A.: Allelopathic effects of Baltic Sea spring bloom dinoflagellates on co-occurring phytoplankton, *Mar. Ecol.-Prog. Ser.*, 439, 45–55, 2011.

Sun, J. and Liu, D.: Geometric models for calculating cell biovolume and surface area for phytoplankton, *J. Plankton Res.*, 25, 1331–1346, 2003.

Tamminen, T.: Nitrate and ammonium depletion rates and preferences during a Baltic spring bloom, *Mar. Ecol.-Prog. Ser.*, 120, 123–133, 1995.

Tamminen, T. and Andersen, T.: Seasonal phytoplankton nutrient limitation patterns as revealed by bioassays over Baltic Sea gradients of salinity and eutrophication, *Mar. Ecol.-Prog. Ser.*, 340, 121–138, 2007.

Coastal carbon pathways and stoichiometry

K. Spilling et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Taylor, A. H., Geider, R. J., and Gilbert, F. J. H.: Seasonal and latitudinal dependencies of phytoplankton carbon-to-chlorophyll *a* ratios: results of a modelling study, *Mar. Ecol.-Prog. Ser.*, 152, 51–66, 1997.

5 Tillmann, U., Alpermann, T., John, U., and Cembella, A.: Allelochemical interactions and short-term effects of the dinoflagellate *Alexandrium* on selected photoautotrophic and heterotrophic protists, *Harmful Algae*, 7, 52–64, 2008.

Toggweiler, J.: Vanishing in Bermuda, *Nature*, 372, 505–506, 1994.

10 Underwood, G. J. C., Boulcott, M., Raines, C. A., and Waldron, K.: Environmental effects on exopolymer production by marine benthic diatoms: dynamics, changes in composition, and pathways of production, *J. Phycol.*, 40, 293–304, doi:10.1111/j.1529-8817.2004.03076.x, 2004.

Vehmaa, A., Kremp, A., Tamminen, T., Hogfors, H., Spilling, K., and Engström-Öst, J.: Copepod reproductive success in spring-bloom communities with modified diatom and dinoflagellate dominance, *ICES J. Mar. Sci.*, 69, 351–357, 2011.

15 Worm, B., Barbier, E. B., Beaumont, N., Duffy, J. E., Folke, C., Halpern, B. S., Jackson, J. B., Lotze, H. K., Micheli, F., and Palumbi, S. R.: Impacts of biodiversity loss on ocean ecosystem services, *Science*, 314, 787–790, 2006.

Coastal carbon pathways and stoichiometry

K. Spilling et al.

Table 1. Summary of the experimental set-up in different years. The treatments were nutrient addition (NPSi), which were additions of nitrogen (N), phosphorus (P) and silicate (Si) in N-P, Si and N-P-Si additions. The light treatment (Light) was a low and high light treatment, 20 and 90 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ respectively. In 2005 there was a diatom or no diatom addition (Diatom) instead of two light levels and in 2007 there was diatoms added in a gradient (Diatom gradient). The diatoms added were: *Thalassiosira levanderi* ($\sim 10 \mu\text{m}$ diameter) and *T. baltica* (20–30 μm diameter), two very typical spring bloom species. The start concentration of NO_3 (Start NO_3) gives the concentration in $\mu\text{g L}^{-1}$ of NO_3 in the control and in the treatments with N addition. The peak Chl *a* values are the minimum and maximum concentration recorded in the control (no nutrient addition) and in treatments with nutrients added respectively.

Year	Start date	Duration days	Treatments	Start NO_3 $\mu\text{g L}^{-1}$	Start Chl <i>a</i> $\mu\text{g L}^{-1}$	Peak Chl <i>a</i> $\mu\text{g L}^{-1}$
2004	24 Mar	44	NPSi, Light	90/260	7.1	21/78
2005	7 Apr	28	NPSi, Diatom	100/250	2.9	20/184
2006	19 Apr	23	NPSi, Light	8/170	35.0	39/199
2007	16 Mar	33	NPSi, Diatom gradient	90/250	4.9	22/77
2008	12 Mar	28	NPSi, Light	100/280	0.6	43/70

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Coastal carbon pathways and stoichiometry

K. Spilling et al.

Table 2. Statistical comparison using 1-way ANOVA on ranks. Tukey's post hoc test of statistical significant differences (*) can be found in Table 3 (exponential growth phase) and 4 (stationary growth phase).

Exponential growth phase					
Parameter	DF	SS	MS	<i>F</i> value	<i>p</i> value
POC	2	5.58	2.79	0.026	0.974
DOC	2	29.3	14.6	0.140	0.870
TGP	2	313	156	1.637	0.211
CAE	2	888	444	5.770	0.007*
RES	2	941	471	6.258	0.005*
Stationary growth phase					
Parameter	DF	SS	MS	<i>F</i> value	<i>p</i> value
POC	2	1395	678	9.246	< 0.001*
DOC	2	958	479	5.399	0.009*
TGP	2	1084	542	6.386	0.005*
CAE	2	739	369	3.874	0.031*
RES	2	564	282	2.804	0.075

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)

[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Coastal carbon pathways and stoichiometry

K. Spilling et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Table 3. Tukey's post hoc tests of carbon parameters during exponential growth phase. Only the statistical significant parameters from Table 2 were tested: carbon assimilation efficiency (CAE) and respiration (RES). The phytoplankton community was categorized according to diatom dominance (Diatoms), mixed community (Mixed) and dinoflagellate dominance (Dinoflagellates). The star (*) indicate statistical significance ($\alpha = 0.05$).

CAE	Diff	St diff	p value
Mixed vs. Diatoms	10.833	3.026	0.013*
Mixed vs. Dinoflagellates	10.517	2.800	0.023*
Dinoflagellates vs. Diatoms	0.317	0.084	0.996
RES			
Mixed vs. Diatoms	11.583	3.272	0.007*
Mixed vs. Dinoflagellates	10.183	2.743	0.026*
Dinoflagellates vs. Diatoms	1.400	0.377	0.925

Coastal carbon pathways and stoichiometry

K. Spilling et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Table 4. Tukey's post hoc tests of carbon parameters during stationary growth phase. Only the statistical significant parameters from Table 2 were tested: particular organic carbon (POC) dissolved organic carbon (DOC), total gross production (TGP) and carbon assimilation efficiency (CAE). The phytoplankton community was categorized according to diatom dominance (Diatoms), mixed community (Mixed) and dinoflagellate dominance (Dinoflagellates). The star (*) indicate statistical significance ($\alpha = 0.05$).

POC	Diff	St diff	p value
Mixed vs. Diatoms	11.071	3.240	0.007*
Mixed vs. Dinoflagellates	4.029	1.120	0.509
Dinoflagellates vs. Diatoms	15.100	4.060	< 0.001*
DOC			
Mixed vs. Diatoms	11.464	3.094	0.011*
Mixed vs. Dinoflagellates	1.414	0.363	0.930
Dinoflagellates vs. Diatoms	10.050	2.492	0.046*
TGP			
Mixed vs. Diatoms	12.381	3.416	0.005*
Mixed vs. Dinoflagellates	2.114	0.554	0.845
Dinoflagellates vs. Diatoms	10.267	2.603	0.036*
CAE			
Mixed vs. Diatoms	2.512	0.654	0.791
Mixed vs. Dinoflagellates	8.671	2.145	0.096
Dinoflagellates vs. Diatoms	11.183	2.675	0.030*

Coastal carbon pathways and stoichiometry

K. Spilling et al.




Figure 1. An example of the data extracted from the mesocosms (data from high light treatment with N and P addition, 2004): dissolved, inorganic nutrients and particulate, organic nutrients **(A)** and carbon parameters **(B)**. The parameters are: nitrate (NO_3), particulate organic nitrogen (PON), phosphate (PO_4), particulate organic phosphorus (POP), dissolved silicate (DSi), biogenic silicate (BSi), particulate organic carbon (POC), dissolved organic carbon (DOC) and total gross production (TGP). All parameters were measured directly except TGP, which was extrapolated from short-term ^{14}C incubations. The growth was divided into exponential and stationary growth phases based on the primary production peak (L^{-1}), indicated with the horizontal bars on top **(B)**. Note the different scales for on the y-axes.




Figure 2. Species evenness plotted against the dinoflagellate proportion of the whole community.

Coastal carbon pathways and stoichiometry

K. Spilling et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Coastal carbon pathways and stoichiometry

K. Spilling et al.




Figure 3. The growth rate during exponential growth of particulate organic carbon (μ_{POC}) and Chlorophyll *a* (μ_{Chla}) at different dinoflagellate proportion of the total phytoplankton community **(A)**, and the relationship between carbon assimilation efficiency and μ_{POC} **(B)**. No significant trend was found for μ_{POC} , but a negative correlation between μ_{Chla} and dinoflagellate proportion. The solid line represents the linear regression (slope = -0.16 ; $R^2 = 0.53$; $p < 0.0001$) and the dashed lines represent the 95% confidence intervals. The carbon assimilation efficiency is the ratio between the measured growth rate in POC and the total gross production (Fig. 1). A positive correlation was found (slope = 1.49 ; $R^2 = 0.12$; $p = 0.04$).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Coastal carbon pathways and stoichiometry

K. Spilling et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 4. Carbon budget parameters: particulate organic carbon (POC), dissolved organic carbon (DOC) and total gross production, during exponential and stationary growth. The phytoplankton community was divided into three categories: diatom dominance (> 80 %), mixed community (20–70 % dinoflagellates) and dinoflagellate dominance (> 70 %). The rationale behind setting the group boundaries was from the apparent difference in species evenness (Fig. 1) between these groups. The stars (*) indicate statistical significance ($\alpha = 0.05$) against one (*) or two groups (**), details can be found in Table 3.

Coastal carbon pathways and stoichiometry

K. Spilling et al.




Figure 5. Carbon budget parameters: carbon assimilation efficiency and respiration, during exponential and stationary growth. The phytoplankton community was divided into three categories: diatom dominance (> 80 %), mixed community (20–70 % dinoflagellates) and dinoflagellate dominance (> 70 %). The rationale behind setting the group boundaries was from the apparent difference in species evenness (Fig. 1) between these groups. The star indicate statistical significance difference ($\alpha = 0.05$) compared to one (*) or two (**) other groups, details can be found in Table 4.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Coastal carbon pathways and stoichiometry

K. Spilling et al.




Figure 6. The C:N, C:P and N:P ratios plotted against the weighted mean of dinoflagellate proportion during exponential (primary production peak) and stationary growth phase (end of experiment). The dashed horizontal line represent the Redfield ratio (molar ratio), the red and blue line represents statistically significant linear regression of exponential and stationary growth phase data. Regression coefficients were for the C:N ratio: slope = -3.3 ; $R^2 = 0.14$; $p = 0.02$ and slope = -9.6 ; $R^2 = 0.29$; $p = 0.0004$ for exponential and stationary growth phase respectively. For the C:P ratio they were: slope = -48.0 ; $R^2 = 0.13$; $p = 0.03$ and slope = -171.4 ; $R^2 = 0.45$ and $p < 0.001$ for exponential and stationary growth phase respectively. For N:P ratio, the regression coefficients for stationary growth phase: slope = -3.15 ; $R^2 = 0.31$; $p = 0.0002$; for the exponential growth phase, the coefficients were: $R^2 = 0.04$; $p = 0.2$.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Coastal carbon pathways and stoichiometry

K. Spilling et al.




Figure 7. The C : Chl *a* ratio at the primary production peak (PP) and at the end of the experiment (End). The phytoplankton community was divided into three categories: diatom dominance (> 80 %), mixed community (20–70 % dinoflagellates) and dinoflagellate dominance (> 70 %). The rationale behind setting the group boundaries was from the apparent difference in species evenness (Fig. 1).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)