

Dear Reviewer,

We would like to thank reviewer for raising very interesting questions and providing us with useful comments to improve the quality of the paper. We have so far tried to address all the issues raised. Author's comments are italicized.

1) In setting up the treatments, the authors compare reed canary grass to bare soil. Is this what would be happening in the field? Bare soil? Given the authors had to remove organic matter and grass from the cores they used that were bare, it seems bare soil is not likely to be what would be occurring in the field perhaps. Therefore, the authors need to justify why they compared to bare soil and perhaps include in the discussion a brief discussion about how these results might have varied if they had not compared to bare soil but instead to whatever native vegetation would have recruited naturally to a rewetted site.

We agree that for evaluation of GHG effects of rewetting under in situ conditions an appropriate reference would be a naturally vegetated (grass) soil. During the present experiment, however, comparison of RCG and bare soil mesocosms (rather than, e.g., grass vegetation) was done in order to tentatively isolate the contribution of RCG in the measured GHG fluxes. In doing this, the GHG emission from the bare soil mesocosm was subtracted from the emission from the vegetated mesocosms. Inclusion of grass-vegetated mesocosms would have been a third experimental treatment, which was not feasible in the present study.

We intended to reflect our approach already in the title of the manuscript, but as both reviewers stresses this point; we obviously failed to justify it properly in the manuscript itself. We have now added such information in the discussion. Page 16, Line no. 318-320.

2) I think in the Introduction it would be worthwhile to mention that reed canary grass is an invasive species in some parts of the world (including the U.S.) and that it may not be prudent to use reed canary grass as a biofuel crop in places where there is concern about this invasive species spreading. Thus there are additional factors that must be taken into account besides the greenhouse gas balance before deciding to plant RCG and this should perhaps be discussed in the conclusion or discussion.

We have now mentioned in the introduction that reed canary grass is considered as an invasive species in some countries. Page 3, Line no. 50.

3) RVI is defined but not explained and this was not an acronym I was familiar with. Please explain this briefly when it is first introduced so the reader knows why it is a useful index.

We have added following section in methodology part:

Page 7, Line no. 141-148: Biomass development was monitored through the non-destructive measurement of ratio vegetation index (RVI). RVI was determined for each mesocosm using a SpectroSense 2+ fitted with SKR1800 sensors (Skype Instruments, Powys, UK). The sensors measured the incident and reflected red light (R) at 656 nm and the incident and reflected infrared light (NIR) at

778 nm. RVI was then calculated as $(NIR_r/NIR_i)/(R_r/R_i)$ where the subscripts i and r denote the incident and reflected radiation. RVI has already been used as a useful predicting factor for modelling ER and CH₄ fluxes (Görres et al., 2014; Kandel et al., 2013a; Kandel et al., 2013b; Karki et al., 2014)

4) In the statistical method section, I think it would be useful to describe more fully what the CorAR1 structure versus compound symmetry represent since these were applied to the CH₄, N₂O or CO₂ fluxes.

Following lines were added in statistical section:

Page 11, Line no. 216-221: Dates were treated as repeated measurements by applying either compound symmetry structure (each dependent variable have constant covariance independent of time) or autocorrelation structure of order 1 (Errors at adjacent time points are correlated) (Maxwell and Delaney, 2004). Best model was selected by use of Akaike's Information Criterion (AIC). For CH₄ and N₂O, autocorrelation structure was selected while compound symmetry was selected for CO₂ fluxes.

5) I would really like to see a diagram/schematic showing GHG flux in and GHG fluxes out of each treatment and at each water level and then also the net balance (a la W. Schlesinger figures). This would visually help me understand the overall net fluxes and would support nicely the authors' premise that RCG can have overall the effect of making a rewetted peatland a sink for CO₂.

We are not familiar with mentioned W. Schlesinger figures. However, we have now added a graph with total GHG balance from each treatment of bare soil and RCG at different GWL in Figure 6d.

6) I would have liked a bit more discussion of the potential policy implications of this study. For example, the findings that increases in CH₄ emissions under RCG were offset by decreased N₂O emissions except when the water levels were at 0 would suggest that it is important if possible when reflooding peatlands to control the degree of wetting. Also, the peaks of N₂O emissions in the RCG treatments that occurred after fertilization suggest that it is critical to only fertilize when absolutely necessary and to keep that fertilization to a minimum. Perhaps some standards need to be set to ensure that fertilization application does not offset the potential benefits of replanting in rewetted peats.

This is indeed a very interesting comment by the reviewer. We agree with the reviewer that good fertilization management could decrease the total emission. We have now acknowledged that further studies are needed to assess the optimum amount and timing of fertilization required for optimum growth of RCG with acceptable N₂O emissions. Page 21, Line no. 442-446.

Technical Comments

1) Add lines in Table 1 to separate SO₄ from NH₄ from NO₃ more easily for the reader.

Done

2) Figure 7: Is this in comparison to the bare treatments? That would explain why there are negative bars for N₂O. But this needs to be clarified in the figure text.

The figure 7 represents the emission only from plants derived from the difference in emissions from RCG treatment and bare soil treatments. We have now made an effort to clarify the figure text.

3) I don't understand the sentence on page 13314 lines 13-14. "...as non-linear increase in gas concentration over time was often observed with non-steady state chambers for used gas measurement." What does "used gas measurement" mean?

We have now changed the sentence as "as non-linear increase in GHG concentration over time was often observed during the non-steady state chambers measurements" Page 7, Line no. 130

4) Page 13323, Line 1. "was" should be "WERE"

Done

5) Page 13324. Line 2. Remove "the" in "...which could suppress CH₄ emissionS."

Done

6) Need to include the scientific name of Reed Canary Grass in the Introduction.

Done

Page 3, Line no. 48

Dear Reviewer,

We would like to thank you for interesting questions and useful comments. We have tried to address all the issues raised by you. The original comment is given first followed by our response.

1. I would like to see some detail in the introduction about the cultivation of RCG. For example, how are crops established, managed and harvested?

We have now added a line in introduction that RCG can be established from seeds and added a reference for detailed management practices. Page 7, Line no. 49-50.

2. Why is bare soil the control? Why not use the vegetation that would grow naturally following rewetting?

We agree that for evaluation of GHG effects of rewetting under in situ conditions an appropriate reference would be a naturally vegetated (grass) soil. During the present experiment, however, comparison of RCG and bare soil mesocosms (rather than, e.g., grass vegetation) was done in order to tentatively isolate the contribution of RCG in the measured GHG fluxes. In doing this, the GHG emission from the bare soil mesocosm was subtracted from the emission from the vegetated mesocosms. Inclusion of grass-vegetated mesocosms would have been a third experimental treatment, which was not feasible in the present study.

We intended to reflect our approach already in the title of the manuscript, but as both reviewers stresses this point; we obviously failed to justify it properly in the manuscript itself. We have now added such information in the discussion. Page 16, Line 318-320.

3. P13313 L24 Why is there not a control with no fertiliser? How were the application rates decided?

Our previous study on effect of fertilization on RCG biomass yield already showed higher biomass yield of RCG with fertilization than without fertilization (Kandel et al. 2013). Similar fertilization rate was applied as applied in reed canary grass field from where the mesocosms were collected. However, higher nitrogen fertilization was given considering the lower mineralization rate at higher GWL. We have now added this information in the methodology section. Page 6, Line no. 110-114.

The optimum amount and timing of fertilization required for optimum growth of RCG could be another experiment and we have now acknowledge this in the discussion section. Page 21, Line no. 442-446.

4. What is RVI? If this is a new method it should be addressed in the introduction.

RVI has now been defined in detail. We have now added a reference where it has been previously used for modelling CO₂ and CH₄ fluxes. Page 7, Line no. 141-148.

5. P13326 L14 Given that biomass yields were similar what is the implication for fertilizer application? Is it necessary?

As mentioned above, RCG showed higher biomass yield with fertilization than without fertilization (Kandel et al. 2013). Nevertheless, in the discussion, we recommend further studies to assess the effect of fertilization on RCG growth. Page 21, Line no. 442-446

6. P13326 L16 Is this uptake of 6.2 kg CO₂/m² an actual CO₂ sink? Will part of this not be removed when the crop is harvested? What proportion of this might be belowground sequestration or how could it be estimated?

We completely agree with the reviewer that the part of CO₂ taken by plants will be removed when the plants are harvested. But we didn't go into further discussion about it is as the uptake of CO₂ was not measured for whole measurement period. Rather we simply made the distinction between ecosystems being C sinks (from a soil C perspective) or CO₂ sinks from an atmospheric perspective. With the data we provide we believe that we can reasonably state that RCG cultivation makes the systems CO₂ sinks from an atmospheric perspective.

Even taking into account the amount of C loss in harvested biomass the total emissions from RCG at 0 cm of GWL was 6.4 kg CO₂. eq. m⁻² year⁻¹ which is almost similar to the growing season GPP. In annual basis the GPP will even be higher. Furthermore, the RCG biomass will additionally reduce the CO₂ emissions by fossil fuel displacement.

Minor comments P13312 L21-22 Chnage to '...60-70 cm early in the 20th century and since then has been used for agricultural purposes.' P13313 L1 Insert 'the' before 'mesocosm'

Done

1 **Effect of reed canary grass cultivation on greenhouse gas emission from peat soil at controlled**
2 **rewetting**

3

4

5 Sandhya Karki^{1*}, Lars Elsgaard¹ and Poul Erik Lærke¹

6

7

8 ¹Aarhus University, Department of Agroecology, Blichers Allé 20, DK-8830 Tjele, Denmark

9 *Corresponding Author: Tel: +45 8715 4785, Email Address: sandhya.karki@agrsci.dk

10 **Abstract**

11 Cultivation of bioenergy crops in rewetted peatland (paludiculture) is considered as a possible land
12 use option to mitigate greenhouse gas (GHG) emissions. However, bioenergy crops like reed canary
13 grass (RCG) can have a complex influence on GHG fluxes. Here we determined the effect of RCG
14 cultivation on GHG emission from peatland rewetted to various extents. Mesocosms were
15 manipulated to three different ground water levels (GWL), i.e., 0, -10 and -20 cm below the soil
16 surface in a controlled semi-field facility. Emissions of CO₂ (ecosystem respiration, ER), CH₄ and
17 N₂O from mesocosms with RCG and bare soil were measured at weekly to fortnightly intervals
18 with static chamber techniques for a period of one year. Cultivation of RCG increased both ER and
19 CH₄ emissions, but decreased the N₂O emissions. The presence of RCG gave rise to 69, 75 and 85%
20 of total ER at -20, -10 and 0 cm GWL, respectively. However, this difference was due to decreased
21 soil respiration at the rising GWL as the plant-derived CO₂ flux was similar at all three GWL. For
22 methane, 70-95% of the total emission was due to presence of RCG, with the highest contribution at
23 -20 cm GWL. In contrast, cultivation of RCG decreased N₂O emission by 33-86% with the major
24 reductions at -10 and -20 cm GWL. In terms of global warming potential, the increase in CH₄
25 emissions due to RCG cultivation was more than off-set by the decrease in N₂O emissions at -10
26 and -20 cm GWL; at 0 cm GWL the CH₄ emissions was offset only by 23%. CO₂ emissions from
27 ER obviously were the dominant RCG-derived GHG flux, but above-ground biomass yields, and
28 preliminary measurements of gross photosynthetic production, showed that ER could be more than
29 balanced due to the photosynthetic uptake of CO₂ by RCG. Our results support that RCG cultivation
30 could be a good land use option in terms of mitigating GHG emission from rewetted peatlands,
31 potentially turning these ecosystems into a sink of atmospheric CO₂.

32
33 **Keywords:** Ecosystem respiration; Ground water level, Methane; Nitrous oxide; Paludiculture

34 1 Introduction

35

36 Peatlands cover 3% of the world's area but contain 30% of the soil organic carbon (Parish et al.,
37 2008), signifying an important role in the global carbon cycle. About 15% of the world's peatlands
38 have been drained for different human purposes mostly for agriculture and forestry and to a lesser
39 extent for peat extraction (Joosten, 2009). Drained peatlands are major sources of CO₂ emissions
40 and estimated to account for about 6% of the total anthropogenic CO₂ emission (Joosten, 2009). In
41 order to reduce the large emissions of CO₂ from drained peatlands, extensive rewetting projects
42 have been implemented in Europe and North America (Höper et al., 2008), and rewetted organic
43 soils have been included in the guidelines for national greenhouse gas (GHG) inventories by the
44 Intergovernmental Panel on Climate Change (IPCC, 2014). In addition, agricultural use of wet and
45 rewetted peatlands for crop growth (paludiculture) is considered as a possible land use option that
46 may indirectly reduce the CO₂ emissions ~~on rewetted organic soils~~ by biomass production for
47 energy purposes (Joosten et al., 2012; Günther et al., 2014).

48 Reed canary grass (RCG) (*Phalaris arundinacea*) is one of the ~~most~~ suitable biomass crops
49 for paludiculture (Wichtmann and Tanneberger, 2011). It can be established from seeds as normal
50 agricultural grass (Kandel et al., 2013b) but in some countries it is considered as an invasive species
51 (Maurer et al., 2003). The plants thrive in wet soils due to aerenchyma tissues (Kercher and Zedler,
52 2004; Askaer et al., 2011) that transport oxygen to the roots in otherwise anaerobic soil
53 compartments. However, cultivating wetland plants like RCG may influence the overall GHG
54 balance by a combination of contrasting effects. First of all, RCG can stimulate the processes of
55 GHG production by increasing the labile soil organic carbon pool, e.g., via root exudates (Ström et
56 al., 2003; Bastviken et al., 2005). Next, the transport of oxygen to anaerobic zones stimulates
57 heterotrophic degradation of organic matter, but at the same time stimulates oxidation of CH₄ (Kao-

58 Kniffin et al., 2010) and suppress CH₄ production due to increase in redox potential (Laanbroek,
59 2010; Sutton-Grier and Megonigal, 2011). RCG may further increase the emissions of reduced soil
60 gasses as the aerenchyma tissues act as a conduit for the direct transport of, e.g., CH₄ and N₂O
61 produced in soil (Joabsson et al., 1999; Jørgensen et al., 2012). Also, RCG can decrease N₂O
62 emissions by assimilation of mineral N which reduces the availability of electron acceptors (nitrate)
63 for denitrifying microorganisms (Roobroeck et al., 2010). In summary, the introduction of RCG at
64 rewetted peatlands may cause a change in the patterns and underlying mechanisms of GHG
65 emission, which is rather complex.

66 | In the natural state, GHG emissions from ~~rewetted~~ peatlands are predominantly controlled by
67 | the position (depth) of the water table (IPCC, 2014). Basically, due to slow diffusion of oxygen in
68 | water (10,000 times slower than in air), ground water level (GWL) has a strong control on the
69 | oxic/anoxic soil boundary and thereby on the biogeochemical processes involved in GHG fluxes
70 | (Dinsmore et al., 2009; Karki et al., 2014). However, the presence of aerenchymatous plants may
71 | strongly interact with GWL in being decisive for the resulting GHG emissions from ~~rewetted-wet~~
72 | peatlands. The objective of the present study was to ~~document the initial effect-~~ quantify the role of
73 | RCG cultivation on the resulting GHG emissions of CO₂, N₂O and CH₄ from peat soils rewetted to
74 | various extents. Such type of information is very important to understand the total GHG balance
75 | from paludiculture and improve the basis for modelling future climate. To accomplish this, the
76 | GHG emissions of all three gases were measured in an annual study with peat soil mesocosms with
77 | RCG and bare soil rewetted to constant GWL of 0, -10 and -20 cm in a controlled semi-field
78 | facility.

79

80 2 Materials and methods

81 2.1 Site description

82 Soil cores were collected from a fen peatland in the Nørre Å river valley, Denmark (56°44'N,
83 9°68'E). The peatland was drained to a depth of 60-70 cm ~~in~~ early in the 20th century and since then
84 used for agricultural purposes. RCG experimental plots were established at the site in 2009 (Kandel
85 et al., 2013b). The top soil layer (0-20 cm) at the study site had the following main properties:
86 highly decomposed peat soil corresponding to H9 on the von Post scale; bulk density, 0.29 g cm⁻³;
87 total organic carbon, 37.8% and total nitrogen, 3.2% (Karki et al., 2014).

88

89 2.2 Experimental design

90 A total of 30 intact soil cores for the mesocosm study were collected in May 2012 by inserting PVC
91 pipes of 60 cm depth and 30 cm diameter into the soil. Half of the soil cores were collected from
92 RCG plots and the other half were collected from a grass field surrounding the RCG plots. The
93 upper 5 cm of the soil and litter layer was removed from the grass field before inserting the PVC
94 pipes and these soil cores were kept bare during the experiment. The soil cores were retrieved with
95 help of a mini excavator and transported to semi-field facilities at AU-Foulum (Karki et al., 2014).
96 The bottom of the PVC pipes were covered with net to allow for free water movement and the pipes
97 were then installed in plastic cylinders (diameter, 37 cm; height, 70 cm). The plastic cylinders were
98 filled with gravel at the bottom 10 cm and the space between the PVC pipes and the wall of the
99 cylinders (ca. 3 cm) was filled with sand. The whole set up was then installed in a trench at the
100 semi-field facility with the soil surface at ground level.

101 Mesocosms with bare soil and RCG were randomly divided into three groups and were
102 manipulated to three different GWL of 0, -10 and -20 cm below the soil surface. Water table was
103 adjusted by fitting a rubber tubing (diameter, 1 cm) to the bottom of each plastic cylinder and

104 placing the other end of the rubber tubing at different heights corresponding to the level of GWL
105 treatment. Water was supplied in the space between the PVC pipes and the wall of the cylinders
106 every day for one hour by a drip irrigation system. Further details on mesocosm incubations and the
107 semi-field facility were given by Karki et al. (2014).

108 | Due to poor regrowth of RCG (both under mesocosms and field conditions), initial weed
109 biomass was uprooted and new RCG seeds were spread on 21 June 2012. RCG was fertilized with
110 surface application of 0.6 g N, 0.1 g P and 0.5 g K per mesocosm on 23 July 2012 (corresponding to
111 80 kg N, 13 kg P and 77 kg K ha⁻¹). This fertilization rate corresponded to the rate applied in a
112 previous study at the RCG field site from where the mesocosms were collected (Kandel et al.,
113 2013a), except that the nitrogen rate was slightly increases in the mesocosm study as lower N
114 mineralization was expected at higher GWLs. After the regrowth of RCG in spring 2013, RCG was
115 fertilized with the same amount of fertilizer on 30 April and again in 28 June 2013. RCG plants
116 were harvested twice, first on 29 October 2012 and then on 27 June 2013. In bare soil mesocosms,
117 emerging weeds were uprooted and mosses were eliminated by application of iron sulphate (FeSO₄)
118 on 29 August 2012. No fertilizer was added to bare soil mesocosms.

119

120 2.3 Gas measurements and flux calculation

121 Dark PVC chambers (diameter, 30 cm; height, 50 cm) equipped with fans and pressure equilibration
122 vents were used for the measurement of CO₂, CH₄ and N₂O (Karki et al., 2014). Gas measurements
123 were carried out between 10:00 and 13:00 at weekly to fortnightly intervals during July 2012 to July
124 2013. Four gas samples (10 mL) were drawn from the chamber headspace with polypropylene
125 syringes during 45 minutes of chamber enclosure and transferred to evacuated 6-mL Exetainers.
126 Gas samples were analysed with an Agilent 7890 gas chromatograph connected to a CTC
127 CombiPAL automatic sample injection system (Agilent, Nærum, Denmark). Fluxes were calculated

128 with the HMR method (Pedersen et al., 2010) in the statistical software R [version 3.0.2](#) (R Core
129 Team, 2013) as non-linear increase in ~~gas-GHG~~ concentration over time was often observed ~~with~~
130 ~~during the~~ non-steady state chambers ~~measurements for used gas measurement~~ (Davidson et al.,
131 2002; Petersen et al., 2012). Thus according to statistical HMR analysis fluxes were calculated
132 either by non-linear or linear models (Pedersen et al., 2010). Out of the total of 435 fluxes for each
133 ~~gasGHG~~, the non-linear approach was applied for 41, 40 and 18% of CO₂, CH₄ and N₂O fluxes
134 from RCG mesocosms, ~~respectively~~ and 22, 16 and 22% of CO₂, CH₄ and N₂O fluxes from bare
135 soil mesocosms, ~~respectively~~. In bare soil at 0 cm GWL, approximately 3% of the CH₄ fluxes were
136 discarded due to episodic release of CH₄ presumably by ebullition.

137

138 2.4 Biomass measurement

139 ~~Biomass development was monitored through the non-destructive measurement of ratio vegetation~~
140 ~~index (RVI). RVI was determined for each mesocosm using a SpectroSense 2+ fitted with~~
141 ~~SKR1800 sensors (Skype Instruments, Powys, UK) as described by Görres et al. (2014). Biomass~~
142 ~~development was monitored through the non-destructive measurement of ratio vegetation index~~
143 ~~(RVI). RVI was determined for each mesocosm using a SpectroSense 2+ fitted with SKR1800~~
144 ~~sensors (Skype Instruments, Powys, UK). The sensors measured the incident and reflected red light~~
145 ~~(R) at 656 nm and the incident and reflected infrared light (NIR) at 778 nm. RVI was then~~
146 ~~calculated as $(NIR_r/NIR_i)/(R_r/R_i)$ where the subscripts i and r denote the incident and reflected~~
147 ~~radiation. RVI has already been used as a useful predicting factor for modelling ER and CH₄ fluxes~~
148 ~~(Kandel et al., 2013a; Kandel et al., 2013b; Görres et al., 2014; Karki et al., 2014)~~

149 RVI measurements were done on the same days as GHG sampling except in winter when the
150 soil was covered with snow or frozen. The total above ground dry biomass from each mesocosm
151 was also determined after each harvest by oven drying ~~the plant material~~ at 60°C to constant weight.

152 After the harvest in 2013, species composition from each mesocosm was determined on dry weight
153 basis to quantify the contribution of volunteer weeds in the total biomass.

154

155 2.5 *Environmental parameters and pore water analysis*

156 Soil temperature at 5 cm depth and soil moisture was measured by temperature and time domain
157 reflectometry (TDR) probes installed permanently in one of the five replicates for each GWL
158 treatment. Soil temperature was measured automatically every hour while soil moisture
159 measurements with TDR (volumetric water content, VWC) were done on every gas sampling
160 occasion. The instrumented mesocosms also had Pt probes installed at 20 cm depth to measure soil
161 redox potential. Soil redox potential was measured at fortnightly intervals from mid-April to July
162 2013 with a portable pH meter (PHM220, Radiometer) by gently pushing a double junction calomel
163 reference electrode (REF251, Hach Lange) into the soil. Measured redox potential were converted
164 to standard hydrogen electrode potential (Eh) by addition of +245 mV (Kjaergaard et al., 2012).

165 A piezometer (length, 65 cm; diameter, 2 cm) with the screen all the way down was installed
166 in the instrumented mesocosms. Approximately 30 mL of soil water was sampled monthly from
167 these piezometers except for February to April 2013 ~~where~~-when water inside the piezometers was
168 frozen. Water samples were analysed for ammonium, nitrate, and sulphate content. Ammonia and
169 nitrate content were measured using an auto-analyzer (Bran+Luebbe GmbH; Norderstedt,
170 Germany) and sulphate was determined by ion chromatography on a Dionex ICS-1500 IC-system
171 (Dionex Corp.; Sunnyvale, CA, USA).

172

173 2.6 Cumulative GHG fluxes

174 For the mesocosms with RCG, CO₂ emissions from ecosystem respiration (ER) were modelled as a
175 function of GWL, temperature, and biomass (RVI) by Model 1 (Karki et al. 2014); for bare soil
176 mesocosms Model 2 excluding RVI was applied:

177

$$178 \quad ER = (b_1 + b_2 \text{GWL}) \times \exp \left(b_3 \left(\frac{1}{10 - T_0} - \frac{1}{T - T_0} \right) \right) \times (b_4 + \text{RVI}) \quad [1]$$

$$179 \quad ER = (b_1 + b_2 \text{GWL}) \times \exp \left(b_3 \left(\frac{1}{10 - T_0} - \frac{1}{T - T_0} \right) \right) \quad [2]$$

180

181 where T_0 is a notional zero respiration temperature, here fixed to -46.02°C (Lloyd and Taylor,
182 1994), T is the air or soil temperature ($^\circ\text{C}$), RVI is the ratio vegetation index, GWL is water table
183 depth below the soil surface (cm) and b_1 , b_2 , b_3 and b_4 are model parameters.

184 All model parameters were estimated by non-linear regression in SigmaPlot 11 (Systat
185 Software, Chicago, IL, USA). Using the obtained model parameters, continuous temperature data
186 and linearly interpolated RVI data, hourly rates of CO₂ emissions were reconstructed for each
187 GWL. These hourly emissions values were summed to yield the annual flux from 10 July 2012 to 9
188 July 2013. The uncertainty of annual fluxes were addressed by deriving the minimum and
189 maximum cumulative fluxes from upper and lower values of model parameters \pm standard errors
190 (SE) (Elsgaard et al., 2012). For model evaluation the Nash–Sutcliffe modelling efficiency (ME)
191 was calculated according to:

192

$$ME = 1 - \frac{\sum_{i=1}^n (Mes_i - Mod_i)^2}{\sum_{i=1}^n (Mes_i - \overline{Mes})^2} \quad [3]$$

194

195 where Mes_i and Mod_i are measured and modelled values, respectively, and \overline{Mes} is the mean of
 196 measured vales (Haefner, 2005).

197 Cumulative CH₄ and N₂O fluxes were calculated by ~~the~~ linear interpolation ~~method~~ between
 198 the sampling dates using the trapezoidal rule (Petersen et al., 2012). ~~The~~ ~~l~~ linear interpolation
 199 method was used as there were no common models to predict CH₄ and N₂O fluxes for vegetated
 200 and bare soil plots. Cumulative fluxes were calculated for each individual mesocosm and then
 201 averaged for each GWL treatment ($n = 5$). Total GHG emissions were calculated by summing
 202 annual CO₂, CH₄ and N₂O emissions at each GWL; CH₄ and N₂O emissions were converted to CO₂
 203 equivalents by multiplying with 28 and 265, respectively, according to the revised global warming
 204 potential (GWP) of the three GHG (Myhre et al., 2013). The plant-derived total GHG emission at
 205 each GWL was estimated as the difference between the total GHG emissions from RCG mesocosms
 206 and bare soil mesocosms. The uncertainty of annual ~~plant-plant~~-derived GHG emissions was
 207 derived from the uncertainty of plant and bare soil emissions added in quadrature.

208

209 2.7 Statistical analysis

210 Statistical analyses were done using R version 3.0.2 (R Core Team, 2013). Data were analyzed
 211 using a linear mixed model including the fixed effect of vegetation (bare soil/RCG), GWL, date and
 212 their two-way interactions. The model also included the random effect of each experimental unit.
 213 Prior to analysis, ~~gas~~-CH₄ and N₂O flux data were log-transformed after addition of a constant
 214 (minimum fluxes of CH₄ and N₂O) to obtain normal distribution and variance homogeneity. ~~Dates~~

215 ~~were treated as repeated measurements and autocorrelation of CorAR1 structure was applied to CH₄~~
216 ~~and N₂O fluxes while compound symmetry was applied to CO₂ fluxes (Oehlert, 2000). Dates were~~
217 ~~treated as repeated measurements by applying either compound symmetry structure (each~~
218 ~~dependent variable have constant covariance independent of time) or autocorrelation structure of~~
219 ~~order 1 (errors at adjacent time points are correlated) (Maxwell and Delaney, 2004). Best model was~~
220 ~~selected by use of Akaike's Information Criterion (AIC). For CH₄ and N₂O, autocorrelation~~
221 ~~structure was selected while compound symmetry was selected for CO₂ fluxes.~~

222
223 A similar linear mixed model was run to determine the effect of GWL on RVI development. One-
224 way ANOVA was used to test the difference in mean yield between the treatments. Significance of
225 all tests was accepted at $P < 0.05$.

226

227 **3 Results**

228 *3.1 Environmental conditions*

229 The average air temperature during the study period was 6.9°C and total precipitation was 667 mm
230 (Fig. 1). Snowfall started in early December 2012 and was observed till end of March 2013 with
231 intermittent freezing and thawing events. The soil was frozen and covered with ice till mid-April
232 2013. The annual average soil temperature (5 cm depth) in RCG treatments was 7.4, 7.7 and 7.6°C
233 at 0, -10 and -20 cm GWL, respectively; for bare soil treatments it was 7.5, 7.4 and 7.9°C at 0, -10
234 and -20 cm GWL, respectively. The average volumetric soil water content during the measurement
235 period was $82 \pm 5\%$, $67 \pm 3\%$, and $58 \pm 3\%$ from RCG treatments at 0, -10 and -20 cm GWL,
236 respectively, and $83 \pm 4\%$, $62 \pm 6\%$, and $55 \pm 7\%$ from bare soil treatments at 0, -10 and -20 cm
237 GWL, respectively (mean \pm standard deviation, $n = 22$). Average soil redox potential was -115, -27,

238 and 40 mV from RCG treatments at 0, -10 and -20 cm GWL, respectively, and -118, -51, 151 mV
239 from bare soil treatments at 0, -10, and -20 cm GWL, respectively (Fig. 2).

240

241 3.2 Biomass yield and RVI

242 The mean biomass yield was 6.0 and 6.6 Mg ha⁻¹ across all GWL in 2012 and 2013, respectively
243 (Fig. 3). During the first year there was a good stand of RCG but during the second year weed
244 biomass became established especially at 0 cm GWL; this was notably meadow foxtail (*Alopecurus*
245 *pratensis* L.) and grasses (*Poa* sp.) which made an important contribution to the total biomass at the
246 time of harvest.

247 The pattern of RVI development was similar among the different GWL treatments; peak
248 values of RVI occurred in late August 2012, where after RVI started to decline due to plant
249 senescence. RVI started to increase again during the regrowth of biomass in spring 2013 (Fig. 4).

250

251 3.3 Pore water properties

252 The annual variation in soil water sulphate concentrations ranged from 1.3 to 56.9 mg L⁻¹.
253 Generally, similar SO₄²⁻ concentrations were found in bare soil and RCG mesocosms at 0 and -10
254 cm GWL, but at -20 cm GWL consistently higher SO₄²⁻ concentrations were found in the bare soil
255 mesocosms (Table 1). For ammonium the concentrations ranged from 0.1 to 10.2 mg L⁻¹ and higher
256 NH₄⁺ concentrations were generally found in bare soil mesocosms than in RCG mesocosms at 0 and
257 -10 cm GWL. In the bare soil treatments the level of NH₄⁺ was lower at -20 cm GWL than at 0 and
258 -10 cm GWL, but in RCG treatments NH₄⁺ concentrations were similar at all the three GWLs
259 (Table 1). The concentration of nitrate was low (<3.1 mg L⁻¹) across all treatments; the highest NO₃⁻
260 levels were generally seen at bare soil treatments at -20 cm GWL (Table 1).

261

262 3.4 Measured GHG fluxes

263 The emission of CO₂ was measured as ER in RCG and bare-soil treatments in order to evaluate the
264 contribution of RCG in the total ER at the different GWLs. The emissions of CO₂ were different
265 between RCG and bare soil mesocosms ($p < 0.001$) and also between the three GWL treatments (p
266 < 0.001) (Table 2). CO₂ emissions decreased consistently with higher GWL both from RCG and
267 bare soil mesocosms. The emissions showed expected seasonal variation with highest CO₂ fluxes
268 during summer time ($p < 0.001$) (Fig. 5a, b). CO₂ emissions ranged from 20 to 485 mg m⁻² h⁻¹
269 across all GWL in bare soil and from 55 to 1700 mg m⁻² h⁻¹ in RCG treatments. Among the air and
270 soil temperature at 5 cm, CO₂ emissions were better correlated with soil temperature in bare soil,
271 but with the air temperature in RCG treatments.

272 Methane fluxes were significantly affected both by vegetation and GWL (Table 2). CH₄
273 emissions were highest at 0 cm GWL both from RCG and bare soil treatments (Fig. 5c, d). CH₄
274 emissions from RCG treatments showed temporal variation ($P < 0.001$) with highest emissions
275 during summer time (Fig. 5c). Peak emissions of CH₄ from RCG treatments were observed in
276 August 2012 across all GWL levels, ranging from 4.4 to 8.9 mg CH₄ m⁻² h⁻¹. During November to
277 early April (i.e., winter season), CH₄ emissions from RCG treatments were below 0.1 mg m⁻² h⁻¹
278 and even occasional uptake (25% of total fluxes measured) of CH₄ was recorded. From bare soil
279 treatments, CH₄ fluxes were generally low and fluctuated between apparent net emission and net
280 uptake except for few episodic peak events, generally from 0 cm GWL. These peak events were
281 considered to represent unsystematic ebullition events.

282 N₂O fluxes from RCG treatments were generally low, fluctuating in a range between -0.02 to
283 0.07 mg m⁻² h⁻¹ except for peak events after fertilizer application (Fig. 5e). Emission peaks of 0.4,
284 0.7 and 0.4 mg N₂O m⁻² h⁻¹ were observed at 0 cm GWL immediately after the first, second and
285 third fertilization events, respectively. Smaller peak emissions of 0.4 and 0.2 mg N₂O m⁻² h⁻¹ were

286 observed at -10 cm GWL after the first and second fertilization event, but at -20 cm GWL, peak
287 emission after the fertilizer application was absent. N₂O emissions from bare soil treatments
288 generally were higher and ranged from -0.02 to 1.9 mg m⁻² h⁻¹. Most of the N₂O emission in bare
289 soil mesocosms was measured during the winter period from November 2012 to April 2013
290 accounting for more than 70% of the cumulative emission at 0 and -10 cm GWL and more than
291 50% at -20 cm GWL.

292

293 3.5 Annual GHG emissions and contribution of plants to annual GHG emissions

294 The estimated parameters for CO₂ flux models are presented in Table 3 showing also that the
295 modelling efficiency was considerably higher for the RCG treatments than the bare soil treatments.
296 Annual CO₂ emissions decreased consistently with raising GWL towards the soil surface both in
297 RCG and bare soil treatments (Fig. 6). In contrast, CH₄ emissions increased systematically both
298 from RCG and bare soil treatments in response to raising GWL (Fig. 6). The annual N₂O emissions
299 showed a contrasting response to raising GWL in bare soil and RCG treatments; in bare soil
300 treatments lower N₂O emissions occurred in response to ~~raising~~raised GWL, but in RCG treatments
301 there was a tendency of higher N₂O emissions in response to ~~raising~~raised GWL (Fig. 6).

302 The presence of plants contributed 69-85% of the total CO₂ emissions from the RCG
303 mesocosms (Fig. 6). The highest contribution was observed at 0 cm GWL and the contribution
304 decreased at lower GWL. RCG likewise ~~contributed~~accounted for more than 70% of total CH₄
305 emissions with the highest contribution of 95% observed at -20 cm GWL. Thus at this GWL (-20
306 cm) CH₄ emission was negligible from bare soil treatments (0.2 g CH₄ m⁻² yr⁻¹) whereas the
307 emissions was substantial from RCG treatments (4.1 g CH₄ m⁻² yr⁻¹). In contrast to CO₂ and CH₄
308 emissions, cultivation of RCG reduced the annual N₂O emissions despite the application of mineral
309 N fertilizer in RCG mesocosms (Fig. 6). At -10 and -20 cm GWL, RCG eliminated 82-86% of the

310 N₂O emissions as compared to bare soil treatments; from 0 cm GWL the reduction corresponded to
311 33% of the N₂O emissions. In terms of GWP, the increase in CH₄ emissions due to RCG cultivation
312 was more than off-set by the decrease in N₂O emissions at -10 and -20 cm GWL, but apparently not
313 at 0 cm GWL where CH₄ emissions were off-set by only 23% by the decreased N₂O emission (Fig.
314 7). CO₂ emissions from ER, though, were the dominant RCG-derived GHG fluxes (Fig. 7).

315 4 Discussion

316

317 During the present study, the effects of RCG cultivation on GHG emission from rewetted peatland
318 was evaluated by comparison of planted and unplanted (bare-soil) mesocosms. Mesocosms with
319 RCG and bare soil (rather than, e.g mesocosms with native grasses) were compared in order to
320 tentatively isolate the contribution of RCG in the measured GHG fluxes. One concern of using this
321 plant exclusion method for GHG studies is the difference in soil moisture regime and temperature
322 that may develop between planted and bare soil treatments which may result in different
323 decomposition rates of soil organic matter (Kuzyakov, 2006). With our experimental setup, we were
324 able to control the GWL throughout the measurement period and this resulted in soil moisture
325 contents (VWC) that were similar between RCG and bare soil treatments at each GWL; this was
326 generally also seen for the soil redox potential and pore water sulphate concentration at least at 0
327 and -10 cm GWL. The average soil temperature difference between the RCG and bare soil
328 treatments was found to be less than 1°C; however during the annual study we observed some
329 seasonal difference in soil temperature especially during spring days (higher temperature in RCG
330 treatments) and summer days (lower temperature in RCG treatments) which was attributed to the
331 RCG cultivation. Yet, the differences in moisture and temperature regime between the planted and
332 bare-soil mesocosms were considered to be modest and pertinent for an evaluation of the effects of
333 RCG on total GHG emissions.

334 Monitoring of environmental variables was achieved by instrumentation of one out of five
335 replicate mesocosms at each GWL. We assumed that the measured variables were representative for
336 all replicates and that the instrumentation did not lead to any bias. This was substantiated by the
337 absence of any systematic deviations in measured GHG fluxes from the instrumented and non-

338 instrumented replicates. Thus, the average difference in annual fluxes with and without
339 instrumentation was less than 15%.

340

341 4.1 *CO₂ emissions*

342 Plants can enhance CO₂ flux from ER directly by above- and below-ground respiration and
343 indirectly by enhancing the decomposition of soil organic matter by the supply of easily degradable
344 root exudates to the soil (priming effect) (Kuzyakov et al., 2001; Van Huissteden et al., 2006). In
345 vegetated soils ER is essentially balanced by photosynthetic CO₂ uptake, and therefore CO₂
346 emissions from ER does not represent the net ecosystem exchange (NEE) of CO₂. Rather than
347 quantifying NEE, an important result of the present study was that plant-derived ER from RCG
348 mesocosms (the major part of total CO₂ emissions) was similar at all three GWL (Fig. 7)
349 substantiating the results of Lafleur et al. (2005) and Riutta et al. (2007) who reported autotrophic
350 respiration to be independent of water table depth. Thus, the observed increase (from 69 to 85%) ~~in~~
351 ~~the contribution of plants~~ in total ER with rising GWL was promoted mainly by decreasing soil
352 respiration at the higher GWL (Fig. 6). The observed contribution of RCG to total CO₂ emissions
353 (on average 76%) was higher than the values of 55% previously reported by Shurpali et al. (2008).
354 Yet, the ~~studies results of by~~ Shurpali et al. (2008) ~~was carried out in~~ were obtained for a drained
355 peatland with an average GWL of -65 cm which would favor aerobic soil respiration to a larger
356 extent than in our soils with GWL no deeper than -20 cm. In accordance with this we also observed
357 a larger soil respiration at -20 cm than at 0 cm GWL.

358

359 4.2 *Methane emissions*

360 Methane fluxes from soil is the result of CH₄ production, consumption and transport (Lai, 2009).

361 Plants play a key role on CH₄ fluxes as they have potential to influence all three processes (Joabsson

362 et al., 1999). CH₄ emissions were higher from RCG than bare soil treatments even though the GWL
363 was raised to the soil surface. Plant roots release organic compounds to soil, which are easily
364 available carbon sources to anaerobic microbial consortia eventually producing the precursors
365 (acetate or H₂/CO₂) for methanogenesis (Ström et al., 2003). Such fresh organic carbon is suggested
366 to be important substrates for methanogenesis as peat carbon is shown to be more recalcitrant to
367 anaerobic decomposition (Tuittila et al., 2000; Hahn-Schöfl et al., 2011).

368 Methane produced in soil can be emitted to the atmosphere by diffusion, ebullition (release of
369 gas bubbles) and plant-mediated transport (Whalen, 2005; Lai, 2009). Indeed, RCG can transport
370 CH₄ from soil to the atmosphere directly through its aerenchyma tissue, thereby bypassing the
371 microbial methane oxidation layer in the soil. On an annual basis it has been estimated that RCG
372 may actually transport 70% of the total CH₄ emissions from a natural wetland in Denmark (Askaer
373 et al., 2011). In the absence of plant-mediated transport, diffusion expectedly would be the
374 dominant pathway of CH₄ emissions in bare soil treatments. CH₄ transport through diffusion is a
375 slow but important process for bringing CH₄ in contact with the CH₄ oxidizing microbial
376 community (Whalen 2005; Lai 2009). In our study there was negligible CH₄ emissions from bare
377 soil at -20 cm GWL, aligning with the results of Schäfer et al. (2012) who reported this drainage
378 depth to be sufficient to suppress diffusive CH₄ emissions due to methane oxidation and reduced
379 methanogenesis.

380 The transport of oxygen by aerenchyma plants to anoxic soil compartments has been reported
381 to increase the redox potential which could suppress ~~the~~ CH₄ emission (Sutton-Grier and
382 Megonigal, 2011). However, in our study neither the redox potential nor the sulphate content was
383 consistently increased by the presence of plants suggesting the role of substrate availability and
384 transport of CH₄ through RCG to be the important factors for controlling CH₄ emissions from the
385 RCG treatments.

386 It is possible that we could have underestimated the total CH₄ emission from bare soil
387 treatments at 0 cm GWL as episodic CH₄ release through ebullition was not taken into account in
388 the annual balance. Ebullition events were identified by occasional erratic time courses of CH₄
389 concentrations during the flux measurements; however as these events were generally associated
390 with the initial (time 0 and 15 min) CH₄ measurements chamber gas samplings it was believed to
391 represent artifacts created during chamber deployment. Yet, episodic release of CH₄ may be more
392 important in bare soil than in vegetated soil as plants may reduce the soil concentration of CH₄ by
393 mediating CH₄ transport and also by rhizospheric oxidation of CH₄; these processes reduce the
394 potential formation of CH₄ bubbles (Chanton, 2005). Tentatively accounting for the observed
395 episodic CH₄ release, a total of 0.04 g m⁻² of CH₄ was released during the study; this was a
396 negligible contribution (< 1%) to the annual CH₄ flux from bare soil at 0 cm GWL. However, as
397 ebullition events are short-lived and unsystematic they could easily be missed by the chamber
398 measurements (Coulthard et al., 2009).

399

400 4.3 N₂O emissions

401 Annual fluxes of N₂O (0.2 to 0.4 g N₂O m⁻² yr⁻¹) from RCG mesocosms were within the range (-0.4
402 to 0.8 g N₂O m⁻² yr⁻¹) reported for undisturbed Danish riparian wetland (Audet et al., 2014).
403 However, annual fluxes were higher in bare soil (0.6 to 2.0 g N₂O m⁻² yr⁻¹) as compared to RCG
404 treatments. Thus, RCG decreased the annual N₂O emissions, contradictory to the finding of
405 Hyvönen et al. (2009) where fertilization in RCG increased the N₂O emissions by 90% as compared
406 to bare soil. However, in the study by Hyvönen et al. (2009) N₂O emissions were quite low (0.01 g
407 N₂O m⁻² yr⁻¹) from the soil without vegetation. Their site was an abandoned peatland (Hyvönen et
408 al. 2009) probably with limited nitrification because of a high C/N ratio (42.3) (Klemedtsson et al.,
409 2005) compared to our peat soil with rich N content (3.2%) and a low C/N ratio (11.6). Thus, the

410 ecosystem studied by Hyvönen et al. (2009) might have been more N limited at the unfertilized sites
411 than was the case for our study site. ~~This, in turn, could cause a more pronounced response to N~~
412 ~~fertilization in the studies of Hyvönen et al. (2009).~~

413 The effect of RCG cultivation on N₂O emissions was highly dependent upon the GWL. At 0
414 cm GWL, there was least effect of RCG cultivation on N₂O emissions due to peak emissions
415 observed after fertilization. Peak emission observed after fertilization events suggest that N₂O
416 emission was limited by mineral N content at 0 cm GWL. Saari et al. (2013) and Silvan et al. (2002)
417 also reported a significant increase in N₂O emission after addition of inorganic nitrogen in riparian
418 wetland due to favorable conditions for denitrification.

419 Previous studies have reported that winter emissions significantly contributed to annual N₂O
420 emissions (Maljanen et al., 2004; Regina et al., 2004). Such emissions in winter have been related
421 to the physical release of N₂O that is produced and trapped under frozen surface layers ~~and as well~~
422 ~~as~~ the emissions of newly produced N₂O (*de novo* emissions) at the onset of thaw stimulated by
423 increased biological activity and changes in physical and chemical soil conditions (Risk et al.,
424 2013). Significant emissions at all GWL were observed in winter from bare soil treatments, but not
425 from RCG treatments. After harvesting, there was regrowth of RCG and also other volunteer
426 grasses which survived throughout the winter and which may have competed with microorganisms
427 for available N. Maljanen et al. (2004) also observed higher N₂O emissions from bare soil as
428 compared to vegetated plots during winter and likewise related the low emission in vegetated plots
429 to low mineral N content due to uptake of nitrate by plants. Bare soil treatments indeed had a higher
430 availability of mineral N (Table 1), and could be more prone to physical damage by freeze and thaw
431 cycles due to lack of plant cover; both these factors stimulate the biological activities related to N₂O
432 emissions as substantiated by the observed slight increase also in CO₂ emissions coinciding with
433 increased N₂O emission especially at 0 and -10 cm GWL.

434

435 *4.4 Effect of RCG cultivation on GHG balance from rewetted peatland*

436 Two of the major concerns of growing wetland plants like RCG in rewetted peatland are the
437 possible increase in CH₄ emissions due to supply of fresh plant material and transport of CH₄ by
438 aerenchyma tissue (Ström et al., 2003; Askaer et al., 2011) and the possible increase in N₂O due to
439 application of N fertilizers (Maljanen et al., 2010). However, in the present experiment, cultivation
440 of RCG decreased N₂O emission to an extent that could offset the increase in CH₄ emission at -10
441 and -20 cm GWL, but apparently not at 0 cm GWL; the latter case being due to peak emissions in
442 N₂O after fertilization events in RCG. This result suggests that emissions at 0 cm GWL can be
443 reduced by reducing the N fertilization rate. Further studies are needed to assess the optimum
444 amount and timing of fertilization required for optimum growth of RCG with acceptable N₂O
445 emissions. Emissions of N₂O caused by N fertilization should not off-set the benefit of fossil fuel
446 substitution obtained by the fertilizer induced increase of biomass production (Kandel et al. 2013a).
447 Regarding the overall GHG emission, the CO₂ emissions from ER was clearly the dominant RCG-
448 derived GHG flux. Yet, CO₂ flux from ER would to a large extent be counterbalanced by gross
449 photosynthesis which expectedly was similar at all GWL treatments (based on the similar biomass
450 yields), though CO₂ flux from photosynthesis was not measured in this annual study. Yet, a
451 photosynthetic uptake of 6.2 kg CO₂ m⁻² was measured from RCG mesocosms at 0 cm GWL during
452 the growing season from May to September 2013 (S. Karki, unpublished results) reflecting that
453 RCG potentially can turn the rewetted ecosystem into a sink of CO₂ from an atmospheric
454 perspective. Adaptation or selection of RCG varieties that thrive especially well under distinct
455 climate and shallow GWL conditions could further help to ~~fulfill this potential~~ improve the GHG
456 balance of paludiculture with RCG.

457

458

459 **5 Conclusion**

460

461 The present study to our knowledge is the first to compare the annual GHG emission from RCG and
462 bare soil treatments of rewetted peatland at controlled GWL. The following conclusions were
463 derived: (i) soil respiration decreased with increasing GWL from -20 to -10 to 0 cm, but RCG-
464 derived ER was similar at all three GWL resulting in the highest contribution of RCG to total ER
465 (85%) at 0 cm GWL, (ii) cultivation of RCG increased CH₄ emission at all GWLs, but relatively
466 most at -20 cm GWL, (iii) N₂O emissions decreased ~~at -10 and -20 cm of GWLs~~ due to RCG
467 cultivation especially during winter; winter emissions were an important component of annual
468 emission from bare soil, (iv) in terms of GWP, the increase in CH₄ emissions due to RCG
469 cultivation was more than off-set by the decrease in N₂O emissions at -10 and -20 cm GWL, (v)
470 CO₂ emissions from ER (the dominant RCG-derived GHG flux) could be balanced by
471 photosynthetic CO₂ uptake at all three GWL as indicated by the large and similar above-ground
472 biomass yields at all GWL, signifying a potential of RCG cultivation to turn the rewetted peatland
473 into a sink of atmospheric CO₂.

474

475 *Acknowledgement*

476

477 The study was partly supported by the Danish Centre for Food and Agriculture (DCA) as a part of
478 the BioBase programme. The authors like to thank Sanmohan Baby for statistical advice. We are
479 also grateful for the technical support we received from Bodil Stensgaard, Finn Henning
480 Christensen, Holger Bak, Jørgen M. Nielsen and Stig T. Rasmussen.

References

- Askaer, L., Elberling, B., Friberg, T., Jørgensen, C. J., and Hansen, B. U.: Plant-mediated CH₄ transport and C gas dynamics quantified in-situ in a *Phalaris arundinacea*-dominant wetland, *Plant Soil*, 343, 287-301, 2011.
- Audet, J., Hoffmann, C. C., Andersen, P. M., Baattrup-Pedersen, A., Johansen, J. R., Larsen, S. E., Kjaergaard, C., and Elsgaard, L.: Nitrous oxide fluxes in undisturbed riparian wetlands located in agricultural catchments: Emission, uptake and controlling factors, *Soil Biol. Biochem.*, 68, 291-299, 2014.
- Bastviken, S. K., Eriksson, P. G., Premrov, A., and Tonderski, K.: Potential denitrification in wetland sediments with different plant species detritus, *Ecol. Eng.*, 25, 183-190, 2005.
- Chanton, J. P.: The effect of gas transport on the isotope signature of methane in wetlands, *Org. Geochem.*, 36, 753-768, 2005.
- Coulthard, T. J., Baird, A. J., Ramirez, J., and Waddington, J. M.: Methane dynamics in peat: Importance of shallow peats and a novel reduced-complexity approach for modeling ebullition, in: *Carbon Cycling in Northern Peatlands*, edited by: Baird, A. J., Belyea, L. R., Comas, X., Reeve, A. S., and Slater, L. D., American Geophysical Union, Washington, D. C, 173-185, 2009.
- Davidson, E. A., Savage, K., Verchot, L. V., and Navarro, R.: Minimizing artifacts and biases in chamber-based measurements of soil respiration, *Agric. For. Meteorol.*, 113, 21-37, 2002.
- Dinsmore, K. J., Skiba, U. M., Billett, M. F., and Rees, R. M.: Effect of water table on greenhouse gas emissions from peatland mesocosms, *Plant Soil*, 318, 229-242, 2009.
- Elsgaard, L., Görres, C.-M., Hoffmann, C. C., Blicher-Mathiesen, G., Schelde, K., and Petersen, S. O.: Net ecosystem exchange of CO₂ and carbon balance for eight temperate organic soils under agricultural management, *Agr. Ecosyst. Environ.*, 162, 52-67, 2012.

- Görres, C. M., Kutzbach, L., and Elsgaard, L.: Comparative modeling of annual CO₂ flux of temperate peat soils under permanent grassland management, *Agr. Ecosyst. Environ.*, 186, 64-76, 2014.
- Günther, A., Huth, V., Jurasinski, G., and Glatzel, S.: The effect of biomass harvesting on greenhouse gas emissions from a rewetted temperate fen, *GCB Bioenergy*, doi:10.1111/gcbb.12214, 2014.
- Haefner, J. W.: *Modelling Biological Systems: Principles and Applications*, 2nd ed., Springer, New York, NY, 2005.
- Hahn-Schöfl, M., Zak, D., Minke, M., Gelbrecht, J., Augustin, J., and Freibauer, A.: Organic sediment formed during inundation of a degraded fen grassland emits large fluxes of CH₄ and CO₂, *Biogeosciences*, 8, 1539-1550, 2011.
- Höper, H., Augustin, J., Cagampan, J. P., Drösler, M., Lundin, L., Moors, E., Vasander, H., Waddington, J. M., and Wilson, D.: Restoration of peatlands and greenhouse gas balances, in: *Peatlands and Climate Change*, edited by: Starck, M., International Peat Society, Saarijärvi, Finland, 182-210, 2008.
- Hyvönen, N. P., Huttunen, J. T., Shurpali, N. J., Tavi, N. M., Repo, M. E., and Martikainen, P. J.: Fluxes of nitrous oxide and methane on an abandoned peat extraction site: effect of reed canary grass cultivation, *Bioresour. Technol.*, 100, 4723-4730, 2009.
- IPCC: 2013 Supplement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories: Wetlands in, edited by: Hiraishi, T., Krug, T., Tanabe, K., Srivastava, N., Baasansuren, J., Fukuda, M., and Troxler, T. G., IPCC, Switzerland, 2014.
- Joabsson, A., Christensen, T. R., and Wallen, B.: Vascular plant controls on methane emissions from northern peatforming wetlands, *Trends Ecol. Evol.*, 14, 385-388, 1999.
- Joosten, H.: *The Global Peatland CO₂ Picture: peatland status and drainage related emissions in all countries of the world*, Wetland International, Ede., 2009.

- Joosten, H., Tapio-Biström, M.-L., and Tol, S.: Peatlands-guidance for climate change mitigation through conservation, rehabilitation and sustainable use, Food and Agriculture Organisation of the United Nations and Wetlands International, Rome, Italy, 2012.
- Jørgensen, C. J., Struwe, S., and Elberling, B.: Temporal trends in N₂O flux dynamics in a Danish wetland - effects of plant-mediated gas transport of N₂O and O₂ following changes in water level and soil mineral-N availability, *Glob. Change Biol.*, 18, 210-222, 2012.
- Kandel, T. P., Elsgaard, L., Karki, S., and Lærke, P. E.: Biomass yield and greenhouse gas emissions from a drained fen peatland cultivated with reed canary grass under different harvest and fertilizer regimes, *Bioenergy Res.*, 6, 883-895, 2013a.
- Kandel, T. P., Elsgaard, L., and Lærke, P. E.: Measurement and modelling of CO₂ flux from a drained fen peatland cultivated with reed canary grass and spring barley, *GCB Bioenergy*, 5, 548-561, 2013b.
- Kao-Kniffin, J., Freyre, D. S., and Balsler, T. C.: Methane dynamics across wetland plant species, *Aquat. Bot.*, 93, 107-113, 2010.
- Karki, S., Elsgaard, L., Audet, J., and Lærke, P.: Mitigation of greenhouse gas emissions from reed canary grass in paludiculture: effect of groundwater level, *Plant Soil*, 383,217-230, 2014.
- Kercher, S. M., and Zedler, J. B.: Flood tolerance in wetland angiosperms: a comparison of invasive and noninvasive species, *Aquat. Bot.*, 80, 89-102, 2004.
- Kjaergaard, C., Heiberg, L., Jensen, H. S., and Hansen, H. C. B.: Phosphorus mobilization in rewetted peat and sand at variable flow rate and redox regimes, *Geoderma*, 173, 311-321, 2012.
- Klemedtsson, L., Von Arnold, K., Weslien, P., and Gundersen, P.: Soil CN ratio as a scalar parameter to predict nitrous oxide emissions, *Glob. Change Biol.*, 11, 1142-1147, 2005.
- Kuzyakov, Y., Ehrensberger, H., and Stahr, K.: Carbon partitioning and below-ground translocation by *Lolium perenne*, *Soil Biol. Biochem.*, 33, 61-74, 2001.

- Kuzyakov, Y.: Sources of CO₂ efflux from soil and review of partitioning methods, *Soil Biol. Biochem.*, 38, 425-448, 2006.
- Laanbroek, H. J.: Methane emission from natural wetlands: interplay between emergent macrophytes and soil microbial processes. A mini-review, *Ann. Bot.*, 105, 141-153, 2010.
- Lafleur, P. M., Moore, T. R., Roulet, N. T., and Frolking, S.: Ecosystem respiration in a cool temperate bog depends on peat temperature but not water table, *Ecosystems*, 8, 619-629, 2005.
- Lai, D. Y. F.: Methane dynamics in northern peatlands: A review, *Pedosphere*, 19, 409-421, 2009.
- Lloyd, J., and Taylor, J. A.: On the temperature dependence of soil respiration, *Funct. Ecol.*, 8, 315-323, 1994.
- Maljanen, M., Komulainen, V.-M., Hytönen, J., Martikainen, P. J., and Laine, J.: Carbon dioxide, nitrous oxide and methane dynamics in boreal organic agricultural soils with different soil characteristics, *Soil Biol. Biochem.*, 36, 1801-1808, 2004.
- Maljanen, M., Sigurdsson, B. D., Guðmundsson, J., Óskarsson, H., Huttunen, J. T., and Martikainen, P. J.: Greenhouse gas balances of managed peatlands in the Nordic countries-present knowledge and gaps, *Biogeosciences*, 7, 2711-2738, 2010.
- Maurer, D. A., Lindig-Cisneros, R., Werner, K. J., Kercher, S., and Miller, R.: The Replacement of Wetland Vegetation by Reed Canarygrass (*Phalaris arundinacea*), *Ecol. Rest.*, 21, 116-119, , 2003.
- Maxwell, S. E., and Delaney, H. D.: Designing experiments and analyzing data: A model comparison perspective, Psychology Press, 2004.
- Myhre, G., Shindell, D., Bréon, F. M., Collins, W., Fuglestedt, J., Huang, J., Koch, D., Lamarque, J. F., Lee, D., Mendoza, B., Nakajima, T., Robock, A., Stephens, G., Takemura, T., and Zhang, H.: Anthropogenic and natural radiative forcing, in: *Climate change 2013: The physical science basis. Contribution of working group I to the fifth assessment report of the Intergovernmental Panel on Climate Change*, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K.,

- Boschung, J., A. N., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 658-740, 2013.
- Oehlert, G. W.: A First Course in Design and Analysis of Experiments, University of Minnesota, 2000.
- Parish, F., Sirin, A., Charman, D., Joosten, H., Minayeva, T., Silvius, M., and Stringer, L.: Assessment on peatlands, biodiversity and climate change: main report, Global Environment Centre, Kuala Lumpur and Wetlands International, Wageningen, 179 pp., 2008.
- Pedersen, A. R., Petersen, S. O., and Schelde, K.: A comprehensive approach to soil-atmosphere trace-gas flux estimation with static chambers, *Eur. J. Soil Sci.*, 61, 888-902, 2010.
- Petersen, S. O., Hoffmann, C. C., Schäfer, C.-M., Blicher-Mathiesen, G., Elsgaard, L., Kristensen, K., Larsen, S. E., Torp, S. B., and Greve, M. H.: Annual emissions of CH₄ and N₂O, and ecosystem respiration, from eight organic soils in Western Denmark managed by agriculture, *Biogeosciences*, 9, 403-422, 2012.
- R. Core Team: R: A language and environment for statistical computing. R foundation for statistical computing, Vienna, Austria, 2013
- Regina, K., Syväsalo, E., Hannukkala, A., and Esala, M.: Fluxes of N₂O from farmed peat soils in Finland, *Eur. J. Soil Sci.*, 55, 591-599, 2004.
- Risk, N., Snider, D., and Wagner-Riddle, C.: Mechanisms leading to enhanced soil nitrous oxide fluxes induced by freeze-thaw cycles, *Can. J. Soil Sci.*, 93, 401-414, 2013.
- Riutta, T., Laine, J., and Tuittila, E.-S.: Sensitivity of CO₂ exchange of fen ecosystem components to water level variation, *Ecosystems*, 10, 718-733, 2007.
- Roobroeck, D., Butterbach-Bahl, K., Brüggemann, N., and Boeckx, P.: Dinitrogen and nitrous oxide exchanges from an undrained monolith fen: short-term responses following nitrate addition, *Eur. J. Soil Sci.*, 61, 662-670, 2010.

- Saari, P., Saarnio, S., Heinonen, J., and Alm, J.: Emissions and dynamics of N₂O in a buffer wetland receiving water flows from a forested peatland, *Boreal Environ. Res.*, 18, 164-180, 2013.
- Schäfer, C. M., Elsgaard, L., Hoffmann, C. C., and Petersen, S. O.: Seasonal methane dynamics in three temperate grasslands on peat, *Plant Soil*, 357, 339-353, 2012.
- Shurpali, N. J., Hyvönen, N. P., Huttunen, J. T., Biasi, C., Nykänen, H., Pekkarinen, N., and Martikainen, P. J.: Bare soil and reed canary grass ecosystem respiration in peat extraction sites in Eastern Finland, *Tellus B*, 60, 200-209, 2008.
- Silvan, N., Regina, K., Kitunen, V., Vasander, H., and Laine, J.: Gaseous nitrogen loss from a restored peatland buffer zone, *Soil Biol. Biochem.*, 34, 721-728, 2002.
- Ström, L., Ekberg, A., Mastepanov, M., and Røjle Christensen, T.: The effect of vascular plants on carbon turnover and methane emissions from a tundra wetland, *Glob. Change Biol.*, 9, 1185-1192, 2003.
- Sutton-Grier, A. E., and Megonigal, J. P.: Plant species traits regulate methane production in freshwater wetland soils, *Soil Biol. Biochem.*, 43, 413-420, 2011.
- Tuittila, E.-S., Komulainen, V.-M., Vasander, H., Nykänen, H., Martikainen, P. J., and Laine, J.: Methane dynamics of a restored cut-away peatland, *Glob. Change Biol.*, 6, 569-581, 2000.
- Van Huissteden, J., van den Bos, R., and Alvarez, I. M.: Modelling the effect of water-table management on CO₂ and CH₄ fluxes from peat soils, *Neth. J. Geosci*, 85, 3-18, 2006.
- Whalen, S. C.: Biogeochemistry of methane exchange between natural wetlands and the atmosphere, *Environ. Eng. Sci*, 22, 73-94, 2005.
- Wichtmann, W., and Tanneberger, F.: Landuse options for rewetted peatlands, in: Carbon credits from peatland rewetting. Climate, biodiversity, land use, edited by: Tanneberger, F., and Wichtmann, W., Stuttgart, Schweizerbart Science Publishers, 107-132, 2011.

Table 1. Concentration of sulphate, ammonium, and nitrate (mg L^{-1}) in ground water samples collected from piezometers from bare soil and reed canary grass (RCG) mesocosms at different ground water levels (GWL).

Treatment and date	SO_4^{2-} (mg L^{-1}) at GWL			NH_4^+ (mg L^{-1}) at GWL			NO_3^- (mg L^{-1}) at GWL		
	0 cm	-10 cm	-20 cm	0 cm	-10 cm	-20 cm	0 cm	-10 cm	-20 cm
<i>Bare soil</i>									
26/07/2012	20.7	33.4	54.2	0.9	0.7	0.0	0.4	0.6	0.7
24/08/2012	6.6	10.0	56.9	1.4	2.0	0.1	0.2	0.1	2.0
26/09/2012	41.7	13.5	52.9	3.4	4.3	0.3	0.1	<0.1	1.8
05/11/2012	2.7	1.8	46.2	3.6	3.5	0.7	<0.1	0.1	0.5
30/11/2012	2.1	4.7	41.9	4.0	3.2	0.8	<0.1	<0.1	0.1
04/01/2013	2.3	4.0	29.7	2.9	3.8	0.4	0.1	0.3	1.0
06/05/2013	1.3	1.4	22.7	5.1	4.6	0.7	0.7	0.1	0.6
11/06/2013	2.0	2.1	18.1	5.8	3.8	0.2	<0.1	<0.1	0.6
16/07/2013	1.3	1.4	17.2	10.2	5.9	0.6	<0.1	<0.1	2.5
<i>RCG</i>									
26/07/2012	10.4	11.4	9.9	2.9	1.2	2.0	2.9	0.4	0.1
24/08/2012	3.8	2.4	9.2	0.2	0.1	0.5	<0.1	<0.1	0.1
26/09/2012	2.0	10.6	3.9	0.6	0.2	0.9	0.1	0.1	0.1
05/11/2012	5.6	3.6	2.3	0.8	0.1	1.0	<0.1	<0.1	<0.1
30/11/2012	4.1	3.2	3.9	0.6	0.4	0.8	0.5	0.2	0.1
04/01/2013	2.3	3.5	5.3	1.5	0.1	0.7	0.1	0.5	0.1
06/05/2013	2.0	1.8	3.5	1.7	0.4	0.2	<0.1	3.1	<0.1
11/06/2013	3.4	4.5	4.7	0.3	0.8	0.1	<0.1	0.1	0.3
16/07/2013	3.3	1.8	10.6	1.5	1.5	0.1	0.1	<0.1	0.3

Table 2. Statistical main effects of vegetation (i.e., reed canary grass cultivation or bare soil), ground water level (GWL), and date on fluxes of CO₂, CH₄ and N₂O as explored with linear mixed models.

Variables	CO ₂			CH ₄			N ₂ O		
	<i>DF</i>	<i>F</i> value	<i>P</i> value	<i>DF</i>	<i>F</i> value	<i>P</i> value	<i>DF</i>	<i>F</i> value	<i>P</i> value
Vegetation	1	956.2	<0.001	1	165.8	<0.001	1	0.5	<0.001
GWL	2	32.2	<0.001	2	15.4	<0.001	2	3.1	0.02
Date	28	75.6	<0.001	28	25.8	<0.001	28	<0.1	<0.001

Table 3. Parameter estimates (b_1 , b_2 , b_3 and b_4) for CO₂ flux (ecosystem respiration) models.

Uncertainties shown in parentheses are standard error of parameter estimates. Also shown are correlation coefficients (r) between observed and modelled data and modeling efficiencies (ME).

Treatment	CO ₂ flux model	b_1 (mg CO ₂ m ⁻² h ⁻¹)	b_2 (mg CO ₂ m ⁻² h ⁻¹ cm ⁻¹)	b_3 (K)	b_4	r	ME
Reed canary grass	Model 1	49.6 (3.8)	0.4 (0.1)	259.1 (15.5)	5.0 (0.7)	0.90	0.82
Bare soil	Model 2	79.1 (6.9)	5.7 (0.5)	286.4 (24.1)	na	0.68	0.46

na, not applicable

Figure captions

Fig. 1. (a) Hourly air temperature at 2 m height at the semi-field facility and hourly average soil temperature at 5 cm depth across all mesocosm treatments, and (b) daily precipitation at the semi-field facility during the study period (July 2012 to July 2013).

Fig. 2. Redox potential (Eh) at different ground water levels (GWL) from reed canary grass (RCG) and bare soil mesocosms. Eh was measured at 20 cm soil depth from April to July 2013.

Fig. 3. Mean dry biomass yield (Mg ha^{-1}) from mesocosms at different ground water level in 2012 and 2013. Error bars show standard error ($n = 5$).

Fig. 4. Average ratio vegetation index (RVI) development during the measurement period across all ground water levels. Error bars show standard error ($n = 15$). Dotted line represent the winter period when RVI was not measured due to ice and snow.

Fig. 5. Time course of greenhouse gas fluxes from the rewetted peat soil mesocosms during July 2012 to July 2013 in treatments with RCG cultivation (left panels) and bare soil (right panels). Data are shown for (a, b) CO_2 fluxes from ecosystem respiration, (c, d) CH_4 fluxes, and (e, f) N_2O fluxes. All data are mean and standard error of five replicates from each of the three ground water levels (GWL) at 0, -10 and -20 cm. Arrows marked by H indicate the times of harvest and arrows marked by F indicate the times of mineral fertilization.

Fig. 6. Annual fluxes of (a) CO_2 from ecosystem respiration, (b) CH_4 , (c) N_2O and (d) total global warming potential (Σ GWP) from the rewetted peat soil mesocosms from July 2012 to July 2013 in treatments with RCG cultivation (gray bars) and bare soil (white bars) at ground water levels of 0, -

10 and -20 cm. Error bars for CO₂ data show the standard error (SE) derived from SE of model parameters. For CH₄ and N₂O, data are shown as mean and SE of individual mesocosms ($n = 5$). Numbers in parentheses indicate the contribution of RCG in total emission at the different GWLs.

Fig. 7. Plant-derived CO₂, CH₄ and N₂O emissions ~~for RCG treatments~~ at different ground water levels as compared in terms of CO₂ equivalents (CO₂ eq.). Plant derived emissions were estimated as the difference between the total emissions from RCG treatments and bare soil treatments.




Fig. 1.




Fig. 2.





Fig. 3.






Fig. 5.




Fig. 6.




Fig. 6.




Fig. 7.