

Technical Note: Hyperspectral Lidar Time Series of Pine Canopy Chlorophyll Content

T. Hakala¹, O. Nevalainen¹, S. Kaasalainen¹, and R. Mäkipää²

[1]{Finnish Geodetic Institute, Masala, Finland}

[2]{Finnish Forest Research Institute (METLA), Vantaa, Finland}

Correspondence to: T. Hakala (Teemu.Hakala@nls.fi)

Abstract

We present an empirical application of hyperspectral lidar for monitoring the seasonal and spatial changes in pine chlorophyll content and upscaling the accurate leaf-level chlorophyll measurements into branch and tree level. The results show the capability of the new instrument for monitoring the changes in the shape and physiology of tree canopy: the spectral indices retrieved from the hyperspectral point cloud agree with laboratory measurements of the chlorophyll content. The approach opens new prospects for replacing destructive and labor-intensive manual sampling with remote observations of tree physiology.

1 Introduction

The photosynthetic activity in tree canopy is an indicator of tree health. Vigorous trees with high foliar biomass and chlorophyll content have high carbon assimilation capacity. Stress in vegetation has been shown to induce changes in the photosynthetically-active pigments such as chlorophyll a and b. Therefore, the leaf chlorophyll content is an important indicator of the photosynthetic capacity as well as tree productivity and stress (Coops et al., 2003, Lausch et al. 2013).

The leaf properties and the distribution of chlorophyll and nutrients within a canopy vary as a function of time and space, and depending on the resource availability (Wang and Schjoerring, 2012, Peltoniemi et al., 2012). Plant phenology and seasonal chlorophyll content cycle are correlated to the CO₂ flux. For monitoring these seasonal variations, methods are needed for accurate and nondestructive chlorophyll estimation, both at the leaf and canopy level (e.g., Gond et al., 1999). Chlorophyll estimation with spectral remote-sensing has been

implemented increasingly in a number of studies (e.g., Coops et al., 2003, Lausch et al., 2013), but improved resolution and more accurate 3D position for the spectra are still being called for, to extend the accurate leaf-level measurement into canopy and stand level (cf. Gaulton et al., 2013). To investigate the spatial variation of the photosynthetic capacity and self-shading of photosynthetically active tissue, the canopy and branch structure must also be included in the measurement.

One way to provide simultaneous structural and spectral information is lidar combined with hyperspectral passive sensing (e.g., Thomas et al., 2006, Asner et al., 2007, Jones et al., 2010), but new applications using multi or hyperspectral laser scanning have increased quite recently. Hancock et al., (2012) demonstrated the potential of dual wavelength, large-footprint, spaceborne lidar to separate ground and canopy returns using the extra information contained in a spectral ratio to complement the canopy height from laser scanning. Three-dimensional (3D) distributions of vegetation biochemical properties were measured with spectral indices developed for the Salford Advanced Laser Canopy Analyser (SALCA), which is also a dual-wavelength lidar (Gaulton et al., 2013). A similar approach was used in the Dual-Wavelength Echidna Lidar (DWEL) (Douglas et al., 2012). A multispectral canopy lidar has also been introduced for simultaneous retrieval of vegetation structure and spectral indices (Woodhouse et al., 2011). In this approach, a tunable laser operating at four wavelengths was used. The limitation of empirical vegetation indices estimating chlorophyll content is that they are also affected by the canopy structural properties. In addition, they can be affected by the internal structure, size, surface and shape of leaves and can thus be species-specific, requiring calibration when applied to specific species (Zhang et al., 2008).

In this technical note, an application of the recently developed hyperspectral lidar instrument (Hakala et al., 2012) is presented for monitoring the seasonal and spatial changes in pine chlorophyll content. As a non-destructive method, the capability of the instrument to upscale the accurate leaf-level chlorophyll content measurements into branch and tree level has been investigated and validated with chemical analysis of chlorophyll content. In this study we used three spectral indices showing good correlation with Scots Pine shoot chlorophyll concentration using the HSL instrument in Nevalainen et al. (2014).

2 Materials and methods

Hyperspectral lidar (HSL) is a prototype laser scanning instrument (Hakala et al., 2012) utilizing a supercontinuum laser. White laser pulses are transmitted to a target and the

distances of reflected echoes are determined from time of flight. A spectrograph and an avalanche photodiode (APD) array connected to a high-speed digitizer are used to determine the spectrum of each returning echo by measuring the intensity of the echo at multiple wavelengths. Also the intensity of each transmitted laser pulse is measured and used to normalize the echo intensity. Current prototype configuration uses a 16 element APD array and an 8 channel digitizer, enabling us to measure at 8 wavelength bands: 545, 641, 675, 711, 742, 778, 978, 1292 nm, full width at half maximum about 20 nm. Before the target is measured a reference target with known reflectance (Spectralon) is measured at distance intervals of about 30 cm and these data are used to calibrate the reflectance over the whole measurement range. Additionally the Spectralon is placed in the scanned area during the actual measurement to validate the calibration. The instrument and data processing presented in more detail in Hakala et al., 2012.

A Scots pine (*Pinus sylvestris* L.) was scanned five times during the 2013 growth season. The tree was approximately 13 years old, 5.5 m high and it was growing in a small forest stand near the institute building. The HSL was mounted on a portable cart, and the tree was scanned from two directions. The scans were co-registered using white spherical reference targets placed on fixed locations on the target area. The distance between the scanner and the tree was about 5 m. The tree was scanned with 0.1° horizontal and about 0.02° vertical resolution and the resulting point clouds contained 200 000- 470 000 echoes from the tree. The beam diameter at the target was about 5 mm.

Needle samples were taken immediately after the scan for laboratory analysis. Six branches were selected and the samples were taken from these branches according to needle cohorts (current year needles, and 1-, 2, and 3-year old needles). Two needle pairs were taken from each cohort of each selected branch. Analysis of the chlorophyll contents followed the protocol described in Wellburn (1994) for extraction with dimethyl-sulfoxide (DMSO). After extraction, the chlorophyll concentrations were determined from solvents spectrophotometrically using wave-lengths 480.0, 649.1 and 665.1 nm (resolution 0.1 – 0.5 nm).

Two of the six sampled branches were clearly identifiable from the HSL point cloud, having enough point density and long enough growth of the branch. Previous year cohorts were selected for further analysis, since they had needles present during all measurements. Therefore the following analysis is performed for two cohorts and five measurement dates.

The parts of the point cloud containing the selected cohorts were isolated in post processing. Three spectral indices were tested for determining chlorophyll content of the needles. Since it was not possible to tune all required wavelengths to optimal positions for every index, we used the nearest available band.

The Modified Chlorophyll Absorption Ratio Index using reflectance at 705 and 750 nm (referred here as MCARI750) was first presented by Wu et al. (2008). Contrary to the original MCARI (Daughtry et al. 2000), MCARI750 uses reflectance at 705 and 750 nm, which have shown better sensitivity to high chlorophyll contents (Wu et al. 2008). MCARI has been designed to measure the depth of the maximum chlorophyll absorption at 670 nm relative to green reflectance peak at 550 nm and reflectance at 700 nm, at canopy scale (Daughtry et al., 2000).

$$MCARI750 = [(R_{750} - R_{705}) - 0.2 * (R_{750} - R_{550})] * (R_{750}/R_{705}) \quad (1)$$

The Modified Simple Ratio (MSR), developed by Chen (1996), strives to have low noise effect and good linearity to vegetation biophysical parameters. MSR has been used to estimate chlorophyll and Leaf Area Index (LAI) at canopy scale. Wu et al. (2008) also developed MSR using reflectance at 705 and 750 nm, referred here as MSR2.

$$MSR2 = \frac{R_{750}/R_{705}-1}{\sqrt{R_{750}/R_{705}+1}} \quad (2)$$

The Simple Ratio (SR) indices directly compare the reflectance and absorbance peaks of chlorophyll pigments, which make them sensitive to changes in chlorophyll content (Wu et al., 2008). Variety of wavelength combinations are used with simple ratio indices, but the one selected for this study is SR6 (Zarco-Tejada et al., 2001). It has been used to estimate chlorophyll at leaf level.

$$SR6 = \frac{R_{750}}{R_{710}} \quad (3)$$

Additionally, normalised difference vegetation index (NDVI) (Rouse et al., 1973) was used to separate needles from branches. NDVI is the most widely used vegetation index. It is based on the contrast between high absorption at red and high reflectance at near-infrared (NIR). NDVI has been developed for canopy scale and it has been used for both chlorophyll and LAI estimation.

$$NDVI = \frac{R_{800}-R_{670}}{R_{800}+R_{670}} \quad (4)$$

As the channels of the prototype HSL are limited to eight separate spectral bands, these indices had to be used with the closest available spectral band.

3 Results

The overall shape of the tree and changes in shape from May to November can be observed in Figure 1 where no spectral information is used. The changes in the shape and the spectra of tree parts are visible in the spectral point clouds. To demonstrate this, we plot the time series of the NDVI over the pine branch from May 15 to Nov 6, 2013 in Figure 2. The outbreak and growth of new shoots (May/Jun 2013) can be observed, as well as the year 2 parts defoliating (Sep/Oct 2013) and falling off completely (Nov 2013).

To validate the capability of the HSL to estimate the chlorophyll content using spectral indices, we compared the HSL data with laboratory analysis over the growing season. We present data for two branch cohorts, denoted M2_1 and M3_1 (one year old part of M2 and M3), which were best visible in the HSL point clouds. The trends in the chlorophyll content and the indices MCARI750, MSR2, and SR6 from HSL data are well reproduced for the individual branches (Figures 3-5). For all three indices, the sample branch M2_1 was best correlated with the laboratory measurements with R^2 0.8-0.9. The R^2 for MCARI750 and MSR2 for M3_1 was 0.7, whereas SR6 performed worse for M3_1 (R^2 0.4). When the data from M2_1 and M3_1 were combined for regression all indices correlated with the chlorophyll content measured in the laboratory (Figure 6). The results were worse for indices averaged over the entire tree point cloud (the right column in Figures 3-5), compared with the average of all year 1 needles measured in the laboratory. This is very likely a result of the variation of the physiological conditions between the tree parts, which is more pronounced when the sampling has been carried out over the entire tree (i.e., the point cloud), rather than just a few needle samples (as in the laboratory experiment). All in all, the analysis of branch parts shows that the spatial distribution of the HSL spectral indices describes the chlorophyll content within the branch, although more measurements are needed to better validate the results.

In figures 3-5, branch M2_1 and M3_1 laboratory measurements consist of two separate needles only. More sampling should have been performed, however, the number of needles in each branch part is limited and the tree had to be sampled several times during the year (this emphasizes the need for non-destructive methods). The number of laser echoes from year 0 and 2 were highly varying; in the spring lidar point clouds the year 0 growths were very small

providing very few echoes. The year 2 and older cohorts started dropping needles before September measurement thus reducing the number of echoes during autumn compared to spring. Therefore we only used year 1 laboratory measurement of needles in plots 3-5 for whole tree (right column), since the weight of the year 0 and 2 laboratory measurements would have been higher compared to the lidar point cloud (lidar point density variable and laboratory sample number constant). Some lidar echoes still originate from the year 0 and 2 needles, reducing the overall correlation between laboratory and lidar data for the whole tree.

The change in the shape of the tree point cloud is visible in Figure 1. The fact that tree shape can be retrieved from HSL point clouds has been shown before in numerous studies (see Kaasalainen et al., 2014 and Refs. therein). We have also shown in our previous study that the tree shape and its changes can be quantified from laser scanner point clouds using quantitative tree structure modelling (Kaasalainen et al., 2014). As the scope of this note was to show the added value of spectral data in the chlorophyll distribution monitoring, the changes in tree shape will be an object of our future study.




Figure 1. Co-registered point clouds from 2013-05-15 scan (grey) and 2013-11-06 scan (red). Growth of the tree is visible and also some movement of the branches can be observed. The height of the tree is about 5.5 m.





Figure 2. NDVI (see the colour bar for values) point clouds of a sample branch M2. The growth of new needles (starting 05-27), already clearly visible new branch tips 06-19, fully grown new needles 09-12 and dying and falloff of old needles (shown in bluish green, low NDVI, colours in 09-12 and 10-03) are visible in the data measured at different times. The measurement dates are shown in the plot titles.



179

180 Figure 3. Top row: distribution of MCARI750 spectral index during separate HSL
 181 measurements, the central mark is the median, the edges of the box are the 25th
 182 percentiles, the whiskers extend to the most extreme data points not considered outliers.
 183 Middle row: Laboratory measurements chlorophyll a+b. Bottom row: Correlation of the
 184 spectral index and laboratory measurement. Subplot columns left to right: sample branch 3
 185 year 1, sample branch 2 year 1, spectral index of whole tree and laboratory measurements of
 186 all year 1 samples.




Figure 4. Same as previous figure (top and bottom rows, laboratory data is the same as in previous figure), this time using MSR2 spectral index.




Figure 5. Same as previous figure, this time using SR6 spectral index.




Figure 6. Correlation of spectral index and laboratory measurement for combined M2_1 and M3_1 data. Left: MSR2, middle: MCARI750, right: SR6. Blue x: M3_1, red circle: M2_1.

4 Conclusions and discussion

We have shown that the hyperspectral lidar provides an empirical approach for efficient mapping the spatial distributions of tree physiological parameters that are correlated to reflectance of the foliage (such as chlorophyll a and b). Because the measurement is non-destructive, it can be repeated for the same target to produce time series of important tree functions, such as moisture condition, photosynthetic capacity, or physiological status.

We demonstrated that the seasonal changes in the shape and physiology of tree parts are visible in 3D; parameters affecting tree physiology can be quantified with spectral indices and linked to a specific location in the tree canopy using the HSL point cloud. We validated the method with reference measurements of chlorophyll a and b concentration in a laboratory. According to our results hyperspectral lidar can be used for the monitoring of the chlorophyll content, but similarly, the approach has potential in the monitoring of the water, carotenoid or lignin content, which all affect reflectance of the foliage (Austin and Ballare 2010).

The benefit of active measurement system, such as HSL, is that they measure backscattered signal that has the potential to eliminate many of the multiple scattering and geometric viewing effects caused by the canopy structure (Gaulton et al., 2013; Morsdorf et al., 2009). The major factors affecting the backscattered signal are the local incidence angle of the target and the area of effective backscattering surface (Gaulton et al., 2013). These factors are also present in this study as one 5mm footprint may include one or several needles with varying incidence angles. However, the influence of these factors is similar with different wavelengths measured at the same optical path. Thus by calculating spectral ratios (i.e. vegetation indices), the influence of the incidence angle and target area can be reduced (Eitel et al., 2011; Gaulton et al., 2013).

However, the influence of multiple scattering effects to the measured backscattered reflectance is not completely removed. Further study would be required to produce physically based model that would properly account for the multiple scattering of needles within single laser footprint and its effect to the measured backscattered reflectance. Some of the limitations of vegetation indices in chlorophyll estimation could be overcome by using inversion of radiative transfer models, such as LIBERTY (Leaf Incorporating Biochemistry

Exhibiting Reflectance and Transmittance Yields) (Dawson et al., 1998) which is specifically developed for needles, or PROSPECT model (Féret et al., 2011).

The tree was scanned from two directions only. Increasing the number of scans from different directions around the tree will improve the results by increasing the point coverage. This will require some instrument development to allow a more efficient field use. Increasing the point density is also an important object of instrument improvement. However, the prototype instrument was capable of showing the potential of 3D spectral measurements.

A major factor causing error and uncertainty in this research was the use of nearest possible channel in vegetation index calculation instead of the band the index was designed to use. Especially close to the vegetation red-edge region even small shift in channel wavelength causes high change in reflectance. This affects the performance of the vegetation indices, especially with indices requiring channels at red edge. However, this was not considered as a major problem as the aim of this study was to test the ability of the HSL in chlorophyll estimation and not to optimize the performance of the indices.

Further work is needed to find the best spectral indices for different applications (e.g., monitoring the effects of drought or limited amount of light on the physiology of different tree parts), and then optimize the spectral channels to match with these indices. This will improve the precision of the results. Increasing the number of spectral channels would also improve the channel optimization and efficiency. Once the approach is well established and calibrated, it has potential for replacing a number of laborious and destructive manual experiments, and hence providing a new tool for remote observations of tree physiology. Although the first results show the potential of the approach, further studies on the laser interaction with the canopy are needed to establish the method physically.

Acknowledgements

This study was funded by the Academy of Finland research projects “New techniques in active remote sensing: hyperspectral laser in environmental change detection” and “Mobile hyperspectral laser remote sensing”.

References

- Asner, G. P.; Knapp, D. E.; Kennedy-Bowdoin, T.; Jones, M. O.; Martin, R. E. et al. Carnegie Airborne Observatory: in-flight fusion of hyperspectral imaging and waveform light detection and ranging for three-dimensional studies of ecosystems, *Journal of Applied Remote Sensing*, 1(1), 013536; doi:10.1117/1.2794018, 2007.
- Austin, A. T. and Ballaré, C. S.: Dual role of lignin in plant litter decomposition in terrestrial ecosystems. *PNAS* 107(10), 4618–4622, doi: 10.1073/pnas.0909396107.
- Chen, J., Evaluation of vegetation indices and modified simple ratio for boreal applications. *Canadian Journal of Remote Sensing* 22, 229–242, 1996.
- Coops, N. C., Stone, C., Culvenor, D. S., Chisholm, L. A. and Merton, R. N.: Chlorophyll content in eucalypt vegetation at the leaf and canopy scales as derived from high resolution spectral data, *Tree Physiology*, 23(1), 23–31, doi:10.1093/treephys/23.1.23, 2003.
- Daughtry, C.: Estimating Corn Leaf Chlorophyll Concentration from Leaf and Canopy Reflectance, *Remote Sensing of Environment*, 74(2), 229–239, doi:10.1016/S0034-4257(00)00113-9, 2000.
- Dawson, T.P., Curran, P.J., Plummer, S.E.: LIBERTY—modeling the effects of leaf biochemical concentration on reflectance spectra. *Remote Sensing of Environment*, 65 (1), 50–60, 1998.
- Douglas, E. S., Strahler, A., Martel, J., Cook, T., Mendillo, C., Marshall, R., Chakrabarti, S., Schaaf, C., Woodcock, C., Li, Z., Yang, X., Culvenor, D., Jupp, D., Newnham, G. and Lovell, J.: DWEL: A Dual-Wavelength Echidna Lidar for ground-based forest scanning, pp. 4998–5001, *IEEE*, 2012.
- Eitel, J.U.H., Vierling, L.A., Long, D.S., Hunt, E.R.: Early season remote sensing of wheat nitrogen status using a green scanning laser. *Agricultural and Forest Meteorology*, 151, 1338–1345, doi:10.1016/j.agrformet.2011.05.015, 2011.
- Féret, J. B., François, C., Gitelson, A., Asner, G., P., Barry, K., M., Panigada, C., Richardson, A., D, Jacquemoud, S.: Optimizing spectral indices and chemometric analysis of leaf chemical properties using radiative transfer modeling, *Remote Sensing of Environment*, 115(10), 2742–2750, doi:10.1016/j.rse.2011.06.016, 2011.

281 Gaulton, R., Danson, F. M., Ramirez, F. A. and Gunawan, O.: The potential of dual-
 282 wavelength laser scanning for estimating vegetation moisture content, *Remote Sensing of*
 283 *Environment*, 132, 32–39, doi:10.1016/j.rse.2013.01.001, 2013.

284 Gond, V., de Pury, D. G. G., Veroustraete, F. and Ceulemans, R.: Seasonal variations in leaf
 285 area index, leaf chlorophyll, and water content; scaling-up to estimate fAPAR and carbon
 286 balance in a multilayer, multispecies temperate forest, *Tree Physiology*, 19(10), 673–679,
 287 doi:10.1093/treephys/19.10.673, 1999.

288 Hakala, T., Suomalainen, J., Kaasalainen, S. and Chen, Y.: Full waveform hyperspectral
 289 LiDAR for terrestrial laser scanning, *Optics Express*, 20(7), 7119,
 290 doi:10.1364/OE.20.007119, 2012.

291 Hancock, S., Lewis, P., Foster, M., Disney, M. and Muller, J.-P.: Measuring forests with dual
 292 wavelength lidar: A simulation study over topography, *Agricultural and Forest Meteorology*,
 293 161, 123–133, doi:10.1016/j.agrformet.2012.03.014, 2012.

294 Jones, T. G., Coops, N. C. and Sharma, T.: Assessing the utility of airborne hyperspectral and
 295 LiDAR data for species distribution mapping in the coastal Pacific Northwest, Canada,
 296 *Remote Sensing of Environment*, 114(12), 2841–2852, doi:10.1016/j.rse.2010.07.002, 2010.

297 Kaasalainen, S., Krooks, A., Liski, J., Raunonen, P., Kaartinen, H., Kaasalainen, M.,
 298 Puttonen, E., Anttila, K., and Mäkipää, R., Change Detection of Tree Biomass with Terrestrial
 299 Laser Scanning and Quantitative Structure Modelling. *Remote Sensing* 6(5), 3906–3922.
 300 doi:10.3390/rs6053906, 2014

301 Lausch, A., Heurich, M., Gordalla, D., Dobner, H.-J., Gwilym-Margianto, S. and Salbach,
 302 C.: Forecasting potential bark beetle outbreaks based on spruce forest vitality using
 303 hyperspectral remote-sensing techniques at different scales, *Forest Ecology and Management*,
 304 308, 76–89, doi:10.1016/j.foreco.2013.07.043, 2013.

305 Morsdorf, F., Nichol, C., Malthus, T., Woodhouse, I.H.: Assessing forest structural and
 306 physiological information content of multi-spectral LiDAR waveforms by radiative transfer
 307 modelling. *Remote Sensing of Environment*, 113 (10), 2152–2163,
 308 doi:10.1016/j.rse.2009.05.019, 2009.

309 Nevalainen, O., Hakala, T., Suomalainen, J., Mäkipää, R., Peltoniemi, M., Krooks, A.,
 310 Kaasalainen, S.: Fast and nondestructive method for leaf level chlorophyll estimation using

311 hyperspectral LiDAR, *Agricultural and Forest Meteorology*, 198–199, 250–258,
 312 doi:10.1016/j.agrformet.2014.08.018, 2014.

313 Peltoniemi, M. S., Duursma, R. A. and Medlyn, B. E.: Co-optimal distribution of leaf nitrogen
 314 and hydraulic conductance in plant canopies, *Tree Physiology*, 32(5), 510–519,
 315 doi:10.1093/treephys/tps023, 2012.

316 Rouse, J.W., Haas, R.H., Schell, J.A., Deering, W.D., Monitoring vegetation systems in the
 317 Great Plains with ERTS. In: Third ERTS Symposium, NASA SP-351, 309–317, 1973

318 Thomas, V., Finch, D. A., McCaughey, J. H., Noland, T., Rich, L. and Treitz, P.: Spatial
 319 modelling of the fraction of photosynthetically active radiation absorbed by a boreal
 320 mixedwood forest using a lidar–hyperspectral approach, *Agricultural and Forest Meteorology*,
 321 140(1-4), 287–307, doi:10.1016/j.agrformet.2006.04.008, 2006.

322 Wang, L. and Schjoerring, J. K.: Seasonal variation in nitrogen pools and $^{15}\text{N}/^{13}\text{C}$ natural
 323 abundances in different tissues of grassland plants, *Biogeosciences*, 9(5), 1583–1595,
 324 doi:10.5194/bg-9-1583-2012, 2012.

325 Wellburn, A. R.: The Spectral Determination of Chlorophylls a and b, as well as Total
 326 Carotenoids, Using Various Solvents with Spectrophotometers of Different Resolution,
 327 *Journal of Plant Physiology*, 144(3), 307–313, doi:10.1016/S0176-1617(11)81192-2, 1994.

328 Woodhouse, I. H., Nichol, C., Sinclair, P., Jack, J., Morsdorf, F., Malthus, T. J. and
 329 Patenaude, G.: A Multispectral Canopy LiDAR Demonstrator Project, *IEEE Geoscience and*
 330 *Remote Sensing Letters*, 8(5), 839–843, doi:10.1109/LGRS.2011.2113312, 2011.

331 Wu, C., Niu, Z., Tang, Q. and Huang, W.: Estimating chlorophyll content from hyperspectral
 332 vegetation indices: Modeling and validation, *Agricultural and Forest Meteorology*, 148(8-9),
 333 1230–1241, doi:10.1016/j.agrformet.2008.03.005, 2008.

334 Zarco-Tejada, P. J., Miller, J. R., Noland, T. L., Mohammed, G. H. and Sampson, P. H.:
 335 Scaling-up and model inversion methods with narrowband optical indices for chlorophyll
 336 content estimation in closed forest canopies with hyperspectral data, *IEEE Transactions on*
 337 *Geoscience and Remote Sensing*, 39(7), 1491–1507, doi:10.1109/36.934080, 2001.

338 Zhang, Y., Chen, J., Miller, J., Noland, T.: Leaf chlorophyll content retrieval from airborne
 339 hyperspectral remote sensing imagery. *Remote Sensing of Environment*, 112 (7), 3234–3247,
 340 doi:10.1016/j.rse.2008.04.005, 2008.