

This discussion paper is/has been under review for the journal Biogeosciences (BG).  
Please refer to the corresponding final paper in BG if available.

# Sensitivity of the air–sea CO<sub>2</sub> exchange in the Baltic Sea and Danish inner waters to atmospheric short term variability

A. S. Lansø<sup>1</sup>, J. Bendtsen<sup>2</sup>, J. H. Christensen<sup>1,3</sup>, L. L. Sørensen<sup>1,3</sup>, H. Chen<sup>4,5</sup>,  
H. A. J. Meijer<sup>4</sup>, and C. Geels<sup>1</sup>

<sup>1</sup>Department of Environmental Science, Aarhus University, 4000 Roskilde, Denmark

<sup>2</sup>Climate Lab, Symbion Science Park, Fruebjergvej 3, 2100 Copenhagen, Denmark

<sup>3</sup>Arctic Research Centre (ARC), Aarhus University, 8000C Aarhus, Denmark

<sup>4</sup>Centre for Isotope Research (CIO), Energy and Sustainability Research Institute Groningen (ESRIG), University of Groningen, Groningen, the Netherlands

<sup>5</sup>Cooperative Institute for Research in Environmental Sciences (CIRES), University of Colorado, Boulder, CO, USA

Received: 4 September 2014 – Accepted: 4 November 2014 – Published: 9 December 2014

Correspondence to: A. S. Lansø (asla@envs.au.dk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

## Abstract

Minimising the uncertainties in estimates of air-sea  $\text{CO}_2$  exchange is an important step toward increasing the confidence in assessments of the  $\text{CO}_2$  cycle. Using an atmospheric transport model makes it possible to investigate the direct impact of atmospheric parameters on the air-sea  $\text{CO}_2$  flux along with its sensitivity to e.g. short-term

5 temporal variability in wind speed, atmospheric mixing height and the atmospheric  $\text{CO}_2$  concentration. With this study the importance of high spatiotemporal resolution of atmospheric parameters for the air-sea  $\text{CO}_2$  flux is assessed for six sub-basins within the Baltic Sea and Danish inner waters. A new climatology of surface water partial pressure of  $\text{CO}_2$  ( $p\text{CO}_2$ ) has been developed for this coastal area based on available data from monitoring stations and underway  $p\text{CO}_2$  measuring systems. Parameterisations depending on wind speed were applied for the transfer velocity to calculate the air-sea  $\text{CO}_2$  flux. Two model simulations were conducted – one including short term variability in atmospheric  $\text{CO}_2$  (VAT), and one where it was not included (CAT).

15 A seasonal cycle in the air-sea  $\text{CO}_2$  flux was found for both simulations for all sub-basins with uptake of  $\text{CO}_2$  in summer and release of  $\text{CO}_2$  to the atmosphere in winter. During the simulated period 2005–2010 the average annual net uptake of atmospheric  $\text{CO}_2$  for the Baltic Sea, Danish Straits and Kattegat was 287 and 471  $\text{Gg C yr}^{-1}$  for the VAT and CAT simulations, respectively. The obtained difference of 184  $\text{Gg C yr}^{-1}$  was 20 found to be significant, and thus ignoring short term variability in atmospheric  $\text{CO}_2$  does have a sizeable effect on the air-sea  $\text{CO}_2$  exchange. The combination of the atmospheric model and the new  $p\text{CO}_2$  fields has also made it possible to make an estimate of the marine part of the Danish  $\text{CO}_2$  budget for the first time. A net annual uptake of 2613  $\text{Gg C yr}^{-1}$  was found for the Danish waters.

25 A large uncertainty is connected to the air-sea  $\text{CO}_2$  flux in particular caused by the transfer velocity parameterisation and the applied  $p\text{CO}_2$  climatology. However, the present study underlines the importance of including short term variability in the atmo-

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

spheric CO<sub>2</sub> concentration in future model studies of the air–sea exchange in order to minimise the uncertainty.

## 1 Introduction

The capacity of ocean and land to take up and re-emit atmospheric CO<sub>2</sub> has a dominating effect on the greenhouse gas balance, and hence changes in climate. Currently, the land areas and the global oceans are estimated to take up about 27 and 28 %, respectively, of the CO<sub>2</sub> emitted by anthropogenic sources (Le Quéré et al., 2013).

In recent years the biogeochemical active coastal seas have been given increased attention (Borges et al., 2006; Chen et al., 2013; Mørk et al., 2014). Although such coastal waters only amount to 7 % of the global oceans, high inputs, production, degradation and export of organic matter might result in coastal air–sea CO<sub>2</sub> fluxes contributing a great deal more than 7 % to the global air–sea flux (Gattuso et al., 1998). Due to the high heterogeneity of these areas, coastal CO<sub>2</sub> fluxes are prone to large uncertainties. Several studies agree that continental shelves, in general, act as sinks, while estuaries act as sources of CO<sub>2</sub> to the atmosphere. However, global estimates vary in size according to applied methodology, with oceanic uptake in shelf areas between 0.21 and 0.40 PgCyr<sup>-1</sup>, and release from estuaries in the range of 0.10 to 0.50 PgCyr<sup>-1</sup> (Cai, 2011; Chen et al., 2013; Chen and Borges, 2009; Laruelle et al., 2010). The poor coverage of observations in both space and time makes validation of these global estimates difficult.

In order to better quantify the impact of coastal regions on the global carbon budget, detailed studies of the processes at the regional scale are necessary (Kulinski and Pempkowiak, 2011). A coastal region that has been well studied is the Baltic Sea. The Baltic Sea is a high latitude inner shelf sea connected to the North Sea through the shallow transition zone of the Danish Straits, and enclosed by land with various terrestrial ecosystems and densely populated areas. Seasonal amplitudes of up to 400  $\mu\text{atm}$  are observed in the partial pressure of CO<sub>2</sub> ( $p\text{CO}_2$ ) in the Baltic Sea (Thomas and Schnei-

BGD

11, 16993–17042, 2014

Sensitivity of the air–sea CO<sub>2</sub> exchange in the Baltic Sea and Danish inner waters

A. S. Lansø et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



## Sensitivity of the air-sea CO<sub>2</sub> exchange in the Baltic Sea and Danish inner waters

A. S. Lansø et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

der, 1999) with maximum values of  $p\text{CO}_2$  found in winter and minimum during summer. Since the difference between the  $p\text{CO}_2$  level in the ocean and the atmosphere controls the direction of the air–sea CO<sub>2</sub> flux, this is an indication of the pronounced seasonal variation of the flux in the Baltic Sea, with outgassing of CO<sub>2</sub> to the atmosphere during winter and uptake during summer (Thomas et al., 2004; Thomas and Schneider, 1999). Despite numerous studies, it is still uncertain, whether the Baltic Sea currently acts as a net sink or source of atmospheric CO<sub>2</sub>, as previous studies have given ambiguous results varying from  $-4.3$  to  $2.7 \text{ g C m}^{-2} \text{ yr}^{-1}$  for the entire Baltic Sea region (Gustafsson et al., 2014; Kulinski and Pempkowiak, 2011; Norman et al., 2013). Thereby, it is also difficult to project how the Baltic Sea will contribute to the global carbon budget in the future. Moreover, the region may possibly have changed from being a net source to a net sink of atmospheric CO<sub>2</sub>, due to the industrialization and the enormous input of nutrients (Omstedt et al., 2009). These inputs will, however, likely change in the future due to changes in climate and anthropogenic activities (Geels et al., 2012; Langner et al., 2009).

As the Baltic Sea is bordered by land areas, the atmospheric CO<sub>2</sub> concentration found here will be directly affected by continental air leading to a greater temporal and spatial variability in the CO<sub>2</sub> level, than what is found over open oceans. The impact of temporal variations in atmospheric CO<sub>2</sub> on the air–sea CO<sub>2</sub> exchange has been discussed by Rutgersson et al. (2008, 2009). They show an overestimation in the amplitude of the seasonal cycle for calculated air–sea CO<sub>2</sub> fluxes, when using a constant annual mean value of atmospheric CO<sub>2</sub> concentration instead of daily levels of the atmospheric concentration. Annually, the difference was less than 10 % between the two cases, but weekly flux deviations of 20 % were obtained. This indicates how synoptic variability in the atmosphere not always can be ignored (Rutgersson et al., 2009). Further, Rutgersson et al. (2008) note that the uncertainties connected with the transfer velocity are much greater. However, it is still worthwhile to minimise the bias in the estimation of the flux by including detailed information of the atmospheric CO<sub>2</sub> concentration. The short term variability (hourly) of both meteorology and atmospheric

CO<sub>2</sub> concentrations is not always accounted for or has not been discussed in previous estimates of the air–sea CO<sub>2</sub> fluxes in the Baltic Sea (Algesten et al., 2006; Gustafsson et al., 2014; Kulinski and Pempkowiak, 2011; Löffler et al., 2012; Norman et al., 2013; Wesslander et al., 2010).

5 To enlighten this matter, the present study aims to determine the importance of the short-term variability in atmospheric CO<sub>2</sub> concentrations for the net air–sea CO<sub>2</sub> flux. The Baltic Sea and Danish inner waters (consisting of Kattegat and the Danish straits; Øresund and the Belt Seas) are chosen as focus areas for the present study. A modelling approach is applied, which includes both short-term (hourly to synoptic) and 10 long-term (seasonal to inter-annual) variability in the atmospheric CO<sub>2</sub> concentrations. The analysis is carried out by constructing a meso-scale model framework based on an atmospheric transport model covering the study region with high resolutions in both space and time. The model includes a new spatial  $p\text{CO}_2$  climatology developed especially for the investigated marine area, as existing climatologies do not cover this area. 15 The advantages of the present study are that the same and consistent method is applied to the entire Baltic Sea and Danish inner waters, and that the impact of spatial and temporal short term variability in atmospheric parameters will be investigated in more details than in the previous studies of this region.

Recently, national CO<sub>2</sub> budgets that include both anthropogenic and natural components have been estimated in various countries (Meesters et al., 2012; Smallman et al., 2014). The present study is likewise part of a national project: Ecosystem Surface Exchange of Greenhouse Gases in an Environment of Changing Anthropogenic and Climate forcing (ECOCLIM) that is to determine the CO<sub>2</sub> budget for Denmark. For that reason the present study will also estimate the marine component of the Danish 20 CO<sub>2</sub> budget.

In Sect. 2 the study area, the applied surface fields of  $p\text{CO}_2$  and the model framework are described. Results are presented in Sect. 3, leading to a discussion in Sect. 4, and concluding remarks in Sect. 5.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## 2 Study setup

### 2.1 Study area

The marine areas investigated in this study comprise of the Baltic Sea, Kattegat and the Danish Straits, with the two latter also referred to as the Danish inner waters (see 5 Fig. 1). In the following a short introduction to these heterogeneous marine areas is given, as well as a description of the overall atmospheric CO<sub>2</sub> field in the region.

The Baltic Sea is a semi-enclosed continental shelf sea area with a large volume of river runoff adding a substantial amount of nutrients and terrestrial carbon to the Baltic Sea (Kulinski and Pempkowiak, 2011). The circulation in the Baltic Sea is influenced 10 by a relative large runoff from the surrounding drainage areas, and this causes a low-saline outflowing surface water mass from the area. The Baltic Sea can, therefore, be considered as a large estuary. Inflow of high-saline water from the North Sea ventilates the bottom waters of the Baltic Sea, and the exchange between these water masses occurs through the shallow North Sea/Baltic Sea transition zone centred around the 15 Danish Straits (Bendtsen et al., 2009). Ice coverage is observed in the northern part of Baltic Sea during winter (Löffler et al., 2012), which has implications for the air-sea exchange of CO<sub>2</sub>.

Atmospheric concentrations of CO<sub>2</sub> in the Baltic region have a greater seasonal amplitude than at e.g. Mauna Loa, Hawaii, which often is referred to as a global reference 20 for the atmospheric CO<sub>2</sub> background, due to the remoteness of the site. The larger seasonal amplitude over the Baltic can be explained by the difference in latitude between the studied area (54–66° N) and Mauna Loa (20° N), and the undisturbed air at the high altitude site of Mauna Loa compared to the semi-enclosed Baltic Sea (Rutgersson et al., 2009). The study by Rutgersson and colleagues also showed that the 25 atmospheric CO<sub>2</sub> concentration in the southern part of the Baltic Sea is more affected by regional anthropogenic and terrestrial sources and sinks, than the more remote northern part of the Baltic Sea area.

BGD

11, 16993–17042, 2014

Sensitivity of the air-sea CO<sub>2</sub> exchange in the Baltic Sea and Danish inner waters

A. S. Lansø et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

## 2.2 Surface water $p\text{CO}_2$ climatology

Model calculations of the surface air-sea gas exchange of  $\text{CO}_2$  are parameterised in terms of the difference in partial pressure of  $\text{CO}_2$  (i.e.  $\Delta p\text{CO}_2$ ) between the atmosphere and the ocean surface. The global climatology of oceanic surface  $p\text{CO}_2$  by Takahashi et al. (2009) is commonly used in atmospheric transport models of  $\text{CO}_2$  (e.g. Geels et al., 2007; Sarrat et al., 2009), and is also applied here for areas outside the Baltic Sea and the Danish inner waters. However, this climatology does not cover the Baltic Sea area, and therefore, a new Baltic Sea climatology has been created and merged with the climatology of Takahashi et al. (2009) in the model domain towards the North Sea and the Northern North Atlantic.

Available  $p\text{CO}_2$  surface measurements and water chemistry data from the Baltic Sea and the Danish inner waters are combined in six sub-domains of the Baltic Sea to provide monthly averaged  $p\text{CO}_2$  values for this new climatology. The sub-domains cover Skagerrak, Kattegat and the Belt Sea (henceforth referred to just as Kattegat), the Western Baltic Sea, the Baltic Proper, the Gulf of Finland, the Bothnian Sea and the Bothnian Bay. Two data sets are analysed; one from marine stations (stationary) and the other obtained from ships (underway). All available data collected since year 2000 is included in the analysis (Fig. 1). From the two data sets monthly mean values for each sub-domain are determined.

Surface measurements of salinity, temperature, alkalinity and pH from six marine measuring stations (operated by the Swedish Meteorological and Hydrological Institute, SMHI; Shark Data Base, 2013) are applied to calculate the surface  $p\text{CO}_2$  values by a similar approach as described in Wesslander et al. (2010). The six stations are located from the central Skagerrak to the Bothnian Bay (Fig. 1), but no measurements are available from the Gulf of Finland. A relatively high frequency of observations is obtained at the six monitoring stations with the number of observations in each month ranging between 4–8 at station A17, 15–36 at station Anholt E, 6–18 at station BY5,

BGD

11, 16993–17042, 2014

Sensitivity of the air-sea  $\text{CO}_2$  exchange in the Baltic Sea and Danish inner waters

A. S. Lansø et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

7–17 at station BY15, 1–5 at station C3 (but no data representing November) and 2–10 at station F9 (but no data representing January, February and November).

Surface levels of  $p\text{CO}_2$  from the central Baltic Sea (Schneider and Sadkowiak, 2012) have been measured by underway- $p\text{CO}_2$  systems (Körtzing et al., 1996; Schneider et al., 2006) from cargo and research ships. In particular, a route between Germany (Kiel) and Finland (Helsinki) has regularly been monitored from cargo-ships, whereas no measurements are available in the northern part of the Baltic Sea, the Danish straits, Kattegat and Skagerrak. Good data coverage of underway  $p\text{CO}_2$  measurements is obtained in the sub-domain of the Western Baltic Sea, with the number of observations in each month ranging between 9000 and 55 000, and in the Baltic Proper, where the corresponding number of observations ranges from 20 000 to 116 000. In the Bothnian Sea the number of observations ranges from 2000 to 77 000, but there are no observations in December. Only a single month (March) is represented in the Bothnian Bay with about 5000 observations. The Gulf of Finland is represented with observations ranging from 3000 and 18 000 each month.

The stationary data from the monitoring stations and the underway data have been combined in such a manner that if underway data exists for a sub-domain, these data is used for the  $p\text{CO}_2$  fields in the given subdomain. Otherwise, measurements from the monitoring stations are used to calculate the  $p\text{CO}_2$  fields. Thus,  $p\text{CO}_2$  fields for Skagerrak, Kattegat, and the Bothnian Bay are calculated solely based on data from the SMHI stations. The  $p\text{CO}_2$  fields for the Western Baltic Sea, the Baltic Proper, the Gulf of Finland and the Bothnian Sea are obtained from the underway measurements of  $p\text{CO}_2$ , except for December in the Bothnian Sea, which is represented by the monitoring station C3. The data used to obtain the monthly averages of surface  $p\text{CO}_2$  in each sub-domain have all been normalised to year 2000 using an annual increase in  $\text{CO}_2$  of  $1.9 \mu\text{atm yr}^{-1}$  found for the central Baltic Sea (Wesslander et al., 2010).

The resulting  $p\text{CO}_2$  climatology for the Baltic Sea and Danish inner waters is combined with the global open ocean  $p\text{CO}_2$  climatology from Takahashi et al. (2009). This climatology is calculated for a global oceanic grid with a horizontal resolution of  $5^\circ \times 4^\circ$  in

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

longitude and latitude, respectively. Consequently, this field has an even coarser spatial resolution than the sub-domains defined in the Baltic Sea area. The global climatology is by Takahashi and co-workers referenced to the year 2000 with an annual trend of  $1.5 \mu\text{atm yr}^{-1}$ . This trend is also used to extrapolate the global data for year 2000 to the proceeding years covered in this study. Note that the trend used for the Baltic Sea and Danish inner waters is  $1.9 \mu\text{atm yr}^{-1}$ , as this trend is shown to match this particular area. However, the difference in annual trends between the two climatologies is so small compared to the absolute  $p\text{CO}_2$  values, thus it is reasonable to assume that the impact on the current results will be insignificant.

The monthly averaged  $p\text{CO}_2$  values show a characteristic seasonal pattern at all monitoring stations and for the underway  $p\text{CO}_2$  data (Fig. 2, Table S1 in the Supplement). The surface  $p\text{CO}_2$  is under-saturated during spring and summer and super-saturated during fall and winter (Fig. 3a). However, there is a large spatial gradient in the seasonal amplitude from Skagerrak and into the Baltic Sea. A seasonal amplitude of about  $140 \mu\text{atm}$  characterises the variation in Skagerrak and Kattegat, where the  $p\text{CO}_2$  varies between  $275$  and  $420 \mu\text{atm}$ , and the surface water is only slightly super-saturated during the winter months. In the Baltic Sea a relatively large seasonal amplitude of up to  $400 \mu\text{atm}$  is observed, as primary production during the growing season, i.e. spring and summer, causes a large uptake of total dissolved inorganic carbon in the surface layer and contributes to lowering the surface  $p\text{CO}_2$  values. The data shows how biological uptake causes a reduction of surface  $p\text{CO}_2$ , despite the general warming during the summer month, which normally tends to increase the  $p\text{CO}_2$  in the surface water. During fall and winter, the surface  $p\text{CO}_2$  value increase, because sub-surface waters enriched in total dissolved inorganic carbon from remineralisation of organic matter during the summer are mixed into the surface layer. In the areas north-east of the Western Baltic Sea in particularly, this allows for high monthly averaged surface  $p\text{CO}_2$  values of  $460$ – $530 \mu\text{atm}$  during winter with the largest average winter values observed in the Gulf of Finland.

<a href="#">Title Page</a>	
<a href="#">Abstract</a>	<a href="#">Introduction</a>
<a href="#">Conclusions</a>	<a href="#">References</a>
<a href="#">Tables</a>	<a href="#">Figures</a>
<a href="#">◀</a>	<a href="#">▶</a>
<a href="#">◀</a>	<a href="#">▶</a>
<a href="#">Back</a>	<a href="#">Close</a>
<a href="#">Full Screen / Esc</a>	
<a href="#">Printer-friendly Version</a>	
<a href="#">Interactive Discussion</a>	

In general a good accordance between the  $p\text{CO}_2$  values calculated from the monitoring stations and underway  $p\text{CO}_2$  data from the sub-domains exists. The underway  $p\text{CO}_2$  data includes both temporal and spatial variability within each sub-domain during the period since 2000. Therefore, their standard deviations (SD) are larger than the SDs from the monitoring stations, which mainly arise due to inter-annual variability in the period. Two sub-domains, the Western Baltic Sea and the Baltic Proper, have good data coverage from both the monitoring stations and underway  $p\text{CO}_2$  data. The stations, BY5 and BY15, that represent the Western Baltic Sea and the Baltic Proper, respectively, have lower surface  $p\text{CO}_2$  values during the summer period than the underway  $p\text{CO}_2$  data, but the difference between the two data sets are within their SD.

### 2.3 Model framework

The model framework is based upon the Danish Eulerian Hemispheric Model (DEHM) – a well validated three dimensional large scale atmospheric chemical transport model (Brandt et al., 2012; Christensen, 1997). DEHM is based on the equation of continuity and use terrain following sigma levels as vertical coordinates. Here, 29 vertical levels are distributed between the surface and 100 hPa with a higher density of vertical levels in the lower part of the atmosphere. The main domain of DEHM covers the Northern Hemisphere with a horizontal grid resolution of  $150\text{ km} \times 150\text{ km}$  using a polar stereographic projection true at  $60^\circ\text{ N}$ . Furthermore, DEHM has nesting capabilities allowing for a nest over Europe with a resolution of  $50\text{ km} \times 50\text{ km}$ , a nest of Northern Europe with an approximate resolution of  $16.7\text{ km} \times 16.7\text{ km}$ , and a  $5.6\text{ km} \times 5.6\text{ km}$  nest covering Denmark. In order to cover the Baltic Sea and the Danish marine areas in focus, a setup with two nests is applied in the current study (the European and the Northern European nests). The main domain and the nests each comprise of  $96 \times 96$  grid points. This study uses a modified version of DEHM solely simulating transport and exchange of  $\text{CO}_2$  (Geels et al., 2002, 2004, 2007), but in the present study with an updated description of the surface exchange of  $\text{CO}_2$  (described in Sect. 2.2.1).

DEHM is driven by meteorological data from the meteorological model MM5v3.7 (Grell et al., 1995) using National Centers for Environmental Prediction, NCEP, data as input.

### 2.3.1 Model inputs

5 To accurately simulate the atmospheric content of CO<sub>2</sub>, a number of CO<sub>2</sub> sources and sinks within the model domain as well as inflow at the lateral boundaries are required together with a background concentration. The atmospheric concentration of CO<sub>2</sub> ( $X_{\text{atm}}$ ) can be described by

$$X_{\text{atm}} = X_{\text{ff}} + X_{\text{bio}} + X_{\text{fire}} + X_{\text{ocn}} + X_{\text{background}} \quad (1)$$

10 where  $X_{\text{ff}}$  is the contribution of CO<sub>2</sub> from fossil fuel emissions,  $X_{\text{fire}}$  from vegetation fires, and  $X_{\text{bio}}$  and  $X_{\text{ocn}}$  are the contribution to the atmospheric concentration from exchange of CO<sub>2</sub> with the terrestrial biosphere and ocean, respectively.  $X_{\text{background}}$  is the atmospheric background of CO<sub>2</sub>.

#### Fossil Fuel ( $X_{\text{ff}}$ )

15 Fossil fuel emissions for the domain covering the Northern Hemisphere are implemented in DEHM from the Carbon Tracker (hereafter referred to as CT) simulation system (CarbonTracker CT2011\_oi, 2013; Peters et al., 2007). This emission map has a three hourly temporal resolution on a 1° × 1° grid.

20 CT is for the European area substituted by a fossil fuel emission inventory with a higher spatiotemporal resolution (hourly, 10 km × 10 km) developed by the Institute of Energy Economics and the Rational Use of Energy (Pregger et al., 2007).

25 For the area of Denmark, emissions with an even finer spatial resolution of 1 km × 1 km are applied obtained from the Department of Environmental Science, Aarhus University. These are based on the Danish national inventory submitted yearly to UNFCCC (United Nations Framework Convention on Climate Change) and constructed from

energy statistics, point source and statistic sub-models (Plejdrup and Gyldenkærne, 2011).

As the European and Danish emission inventories are for the years 2005 and 2011, respectively, these inventories are scaled to total yearly national estimates of carbon emissions from fossil fuel consumption conducted by the Carbon Dioxide Information Analysis Center, CDIAC, in order to account for the year to year change in emissions (Boden et al., 2013).

### Biosphere ( $X_{\text{bio}}$ )

Terrestrial biosphere fluxes from the CT system, with a spatial resolution of  $1^\circ \times 1^\circ$  and a temporal resolution of three hours, are applied in DEHM. In the CT assimilation system the Carnegie-Ames Stanford Approach (CASA) biogeochemical model is used for prior fluxes (Giglio et al., 2006; van der Werf et al., 2006). The prior terrestrial biosphere fluxes are optimised in the CT assimilation system by atmospheric observations of  $\text{CO}_2$ . Via this atmospheric inversion a best guess of surface fluxes is obtained, and the optimised fluxes are implemented in DEHM.

### Fires ( $X_{\text{fire}}$ )

$\text{CO}_2$  emissions due to vegetation fires are obtained from the CT fire module and applied in DEHM. The CT fire module is based on the Global Fire Emission Database, GFEDv3.1, and CASA, while the burned area from GFED is based on MODIS satellite observations of fire counts. The resolution is likewise three hourly on a  $1^\circ \times 1^\circ$  grid.

### Ocean ( $X_{\text{ocn}}$ )

The  $\text{CO}_2$  flux ( $F$ ) at the air-sea interface is calculated using the relation:  $F = k\alpha\Delta p\text{CO}_2$ , where,  $k$  is the exchange coefficient,  $\alpha$  is the gas solubility and  $\Delta p\text{CO}_2$  is the difference in partial pressure of  $\text{CO}_2$  between the surface water and the overlying air. The gas solubility of  $\text{CO}_2$  is determined from Weiss (1974) and depends on the

BGD

11, 16993–17042, 2014

Sensitivity of the air-sea  $\text{CO}_2$  exchange in the Baltic Sea and Danish inner waters

A. S. Lansø et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



water temperature and salinity. A  $0.25^\circ \times 0.25^\circ$  salinity map is implemented in DEHM for the calculation of  $\text{CO}_2$  solubility (Boyer et al., 2005). To calculate  $\Delta p\text{CO}_2$  the surface  $p\text{CO}_2$  fields described in Sect. 2.2 are applied together with the concentration of  $\text{CO}_2$  in the lowest atmospheric layer in DEHM.

5 No standardised parameterisation of the transfer velocity,  $k$ , exists, but  $k$  is most often parameterised as a power function of the wind speed (Garbe et al., 2014; Rutgersson et al., 2008) normalised to the Schmidt's number ( $Sc$ ) according to Wanninkhof (1992). In the present study we use the parameterisation of Wanninkhof (1992) (hereafter referred to as W92), which is the parameterisation most frequently applied.

10 This transfer velocity is a function of the wind speed at 10 m above the surface ( $u_{10}$ ) and when normalised to  $Sc$  at 20 °C in salt water it has the form:

$$k_{660} = \left(0.31u_{10}^2\right) \sqrt{\frac{660}{Sc}} \quad (2)$$

15 However, a few additional parameterisations that could be more representative for the study area are also tested. One is Nightingale et al. (2000), who estimate a transfer velocity based on tracer gas measurements in the North Sea of:

$$k_{660} = \left(0.333u_{10} + 0.222u_{10}^2\right) \sqrt{\frac{660}{Sc}} \quad (3)$$

Another is by Weiss et al. (2007), who have carried out measurements using eddy covariance techniques in the Arkona basin located within the Baltic Sea to estimate an accurate  $k$  for this particular area. This parameterisation takes the form:

$$20 \quad k_{660} = \left(0.365u_{10}^2 + 0.46u_{10}\right) \sqrt{\frac{660}{Sc}} \quad (4)$$

Sea ice coverage is in DEHM obtained from NCEP. The sea ice coverage is implemented in the calculations of the air–sea  $\text{CO}_2$  exchange, such that the flux in a grid

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

cell is reduced by the fraction of sea ice. If the fraction of sea ice coverage is 1, the entire grid cell will be covered with ice, and no exchange of CO<sub>2</sub> will take place between the ocean and atmosphere. Recent studies have shown that CO<sub>2</sub> exchange between ice-covered sea and the atmosphere does take place, but to what extent has not yet 5 been quantified (Parmentier et al., 2013; Sørensen et al., 2014). For that reason the exchange over sea ice is not accounted for here.

$k_{660}$ ,  $\alpha$  and  $\Delta p\text{CO}_2$  are calculated at each time step of the model simulation (the time step of the model varies between ca. 3–20 min depending of e.g. the nest). Consequently, the air–sea CO<sub>2</sub> flux has the same temporal resolution as the simulated 10 atmospheric CO<sub>2</sub>.

### Atmospheric background ( $X_{\text{background}}$ )

The level of atmospheric CO<sub>2</sub> has been increasing since pre-industrial times. It is not feasible to simulate this entire time period with the model system as to replicate this build-up. Therefore, an atmospheric background of CO<sub>2</sub> is needed. The atmospheric 15 background of CO<sub>2</sub> is established on the basis of the NOAA ESRL GLOBALVIEW-CO<sub>2</sub> data product using observations from the Baltic Station, BAL (lat = 55°35' N, lon = 17°22' E) (GLOBALVIEW-CO<sub>2</sub>, 2013). BAL lies within the area of interest, but far from local sources and sinks. It can therefore be assumed to represent the atmospheric 20 background level in the study area. The atmospheric background of CO<sub>2</sub> is calculated based on the following equation:

$$X_{\text{background}} = X_{\text{CO}_2 \text{ 2000}} + 1.91(\text{year} - 2000) + 0.16\text{month} \quad (5)$$

Here  $X_{\text{CO}_2 \text{ 2000}} = 370.15$  ppm is the mean CO<sub>2</sub> concentration at the station in 2000, year and month is the simulated year and month, and 1.91 and 0.16 represent the yearly 25 and monthly trend of atmospheric CO<sub>2</sub>. The trends are based on the times series at BAL for the period 2000–2010, in order to get a representative overall trend for the period in focus here (2005–2010).

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

DEHM only covers the Northern Hemisphere; hence boundary conditions for the main domain are needed at the lateral boundaries towards the Southern Hemisphere, as to account for inflow from the Southern Hemisphere. Three dimensional atmospheric mole fractions of CO<sub>2</sub> from the CT system are applied at these boundaries.

## 3 Results

### 3.1 Model evaluation

The period 2005–2010 is simulated by DEHM with setup and fluxes as described in Sect. 2. The performance of the model for this period is evaluated by comparing simulated atmospheric CO<sub>2</sub> concentrations against observed. The comparison is made at six stations within the study area, where both remote continental (PAL), marine (F3, MHD, OST, WES) and anthropogenic (LUT) influenced stations are represented.

Measured and simulated atmospheric CO<sub>2</sub> from the marine site Östergarnsholm, Sweden (OST, 57°27' N, 18°59' E) and the anthropogenic continental site Lutjewad, the Netherlands (LUT, 53°40' N, 6°31' E) (van der Laan et al., 2009) are shown for year 2007 in Fig. 4. The Östergarnsholm marine micrometeorological field station has been running semi-continuously since 1995, measuring atmospheric CO<sub>2</sub> since 2005. The site has been shown to represent marine conditions and is described further in Rutgersson et al. (2008) and Höglström et al. (2008). The site at Lutjewad is on the other hand both influenced by background air from the North Sea and polluted air from the continent, depending on the wind direction. Hourly mean concentrations are plotted for simulated and measured atmospheric CO<sub>2</sub>, and at both sites a large diurnal variability is seen in the observations. The model is not able to capture the large amplitude in the diurnal cycle, but correlations of 0.75 and 0.71 are obtained for LUT and OST, respectively. The root mean square errors, RMSE, are 9.6 and 8.8 ppm, respectively.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

These high RMSEs are linked to the underestimation of the diurnal cycle in the model. Earlier model studies have shown the same tendency to underestimate the observed variability (e.g. Geels et al., 2007). The underestimation of the diurnal cycle by DEHM is most likely caused by the coarse spatial resolution of the biosphere fluxes. Further,

5 weekly averages are made for both observed and modelled concentrations of atmospheric CO<sub>2</sub> (see Fig. 4). Improvements are obtained in both correlation and RMSE to 0.89 and 5.3 ppm for LUT, and 0.91 and 5.6 ppm for OST. Synoptic scale variability is seen in the atmospheric CO<sub>2</sub> concentration in both the simulated and observed time-series. Especially at LUT large positive spikes are seen due to the influence of air from 10 densely populated and industrialised regions.

Flask measurements of CO<sub>2</sub> at F3, the Netherlands (54°51' N, 4°44' E) (van der Laan-Luijkx et al., 2010) are compared to hourly modelled averages (Fig. 5) during the six year simulated period. This results in a correlation of 0.64, and a RMSE of 5.7 ppm. F3 is positioned on an oil platform in the North Sea, where local sources 15 can influence the measured CO<sub>2</sub> concentration under certain wind conditions. Consequently, the most extreme outliers were filtered out with the help of simultaneous CH<sub>4</sub> and CO measurements, when the influence from the local source was obvious. As to examine the model performance on a longer time scale, weekly averages are made for the three remaining stations Mace Head, Ireland (MHD, 53°20' N, 9°54' W) (Biraud 20 et al., 2000), Pallas-Sammaltunturi, Finland (PAL, 67°58' N, 24°07' E) (FMI, 2013) and Westerland, Germany (WES, 54°56' N, 8°19' E) (UBA, 2014) for the six year period (Fig. 5). In general a reasonable correspondence between model and observations is seen during this period with correlations of 0.96, 0.98 and 0.89, and RMSEs of 1.8, 1.9 and 3.8 ppm for MHD, PAL and WES, respectively. The ability of the model to capture 25 the seasonal cycle is contributing to the very high correlation, but the model is also capable of capturing weekly variability and transport events especially during winter.

To conclude, this evaluation shows that the DEHM model captures the overall CO<sub>2</sub> pattern across the marine region in focus in the current study.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## 3.2 Air-sea CO<sub>2</sub> fluxes

In order to investigate the effect on the air-sea CO<sub>2</sub> flux from variations in the atmospheric CO<sub>2</sub> concentration at the Baltic Sea and Danish inner waters, two different model simulations are conducted. One model simulation has atmospheric CO<sub>2</sub> concentrations that vary from time step to time step according to the fluxes and atmospheric transport in DEHM. This is in the following referred to as the VAT ("Variable ATmosphere") simulation. The other simulation contains at each time step and grid cell the monthly mean CO<sub>2</sub> concentration for the given month. This is in the following referred to as CAT ("Constant ATmosphere"). All other settings are identical in the two simulations. The simulations are made for the period 2005 to 2010 using the transfer velocity parameterisation by W92.

First, the results of atmospheric CO<sub>2</sub> concentrations and air-sea CO<sub>2</sub> fluxes from the VAT simulation will be presented. These results can be used to get an understanding of how the atmospheric CO<sub>2</sub> concentrations vary, and of how the air-sea CO<sub>2</sub> fluxes behave in terms of size and direction in the different sub-basins of the Baltic Sea and Danish inner waters. This will be followed by the comparison of the VAT and CAT simulation.

### 3.2.1 Variable atmospheric CO<sub>2</sub> concentration

The variability of atmospheric CO<sub>2</sub> in the Baltic area is illustrated in Fig. 6, which shows a few examples of the hourly simulated surface concentration. The top panel shows the variability in February 2007, where synoptic scale variability influence transport of CO<sub>2</sub>, and hence the surface concentrations. On 1 February 2007 at 4:00 GMT, a low pressure system has during the last days moved through Southern Scandinavia and is now located over Poland. This system has rotated continental air with high levels of CO<sub>2</sub> from east towards the Baltic Sea. On 3 February 2007, the prevailing winds are now westerly, where marine air masses with lower CO<sub>2</sub> concentrations are transported towards the Baltic Sea. The lower panel of Fig. 6 shows the diurnal variability on 14

BGD

11, 16993–17042, 2014

Sensitivity of the air-sea CO<sub>2</sub> exchange in the Baltic Sea and Danish inner waters

A. S. Lansø et al.

<a href="#">Title Page</a>	
<a href="#">Abstract</a>	<a href="#">Introduction</a>
<a href="#">Conclusions</a>	<a href="#">References</a>
<a href="#">Tables</a>	<a href="#">Figures</a>
<a href="#">◀</a>	<a href="#">▶</a>
<a href="#">◀</a>	<a href="#">▶</a>
<a href="#">Back</a>	<a href="#">Close</a>
<a href="#">Full Screen / Esc</a>	
<a href="#">Printer-friendly Version</a>	
<a href="#">Interactive Discussion</a>	

July 2007. At 2:00 GMT, air masses with high CO<sub>2</sub> concentrations are transported from land towards the marine areas – most evident in near-coastal areas. The same is the case at 14:00 GMT, but with lower concentrations due to the uptake of CO<sub>2</sub> by the terrestrial biosphere at this time of the day. These examples show that large spatial gradients of up to 20 ppm can develop across the Baltic Sea during summer.

The seasonal averaged air–sea CO<sub>2</sub> fluxes estimated by DEHM in the VAT simulation are shown in Fig. 3b. In winter, a gradient is seen from the North Sea through the Danish inner Straits towards the Baltic Sea, indicating a large release of CO<sub>2</sub> to the atmosphere in the Baltic, and uptake in the North Sea. Progressing to spring, the gradient towards the Baltic ceases and all areas now have marine uptake of atmospheric CO<sub>2</sub>, which continues throughout the summer. In fall, the gradient starts to build up again, and the Baltic Sea becomes a source of CO<sub>2</sub> to the atmosphere.

The monthly mean 2005–2010 sub-basin averaged fluxes likewise depict this seasonality (Fig. 7). The highest seasonal amplitudes are found in the Baltic Sea area stretching from the Baltic Proper and northwards with the greatest seasonal amplitude of 12 g C m<sup>-2</sup> month<sup>-1</sup> found in the Bothnian Sea. Less seasonal variation in the CO<sub>2</sub> flux is obtained for Kattegat and the Danish Straits, which yearly experience a variability of just 4.3 g C m<sup>-2</sup> month<sup>-1</sup>.

The total sub-basin monthly mean fluxes of CO<sub>2</sub> between the atmosphere and ocean also show a seasonal variation for all areas with release in winter and uptake of atmospheric CO<sub>2</sub> in summer (Table 1). The entire area comprising of the six sub-basins has for the period 2005–2010 an average annual net uptake of atmospheric CO<sub>2</sub> of 287 Gg C yr<sup>-1</sup>. However, the net exchange varies greatly from sub-basin to sub-basin. Kattegat, the Western Baltic Sea and the Baltic Proper all have annual net uptake of atmospheric CO<sub>2</sub> averaged over 2005 to 2010, while the remaining three sub-basins release CO<sub>2</sub> to the atmosphere. The Baltic Proper contributes the most to the total annual averaged flux with an uptake of 254 g C yr<sup>-1</sup>, but during the individual months the fluxes in the Baltic Proper are even larger (up to 900 g C month<sup>-1</sup>). Monthly fluxes of this considerable size are not obtained in any of the other sub-domains. This is of

**Sensitivity of the air–sea CO<sub>2</sub> exchange in the Baltic Sea and Danish inner waters**

A. S. Lansø et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

course related to the fact that the Baltic Proper has the greatest spatial extent of all the six sub-basins.

To estimate the marine contribution in the Danish national CO<sub>2</sub> budget, the air-sea CO<sub>2</sub> flux in the Danish Exclusive Economic Zone (EEZ) is calculated. The EEZ is a zone adjacent to the territorial waters extending up to 200 nautical mile off shore, and in the EEZ the coastal state has the right to explore, exploit and manage all resources within this area (United Nations Chapter XXI Law of the Sea, 1984). The Danish EEZ has an area of approximately 105 000 km<sup>2</sup> (Fig. S2 in the Supplement). During the six years simulated an average annual uptake in the Danish EEZ of 2613 Gg C yr<sup>-1</sup> is obtained. The main part of the uptake in the Danish EEZ occurs in the North Sea. The North Sea has the largest extent in the Danish EEZ and combined with a small seasonal amplitude in  $p\text{CO}_2$ , this results in a constant uptake throughout the year. The other sub-basins within the Danish EEZ all release CO<sub>2</sub> in winter and take up CO<sub>2</sub> during summer. Compared to the Danish national emissions of anthropogenic CO<sub>2</sub>, the marine uptake by the Danish EEZ corresponds to 18 % per year of these emissions (Table 2). For the six year period investigated, the annual mean inventory in CO<sub>2</sub> excluding land use and land use change is 14.6 Tg C (Nielsen et al., 2013).

### 3.2.2 Constant atmospheric CO<sub>2</sub> concentration

The impact of variations in the atmospheric CO<sub>2</sub> concentration is analysed in the following by comparing the results of the net annual averaged air-sea CO<sub>2</sub> fluxes for the VAT and CAT simulations in the six sub-basins. A total annual difference of 184 Gg C yr<sup>-1</sup> is obtained, which corresponds to a 64 % difference (calculated with VAT as the reference). CAT gives a total annual uptake of 471 Gg C yr<sup>-1</sup>, while VAT only has an annual uptake of 287 Gg C yr<sup>-1</sup>. The seasonal difference between VAT and CAT across the study area is seen in Fig. 8. The monthly fluxes in the sub-basins maintain the same direction in both VAT and CAT. However, for months where the different sub-basins experience outgassing, the outgassing is reduced in the CAT simulation as compared

to in the VAT simulation. For months with uptake of CO<sub>2</sub> in the individual sub-basins, a higher uptake is simulated with the CAT setup than with the VAT setup.

In order to further analyse the difference between the VAT and the CAT simulations, times series of the driving parameters are compared. Examples of the atmospheric pCO<sub>2</sub> ( $p\text{CO}_2^a$ ) in the lowest model layer in the VAT and CAT simulations are shown for a coastal site south of Sweden (55°18' N, 13°55' E) in Figs. 9 and 10 for February and July 2007, respectively. February represents a case of outgassing, and July a case of marine uptake of atmospheric CO<sub>2</sub>. Time series of wind velocity at 10 m,  $u_{10}$ , and the atmospheric mixing height,  $h_{\text{mix}}$ , are also plotted. Further, the differences in the atmospheric partial pressure of CO<sub>2</sub> ( $\Delta p\text{CO}_2^a$ ) and in the air-sea CO<sub>2</sub> flux ( $\Delta F_{\text{CO}_2}$ ) between the two simulations are shown (calculated as VAT – CAT). Differences in the  $p\text{CO}_2^a$  in the two simulations determines the difference in  $\Delta p\text{CO}_2^a$  between the two simulations, as the partial pressure of CO<sub>2</sub> in the water is the same in the two simulations.  $p\text{CO}_2^a$  is the only variable allowed to vary in the air-sea CO<sub>2</sub> flux calculations between VAT and CAT, and is thus responsible for the obtained flux differences.

For both months  $p\text{CO}_2^a$ \_VAT fluctuates around the constant  $p\text{CO}_2^a$ \_CAT. During the first half of February, a period of anti-correlation between  $p\text{CO}_2^a$ \_VAT and  $u_{10}$  is seen. This anti-correlation is greatest during the second week with a weekly correlation coefficient ( $r$ ) equal to  $-0.69$ . During the last week of February, a positive correlation of  $r = 0.62$  between the two is obtained with wind speeds above 10 m s<sup>-1</sup> and high  $p\text{CO}_2^a$  levels in the atmosphere. In February no clear diurnal cycle is seen in the mixing height, but it seems to follow the pattern of the wind speed with decreases in  $h_{\text{mix}}$  during periods with low wind speeds and increases in  $h_{\text{mix}}$  during high wind speeds. The correlation between these two parameters in February is  $r = 0.72$ .

In July a clear diurnal variability is seen in  $p\text{CO}_2^a$ \_VAT, and an anti-correlation between  $h_{\text{mix}}$  and  $p\text{CO}_2^a$ \_VAT is evident throughout the month with the highest anti-correlation during the last week (with  $r = -0.72$ ). In the VAT simulation the so-called diurnal rectifier effect is modelled. The collaboration between terrestrial ecosystems and boundary layer dynamics that act towards lowering  $p\text{CO}_2^a$  during the day and in-

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

crease it during night is known as the rectifier effect. In particularly during the growing season the rectifier effect is apparent (Denning et al., 1996).

An anti-correlation between  $\Delta p\text{CO}_2^a$  and  $\Delta F_{\text{CO}_2}$  is seen in both February and July, which will be discussed in more detail in Sect. 4.3. During winter the largest difference in the air-sea  $\text{CO}_2$  flux between VAT and CAT coincides with high wind speeds or large differences in the atmospheric  $\text{CO}_2$  concentrations (hence large  $\Delta p\text{CO}_2^a$  values). In summer the diurnal cycle in the atmospheric  $\text{CO}_2$  levels are translated into the flux difference.

Vertical profiles of atmospheric  $\text{CO}_2$  at the site south of Sweden have been plotted together with  $h_{\text{mix}}$  in Fig. 11. Note that the unit in Fig. 11 is ppm and not  $\mu\text{atm}$ . The variability of  $\text{CO}_2$  is also evident in the vertical profile, where air masses with low or high  $\text{CO}_2$  concentrations are being transported to and from the site ( $55^{\circ}18' \text{N}$ ,  $13^{\circ}55' \text{E}$ ). Continental air is represented by high levels of  $\text{CO}_2$  that extend up to 2 km into the atmosphere, while marine air masses have lower levels of  $\text{CO}_2$  corresponding to the levels above 2 km. The shift between the two types of air masses is clearly seen in the vertical profile e.g. on 2 February. Here higher wind speed leads to transport of marine air masses to the site (see Fig. 9). Like Fig. 9, the vertical profile in February shows no clear connection between surface concentrations of  $\text{CO}_2$  and  $h_{\text{mix}}$ . In July, the vertical profile depicts the rectifier effect. Low surface values of  $\text{CO}_2$  coincide with the greatest boundary layer heights found during the day time, and high surface levels of  $\text{CO}_2$  concur during night time with the nocturnal boundary layer. It is remarkable how the vertical profile during July 2007 represents a much more mixed atmosphere as compared to February 2007, where the marine and continental air masses clearly are distinguished from each other.

BGD

11, 16993–17042, 2014

---

## Sensitivity of the air-sea $\text{CO}_2$ exchange in the Baltic Sea and Danish inner waters

A. S. Lansø et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

## 4 Discussion

### 4.1 Surface water $p\text{CO}_2$ climatology

A representative map of surface  $p\text{CO}_2$  has been created for Skagerrak and six sub-domains in the Baltic using two data sets, one obtained from monitoring stations and 5 one using underway measurements of surface  $p\text{CO}_2$  (see Sect. 2.2).

Previous estimates of  $p\text{CO}_2$  at two positions within the Baltic Sea have shown an inter-annual variability of up to 25 % in winter and almost 140 % in summer (Wesslander et al., 2010). Likewise, large short term variability has been measured in different coastal systems (Dai et al., 2009; Leinweber et al., 2009; Wesslander et al., 2011).

10 The representation of surface  $p\text{CO}_2$  values in the sub-domains by a monthly averaged value does not account for the temporal variability during each month and the spatial variability in the relatively large areas. Also, the inter-annual variability is not accounted for by the average values. However, the simplified description of the conditions in the Baltic Sea in a number of sub-domains is currently the best solution in order to obtain 15 a surface field of  $p\text{CO}_2$  that spatially covers the whole model domain for this study. The estimated surface fields of  $p\text{CO}_2$  are based on relatively few observations, and therefore, it has not been possible to include neither spatial, nor short-term or inter-annual variability. Although, underway  $p\text{CO}_2$  measurements (Schneider and Sadkowiak, 2012) have increased the data coverage in the central Baltic Sea significantly in the most 20 recent years.

### 4.2 Air-sea $\text{CO}_2$ fluxes

The atmospheric  $\text{CO}_2$  concentration is seen to vary greatly within the study area (Figs. 6 and 11). The dynamics of the fluxes and the atmospheric transport and mixing lead to short-term variations and spatial gradients in the atmospheric  $\text{CO}_2$  level across 25 the marine areas in focus here. Pressure systems move through the region transporting air masses with different characteristics and  $\text{CO}_2$  levels to and from the Baltic and the

BGD

11, 16993–17042, 2014

Sensitivity of the air-sea  $\text{CO}_2$  exchange in the Baltic Sea and Danish inner waters

A. S. Lansø et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



## Sensitivity of the air-sea $\text{CO}_2$ exchange in the Baltic Sea and Danish inner waters

A. S. Lansø et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Danish inner waters. In the growing season, the effect from the terrestrial biosphere is apparent with a clear diurnal cycle in the atmospheric  $\text{CO}_2$  caused by respiration during night time and photosynthesis during the day. Even these short term variations in atmospheric  $\text{CO}_2$  concentrations over land can be transported to marine areas, indicating why atmospheric short term variability is important to include in the air-sea flux estimations.

For the six year period an annual average uptake of  $287 \text{ Gg C} \text{yr}^{-1}$  is obtained with the VAT-setup as a total for the six sub-basins. A statistical analysis of the simulated fluxes shows that Kattegat and the Western Baltic Sea are annual sinks (at a significance level of 0.05), while the Gulf of Finland and the Bothnian Bay are annual sources of atmospheric  $\text{CO}_2$ . In the transition zone between these areas, i.e. the Baltic Proper and the Bothnian Sea, large variations in the annual flux are seen in this study. During the six years simulated, these sub-domains change annually between being sources and sinks of  $\text{CO}_2$  to the atmosphere. This also affects the total flux for the entire investigated area, which also shifts between being an annual source and sink. A significant test (Student's  $t$  test with a significance level of 0.05) show that the variability from year to year is so large that we cannot conclude that the area is a net sink, despite the estimated averaged uptake of  $287 \text{ Gg C} \text{yr}^{-1}$ .

The air-sea  $\text{CO}_2$  fluxes obtained from the VAT simulation for six sub-basins are compared to previous results from the area as to assess the consistency. Previous studies of the air-sea  $\text{CO}_2$  flux in the Baltic Sea area are ambiguous as to the Baltic Sea's role in the carbon cycle. This is partly caused by the various techniques used, ranging from in-situ measurements using eddy covariance method to model simulations (Kulinski and Pempkowiak, 2011; Rutgersson et al., 2009; Weiss et al., 2007; Wesslander et al., 2010), and partly by the different spatial areas investigated. Some of the previous studies are site specific (Algesten et al., 2006; Kuss et al., 2006; Löffler et al., 2012; Rutgersson et al., 2008; Wesslander et al., 2010), and other studies cover the entire area (Gustafsson et al., 2014; Kulinski and Pempkowiak, 2011; Norman et al., 2013). None of the previous regional studies have based their estimates of the air-sea  $\text{CO}_2$

flux on results from an atmospheric transport model capable of combining large spatial coverage with high spatiotemporal resolution of the entire Baltic region as in the present study. Results from previous studies and the present study have been converted to the same unit of  $\text{gC m}^{-2} \text{yr}^{-1}$ , as to allow for a direct comparison (Table 3).

5 Table 3 reveals that in terms of the direction of the flux, the present study corresponds well with some of the previous studies while contradicts others. As the results obtained from the VAT simulation lie within in the range of previous estimates, it seems reasonable to use the current model setup for sensitivity analysis of the air-sea  $\text{CO}_2$  flux in the region. Additional, it can be concluded that the obtained results from the  
10 VAT simulation together with recent studies converge towards the Baltic Sea and the Danish inner waters being annual sinks of atmospheric  $\text{CO}_2$ .

### 4.3 Impact of atmospheric short term variability

The difference of  $184 \text{ Gg C yr}^{-1}$  between the annual air-sea flux in the CAT and VAT simulations was tested to be significantly different from zero at a 0.05 significance level.

15 Therefore, it can be concluded that using a constant level of atmospheric  $\text{CO}_2$  has a significant impact on the estimated annual air-sea  $\text{CO}_2$  flux. The greatest differences are found in winter and fall in the Baltic Sea area (Fig. 8). But large differences are also found over open water areas in spite of a less variable atmospheric  $\text{CO}_2$  concentration here. The generally higher wind speeds over open water might lead to the large flux  
20 difference here.

The deviation between the two simulations in the study region is mainly caused by a reduction in the winter uptake in the CAT simulation. An optimal situation where higher uptake during winter occurs in the VAT simulation is seen, when high wind speeds coincide with lower  $p\text{CO}_2^a$  in the VAT than the CAT simulation. Thereby,  $\Delta p\text{CO}_2$  is greater in the VAT simulation than in the CAT simulation. In combination with the non-linearity of the wind speed in the parameterisation of the transfer velocity, this leads to a larger outgassing of  $\text{CO}_2$  in the VAT simulation. Such a situation is seen at the coastal site south of Sweden ( $55^{\circ}18' \text{N}, 13^{\circ}55' \text{E}$ ) in Fig. 9, where episodes with high

BGD

11, 16993–17042, 2014

## Sensitivity of the air-sea $\text{CO}_2$ exchange in the Baltic Sea and Danish inner waters

A. S. Lansø et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

wind speeds in the beginning of the month transport air masses with lower CO<sub>2</sub> levels to the site and increase the flux in VAT as compared to CAT (i.e. a positive  $\Delta F_{\text{CO}_2}$ ).

The higher marine CO<sub>2</sub> uptake in summer by the CAT simulation is a result of diurnal boundary layer dynamics and the diurnal cycle or lack of it in the atmospheric CO<sub>2</sub> concentrations. The rectifier effect is not accounted for in the CAT simulation, and the constant  $p\text{CO}_2^a$  in CAT is higher during the day and lower during the night than in the VAT simulation. This allows for a greater air-sea  $\Delta p\text{CO}_2$  in the CAT simulation during day, which together with a tendency of higher wind speeds during day time increases the oceanic uptake in CAT. This is illustrated by  $\Delta F_{\text{CO}_2}$ , where positive values indicate how the flux is numerical larger in CAT than VAT (see Fig. 10). As described in Sect. 3.1, the diurnal cycle of atmospheric CO<sub>2</sub> is generally underestimated by the DEHM model. Due to this underestimation the difference between the VAT and CAT simulations found during the growing season is likely a conservative estimate.

While Rutgersson et al. (2008) find a slightly overestimated seasonal amplitude, when using a constant atmospheric CO<sub>2</sub> concentration, the present study finds that the seasonal cycle of the CAT simulation is displaced downwards as compared to the VAT simulation. This displacement results in a greater annual uptake in the CAT simulation.

#### 4.4 Uncertainties

The estimated air-sea CO<sub>2</sub> flux is controlled by several parameters in the applied model setup: choice of transfer velocity parameterisation, wind speed, temperature, salinity, atmospheric CO<sub>2</sub> concentration and marine  $p\text{CO}_2$  surface values. Each of these is connected with some uncertainty and errors.

Takahashi et al. (2009) estimate the combined precision on the global air-sea flux to be on the order of  $\pm 60\%$ , when including a possible climatology bias due to interpolation and under-sampling. The uncertainty might be higher in the current study, as the climatology for the  $p\text{CO}_2$  in surface waters used here covers areas, where the spatiotemporal variability in the measured  $p\text{CO}_2$  is higher than in open waters. The natural

BGD

11, 16993–17042, 2014

Sensitivity of the air-sea CO<sub>2</sub> exchange in the Baltic Sea and Danish inner waters

A. S. Lansø et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

variability within the sub-domains is represented by the SDs in Fig. 2 and it reflects both the spatial and temporal variation in the domains during the period of sampling, i.e. the last decade. The Baltic Sea domains (i.e. except the Kattegat sub-domain) are all characterised by a significant under-saturation of the surface water during spring and summer. During winter these stations are in general supersaturated with respect to the atmospheric  $p\text{CO}_2$ . Thus, the sign of the  $\text{CO}_2$  flux during the seasons is assumed to be well-determined in the Baltic Sea sub-domains due to the large seasonal amplitudes. However, during the seasonal change between summer and winter, where typical SDs in the climatology of 50 ppm are seen, we estimate that the uncertainty due to the ocean surface  $p\text{CO}_2$ -values is in the order of 50 % in the Baltic Sea. The uncertainty in the Kattegat sub-domain is estimated to be up to 50–100 % because of the relatively small seasonal amplitude.

Atmospheric  $\text{CO}_2$ , wind speed and temperature all vary in each model time step and grid cell. The uncertainties of wind speed and temperature are small compared to the uncertainties of the  $p\text{CO}_2$  fields. Figures 4 and 5 show how well the DEHM model captures the weekly and seasonal variability in the atmospheric  $\text{CO}_2$  concentrations. However, some problems arise in capturing the variability on shorter time scales (e.g. diurnal). The diurnal cycle is under-estimated in this model setup, which is related to the coarse resolution of the biosphere fluxes, and of the model itself.

Short term variability does not only exist in the atmospheric concentration of  $\text{CO}_2$ , it has also been detected in the  $p\text{CO}_2$  of surface water (Dai et al., 2009; Leinweber et al., 2009; Rutgersson et al., 2008; Wesslander et al., 2011). The magnitude of the short term variability is site dependent with smallest variability found in open oceans (Dai et al., 2009) and greatest at near-coastal sites (Leinweber et al., 2009; Wesslander et al., 2011). Off the Californian coast, Leinweber et al. (2009) find a diurnal cycle of  $p\text{CO}_2$  with an average amplitude of 20  $\mu\text{atm}$  – a diurnal amplitude double of what they find in the atmosphere. Short term variability of marine  $p\text{CO}_2$ , could potentially alter the annual estimate of the coastal air–sea  $\text{CO}_2$  flux. Thus, in the present study the fluxes at the near-coastal areas within the sub-domains could be affected by this

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

short term variability and could possibly modify the total flux for the sub-domains. However, the short term variability in marine  $p\text{CO}_2$  is not included in this study, and it is therefore difficult to estimate how this might affect the estimated flux. Further, the short term variability in the air and water might be correlated, thus it is not possible to make a deduction of the combined effect in the present model study.

The largest uncertainty connected to the estimated air-sea flux is related to the choice of transfer velocity. The results from the VAT simulation presented here were calculated using the parameterisation by Wanninkhof (1992), but model simulations using parameterisation of Nightingale et al. (2000) and Weiss et al. (2007) have also been conducted. With these parameterisations the annual flux for the study area is changed to  $-667$  and  $-858\text{ Gg Cyr}^{-1}$ , respectively. The present study supports the findings briefly touched upon by Rutgersson et al. (2009), who conclude that the uncertainty due to the value of atmospheric CO<sub>2</sub> is small compared to uncertainty in transfer velocity. We, however, stress that ignoring short term variability in marine and atmospheric  $p\text{CO}_2$  introduces a bias in the estimates of the air-sea CO<sub>2</sub> flux.

## 5 Conclusion

Using an atmospheric CO<sub>2</sub> model with a relatively high spatial and temporal resolution we have estimated the air-sea flux of CO<sub>2</sub> in the Danish inner waters and the Baltic Sea region. More specifically we have made a detailed analysis of the sensitivity to temporal variability in atmospheric CO<sub>2</sub> and the related impact of driving parameters like wind speed and atmospheric mixing height.

In the process of this study new monthly marine  $p\text{CO}_2$  fields have been developed for the region combining existing data from monitoring stations and measurements from ships. Due to the sparseness of these data, only seasonal variations are included in the  $p\text{CO}_2$  fields.

The atmospheric concentration of CO<sub>2</sub> is often assumed to be constant or only vary by season in many marine model studies, but according to this novel sensitivity analy-

sis, neglecting e.g. the diurnal variability in atmospheric CO<sub>2</sub> concentrations could lead to a systematic bias in the annual net air–sea flux. Previous studies have looked at the entire Baltic region (Gustafsson et al., 2014; Kulinski and Pempkowiak, 2011; Norman et al., 2013; Thomas and Schneider, 1999), but not with the same approach as in the present study.

In all the included sub-basins a seasonal cycle was detected in the air–sea CO<sub>2</sub> flux with release of CO<sub>2</sub> during winter and fall, and uptake of atmospheric CO<sub>2</sub> in the remaining months. An annual flux for the study area of  $-287 \text{ Gg C yr}^{-1}$  ( $-0.7 \text{ g C m}^2 \text{ yr}^{-1}$ ) was obtained for the six years simulated. This agrees with the previous findings of Norman et al. (2013) and Gustafsson et al. (2014), who estimate annual air–sea CO<sub>2</sub> fluxes of  $-2.6$  and  $-4.3 \text{ g C m}^2 \text{ yr}^{-1}$ , respectively. Significant testing of the flux for the study area shows that for the six years simulated here, we cannot conclude that the total Baltic Sea region is an annual sink. Instead our results indicate that the region alternates between being an annual source and sink. If examining the results at sub-basin scale, this alternation is caused by two sub-basins: the Baltic Proper and the Bothnian Sea. Furthermore, our results suggest that Kattegat and the Western Baltic Sea act as annual sinks of atmospheric CO<sub>2</sub>, and the Gulf of Finland and the Bothnian Bay act as sources.

In order to test the importance of short term (hourly) variations in the atmospheric CO<sub>2</sub> in relation to the yearly air–sea flux, two different model simulations have been made. The first simulation includes the short-term variations (the VAT simulation), while the other simulation includes a monthly constant atmospheric CO<sub>2</sub> concentration (the CAT simulation). A significant difference of  $184 \text{ Gg C yr}^{-1}$  (corresponding to 67 %) was obtained for the air–sea CO<sub>2</sub> flux for the Baltic Sea and Danish inner waters between the two model simulations. The seasonal amplitude of the air–sea CO<sub>2</sub> flux was in the CAT simulation shifted downwards as compared to the VAT simulation, resulting in a reduced winter release of CO<sub>2</sub> in the CAT simulation and an increased summer uptake. The difference occurs solely due to the difference in the atmospheric CO<sub>2</sub> concentrations.

## Sensitivity of the air–sea CO<sub>2</sub> exchange in the Baltic Sea and Danish inner waters

A. S. Lansø et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

As a part of the Danish project ECOCLIM with focus on the Danish CO<sub>2</sub> budget, the natural marine annual flux of CO<sub>2</sub> was for the first time estimated by the present study. The Danish waters – in this context defined as the Danish Exclusive Economic Zone – is according to our simulations taking up 2613 Gg C yr<sup>-1</sup> with the majority taken up in the North Sea. This is comparable to approximately 18 % of the Danish anthropogenic CO<sub>2</sub> emissions.

The uncertainty in estimating the air–sea CO<sub>2</sub> exchange has many contributions. The largest is found to be the parameterisation of the transfer velocity, which supports previous findings. The present study, however, underlines the importance of including short term variability in the atmospheric CO<sub>2</sub> in order to minimise the uncertainties in the air–sea CO<sub>2</sub> flux. This further leads to the deduction that also short term variability in *p*CO<sub>2</sub> of the water, in particular of coastal areas, needs to be included, as short term variability in near coastal surface water *p*CO<sub>2</sub> potentially is greater than in the atmosphere.

To conclude, we recommend that future studies of the air–sea CO<sub>2</sub> exchange include short term variability of CO<sub>2</sub> in the atmosphere. Thereby, the uncertainty related to estimating the marine part of the carbon budget at regional to global scales can be reduced.

The Supplement related to this article is available online at  
doi:10.5194/bgd-11-16993-2014-supplement.

**Acknowledgements.** This study is a part of a Ph.D. project within ECOCLIM founded by the Danish Strategic Research Council (grant number 10-093901). Further, several scientists involved with this work are likewise connected to the Nordic Centre of Excellence, DEFROST. We are grateful to Anna Rutgersson at Uppsala University for sharing atmospheric measurements of CO<sub>2</sub> at Östergarnsholm. We likewise thank the community for making observations available at Lutjewad, Mace Head, Pallas-Sammaltunturi and Westerland. We thank SMHI for sharing marine data from the six monitoring sites. Results from Carbon Tracker CT2011\_oi provided by

BGD

11, 16993–17042, 2014

---

**Sensitivity of the air–sea CO<sub>2</sub> exchange in the Baltic Sea and Danish inner waters**

A. S. Lansø et al.

---

<a href="#">Title Page</a>	
<a href="#">Abstract</a>	<a href="#">Introduction</a>
<a href="#">Conclusions</a>	<a href="#">References</a>
<a href="#">Tables</a>	<a href="#">Figures</a>
<a href="#">◀</a>	<a href="#">▶</a>
<a href="#">◀</a>	<a href="#">▶</a>
<a href="#">Back</a>	<a href="#">Close</a>
<a href="#">Full Screen / Esc</a>	
<a href="#">Printer-friendly Version</a>	
<a href="#">Interactive Discussion</a>	

NOAA ESRL, Boulder, Colorado, USA from the website at <http://carbontracker.noaa.gov> have contributed to this work, as have GLOBALVIEW-CO<sub>2</sub>, 2013 and European and Danish fossil fuel emission inventories from IER and Aarhus University, respectively.

## References

5 Algesten, G., Brydsten, L., Jonsson, P., Kortelainen, P., Löfgren, S., Rahm, L., Räike, A., Sobek, S., Tranvik, L., Wikner, J., and Jansson, M.: Organic carbon budget for the Gulf of Bothnia, *J. Marine Syst.*, 63, 155–161, 2006.

10 Bendtsen, J., Gustafsson, K. E., Söderkvist, J., and Hansen, J. L. S.: Ventilation of bottom water in the North Sea-Baltic Sea transition zone, *J. Marine Syst.*, 75, 138–149, 2009.

15 Biraud, S., Ciais, P., Ramonet, M., Simmonds, P., Kazan, V., Monfray, P., O'Doherty, S., Spain, T. G., and Jennings, S. G.: European greenhouse gas emissions estimated from continuous atmospheric measurements and radon 222 at Mace Head, Ireland, *J. Geophys. Res.-Atmos.*, 105, 1351–1366, 2000.

20 Boden, T. A., Marland, G., and Andres, R. J.: Global, Regional, and National Fossil-Fuel CO<sub>2</sub> Emissions, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, Tenn., USA, 2013.

25 Borges, A. V., Schiettecatte, L. S., Abril, G., Delille, B., and Gazeau, E.: Carbon dioxide in European coastal waters, *Estuar. Coast. Shelf S.*, 70, 375–387, 2006.

30 Boyer, T., Levitus, S., Garcia, H., Locarnini, R. A., Stephens, C., and Antonov, J.: Objective analyses of annual, seasonal, and monthly temperature and salinity for the world ocean on a 0.25 degrees grid, *Int. J. Climatol.*, 25, 931–945, 2005.

35 Brandt, J., Silver, J. D., Frohn, L. M., Geels, C., Gross, A., Hansen, A. B., Hansen, K. M., Hedegaard, G. B., Skjøth, C. A., Villadsen, H., Zare, A., and Christensen, J. H.: An integrated model study for Europe and North America using the Danish Eulerian Hemispheric Model with focus on intercontinental transport of air pollution, *Atmos. Environ.*, 53, 156–176, 2012.

40 Cai, W. J.: Estuarine and coastal ocean carbon paradox: CO<sub>2</sub> sinks or sites of terrestrial carbon incineration?, *Annu. Rev. Marine Sci.*, 3, 123–145, 2011.

45 CarbonTracker: CT2011\_oi, available at: <http://carbontracker.noaa.gov> (last access: 1 August 2013), 2013.

BGD

11, 16993–17042, 2014

Sensitivity of the air-sea CO<sub>2</sub> exchange in the Baltic Sea and Danish inner waters

A. S. Lansø et al.

<a href="#">Title Page</a>	
<a href="#">Abstract</a>	<a href="#">Introduction</a>
<a href="#">Conclusions</a>	<a href="#">References</a>
<a href="#">Tables</a>	<a href="#">Figures</a>
<a href="#">◀</a>	<a href="#">▶</a>
<a href="#">◀</a>	<a href="#">▶</a>
<a href="#">Back</a>	<a href="#">Close</a>
<a href="#">Full Screen / Esc</a>	
<a href="#">Printer-friendly Version</a>	
<a href="#">Interactive Discussion</a>	

Chen, C. T. A. and Borges, A. V.: Reconciling opposing views on carbon cycling in the coastal ocean: continental shelves as sinks and near-shore ecosystems as sources of atmospheric CO<sub>2</sub>, *Deep-Sea Res. Pt. II*, 56, 578–590, 2009.

5 Chen, C.-T. A., Huang, T.-H., Chen, Y.-C., Bai, Y., He, X., and Kang, Y.: Air–sea exchanges of CO<sub>2</sub> in the world's coastal seas, *Biogeosciences*, 10, 6509–6544, doi:10.5194/bg-10-6509-2013, 2013.

Christensen, J. H.: The Danish Eulerian hemispheric model – a three-dimensional air pollution model used for the Arctic, *Atmos. Environ.*, 31, 4169–4191, 1997.

10 Dai, M. H., Lu, Z. M., Zhai, W. D., Chen, B. S., Cao, Z. M., Zhou, K. B., Cai, W. J., and Chen, C. T. A.: Diurnal variations of surface seawater *p*CO<sub>2</sub> in contrasting coastal environments, *Limnol. Oceanogr.*, 54, 735–745, 2009.

Denning, A. S., Randall, D. A., Collatz, G. J., and Sellers, P. J.: Simulations of terrestrial carbon metabolism and atmospheric CO<sub>2</sub> in a general circulation model .2. Simulated CO<sub>2</sub> concentrations, *Tellus B*, 48, 543–567, 1996.

15 FMI: Atmospheric CO<sub>2</sub> Hourly Concentration Data, Pallas-Sammaltunturi – FMI, WMO World Data Centre for Greenhouse Gases, Japan Meteorol. Agency, Tokyo, 2013.

Garbe, C. S., Rutgersson, A., Boutin, J., de Leeuw, G., Delille, B., Fairall, C. W., Gruber, N., Hare, J., Ho, D. H., Johnson, M. T., Nightingale, P. D., Pettersson, H., Piskozub, J., Sahlee, E., Tsai, W., Ward, B., Woolf, D. K., and Zappa, C. J.: Transfer across the air–sea interface, in: *Ocean–Atmosphere Interactions of Gases and Particles*, edited by: Liss, P. S. and Johnson, M. T., Springer Earth System Science, 55–111, 2014.

Gattuso, J. P., Frankignoulle, M., and Wollast, R.: Carbon and carbonate metabolism in coastal aquatic ecosystems, *Annu. Rev. Ecol. Syst.*, 29, 405–434, 1998.

25 Geels, C., Christensen, J. H., Frohn, L. M., and Brandt, J.: Simulating spatiotemporal variations of atmospheric CO<sub>2</sub> using a nested hemispheric model, *Phys. Chem. Earth*, 27, 1495–1505, 2002.

Geels, C., Doney, S. C., Dargaville, R., Brandt, J., and Christensen, J. H.: Investigating the sources of synoptic variability in atmospheric CO<sub>2</sub> measurements over the Northern Hemisphere continents: a regional model study, *Tellus B*, 56, 35–50, 2004.

30 Geels, C., Gloor, M., Ciais, P., Bousquet, P., Peylin, P., Vermeulen, A. T., Dargaville, R., Aalto, T., Brandt, J., Christensen, J. H., Frohn, L. M., Haszpra, L., Karstens, U., Rödenbeck, C., Ramonet, M., Carboni, G., and Santaguida, R.: Comparing atmospheric transport models for

BGD

11, 16993–17042, 2014

Sensitivity of the air–sea CO<sub>2</sub> exchange in the Baltic Sea and Danish inner waters

A. S. Lansø et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

future regional inversions over Europe – Part 1: mapping the atmospheric CO<sub>2</sub> signals, *Atmos. Chem. Phys.*, 7, 3461–3479, doi:10.5194/acp-7-3461-2007, 2007.

5 Geels, C., Hansen, K. M., Christensen, J. H., Ambelas Skjøth, C., Ellermann, T., Hede-  
gaard, G. B., Hertel, O., Frohn, L. M., Gross, A., and Brandt, J.: Projected change in at-  
mospheric nitrogen deposition to the Baltic Sea towards 2020, *Atmos. Chem. Phys.*, 12,  
2615–2629, doi:10.5194/acp-12-2615-2012, 2012.

10 Giglio, L., van der Werf, G. R., Randerson, J. T., Collatz, G. J., and Kasibhatla, P.: Global  
estimation of burned area using MODIS active fire observations, *Atmos. Chem. Phys.*, 6,  
957–974, doi:10.5194/acp-6-957-2006, 2006.

15 10 GLOBALVIEW-CO<sub>2</sub>: Cooperative Global Atmospheric Data Integration Project, Updated  
Annually, Multi-Laboratory Compilation of Synchronized and Gap-Filled Atmospheric  
Carbon Dioxide Records for the Period 1979–2012 (obspack\_co2\_1\_GLOBALVIEW-  
CO<sub>2</sub>\_2013\_v1.0.4\_2013-12-23), Compiled by NOAA Global Monitoring Division, Boulder,  
Colorado, USA, doi:10.3334/OBSPACK/1002, 2013.

20 15 Grell, G. A., Dudhia, J., and Stauffer, D. R.: A Description of the Fifth-Generation Penn  
State/NCAR Mesoscale Model (MM5), National Center for Atmospheric Research, NCAR  
Technical Note NCAR/TN-398+STR, Boulder, Colorado, USA, 1995.

25 Gustafsson, E., Deutsch, B., Gustafsson, B. G., Humborg, C., and Mörth, C. M.: Carbon cycling  
in the Baltic Sea – the fate of allochthonous organic carbon and its impact on air–sea CO<sub>2</sub>  
exchange, *J. Marine Syst.*, 129, 289–302, 2014.

30 Högström, U., Sahlée, E., Drennan, W. M., Kahma, K. K., Smedman, A. S., Johansson, C.,  
Pettersson, H., Rutgersson, A., Tuomi, L., Zhang, F., and Johansson, M.: Momentum fluxes  
and wind gradients in the marine boundary layer – a multi-platform study, *Boreal Environ.  
Res.*, 13, 475–502, 2008.

25 Körtzinget, A., Thomas, H., Schneider, B., Gronau, N., Mintrop, L., and Durr, H. H.: At sea  
intercompariosn of two newly designed underway  $p\text{CO}_2$  systems – encouraging results, *Mar.  
Chem.*, 52, 133–145, 1996.

30 Kuliński, K. and Pempkowiak, J.: The carbon budget of the Baltic Sea, *Biogeosciences*, 8,  
3219–3230, doi:10.5194/bg-8-3219-2011, 2011.

30 Kuss, J., Roeder, W., Wlost, K. P., and DeGrandpre, M. D.: Time-series of surface water CO<sub>2</sub>  
and oxygen measurements on a platform in the central Arkona Sea (Baltic Sea): seasonality  
of uptake and release, *Mar. Chem.*, 101, 220–232, 2006.

## Sensitivity of the air–sea CO<sub>2</sub> exchange in the Baltic Sea and Danish inner waters

A. S. Lansø et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Langner, J., Andersson, C., and Engardt, M.: Atmospheric input of nitrogen to the Baltic Sea basin: present situation, variability due to meteorology and impact of climate change, *Boreal Environ. Res.*, 14, 226–237, 2009.

5 Laruelle, G. G., Durr, H. H., Slomp, C. P., and Borges, A. V.: Evaluation of sinks and sources of CO<sub>2</sub> in the global coastal ocean using a spatially-explicit typology of estuaries and continental shelves, *Geophys. Res. Lett.*, 37, L15607, doi:10.1029/2010GL043691, 2010.

10 Le Quéré, C., Andres, R. J., Boden, T., Conway, T., Houghton, R. A., House, J. I., Marland, G., Peters, G. P., van der Werf, G. R., Ahlström, A., Andrew, R. M., Bopp, L., Canadell, J. G., Ciais, P., Doney, S. C., Enright, C., Friedlingstein, P., Huntingford, C., Jain, A. K., Jourdain, C., Kato, E., Keeling, R. F., Klein Goldewijk, K., Levis, S., Levy, P., Lomas, M., Poulter, B., Rau-  
pach, M. R., Schwinger, J., Sitch, S., Stocker, B. D., Viovy, N., Zaehle, S., and Zeng, N.: The global carbon budget 1959–2011, *Earth Syst. Sci. Data*, 5, 165–185, doi:10.5194/essd-5-165-2013, 2013.

15 Leinweber, A., Gruber, N., Frenzel, H., Friederich, G. E., and Chavez, F. P.: Diurnal carbon cycling in the surface ocean and lower atmosphere of Santa Monica Bay, California, *Geophys. Res. Lett.*, 36, L08601, doi:10.1029/2008GL037018, 2009.

Löffler, A., Schneider, B., Perttilä, M., and Rehder, G.: Air–sea CO<sub>2</sub> exchange in the Gulf of Bothnia, Baltic Sea, *Cont. Shelf Res.*, 37, 46–56, 2012.

20 Meesters, A. G. C. A., Tolk, L. F., Peters, W., Hutjes, R. W. A., Vellinga, O. S., Elbers, J. A., Vermeulen, A. T., van der Laan, S., Neubert, R. E. M., Meijer, H. A. J., and Dolman, A. J.: Inverse carbon dioxide flux estimates for the Netherlands, *J. Geophys. Res.-Atmos.*, 117, L20306, doi:10.1029/2012JD017797, 2012.

Mørk, E. T., Sørensen, L. L., Jensen, B., and Sejr, M. K.: Air–sea CO<sub>2</sub> gas transfer velocity in a shallow estuary, *Bound.-Lay. Meteorol.*, 151, 119–138, 2014.

25 Nielsen, O.-K., Plejdrup, M. S., Winther, M., Nielsen, M., Gyldenkærne, S., Mikkelsen, M. H., Albrektsen, R., Thomsen, M., Hjelgaard, K., Hoffmann, L., Fauser, P., Bruun, H. G., Johannsen, V. K., Nord-Larsen, T., Vesterdal, L., Møller, I. S., Caspersen, O. H., Rasmussen, E., Petersen, S. B., Baunbæk, L., and Hansen, M. G.: Denmark's National Inventory Report 2013, Emission Inventories 1990–2011 – Submitted under the United Nations Framework Convention on Climate Change and the Kyoto Protocol, Aarhus University, DCE – Danish Centre for Environment and Energy, 1–1202, 2013.

BGD

11, 16993–17042, 2014

Sensitivity of the air–sea CO<sub>2</sub> exchange in the Baltic Sea and Danish inner waters

A. S. Lansø et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Nightingale, P. D., Malin, G., Law, C. S., Watson, A. J., Liss, P. S., Liddicoat, M. I., Boutin, J., and Upstill-Goddard, R. C.: In situ evaluation of air–sea gas exchange parameterizations using novel conservative and volatile tracers, *Global Biogeochem. Cy.*, 14, 373–387, 2000.

5 Norman, M., Rutgersson, A., and Sahlée, E.: Impact of improved air–sea gas transfer velocity on fluxes and water chemistry in a Baltic Sea model, *J. Marine Syst.*, 111, 175–188, 2013.

Omstedt, A., Gustafsson, E., and Wesslander, K.: Modelling the uptake and release of carbon dioxide in the Baltic Sea surface water, *Cont. Shelf Res.*, 29, 870–885, 2009.

10 Parmentier, F. J. W., Christensen, T. R., Sørensen, L. L., Rysgaard, S., McGuire, A. D., Miller, P. A., and Walker, D. A.: The impact of lower sea-ice extent on Arctic greenhouse-gas exchange, *Nat. Clim. Change*, 3, 195–202, 2013.

Peters, W., Jacobson, A. R., Sweeney, C., Andrews, A. E., Conway, T. J., Masarie, K., Miller, J. B., Bruhwiler, L. M. P., Pétron, G., Hirsch, A. I., Worthy, D. E. J., van der Werf, G. R., Randerson, J. T., Wennberg, P. O., Krol, M. C., and Tans, P. P.: An atmospheric perspective on North American carbon dioxide exchange: CarbonTracker, *P. Natl. Acad. Sci. USA*, 104, 18925–18930, 2007.

15 Pleijdrup, M. S. and Gyldenkærne, S.: Spatial Distribution of Emissions to Air – the SPREAD model, National Environmental Research Institute, Aarhus University, Denmark, 2011.

Pregger, T., Scholz, Y., and Friedrich, R.: Documentation of the Anthropogenic GHG Emission Data for Europe Provided in the Frame of CarboEurope GHG and CarboEurope IP, University of Stuttgart, IER – Institute of Energy Economics and the Rational Use of Energy, 2007.

20 Rutgersson, A., Norman, M., Schneider, B., Pettersson, H., and Sahlée, E.: The annual cycle of carbon dioxide and parameters influencing the air–sea carbon exchange in the Baltic Proper, *J. Marine Syst.*, 74, 381–394, 2008.

Rutgersson, A., Norman, M., and Åström, G.: Atmospheric CO<sub>2</sub> variation over the Baltic Sea and the impact on air–sea exchange, *Boreal Environ. Res.*, 14, 238–249, 2009.

25 Sarrat, C., Noilhan, J., Lacarrère, P., Masson, V., Ceschia, E., Ciais, P., Dolman, A., Elbers, J., Gerbig, C., and Jarosz, N.: CO<sub>2</sub> budgeting at the regional scale using a Lagrangian experimental strategy and meso-scale modeling, *Biogeosciences*, 6, 113–127, doi:10.5194/bg-6-113-2009, 2009.

30 Schneider, B. and Sadkowiak, B.: Underway *p*CO<sub>2</sub> Measurements in Surface Waters during the VOS and Research Vruises in the Baltic Sea, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, Tennessee, USA, 2012.

## Sensitivity of the air–sea CO<sub>2</sub> exchange in the Baltic Sea and Danish inner waters

A. S. Lansø et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Schneider, B., Kaitala, S., and Maunula, P.: Identification and quantification of plankton bloom events in the Baltic Sea by continuous  $p\text{CO}_2$  and chlorophyll *a* measurements on a cargo ship, *J. Marine Syst.*, 59, 238–248, 2006.

Shark Data Base: SMHI, available at: <http://www.smhi.se/klimatdata/oceanografi/Havsmiljodata> (last access: 1 May 2014), 2013.

Smallman, T. L., Williams, M., and Moncrieff, J. B.: Can seasonal and interannual variation in landscape  $\text{CO}_2$  fluxes be detected by atmospheric observations of  $\text{CO}_2$  concentrations made at a tall tower?, *Biogeosciences*, 11, 735–747, doi:10.5194/bg-11-735-2014, 2014.

Sørensen, L. L., Jensen, B., Glud, R. N., McGinnis, D. F., Sejr, M. K., Sievers, J., Søgaard, D. H., Tison, J.-L., and Rysgaard, S.: Parameterization of atmosphere–surface exchange of  $\text{CO}_2$  over sea ice, *The Cryosphere*, 8, 853–866, doi:10.5194/tc-8-853-2014, 2014.

Takahashi, T., Sutherland, S. C., Wanninkhof, R., Sweeney, C., Feely, R. A., Chipman, D. W., Hales, B., Friederich, G., Chavez, F., Sabine, C., Watson, A., Bakker, D. C. E., Schuster, U., Metzl, N., Yoshikawa-Inoue, H., Ishii, M., Midorikawa, T., Nojiri, Y., Körtzinger, A., Steinhoff, T., Hoppema, M., Olafsson, J., Arnarson, T. S., Tilbrook, B., Johannessen, T., Olsen, A., Bellerby, R., Wong, C. S., Delille, B., Bates, N. R., and de Baar, H. J. W.: Climatological mean and decadal change in surface ocean  $p\text{CO}_2$ , and net sea–air  $\text{CO}_2$  flux over the global oceans, *Deep-Sea Res. Pt. II*, 56, 554–577, 2009.

Thomas, H. and Schneider, B.: The seasonal cycle of carbon dioxide in Baltic Sea surface waters, *J. Marine Syst.*, 22, 53–67, 1999.

Thomas, H., Bozec, Y., Elkalay, K., and de Baar, H. J. W.: Enhanced open ocean storage of  $\text{CO}_2$  from shelf sea pumping, *Science*, 304, 1005–1008, 2004.

Thomas, H., Bozec, Y., de Baar, H. J. W., Elkalay, K., Frankignoulle, M., Schiettecatte, L.-S., Kattner, G., and Borges, A. V.: The carbon budget of the North Sea, *Biogeosciences*, 2, 87–96, doi:10.5194/bg-2-87-2005, 2005.

UBA: Air Monitoring Network of the Federal Environment Agency (UBA), Germany Monitoring Site Westerland, 2014.

United Nations Chapter XXI: Law of the Sea, available at: [https://treaties.un.org/Pages/ViewDetailsIII.aspx?&src=TREATY&mtdsg\\_no=XXI~6&chapter=21&Temp=mtdsg3&lang=en](https://treaties.un.org/Pages/ViewDetailsIII.aspx?&src=TREATY&mtdsg_no=XXI~6&chapter=21&Temp=mtdsg3&lang=en) (last access: May 2014), 1984.

van der Laan, S., Neubert, R. E. M., and Meijer, H. A. J.: A single gas chromatograph for accurate atmospheric mixing ratio measurements of  $\text{CO}_2$ ,  $\text{CH}_4$ ,  $\text{N}_2\text{O}$ ,  $\text{SF}_6$  and  $\text{CO}$ , *Atmos. Meas. Tech.*, 2, 549–559, doi:10.5194/amt-2-549-2009, 2009.

## Sensitivity of the air–sea $\text{CO}_2$ exchange in the Baltic Sea and Danish inner waters

A. S. Lansø et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



van der Laan-Luijkx, I. T., Neubert, R. E. M., van der Laan, S., and Meijer, H. A. J.: Continuous measurements of atmospheric oxygen and carbon dioxide on a North Sea gas platform, *Atmos. Meas. Tech.*, 3, 113–125, doi:10.5194/amt-3-113-2010, 2010.

van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Kasibhatla, P. S., and Arel-Iano Jr., A. F.: Interannual variability in global biomass burning emissions from 1997 to 2004, *Atmos. Chem. Phys.*, 6, 3423–3441, doi:10.5194/acp-6-3423-2006, 2006.

Wanninkhof, R.: Relationship between wind-speed and gas-exchange over the ocean, *J. Geophys. Res-Oceans*, 97, 7373–7382, 1992.

Weiss, A., Kuss, J., Peters, G., and Schneider, B.: Evaluating transfer velocity-wind speed relationship using a long-term series of direct eddy correlation  $\text{CO}_2$  flux measurements, *J. Marine Syst.*, 66, 130–139, 2007.

Weiss, R. F.: Carbon dioxide in water and seawater: the solubility of a non-ideal gas, *Mar. Chem.*, 2, 203–215, 1974.

Wesslander, K., Omstedt, A., and Schneider, B.: Inter-annual and seasonal variations in the air-sea  $\text{CO}_2$  balance in the central Baltic Sea and the Kattegat, *Cont. Shelf Res.*, 30, 1511–1521, 2010.

Wesslander, K., Hall, P., Hjalmarsson, S., Lefevre, D., Omstedt, A., Rutgersson, A., Sahlée, E., and Tengberg, A.: Observed carbon dioxide and oxygen dynamics in a Baltic Sea coastal region, *J. Marine Syst.*, 86, 1–9, 2011.

## Sensitivity of the air-sea $\text{CO}_2$ exchange in the Baltic Sea and Danish inner waters

A. S. Lansø et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



**Table 1.** Monthly mean fluxes for the period 2005–2010 depicting seasonal variation of the air–sea CO<sub>2</sub> exchange. Unit is in Gg C per sub-basin. Positive sign indicates release of CO<sub>2</sub> from the ocean to the atmosphere, negative sign indicates uptake of atmospheric CO<sub>2</sub> by the ocean. This sign convention is used throughout the paper.

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Ann
Kattegat	29	-21	-98	-42	-25	-28	-26	-33	-33	-15	14	43	-235
Western Baltic	125	31	-113	-226	-206	-142	-153	-92	60	137	236	140	-203
Baltic Proper	804	365	92	-654	-808	-718	-844	-784	-178	481	995	993	-254
Gulf of Finland	49	60	8	-61	-92	-68	-74	-67	1	78	151	117	102
Bothnian Sea	207	120	83	-253	-383	-325	-355	-284	-22	439	529	412	167
Bothnian Bay	31	23	10	-7	-50	-91	-118	-18	48	205	94	9	137

<a href="#">Abstract</a>	<a href="#">Introduction</a>
<a href="#">Conclusions</a>	<a href="#">References</a>
<a href="#">Tables</a>	<a href="#">Figures</a>
<a href="#">◀</a>	<a href="#">▶</a>
<a href="#">◀</a>	<a href="#">▶</a>
<a href="#">Back</a>	<a href="#">Close</a>
<a href="#">Full Screen / Esc</a>	
<a href="#">Printer-friendly Version</a>	
<a href="#">Interactive Discussion</a>	

**Table 2.** The top row shows the annual Danish CO<sub>2</sub> emissions as reported to UNFCCC. The middle row contains the annual uptake of CO<sub>2</sub> in the marine area defined as the Danish Exclusive Economic Zone as estimated in this study. The bottom rows give this uptake as a percentage of the Danish anthropogenic CO<sub>2</sub> emissions.

	2005	2006	2007	2008	2009	2010	6 yr avg.
DK CO <sub>2</sub> (Tg C)	14.3	16.5	15.2	14.3	13.6	13.7	14.6
Total Uptake – EEZ (Tg C)	-2.6	-2.4	-2.8	-2.6	-2.6	-2.7	-2.6
Percentage of CO <sub>2</sub>	18	14	18	18	19	20	18

[Title Page](#)  
[Abstract](#) [Introduction](#)  
[Conclusions](#) [References](#)  
[Tables](#) [Figures](#)

[◀](#) [▶](#)  
[◀](#) [▶](#)  
[Back](#) [Close](#)  
[Full Screen / Esc](#)

[Printer-friendly Version](#)  
[Interactive Discussion](#)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Table 3.** Present study compared to previous results within the different sub-domains. Study type indicates the type of previous study (Mod. – model based, MBA – mass balance approach, Meas. – measurement based, Cru. – cruise based) and its spatial extend (s-b – sub-basins, s-s – site specific). All shown results are in  $\text{g C m}^2 \text{ yr}^{-1}$ .

	Previous results	Study	Study type	Presents Study
Kattegat	–40.0	Gustafsson et al. (2014)	Mod., s-b	–7.0
	19.0	Norman et al. (2013)	Mod., s-b	
	–13.9	Wesslander et al. (2010)	Meas., s-s	
Western Baltic	–34	Gustafsson et al. (2014)	Mod., s-b	–3.1
	–36.0	Kuss et al. (2006)	Meas., s-s	
	–14.4 to 17.9	Norman et al. (2013)	Mod., s-b	
	28.1	Wesslander et al. (2010)	Meas., s-s	
Baltic Proper	–4.2 to –4.3	Norman et al. (2013)	Mod., s-b	–1.5
	–10.8	Schneider and Thomas(1999)	Cru., s-b	
	19.7	Wesslander et al. (2010)	Meas., s-s	
Bothnian Sea	2.2	Gustafsson et al. (2014)	Mod., s-b	2.2
	–8.8	Löffler et al. (2012)	Cru., s-b	
	–0.6	Norman et al. (2013)	Mod., s-b	
Bothnian Bay	12.0	Gustafsson et al. (2014)	Mod., s-b	3.8
	1.7	Löffler et al. (2012)	Cru., s-b	
	4.3	Norman et al. (2013)	Mod., s-b	
Gulf of Finland	7.4	Norman et al. (2013)	Mod., s-b	4.3
Total Baltic Sea	2.7	Kulinski and Pempkowiak (2010)	MBA, s-b	–0.7
	–4.3	Gustafsson et al. (2014)	Mod., s-b	
	–2.6	Norman et al. (2013)	Mod., s-b	

## Sensitivity of the air-sea CO<sub>2</sub> exchange in the Baltic Sea and Danish inner waters

A. S. Lansø et al.

**Figure 1.** The locations of the six monitoring stations in the Baltic Sea, where surface  $p\text{CO}_2$  values are calculated (Shark Data Base, 2013). The stations are located in Skagerrak (A17), Kattegat and the Danish straits (Anholt E), the Western Baltic Sea (BY5), Baltic proper (BY15), the Bothian Sea (C3) and the Bothnian Bay (F9). Data from underway measurements of surface  $p\text{CO}_2$  is shown with yellow and covers in particular the area between Kiel and Helsinki. The division of the six sub-domains is indicated with black lines.

## Title Page

## Abstract

## Introduction

## Conclusions

## References

## Tables

## Figures

1

1

## Back

Close


Full Screen / Esc

[Printer-friendly Version](#)

## Interactive Discussion

## Sensitivity of the air–sea $\text{CO}_2$ exchange in the Baltic Sea and Danish inner waters


A. S. Lansø et al.



**Figure 2.** Monthly averaged surface  $\text{pCO}_2$  values from the six monitoring stations and from underway  $\text{pCO}_2$  data in the sub-domains in the study region. Monthly averaged values are shown with bullets together with the SDs. **(a)** Values from monitoring stations in Skagerrak (A17, blue) and Kattegat (Anholt E, green), **(b)** station BY5 (blue) and underway  $\text{pCO}_2$  in the Western Baltic Sea (black), **(c)** station BY15 (blue) and underway  $\text{pCO}_2$  from the Baltic Proper (black), **(d)** underway  $\text{pCO}_2$  data from the Gulf of Finland and **(e)** station F9 (blue), C3 (green) and underway  $\text{pCO}_2$  data from the Bothnian Sea (black) and underway  $\text{pCO}_2$  from March in the Bothnian Bay (red).

## Sensitivity of the air–sea $\text{CO}_2$ exchange in the Baltic Sea and Danish inner waters

A. S. Lansø et al.




**Figure 3.** (a)  $\Delta p\text{CO}_2$  for selected months during 2005. For the calculations of  $\Delta p\text{CO}_2$  the combined surface map of the global  $p\text{CO}_2$  climatology by Takahashi et al. (2009) and the climatology for the Baltic Sea constructed in this study is used. The coarse resolution of the global climatology is clearly visible along the west coast of Norway. Periods of under and over-saturation are seen which indicates the direction of the air–sea  $\text{CO}_2$  flux (positive  $\Delta p\text{CO}_2$  indicates release of  $\text{CO}_2$  to atmosphere, negative values indicate uptake of atmospheric  $\text{CO}_2$ ). (b) The mean seasonal air–sea  $\text{CO}_2$  flux for the years 2005 to 2010 in  $\text{g C m}^{-2} \text{ month}^{-1}$ . Positive sign indicates release of  $\text{CO}_2$  from the ocean to the atmosphere, negative sign indicates uptake of atmospheric  $\text{CO}_2$  by the ocean. This sign convention is used throughout the paper.

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)

## Sensitivity of the air–sea $\text{CO}_2$ exchange in the Baltic Sea and Danish inner waters


A. S. Lansø et al.




**Figure 4.** One hour averages of modelled and continuously measured atmospheric CO<sub>2</sub> concentrations in 2007. Also weekly averages of both modelled and measured CO<sub>2</sub> concentrations are shown.

## Sensitivity of the air–sea $\text{CO}_2$ exchange in the Baltic Sea and Danish inner waters

A. S. Lansø et al.




**Figure 5.** Top panel shows hourly averages of modelled atmospheric  $\text{CO}_2$  concentrations compared to flask measurement at F3. The three panels below include comparisons of weekly averages of modelled and continuous measurements of  $\text{CO}_2$  at MHD, PAL and WES for the period 2005–2011.



**Figure 6.** Examples of the simulated variability of atmospheric CO<sub>2</sub> in the study area shown here as extracted from the European domain in DEHM with a 50 km × 50 km resolution. Top panel: two situations for February 2007. Continental air masses cover the Baltic region on 1 February, while marine air masses are dominating on 3 February. Bottom panel: the diurnal variability on 14 July 2007 (night time (left) and day time (right)).

# Sensitivity of the air-sea $\text{CO}_2$ exchange in the Baltic Sea and Danish inner waters

A. S. Lansø et al.



**Figure 7.** The monthly mean air-sea  $\text{CO}_2$  flux for the years 2005 to 2010 in the six sub-basins in  $\text{g C m}^{-2} \text{ month}^{-1}$ .

## Title Page

## Abstract

## Introduction

## Conclusion

## References

## Tables

## Figures

18

Ba

Close


Full Screen / Esc

www.ijerpi.org | 10

## Interactive Discussion


## Sensitivity of the air–sea $\text{CO}_2$ exchange in the Baltic Sea and Danish inner waters

A. S. Lansø et al.



**Figure 8.** The seasonal flux difference between the VAT and CAT simulations for the period 2005 to 2010 in  $\text{gC m}^{-2} \text{month}^{-1}$  calculated as VAT – CAT.

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)



**Figure 9.** Time series of driving parameters as extracted from the simulations at the site south of Sweden ( $55^{\circ}18'N, 13^{\circ}55'E$ ) February 2007. Top panel:  $pCO_2^a$  for VAT and CAT together with  $u_{10}$ . Middle panel:  $pCO_2^a$  for VAT and CAT together with  $h_{mix}$ . Bottom panel: difference in  $pCO_2^a$  ( $\Delta pCO_2^a$ ) and  $F_{CO_2}$  ( $\Delta F_{CO_2}$ ) between VAT and CAT.

# Sensitivity of the air-sea $\text{CO}_2$ exchange in the Baltic Sea and Spanish inner waters

A. S. Lansø et al.

## Title Page

## Abstract

## Abstract

## Conclusio

## Conclusions References

## Tables

## Tables | Figures

1

◀ ▶

◀

Back

Back  Close


Full Screen / Esc

卷之三十一

Autumn 2018 | Page 10

## Sensitivity of the air–sea $\text{CO}_2$ exchange in the Baltic Sea and Danish inner waters


A. S. Lansø et al.



**Figure 10.** Simulated parameters as in Fig. 9 at the site south of Sweden (55°18'N, 13°55'E), but for July 2007. Top panel:  $\text{pCO}_2^a$  for VAT and CAT together with  $u_{10}$ . Middle panel:  $\text{pCO}_2^a$  for VAT and CAT together with  $h_{\text{mix}}$ . Bottom panel: difference in  $\text{pCO}_2^a$  ( $\Delta\text{pCO}_2^a$ ) and  $F_{\text{CO}_2}$  ( $\Delta F_{\text{CO}_2}$ ) between VAT and CAT.

## Sensitivity of the air–sea $\text{CO}_2$ exchange in the Baltic Sea and Danish inner waters

A. S. Lansø et al.



**Figure 11.** Simulated vertical profiles of atmospheric CO<sub>2</sub> at the site south of Sweden (55°18' N, 13°55' E) in units of ppm. Top panel: 1–10 February 2007. Bottom panel: 11–20 July 2007. The black line represents the mixing height in km. Note the different scales used in the two plots.